Title: Does diversity beget diversity in microbiomes?

Authors: Naïma Madi[1], Michiel Vos[2], Carmen Lia Murall[1], Pierre Legendre[1] and B.

Jesse Shapiro[1,3,4*]

1. Département de sciences biologiques, Université de Montréal, Canada

2. European Centre for Environment and Human Health, University of Exeter,

Penryn, UK

3. Department of Microbiology and Immunology, McGill University, Canada

4. McGill Genome Centre, Canada

*correspondence: jesse.shapiro@umontreal.ca

keywords: microbiome, evolution, ecology, Earth Microbiome Project, 16S rRNA

Abstract

Microbes are embedded in complex communities where they engage in a wide array of

intra- and inter-specific interactions. The extent to which these interactions drive or

impede microbiome diversity is not well understood. Two contrasting hypotheses have

been put forward to explain how species interactions could influence diversity.

'Ecological Controls' (EC) predicts a negative relationship, where the evolution or

migration of novel types is constrained as available niches become filled. In contrast,

'Diversity Begets Diversity' (DBD) predicts a positive relationship, with existing

diversity promoting the accumulation of further diversity via niche construction and other

interactions. Using the Earth Microbiome Project, the largest standardized survey of

global biodiversity to date, we provide evidence that DBD is strongest in less diverse

biomes, such as the animal gut, but weaker in more diverse biomes, such as soil,

presumably because most available niches are filled. Also consistent with niche

adaptation, resident genera that are strongly specialized for a particular environment

show a stronger DBD relationship than do generalists or migrants from another

environment. In contrast with the Black Queen Hypothesis that predicts a negative

correlation between genome size and community metabolic interactions, we find that

genera with larger genomes exhibit a stronger DBD response, which could be due to a

higher potential for interactions and niche construction offered by more diverse gene

repertoires. Our results highlight the importance of species interactions in shaping

microbiome diversity, and the need to account for such interactions in the study of

microbial genomes, physiology, and ecological function.

Introduction

The majority of the genetic diversity on Earth is encoded by microbes (1–3) and

the functioning of all Earth's ecosystems is reliant on diverse microbial communities (4).

High-throughput 16S rRNA gene amplicon sequencing studies continue to yield

unprecedented insight into the taxonomic richness of microbiomes (e.g. (5, 6)), and

abiotic drivers of community composition (e.g. pH; (7, 8)) are increasingly characterized.

Although it is known that biotic (microbe-microbe) interactions can also be important in

determining community composition (9), comparatively little is known about how such

interactions, positive (e.g. cross-feeding; (10)) or negative (e.g. toxin-mediated

interference competition; (11, 12)), shape microbiome diversity.

The dearth of studies exploring how microbial interactions could influence

diversity stands in marked contrast to a long research tradition on biotic controls of plant

and animal diversity (13, 14). In an early study of 49 animal (vertebrate and invertebrate)

community samples, Elton plotted the number of species versus the number of genera and

observed a ~1:1 ratio in each individual sample, but a ~4:1 ratio when all samples were

pooled (14). He took this observation as evidence for competitive exclusion preventing

related species, more likely to overlap in niche space, to co-exist. This concept, more

recently referred to as niche filling or Ecological Controls (EC) (15), predicts speciation

(or, more generally, diversification) rates to decrease with increasing standing species

diversity because of less available niche space (16). In contrast, the Diversity Begets

Diversity (DBD) model predicts that when species interactions create novel niches,

standing biodiversity favors further diversification (17, 18). For example, niche

construction (i.e. the physical, chemical or biological alteration of the environment) could

influence the evolution of the species constructing the niche, as well as that of co-

occurring species (19, 20).

Empirical evidence for the action of EC vs. DBD in natural plant and animal

communities has been mixed (18, 21–24). Laboratory evolution experiments tracking the

diversification of a focal bacterial lineage in communities of varying complexity have

also yielded contradictory results, with support for EC, DBD, or intermediate scenarios

(25, 26). For example, diversification of a focal *Pseudomonas* clone was favored by

increasing community diversity in the range of 0-20 strains or species within the same

genus (18, 27) but diversification was inhibited by very diverse communities (*e.g.*

hundreds or thousands of species in compost; (28)). These experiments are consistent

with interspecific competition initially driving (29), but eventually inhibiting

diversification as niches are filled.

Most laboratory experiments are restricted to relatively short evolutionary time

scales and include only a small number of taxa; it is therefore unclear if they can be

generalized to natural communities consisting of very many more taxa evolving over

much longer periods, spanning more environmental changes, multiple speciation events,

and large-scale genomic changes. Moreover, because many free-living microbes have

high migration rates ("everything is everywhere, but the environment selects" (30)), we

expect that the majority of diversity present within a sample is the result of immigration

from a meta-community pool rather than having evolved *in situ*. As such, here we

broadly define "diversity begets diversity" (DBD) to include the combined effects of

community assembly from a migrant pool ('ecological species sorting') and *in situ*

evolutionary diversification (**Fig. 1**).

Fig. 1. Contrasting the Diversity Begets Diversity (DBD) and Ecological Controls (EC) models. In the top panel, we consider an evolutionary diversification model of a focal lineage as a function of initial diversity present at the time of diversification. In the bottom panel, we consider a community assembly model, in which standing diversity (black points) in a community selects (for or against) new ASVs which arrive via migration (purple points & arrows). **(A)** For example, sample 1 contains one non-focal genus, and two ASVs within the focal genus (point at x=1, y=2 in the plot). Sample 2 contains three non-focal genera, and four ASVs within the focal genus (point at x=3, y=4). Tracing a line through these points yields a positive diversity slope, supporting the DBD model (red). **(B)** Alternatively, a negative slope would support the Ecological Controls (EC) model (blue line).

To test whether patterns of diversity in natural communities conform to EC or DBD dynamics, we used 2,000 microbiome samples from the Earth Microbiome Project (EMP), the largest available repository of biodiversity based on standardized sampling

and sequencing protocols (31). All samples were rarefied to 5,000 observations (counts of

16S rRNA gene sequences classified into Amplicon Sequence Variants; ASVs), as

diversity estimates are highly sensitive to sampling effort (32). Following Elton (14), we

use the equivalent of Species:Genus ratios. We then calculated a range of taxonomic

diversity ratios (ASV:Genus, Genus:Family, Family:Order, Order:Class, and

Class:Phylum) as proxies for diversity within a focal lineage, from shallow to deep

evolutionary time. Finally, we plot each ratio as a function of the number of non-focal

lineages (Genera, Families, Orders, Classes, and Phyla, respectively) with which the focal

lineage could interact. We refer to the slope of these plots as the "diversity slope", with

negative slopes supporting EC and positive slopes supporting DBD (**Fig. 1**).

Results

Quantifying the DBD-EC continuum in prokaryote communities. We used

generalized linear mixed models (GLMMs) to estimate the diversity slope at each

taxonomic level in the EMP data, which revealed a tendency toward positive slopes with

significant variation by lineage, environment, and their interaction (**Table S1, Figures S1**

and S2-S6, Supplementary Data file 1 Section 1). Examples of how the diversity slope

varies across lineages and environments are shown in **Fig. 2** and **S2-S6**. To assess the

significance of these slope estimates in light of potential sampling bias and data structure

(32, 33), we considered two ecologically plausible null models, both of which randomize

the associations between ASVs within a sample, thus breaking any true biotic interactions

(**Materials and Methods**). Simulation results suggest that diversity slopes in the real

EMP data are not trivially explained by null models (**Fig. S7; Table S2**) , and we revisit

these models in testing key hypotheses below. Because taxonomic labels can be

unavailable or inconsistent with phylogenetic relationships (34, 35) we repeated the

analyses using nucleotide sequence identity in the 16S gene instead of taxonomy, and

again recovered generally positive diversity slopes (**SI Materials and Methods**).

Fig. 2. Focal lineage diversity as a function of community diversity in the top two most prevalent taxa at each taxonomic level. Significant positive diversity slopes are shown in red, negative in blue (linear models, $P < 0.05$, Bonferroni corrected for 17 tests), and non-significant in grey. Four representative environments are shown (see Supplementary Figures S2-S6 for plots in all 17 environments).

DBD reaches a plateau at high diversity. It is expected from theory and experimental studies that a positive DBD relationship should eventually reach a plateau, giving way to EC as niches become saturated (25, 28). This expectation is borne out in our dataset, particularly in the nucleotide sequence-based analyses which support quadratic or cubic relationships over linear diversity slopes (**Fig. S8**). For example, in the animal distal gut, a relatively low-diversity biome, we observed a strong linear DBD relationship at most phylogenetic depths; in contrast, the much more diverse soil biome clearly reaches a plateau (**Fig. S9**).

To comprehensively test the hypothesis that more diverse microbiomes experience weaker DBD due to saturated niche space, we used a GLMM including the interaction between diversity and environment as a fixed effect. We considered this

model only for taxonomic ratios with significant diversity slope variation by environment

(**Table S1**): Family:Order, Order:Class, and Class:Phylum. Consistent with a plateau as

niches become filled, diversity slopes were significantly more positive in less diverse

(often host-associated) biomes (**Fig. 3**, **Figure S10, Supplementary Data file 1 Section**

2). Importantly, the interaction observed in the real EMP data between community

diversity and biome type in shaping focal lineage diversity was not significant in null

models (in which each environment has its own characteristic level of diversity) at any

taxonomic ratio (**Supplementary Data file 2 Section 2**). This suggests that the plateau of

DBD at higher levels of community diversity is not easily explained by data structure.

Fig. 3. Diversity slope is higher in low-diversity (often host-associated) microbiomes.
The x-axis shows the mean number of phyla in each biome. On the y-axis, the diversity
slope was estimated by a GLMM predicting focal lineage diversity as a function of the
interaction between community diversity and environment type at the Class:Phylum ratio
(**Supplementary Data file 1 Section 2.3**). The line represents a regression; the shaded
area depicts 95% confidence limits of the fitted values.

Abiotic drivers of diversity. Our results (**Fig. 3**) suggest that community diversity is a

major determinant of the EC-DBD continuum, and by extension that biotic interactions

may override abiotic factors in determining where a community lies on the continuum.

To formally test for the additional role abiotic drivers might play in generating the

observed EC-DBD continuum, we analyzed two data sets in more detail.

First, we analyzed a subset of 192 EMP samples with measurements of four key

abiotic factors shown to affect microbial diversity (pH, temperature, latitude, and

elevation; (7, 8, 15, 36)). We fitted a GLMM with focal lineage-specific diversity as the

dependent variable, and with the number of non-focal lineages, the four abiotic factors

and their interactions as predictors (fixed effects). As in the full EMP dataset (**Table S1**),

focal lineage diversity was positively associated with community diversity at all

taxonomic ratios in the EMP subset (**Table S3**). As expected, certain abiotic factors,

alone or in combination with diversity, had significant effects on focal lineage diversity.

However, the effects of abiotic factors were always weaker than the effect of community

diversity (**Table S3**; **Supplementary Data file 1 Section 3**). Although only a small

subset of abiotic factors was considered, this analysis suggests that positive diversity

slopes are not likely to be primarily driven by the abiotic environment.

Second, we used a global dataset of 237 soil samples associated with more

detailed environmental metadata (36) (**SI Materials and Methods**). This dataset revealed

weaker evidence for DBD and stronger abiotic effects on diversity. Community diversity

generally had significant positive effects on focal-lineage diversity, but the effect was

weak and not detectable at all taxonomic ratios (**Table S4**). Known abiotic drivers of soil

bacterial diversity such as pH (7) and latitude (36) had effects of similar or stronger

magnitude as community diversity (**Table S4, Supplementary Data file 3**). Together,

these results suggest a mixture of biotic and abiotic drivers of diversity, with their relative

contributions varying depending on the environment type. In particular, environments

that have reached a DBD plateau such as soil, appear to have diversity levels controlled

mainly by abiotic factors.

DBD is more pronounced in microbiome residents than in migrants or generalists.

A recent meta-analysis of 16S sequence data from a variety of biomes suggested an

important distinction between generalist lineages found in many environments, compared

to more restricted specialists (37). Generalists were inferred to have higher speciation

rates, suggesting that the DBD-EC balance might differ between generalists and

specialists. To further investigate this difference, we defined microbiome specialists

(which we call "residents") and generalists in the EMP dataset. We first clustered

environmental samples by their genus-level community composition using fuzzy k-means

clustering (**Fig. 4a**), which identified three major clusters: 'animal-associated', 'saline',

and 'non-saline'. The clustering included some outliers (*e.g.* plant corpus grouping with

animals), but was generally consistent with known distinctions between host-associated

vs. free-living (31), and saline vs. non-saline communities (38, 39). Resident genera were

defined as those with a strong preference for a particular environment cluster (whether

due to dispersal limitation or narrow niche breadth) using indicator species analysis

(permutation test, $P<0.05$; **Fig. 4a**; **Fig. S11**; **Supplementary Data file 4**), and genera

without a strong preference were considered generalists. When residents of one

environmental cluster were observed in a different cluster, we defined them as "migrants"

in that sample. For each environment cluster, we ran a GLMM with resident genus-level

diversity (the number of non-focal genera) as a predictor of focal-lineage diversity (the

ASV:Genus ratio) for residents, generalists, or migrants to that sample (**Supplementary**

Data file 1 Section 4).

Resident community diversity had no significant effect on the diversity of

generalists in animal-associated, saline and non-saline clusters (Wald test, $P>0.05$), but

was positively correlated with lineage-specific resident diversity (Wald test, $z=7.1$, $P=$

1.25e-12; $z=3.316$, $P=0.0009$; $z=7.109$, $P=1.17e-12$, respectively). Resident community

diversity significantly decreased migrant diversity in saline ($z=-3.194$, $P=0.0014$) and

non-saline environment clusters ($z=-2.840$, $P=0.0045$), but had no significant effect in the

animal-associated cluster ($P>0.05$) (**Fig. 4b**). These results suggest that, although

generalist lineages may have higher speciation rates and colonize more habitats than

specialists (37), they are at a relative disadvantage in accumulating taxonomic diversity

via community assembly. Migrants (residents of one environment type that arrive in

another) are at an even greater disadvantage, because higher resident diversity apparently

selects against migrant diversity, supporting an EC model in two out of three

environment types (**Fig. 4b**). The accumulation of diversity via migration from foreign

environments may thus be limited, presumably because most niches are already occupied

by residents.

Fig. 4. The DBD relationship varies between resident and non-resident genera. (A) PCA showing genera clustering into their preferred environment clusters. Circles indicate genera and triangles indicate environments (EMPO 3 biomes). The three environment clusters identified by fuzzy *k*-means clustering are: Non-saline (NS, blue), saline (S, green) and animal-associated (purple). Resident genera were identified by indicator species analysis. **(B) DBD in resident versus non-resident genera across environment clusters.** Results of GLMMs modeling focal lineage diversity as a function of the interaction between community diversity and resident/migrant/generalist status. The x-axis shows the standardized number of non-focal resident genera (community diversity); the y-axis shows the number of ASVs per focal genus. Resident focal genera

are shown in orange, migrant focal genera in red, and generalist focal genera in black.

Red stars indicate a significantly positive or negative slope (Wald test, $P<0.005$).

DBD is associated with larger genome size. In soil, bacteria with larger genomes, and

thus larger repertoires of accessory genes, are able to occupy a wider range of niches

(40). A higher proportion (and thus, possibly, a higher diversity) of such broadly-

distributed, metabolically versatile bacteria might be able to survive and thrive when they

disperse into a new location. We might therefore expect lineages with larger genomes to

exhibit stronger DBD. To test this expectation, we assigned genome sizes to 576 genera

for which at least one whole genome sequence was available, and added an interaction

term between genome size and diversity as a fixed effect in the GLMM (**Materials and**

Methods). Consistent with this expectation, we observed a significant positive effect of

genome size on the diversity slope (Wald test, $z=2.5$, $P=0.01$; **Fig. 5, Supplementary**

Data file 1 Section 5). This effect was not observed in any null model, in which the

interaction between community diversity and genome size was never significant

(**Supplementary Data file 2 Section 3**). Therefore the effect of genome size cannot be

trivially explained by data structure. The positive relationship between genome size and

DBD is likely even stronger than estimated, because assigning genome sizes to entire

genera is imprecise (*i.e.* there is substantial variation in genome size within a genus, or

even within species). This result supports a model in which species with larger gene

repertoires and greater opportunity to engage in niche construction (19) are more prone to

interact with other species, and therefore to co-diversify or co-assemble at higher rates.

Fig. 5. Positive effect of genome size on DBD. Results are shown from a GLMM

predicting focal lineage diversity as a function of the interaction between community

diversity and genome size at the ASV:Genus ratio (**Supplementary Data file 1 Section**

5). The x-axis shows the standardized number of non-focal genera (community diversity);

the y-axis shows the number of ASVs per focal genus. Variable diversity slopes

corresponding to different genome sizes are shown in a blue color gradient; the shaded

area depicts 95% confidence limits of the fitted values.

Taken together with the observation that DBD is stronger in habitat residents

(specialists) than generalists (**Fig. 4b**), these results suggest two independent and

opposing effects: generalists tend to have larger genomes (37, 40), increasing their ability

to colonize new locations, while still being at a disadvantage compared to residents. To

investigate this apparent tradeoff, we considered how the effects of generalism and

genome size could interact to shape DBD. However, we did not find strong evidence for

interactions between genome size and generalist lifestyle, and instead found support for a

simpler model in which genome size correlates with DBD independently of

generalist/resident/migrant status (**Supplementary Text; Fig. S12, S13**).

Discussion

Using ~10 million individual marker sequences from the EMP, we demonstrate a

tendency for diversity in focal lineages to be positively associated with overall

community diversity, with significant variation across lineages and environments. The

strength of the DBD relationship dissipates with increasing microbiome diversity, which

we hypothesize is caused by niche saturation. In more diverse biomes such as soil, abiotic

factors may become more important in driving diversity after DBD reaches a plateau. The

effect of DBD is strongest among habitat residents (specialists), suggesting long-term

niche adaptation and selection against the accumulation of migrant diversity in most

environments.

The positive correlation between genome size and DBD observed here is at odds

with the Black Queen hypothesis, which predicts that social conflict between interacting

species leads to the inactivation and loss of genes involved in shareable metabolites

(public goods), eventually leading to a reduction in genome size (41). It thus predicts a

negative correlation between the degree of species interactions (*i.e.* community diversity)

and genome size (41), opposite to what is observed in our analysis. The stronger DBD in

larger genomes observed here could be driven by larger metabolic repertoires encoded by

larger genomes, resulting in more opportunities to benefit from cross-feeding and other

interspecies interactions. This deserves further study.

While most of the DBD literature considers a model of evolutionary

diversification (15, 17), our results pertain mainly to ecological community assembly

dynamics. At the limited resolution of 16S marker gene sequences, we do not expect

measurable diversification within an individual microbiome sample; however community

diversity could still select for (as in DBD) or against (as in EC) standing diversity in a

focal lineage, even if this lineage diversified before the sampled community assembled.

The EMP survey data support theory and experimental data on biotic drivers of

diversity. In particular, our results support the experimental and theoretical results that

DBD is strong when community diversity is low (18, 27), spurring the accumulation of

diversity in a positive feedback until niches are filled and EC predominates (25, 26, 28,

29). However, due to the correlational nature of the EMP data, it is not possible to test

whether DBD is primarily due to the creation of novel niches via biotic interactions and

niche construction (20), or due to increased competition leading to specialization on

underexploited resources (12, 27). Regardless of the underlying mechanisms, our results

demonstrate a general scaling between different levels of community diversity, which has

important implications for modeling and predicting community function and stability (42,

43). The answer to the question 'why are microbiomes so diverse?' might in a large part

be because microbiomes are so diverse (21).

Acknowledgements.

We thank Luke Thompson for assistance obtaining EMP data and Zofia Ecaterina

Taranu, Vincent Fugère and Guillaume Larocque for advice on GLMMs. We are also

grateful to Steven Kembel and Tom Battin for critical comments that improved the

manuscript. **Funding:** This project was made possible by an NSERC Discovery Grant

and Canada Research Chair to BJS.

Author contributions. Conceptualization: BJS, MV. Data curation: NM. Formal

analysis: NM, MV, BJS. Funding acquisition: BJS. Investigation: NM, MV, CLM, PL,

BJS. Methodology: NM, MV, CLM, PL, BJS. Resources: BJS, PL. Supervision: PL, BJS.

Software: NM. Visualization: NM. Writing original draft: NM, MV, BJS. Writing -

review & editing: NM, MV, CLM, PL, BJS.

Competing interests: none to declare.

Data and materials availability: All data is available from the Earth Microbiome

Project (ftp.microbio.me), as detailed in the Methods. All computer code used for

analysis are available at https://github.com/Naima16/dbd.git.

Materials and Methods

Earth Microbiome Project dataset. We used the EMP '2000 subset' of 16S rRNA gene

sequences, rarefied to 5000 sequences per sample. This subset contains 155,002 ASVs

from 2,000 samples with an even distribution across 17 natural environments (EMP

Ontology level 3). Data were downloaded from the EMP FTP server (ftp.microbio.me),

on February 9, 2018. See *SI Extended Materials and Methods* for further details on data

processing.

Generalized linear mixed model (GLMM) analyses. We used GLMMs to determine

how the richness of a focal lineage (*e.g.* its ASV:Genus ratio) is affected by the diversity

of other lineages (*e.g.* non-focal genera) in the community. The effects of environment

(as defined by the EMP Ontology 'level 3 biomes') and the identity of the focal lineage

were included as random effects on the slope and intercept. We also controlled for the

submitting laboratory (identified by the principal investigator) and the EMP unique

sample identifier (i.e. if two taxa were part of the same sample).

   All models were fitted in Rstudio (Version 1.1.442, R Version 3.5.2) using the

glmer function of the lme4 package (44). Data standardization (transformation to a mean

of zero and a standard deviation of one) was applied to all predictors to get comparable

estimates. We used likelihood-ratio tests (LRT) (anova R function from stats package) as

follows: 1) on nested models to assess the significance of random effects (in the nested

models, each effect was dropped one at a time); 2) on the full model and the null model

comprising only random effects, to assess the significance of fixed effects (45); 3) on the

full model and the model without the interaction term, to assess the significance of

interactions. All models reported here were found to be significant ($P<0.05$). See *SI*

Extended Materials and Methods for further details on inclusion of abiotic factors and

variation across environments.

Null models. We considered two null models, both of which randomize the associations

between ASVs within a sample, thus breaking any true biotic interactions. Both models

were randomly generated matrices of the same size as the real EMP dataset. Model 1 was

generated by sampling from a Poisson distribution for each element. To account for

differences in diversity across the 17 EMP environments, Model 2 was generated by

sampling from a separate Poisson distribution for each environment (**SI Extended**

Materials and Methods). In a GLMM analysis of the simulated data, Model 1 produced

a significantly negative slope describing the relationship between focal lineage diversity

(ASV:genus) and community diversity (# of genera), whereas Model 2 produced a

significantly positive slope (**Fig. S7, Table S2**). To estimate the power to detect DBD or

EC effects, we specifically added these effects to data simulated under each of the null

models. As expected, adding DBD reversed the negative slope of Model 1 and rendered it

positive (**Table S2, Supplementary Data file 2 Section 1**), suggesting that a true DBD

effect is easily detectable under null Model 1. The baseline slope of null Model 2 was

already positive, and was increased somewhat with the addition of a DBD effect. In

contrast, the addition of EC to either null model had little effect on the slope. Together,

these null models suggest some caution in interpreting the general positive slopes

observed in the real EMP data (**Fig 2; Table S1**). Positive slopes are consistent with

DBD, and are highly unexpected under null Model 1, but could arise by chance under

null Model 2, in which biomes differ in their levels of diversity. This is because Model 2

is effectively a composite of Model 1, in which different biomes, each with a negative

slope are 'stacked' to yield an overall positive slope. In the real data, most individual

environments tend toward a positive slope (**Fig. S9, S14**), and the variation among

environments is accounted for in the GLMM (**Table S1**). Therefore, the real data are not

trivially explained by either null model.

Nucleotide sequence-based analysis. We clustered ASVs at decreasing levels of

nucleotide identity, from 100% identical ASVs down to 75% identity (roughly equivalent

to phyla (46)). We estimated focal cluster diversity as the mean number of descendants

per cluster (e.g. number of 100% clusters per 97% cluster) and plotted this against the

total number of clusters (97% identity in this example). This approach has the advantage

of including sequences even if they come from unnamed taxa. For each of the six

nucleotide divergence ratios tested, the relationship between diversity and diversification

was positive (**Fig. S8**), consistent with DBD and suggesting that the taxonomic analyses

were qualitatively unbiased. See *SI Extended Materials and Methods* for further details.

Defining residents, generalists, and migrants. We clustered the environmental samples

based on their genus-level community composition using fuzzy *k*-means clustering. The

clustering (cmeans function, package e1071 in R) was done on the 'hellinger'

transformed data (decostand function, vegan R package). To identify resident genera to

each cluster, we used indicator species analysis (47) as implemented in the indval

function (labdsv R package). We defined residents as genera with indval indices between

0.4 and 0.9, with permutation test $P < 0.05$. Genera not associated with any cluster were

considered generalists. We used principal component analysis (PCA) to visualize

clustering and indicator genera (rda function, vegan R package). We then ran a separate

GLMM for each environmental cluster, with resident genus-level diversity (number of

non-focal genera) as a predictor of focal genus diversity (ASV:Genus ratio) for resident,

migrant (residents of one cluster found in a different cluster) and generalist genera. The

fixed effect was specified as the interaction between diversity and a factor defining the

genus-cluster association (with three levels: resident, migrant and generalist). Random

effects on intercept and slope were kept as in the GLMMs described above.

Genome size analysis. We chose a subset of genera represented by one or more

sequenced genomes in the NCBI microbial genomes database

(https://www.ncbi.nlm.nih.gov/genome/browse#!/prokaryotes/). For these genera, a

representative genome size was assigned by selecting the genome with the lowest number

of scaffolds (if no closed genomes were available) (**Supplementary file 5**). If multiple

genomes were available with the same level of completion, the largest genome size was

used, and only genera with two or more ASVs were included in the analysis. Intracellular

symbionts were excluded. We fitted a GLMM on the subset of data with known genome

size (576 genera, ranging from ~1 to 15 Mbp) with the interaction between community

diversity and genome size as a predictor of focal lineage diversity at the ASV:Genus

level. All the other random effects on intercept and slope were kept as in the previous

GLMMs.

References

1.   S. Sunagawa, *et al.*, Ocean plankton. Structure and function of the global ocean
microbiome. *Science* **348**, 1261359 (2015).

2.   P. Lapierre, J. P. Gogarten, Estimating the size of the bacterial pan-genome. *Trends*
Genet. **25**, 107–110 (2009).

3.   Hug, L. A. et al. A new view of the tree and life's diversity. Nature Microbiology **1**,
16048 (2016).

4.   P. G. Falkowski, T. Fenchel, E. F. Delong, The microbial engines that drive Earth's
biogeochemical cycles. *Science* **320**, 1034–1039 (2008).

5.   M. L. Sogin, *et al.*, Microbial diversity in the deep sea and the underexplored "rare
biosphere." *Proc. Natl. Acad. Sci. U. S. A.* **103**, 12115–12120 (2006).

6.   S. Louca, F. Mazel, M. Doebeli, L. W. Parfrey, A census-based estimate of Earth's
bacterial and archaeal diversity. *PLoS Biol.* **17**, e3000106 (2019).

7.   Lauber, C. L., Hamady, M., Knight, R. & Fierer, N. Soil pH as a predictor of soil
bacterial community structure at the continental scale: a pyrosequencing-based
assessment. *Appl. Environ. Microbiol.* **75**, 5111-5120 (2009).

8.   J. F. Power, *et al.*, Microbial biogeography of 925 geothermal springs in New
Zealand. *Nat. Commun.* **9**, 2876 (2018).

9.   Needham, D. M. & Fuhrman, J. A. Pronounced daily succession of phytoplankton ,
archaea and bacteria following a spring bloom. *Nature Microbiology* **1**, 16005
(2016).

10.   E. C. Seth, M. E. Taga, Nutrient cross-feeding in the microbial world. *Front.*
Microbiol. **5**, 350 (2014).

11.   T. L. Czárán, R. F. Hoekstra, L. Pagie, Chemical warfare between microbes
promotes biodiversity. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 786–790 (2002).

12.   M. E. Hibbing, C. Fuqua, M. R. Parsek, S. B. Peterson, Bacterial competition:
surviving and thriving in the microbial jungle. *Nat. Rev. Microbiol.* **8**, 15–25
(2010).

13.   G. F. Gause, *The Struggle for Existence* (Williams & Wilkins, Baltimore, 1934).

14.   C. Elton, Competition and the Structure of Ecological Communities. *J. Anim. Ecol.*
15, 54–68 (1946).

15.   D. Schluter, M. W. Pennell, Speciation gradients and the distribution of
465   biodiversity. *Nature* **546**, 48–55 (2017).

466   16.   D. L. Rabosky, A. H. Hurlbert, Species richness at continental scales is dominated
467   by ecological limits. *Am. Nat.* **185**, 572–583 (2015).

468   17.   R. H. Whittaker, Evolution and Measurement of Species Diversity. *Taxon* **21**, 213–
469   251 (1972).

470   18.   V. Calcagno, P. Jarne, M. Loreau, N. Mouquet, P. David, Diversity spurs
471   diversification in ecological communities. *Nat. Commun.* **8**, 15810 (2017).

472   19.   M. San Roman, A. Wagner, An enormous potential for niche construction through
473   bacterial cross-feeding in a homogeneous environment. *PLoS Comput. Biol.* **14**,
474   e1006340 (2018).

475   20.   K. N. Laland, F. J. Odling-Smee, M. W. Feldman, Evolutionary consequences of
476   niche construction and their implications for ecology. *Proc. Natl. Acad. Sci. U. S.*
477   *A.* **96**, 10242–10247 (1999).

478   21.   B. C. Emerson, N. Kolm, Species diversity can drive speciation. *Nature* **434**, 1015–
479   1017 (2005).

480   22.   M. W. Palmer, T. A. Maurer, Does Diversity Beget Diversity? A Case Study of
481   Crops and Weeds. *J. Veg. Sci.* **8**, 235–240 (1997).

482   23.   Price, T. D. et al. Niche filling slows the diversification of Himalayan songbirds.
483   *Nature* **509**, 222-225 (2014).

484   24.   D. L. Rabosky, *et al.*, An inverse latitudinal gradient in speciation rate for marine
485   fishes. *Nature* **559**, 392–395 (2018).

486   25.   M. A. Brockhurst, N. Colegrave, D. J. Hodgson, A. Buckling, Niche occupation
487   limits adaptive radiation in experimental microcosms. *PLoS One* **2**, e193 (2007).

488   26.   J. R. Meyer, R. Kassen, The effects of competition and predation on diversification
489   in a model adaptive radiation. *Nature* **446**, 432–435 (2007).

490   27.   A. Jousset, N. Eisenhauer, M. Merker, N. Mouquet, S. Scheu, High functional
491   diversity stimulates diversification in experimental microbial communities. *Sci Adv*
492   **2**, e1600124 (2016).

493   28.   P. Gómez, A. Buckling, Real-time microbial adaptive diversification in soil. *Ecol.*
494   *Lett.* **16**, 650–655 (2013).

29.   S. F. Bailey, J. R. Dettman, P. B. Rainey, R. Kassen, Competition both drives and

impedes diversification in a model adaptive radiation. *Proc. Biol. Sci.* **280**,
20131253 (2013).

30. R. de Wit, T. Bouvier, "Everything is everywhere, but, the environment selects";
what did Baas Becking and Beijerinck really say? *Environ. Microbiol.* **8**, 755–758
(2006).

31. Thompson, L. R. et al. A communal catalogue reveals Earth's multiscale microbial
diversity. *Nature* **551**, 457-463 (2017).

32. N. J. Gotelli, R. K. Colwell, Quantifying biodiversity: procedures and pitfalls in the
measurement and comparison of species richness. *Ecol. Lett.* **4**, 379–391 (2001).

33. O. Jarvinen, Species-To-Genus Ratios in Biogeography: A Historical Note. *J.*
Biogeogr. **9**, 363–370 (1982).

34. D. H. Parks, *et al.*, A standardized bacterial taxonomy based on genome phylogeny
substantially revises the tree of life. *Nat. Biotechnol.* **36**, 996–1004 (2018).

35. M. Vos, A species concept for bacteria based on adaptive divergence. *Trends*
Microbiol. **19**, 1–7 (2011).

36. M. Delgado-Baquerizo, *et al.*, A global atlas of the dominant bacteria found in soil.
Science **359**, 320–325 (2018).

37. S. Sriswasdi, C.-C. Yang, W. Iwasaki, Generalist species drive microbial dispersion
and evolution. *Nat. Commun.* **8**, 1162 (2017).

38. C. A. Lozupone, R. Knight, Global patterns in bacterial diversity. *Proc. Natl. Acad.*
Sci. U. S. A. **104**, 11436–11440 (2007).

39. J.-C. Auguet, A. Barberan, E. O. Casamayor, Global ecological patterns in
uncultured Archaea. *ISME J.* **4**, 182–190 (2010).

40. A. Barberán, *et al.*, Why are some microbes more ubiquitous than others?
Predicting the habitat breadth of soil bacteria. *Ecol. Lett.* **17**, 794–802 (2014).

41. Morris, J. J. & Lenski, R. E. The Black Queen Hypothesis: evolution of
dependencies through adaptive gene loss. *MBio* **3**, e00036-12 (2012).

42. K. Z. Coyte, J. Schluter, K. R. Foster, The ecology of the microbiome: Networks,
competition, and stability. *Science* **350**, 663–666 (2015).

43. F. Pennekamp, *et al.*, Biodiversity increases and decreases ecosystem stability.
Nature **563**, 109–112 (2018).

44. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software, Articles* **67**, 1–48 (2015).

45. W. Forstmeier, H. Schielzeth, Cryptic multiple hypotheses testing in linear models:
overestimated effect sizes and the winner's curse. *Behav. Ecol. Sociobiol.* **65**, 47–
55 (2011).

46. K. T. Konstantinidis, J. M. Tiedje, Towards a genome-based taxonomy for
prokaryotes. *J. Bacteriol.* **187**, 6258–6264 (2005).

47. M. Dufrene, P. Legendre, Species Assemblages and Indicator Species: The Need
for a Flexible Asymmetrical Approach. *Ecol. Monogr.* **67**, 345–366 (1997).

536