Beat-based and memory-based temporal expectations in rhythm: similar perceptual effects, different underlying mechanisms

Fleur L. Bouwer, Henkjan Honing, & Heleen A. Slagter

University of Amsterdam

Author Note

Fleur L. Bouwer, Department of Psychology, Amsterdam Brain and Cognition, University of Amsterdam

Henkjan Honing, Institute for Logic, Language and Computation, Amsterdam Brain and Cognition, University of Amsterdam

Heleen A. Slagter, Department of Experimental and Applied Psychology, Vrije Universiteit Amsterdam and Department of Psychology, Amsterdam Brain and Cognition, University of Amsterdam

FLB is supported by an ABC Talent Grant awarded by Amsterdam Brain and Cognition. HAS is supported by a European Research Council (ERC) starting grant (679399). We would like to thank Peter Saalbrink for his assistance with the data collection.

Correspondence concerning this article should be addressed to Dr. Fleur Bouwer, Department of Psychology, University of Amsterdam (bouwer@uva.nl) or Prof. Dr. Heleen Slagter, Department of Applied and Experimental Psychology, Vrije Universiteit Amsterdam (h.a.slagter@vu.nl).

Abstract

Predicting the timing of incoming information allows the brain to optimize information processing in dynamic environments. Behaviorally, temporal expectations have been shown to facilitate processing of events at expected time points, such as sounds that coincide with the beat in musical rhythm. Yet, temporal expectations can develop based on different forms of structure in the environment, not just the regularity afforded by a musical beat. Little is still known about how different types of temporal expectations are neurally implemented and affect performance. Here, we orthogonally manipulated the periodicity and predictability of rhythmic sequences to examine the mechanisms underlying beat-based and memory-based temporal expectations, respectively. Behaviorally and using EEG, we looked at the effects of beat-based and memory-based expectations on auditory processing when rhythms were task relevant or task irrelevant. At expected time points, both beat-based and memory-based expectations facilitated target detection and led to attenuation of P1 and N1 responses, even when expectations were task-irrelevant (unattended). For beat-based expectations, we additionally found reduced target detection and enhanced N1 responses for events at unexpected time points (e.g., off-beat), regardless of the presence of memory-based expectations or task relevance. This latter finding supports the notion that periodicity selectively induces rhythmic fluctuations in neural excitability and furthermore indicates that while beat-based and memory-based expectations may similarly affect auditory processing of expected events, their underlying neural mechanisms may be different.

Keywords: rhythm, temporal expectations, beat, attention, prediction, auditory

Beat-based and memory-based temporal expectations in rhythm: similar perceptual effects,

different underlying mechanisms

To optimize sensory processing and perception in a changing environment, the human brain continuously tries to predict incoming information (Clark, 2013; Friston, 2005). Being able to not only predict the content of sensory input ("what"), but also its timing ("when") allows the system to prepare for and focus on time points when useful information is likely to occur (Large & Jones, 1999; Nobre & van Ede, 2018). Indeed, temporal expectations have been shown to improve processing of events at expected time points (Haegens & Zion Golumbic, 2018; Henry & Obleser, 2012; Nobre & van Ede, 2018; Rohenkohl, Cravo, Wyart, & Nobre, 2012; ten Oever, Schroeder, Poeppel, van Atteveldt, & Zion-Golumbic, 2014). Additionally, temporal expectations allow us to align our actions to sensory input, enabling complex behaviors such as dancing and synchronizing to musical rhythm (Honing & Bouwer, 2019; McGarry, Sternin, & Grahn, 2019; Merchant, Grahn, Trainor, Rohrmeier, & Fitch, 2015).

Temporal expectations are often studied in the context of some form of periodic input, such as a regular beat in music (*beat-based expectations*). However, temporal expectations can be formed based on different types of structure in the environment, which need not necessarily be periodic. For example, temporal expectations can result from learning the relationship between a cue and a particular temporal interval, or from learning (nonperiodic) sequences of temporal intervals (Nobre & van Ede, 2018). In the latter two cases, expectations rely on memory of absolute durations. We will refer to these as *memory-based expectations*. Note that elsewhere, the terms duration-based timing, absolute timing, and interval-based timing have also been used (Breska & Ivry, 2018; Merchant & Honing, 2014; Teki, Grube, Kumar, & Griffiths, 2011).

Temporal expectations have been explained by entrainment models, such as Dynamic Attending Theory (DAT). Such models propose that temporal expectations result from synchronization between internal oscillations and external rhythmic stimulation (Haegens & Zion Golumbic, 2018; Henry & Herrmann, 2014; Jones & Boltz, 1989; Large & Jones, 1999). On a neural level, the internal oscillations can be thought of as fluctuations in low-frequency oscillatory activity, or cortical excitability, such that the high-excitability phase of low frequency neural oscillations coincides with the timing of expected events, facilitating their processing by increasing sensory gain (Lakatos, Karmos, Mehta, Ulbert, & Schroeder, 2008; Schroeder & Lakatos, 2009).

It is unclear whether entrainment, thought to underlie beat-based expectations, can account for memory-based expectations, which do not rely on periodic input (Breska & Deouell, 2017b; Morillon, Schroeder, Wyart, & Arnal, 2016; Rimmele, Morillon, Poeppel, & Arnal, 2018). Arguably, memory-based expectations can better be explained within a predictive processing framework. Notably, beat-based expectations, while often thought to be dependent on entrainment, have also been modeled using hierarchical predictive coding models (Forth, Agres, Purver, & Wiggins, 2016; van der Weij, Pearce, & Honing, 2017). In such models, temporal expectations reflect the learned probability of an event at a given time, similar to learning the probability of content ("what"; Koelsch, Vuust, & Friston, 2019). Thus, whether beat-based and memory-based temporal expectations are based on shared or separate underlying mechanisms – be it entrainment, leading to changes in sensory gain, or predictive processing based on learned probabilistic information – is presently a matter of active debate. In the current EEG study, we addressed this outstanding question by directly comparing the effects of beat-based and memory-based expectations on auditory processing and behavior.

Previously, beat-based and memory-based timing have been differentiated in terms of their occurrence in different species (Honing, Bouwer, Prado, & Merchant, 2018; Honing &

Merchant, 2014), the neural networks involved (Teki et al., 2011), and in how they are affected by neuropsychological disorders (Breska & Ivry, 2018), suggesting that beat-based and memory-based expectations are subserved by separate mechanisms. However, some have argued for one integrated system for timing (Schwartze & Kotz, 2013; Teki, Grube, & Griffiths, 2012), based on neuropsychological evidence (Cope, Grube, Singh, Burn, & Griffiths, 2014), and for reasons of parsimony (Rimmele et al., 2018). Complicating this discussion, to date, many studies examining temporal expectations used isochronous sequences of events to elicit temporal expectations in general, and beat-based expectations in particular (Arnal, Doelling, & Poeppel, 2014; Breska & Deouell, 2016; Breska & Ivry, 2018; Henry & Obleser, 2012; Lakatos et al., 2013; Lawrance, Harper, Cooke, & Schnupp, 2014; Rohenkohl et al., 2012; Schwartze, Farrugia, & Kotz, 2013; Teki et al., 2011). Isochronous sequences are fully predictable, both in terms of their absolute intervals, and in terms of their ongoing periodicity, and therefore do not allow for differentiation between beat-based and memory-based expectations.

A handful of studies attempted to directly compare beat-based and memory-based expectations by using both isochronous stimuli, and stimuli that were predictable, but not periodic. In a behavioral experiment, responses to sequences that were isochronous (affording both beat-based and memory-based expectations), predictably speeding up or slowing down (affording only memory-based expectations), or with random timing (no expectations) were compared (Morillon et al., 2016). Beat-based expectations improved both perceptual sensitivity and response speed, while memory-based expectations only affected perceptual sensitivity, which was suggested to result from a special relationship between beat-based expectations and the motor system (Morillon et al., 2016). However, in another behavioral study, both beat-based and memory-based expectations improved response speed (Breska & Ivry, 2018), and phase coherence of delta oscillations, which is often used as a proxy for

neural entrainment, was shown to be similarly enhanced by memory-based and beat-based expectations (Breska & Deouell, 2017b), suggesting either that phase coherence is not a good measure of entrainment, as suggested by the authors, or that entrainment is a general, rather than a context-specific, mechanism of temporal expectations (Rimmele et al., 2018). Note that in all three studies described above (Breska & Deouell, 2017b; Breska & Ivry, 2018; Morillon et al., 2016), while care was taken to design stimuli that elicited only memory-based but not beat-based expectations, responses to isochronous stimuli were used as a proxy for beat-based expectations. In addition to affording beat-based expectations, isochronous stimuli are by definition more predictable in terms of learning their intervals than any other type of memory-based sequence (e.g., only one interval needs to be learned in an isochronous sequence). Thus, in these studies, the effects of beat-based expectations were always confounded with increases in temporal predictability based on memory. In addition, in the latter two studies (Breska & Deouell, 2017b; Breska & Ivry, 2018), static visual stimuli were used. The auditory system has repeatedly been found to be superior over the visual system in eliciting temporal expectations in humans (Grahn, 2012; Grahn, Henry, & McAuley, 2011; Zarco, Merchant, Prado, & Mendez, 2009). While it is possible to achieve equal synchronization performance (a measure of temporal expectations) with visual and auditory stimuli alike, this requires the use of moving, rather than static visual input (Grahn, 2012; Iversen, Patel, Nicodemus, & Emmorey, 2015). Thus, the stimuli used by previous studies to study beat-based expectations not only allowed for memory-based strategies to form expectations, they were arguably not optimal for creating beat-based expectations at all.

Here, to gain a better understanding of how beat-based and memory-based expectations may (differentially) influence early sensory processing, we orthogonally manipulated beat-based and memory-based expectations using auditory stimuli, and examined their effects on both behavioral and auditory ERP responses. Effects on ERPs have not been

studied before in this context and may provide a potentially fruitful way of differentiating

between beat-based and memory-based expectations. Temporal expectations have been

reported to both enhance and attenuate the auditory P1 and N1 responses. Enhancement of

sensory responses at expected time points (Bouwer & Honing, 2015; Escoffier et al., 2015;

Fitzroy & Sanders, 2015; Hsu, Hämäläinen, & Waszak, 2013; Rimmele, Jolsvai, & Sussman,

2011; Tierney & Kraus, 2013) is in line with entrainment models of temporal expectations,

which assume increased sensory gain at expected time points (Large & Jones, 1999). By

contrast, attenuation of sensory responses at expected time points (Lange, 2009; Paris, Kim,

& David, 2016; Sanabria & Correa, 2013; Schwartze et al., 2013; Sherwell, Garrido, &

Cunnington, 2017; van Atteveldt et al., 2015) is in line with predictive models of brain

function that assert more efficient processing of incoming information when predicted

information is suppressed (Friston, 2005; Marzecová, Widmann, Sanmiguel, Kotz, &

Schröger, 2017; Schröger, Kotz, & SanMiguel, 2015; Schröger, Marzecová, & Sanmiguel,

2015). Whether temporal expectations lead to enhancement or attenuation of sensory

responses may depend on the type of temporal structure that affords an expectation. Periodic

input, affording beat-based expectations, may lead to entrainment and thus increased neural

excitability at expected time points, resulting in enhancement of early sensory responses

(Haegens & Zion Golumbic, 2018). By contrast, learned probabilistic information about

timing, affording memory-based expectations, may lead to the suppression of predicted

information, resulting in attenuation of early sensory responses. Thus, if based on separate

mechanisms, one subserved by entrainment and one by predictive processing, beat-based and

memory-based expectations may have opposing effects on sensory responses, enhancement

and attenuation of responses respectively.

In addition to the effects of beat-based and memory-based expectations on behavioral

and auditory responses, we examined the effects of task relevance on both types of

expectations. Entrainment and beat-based processing have been shown to be somewhat independent of task relevance (Bouwer, Van Zuijen, & Honing, 2014; Bouwer, Werner, Knetemann, & Honing, 2016; Breska & Deouell, 2014; Rohenkohl, Coull, & Nobre, 2011). Predictive processing, however, has been shown to depend on and interact with task relevance in both the visual and the auditory domain (Hsu, Hämäläinen, & Waszak, 2018; Kok, Rahnev, Jehee, Lau, & de Lange, 2012; Paris et al., 2016; Todorovic, Schoffelen, Ede, Maris, & de Lange, 2015). Thus, if beat-based expectations rely on entrainment while memory-based expectations rely on learning probabilistic information, they may be affected by task relevance differently.

In the current study, in two experiments, we compared responses to auditory sequences that were either periodic, affording beat-based expectations, or aperiodic, thus not affording beat-based expectations. Also, sequences could either consist of fully predictable temporal intervals, affording memory-based expectations, or unpredictable, randomly concatenated intervals. We not only examined responses for events on the beat (e.g., in phase with the periodicity, at expected times), but also off the beat (e.g., out of phase with the periodicity, at less expected times), as entrainment theories predict not only increased sensory gain at expected moments, but also reduced sensory gain in between (Breska & Deouell, 2014). Thus, on the beat, when comparing periodic with aperiodic sequences, we could assess the facilitating effects of the presence of beat-based expectations, while off the beat, we could assess the effects of events occurring at times that were "mispredicted" in terms of the expected beat (Hsu, Bars, & Ha, 2015).

In Experiment 1, we examined the effects of beat-based and memory-based expectations at two positions (on and off the beat) on behavioral performance, by measuring the speed and accuracy of the detection of targets in the form of rare softer tones in the rhythmic sequences. In Experiment 2, we additionally recorded ERP responses to all non-

target sounds, to examine the effects of beat-based and memory-based expectations at two positions on P1 and N1 responses. Additionally, in Experiment 2, ERP responses were examined both when sequences were task-relevant and task-irrelevant.

In terms of behavioral outcomes, we expected faster and more accurate detection of targets with expected than unexpected timing, both for beat-based and memory-based expectations. Additionally, beat-based and memory-based expectations may interact in two distinct ways. First, the presence of beat-based expectations may lead to an increase in sensory gain for sounds on the beat, as predicted by entrainment models. Increased gain should increase the precision of memory-based predictions (Feldman & Friston, 2010), leading to enhanced effects of memory-based expectations in the presence of beat-based expectations. Thus, interacting effects of beat-based and memory-based expectations, which are larger than could be expected based on additivity, could indicate separate mechanisms, with beat-based expectations influencing the gain or precision of sensory processing through entrainment, and memory-based expectations affecting the probabilistic predictions themselves. Alternatively, if beat-based and memory-based expectations rely on shared mechanisms, the simultaneous presence of both types of expectations may lead to interference, with smaller effects of either type when both need to be engaged. In terms of ERP responses, qualitative differences between the effects of beat-based and memory-based expectations – the former leading to enhancement, and the latter to attenuation of sensory responses – could indicate different underlying mechanisms. However, if both types of expectations rely on similar mechanisms, be it based on entrainment or predictive processing, we expected them to affect sensory responses similarly.

Methods

Experiment 1

Participants. Thirty-four participants (26 women), aged between 19 and 45 years old (M = 24.6, SD = 5.7) with no history of neurological or hearing disorders took part in the experiment. Data from two participants were removed due to technical problems, leaving 32 participants for the analysis. All participants provided written consent prior to the study, and participants were reimbursed with either a monetary fee or course credit. The study was approved by the Ethics Review Board of the Faculty of Social and Behavioral Sciences of the University of Amsterdam. The statistical analysis of Experiment 1 was preregistered (aspredicted.org/blind.php?x=mt5z2h).

Stimuli. We created sound patterns of five or six consecutive temporal intervals (Figure 1A), marked by identical woodblock sounds of 60 ms length, generated in GarageBand (Apple Inc.). Patterns of five or six intervals are short enough to allow for learning of the temporal intervals (Schultz, Stevens, Keller, & Tillmann, 2013), and to not make too large demands on working memory (Grahn & Schuit, 2012). At the same time, with a total length of 1800 ms, patterns were long enough to avoid the perception of a regular, periodic beat when patterns were concatenated into sequences, as people do not readily perceive a beat with a period of 1800 ms (London, 2012). Patterns were concatenated into sequences of 128 patterns, with a final tone added to each sequence. Each sequence thus lasted for 3 minutes and 51 seconds.

Beat-based expectations. For the periodic, beat-based patterns, temporal intervals were related by the integer ratios of 1:2:2:3:4 (five intervals) and 1:1:1:2:3:4 (six intervals). The shortest interval was set at 150 ms, leading to inter-onset intervals for the other intervals of 300, 450, and 600 ms. In the periodic patterns, temporal intervals were organized to form groups of four units length (600 ms) and grouped such that a perceptually accented tone was

Figure 1. Schematic overview of the stimuli. A) Patterns of 5 (patterns 1 and 2) and 6 (patterns 3 and 4) temporal intervals were created. All patterns were 12 units length, equaling 1800 ms (1 unit = 150 ms). Periodic patterns (left, orange shades) consisted of intervals related by integer ratios 1:2:3:4. In periodic patterns, groups of intervals always added up to unit length 4, creating a beat with an inter-beat onset of 600 ms (red dotted lines). Aperiodic patterns (right, purple shades) consisted of intervals related by non-integer ratios, not allowing for the perception of a regular beat. B) Patterns were concatenated into sequences of either 128 identical patterns (predictable sequences), or 128 randomly chosen patterns (unpredictable sequences). For the unpredictable sequences, not only patterns starting at beat 1 (patterns 1.1, 2.1, 3.1, and 4.1) were used, but also cyclic permutations of these patterns, starting at beats 2 or 3 (patterns 1.2, 1.3, 2.2, etc.). For all analyses, only events after intervals with unit lengths 1 and 3 were used, to control for acoustic context.

present at the start of each group (Grahn & Brett, 2007; Povel & Essens, 1985). In these

patterns a beat could be perceived with an inter-beat interval of 600 ms (100 BPM or 1.7 Hz),

the optimal rate for human beat perception (London, 2012). These patterns could be regarded

as strictly metric, with the periodicity of the pattern always being marked by a sound (Grahn

& Brett, 2007). Each pattern consisted of 12 units length, or three beats of four units length.

To create aperiodic equivalents of the beat-based patterns, we changed the ratios by

which the temporal intervals were related. For the aperiodic patterns, intervals were related by

non-integer ratios of 1:1.4:1.4:3:5.2 (five intervals) and 1:1:1:1.4:3:4.6 (six intervals). The

aperiodic patterns were equal to their periodic counterparts in terms of length, grouping, and

number of tones. However, the aperiodic patterns did not contain a periodic beat at unit length

four. A pilot confirmed that aperiodic patterns were rated as less beat-inducing than periodic patterns.

Note that it is impossible to create sequences of sounds that are not to some extent (quasi-) periodic (Breska & Deouell, 2017a; Obleser, Henry, & Lakatos, 2017). However, the sequences clearly differed in the presence of periodicity at a rate that afforded beat-based expectations. First, in the aperiodic patterns, contrary to the periodic patterns, events did not align with the intended beat at a rate close to the ideal tempo for beat perception (1.6 Hz). This was also apparent from a spectral analysis of the waveforms (results not shown). Peaks at 1.6 Hz (the beat frequency) were larger for the beat-based sequences than the memory-based sequences. Second, while periodicity was present in the aperiodic sequences at the level of concatenated patterns (with a period of 1800 ms, or 33 BPM or 0.6 Hz), this was too slow for humans to readily perceive a beat in (London, 2012). Finally, we performed an informal pilot experiment in which we asked 17 participants to rate on a scale from 1 to 10 how strongly they heard a beat in the aperiodic and periodic patterns. On average, each periodic pattern was rated as containing more beat (on average 9.3) than each aperiodic pattern (on average 6.7), confirming that our manipulation of periodicity indeed affected perception at the level of hearing a beat. Thus, while we are aware that the aperiodic patterns could be classified as quasi or weakly periodic, for clarity, we will refer to them as aperiodic.

Memory-based expectations. Fully predictable sequences were created by concatenating 128 identical patterns into a sequence (Figure 1B). The surface structure of temporal intervals in these sequences could easily be predicted based on probabilistic information alone. Unpredictable sequences were created by concatenating 128 semi-randomly chosen patterns. Patterns were chosen both from the original patterns, which were also used for the predictable sequences (patterns starting at beat 1, see Figure 1), and from cyclic permutations of these (patterns starting at beats 2 or 3, see Figure 1). The cyclic

permutations were identical to the original patterns when looped (as in the predictable sequences), but not when concatenated in random order (as in the unpredictable sequences). Within an unpredictable sequence, only patterns with either five or six intervals were concatenated. The longest interval had a different length in the aperiodic patterns with five and six intervals (unit lengths 5.2 and 4.6 respectively). This was necessary to keep the overall length of each pattern identical. However, combining the two sets would lead to a larger number of possible temporal intervals in the aperiodic than periodic sequences, possibly confounding the effects of beat-based expectations with differences in entropy. Thus, to keep both the number of events per pattern (event density) and the number of possible intervals (entropy) identical between conditions, we did not combine the sets of rhythms with five and six intervals. Within unpredictable sequences, each pattern could occur maximally twice consecutively.

        Position. With manipulations of periodicity and predictability, we were able to compare responses to events that were expected based on a beat or on learned interval structure with responses that could not be predicted based on their timing. However, we also wanted to examine how beat-based expectations affected responses to events with unexpected timing (e.g., not unpredicted, but rather mispredicted, see also Hsu et al., 2018). Therefore, we not only probed events that were in phase with the periodicity (e.g., on the beat), but also events that were out of phase with the periodicity (e.g., off the beat, see Figure 1). The off-beat events fell at quarter-phase (150 or 450 ms) relative to the beat. We did not include events at antiphase (300 ms), as possibly, participants could perceive an additional beat at the faster subdivision rate, making these events ambiguous in terms of their metrical salience.

        We assumed that people would not perceive a beat in the aperiodic patterns, and therefore, that the distinction between on the beat and off the beat would be meaningless for these patterns in terms of temporal expectations. However, grouping effects could lead to

differences in the perception of events on and off the beat. For example, the auditory system

tends to group events in groups of two (Abecasis, Brochard, Granot, & Drake, 2005;

Brochard, Abecasis, Potter, Ragot, & Drake, 2003; Potter, Fenwick, Abecasis, & Brochard,

2009), and perceptual accents are known to arise based on the surface structure of a rhythm

(Povel & Essens, 1985). To be able to assess the effects of temporal expectations at different

positions without confounding these with differences in grouping, we classified events in the

aperiodic patterns as on-beat or off-beat, depending on their grouping in the periodic

counterpart, even when we assumed no beat would be perceived in response to the aperiodic

patterns (i.e., when we refer to an event as "on-beat" in an aperiodic pattern, it is an event that

falls on the beat in the periodic equivalent, and thus is matched to its periodic equivalent in

terms of grouping).

Targets. In the behavioral task, temporal expectations were probed implicitly, by

introducing infrequent intensity decrements as targets. Based on previous experiments, we

expected that temporal expectations would improve the detection of these targets (Bouwer &

Honing, 2015; Bouwer et al., 2014, 2016; Potter et al., 2009). Intensity decrements of 6 dB

were used (Bouwer & Honing, 2015). In each sequence of 128 patterns, 32 patterns (25

percent) contained a target. Half of the targets appeared on the beat, and half of the targets off

the beat. In each sequence, 26 targets were in positions after temporal intervals with unit

lengths 1 and 3, present in both periodic and aperiodic patterns. Only these targets were used

for the analysis, to equate their acoustic context. At least two standard patterns separated a

pattern containing a target.

Procedure. A total of 16 sequences were presented to each participant, four of each

type. Sequences of different types were semi-randomized, with each type appearing once

every four sequences, and therefore a maximum of two sequences of the same type in a row.

Upon arrival, participants completed a consent form and were allowed to practice the task.

They were instructed to avoid movement, listen to the rhythm carefully, and press a button as fast as possible when they heard a target. Participants were allowed breaks between sequences. An entire experimental session lasted for about 2 hours. Participants were tested individually in a dedicated lab at the University of Amsterdam. Sounds were presented at 70 dB SPL with one Logitech speaker positioned in front of the participants, using Presentation software (version 19.0, www.neurobs.com).

Data analysis. In total, each participant was presented with 52 targets for each condition that were included in the analysis. All responses made within 2000 ms of a target were recorded. Responses faster than 150 ms were discarded, as were responses that were more than 2.5 standard deviations from the mean of a participant's reaction time within each condition. Removal of outliers led to the exclusion of 2.9 percent of the responses in Experiment 1 and 3.1 percent of the responses in Experiment 2. Participants who did not achieve a hit rate of higher than 50 percent in any of the conditions were excluded from the analysis. In Experiment 1, on this ground, two participants were excluded, leaving 30 participants for the analysis of hit rates. One additional participant was excluded for the analysis of reaction times, as this participant had less than five valid reaction times in one condition (i.e., less than 5 out of a possible 52 targets were hits) . In Experiment 2, no participants were excluded from the analysis. Hit rates for each condition and participant and average reaction times for each condition and participant were entered into three-way repeated measures ANOVAs, with Periodicity (periodic, aperiodic), Predictability (predictable, unpredictable), and Position (on the beat, off the beat) as within-subject factors. For significant interactions ($p<0.05$), post hoc tests of simple effects were performed. For all simple effects, we report uncorrected p-values. Effect sizes are reported as partial eta squared. All statistical analyses were performed in SPSS 24.

Experiment 2

 Participants. Thirty-two participants (22 women), aged between 19 and 44 years old (M = 23.4, SD = 4.9) with no history of neurological or hearing disorders took part in Experiment 2. Data of one participant were removed due to excess noise in the EEG signal, leaving thirty-one participants for the analysis. All participants provided written consent prior to the study, and participants were reimbursed with either a monetary fee or course credit. The study was approved by the Ethics Review Board of the Faculty of Social and Behavioral Sciences of the University of Amsterdam.

 Stimuli and Procedure. The materials and procedure for Experiment 2 were identical to those for Experiment 1. In Experiment 2, participants additionally completed an unattended version of the experiment. In the unattended condition, participants were asked to ignore the rhythmic sequences and focus on a self-selected muted movie, rendering the stimuli task irrelevant. All participants first completed the unattended EEG experiment, and subsequently the attended EEG experiment. The EEG recording lasted for around 2 hours. Participants were encouraged to take breaks if needed. During the recording, data quality was assessed online by the experimenter, and if needed, channels were refitted and extra electrode gel was applied. For Experiment 2, one experimental session was about 4 hours, including breaks, practice, and setting up equipment.

 EEG recording. EEG was recorded using a 64-channel Biosemi Active-Two acquisition system (Biosemi, Amsterdam, The Netherlands), with a standard 10/20 configuration and additional electrodes for EOG channels, on the nose, on both mastoids, and on both earlobes. The EEG signal was recorded at 1 kHz.

 EEG analysis. Preprocessing was performed in MATLAB and EEGLAB (Delorme & Makeig, 2004). Data were offline re-referenced to linked mastoids, bad channels were removed, and independent component analysis was used to remove eye-blinks. Subsequently,

bad channels were replaced by values interpolated from the surrounding channels. Visual inspection of the ERPs revealed a postauricular muscle response (PAM) in several subjects. The auditory evoked potential can be easily contaminated by the PAM response (Bell, Smith, Allen, & Lutman, 2004; Picton, Hillyard, Krausz, & Galambos, 1974). To avoid contamination, we re-referenced the data to earlobes for all further analyses. For completeness, we also report results from the mastoid-referenced data.

Data were offline down-sampled to 512 Hz, and filtered using 0.1 Hz high-pass and 40 Hz low-pass finite impulse response filters. Epochs for each condition separately were extracted for non-target sounds, from 200 ms preceding the onset of each event till 500 ms after the onset of each event. Only epochs for events following an interval of unit length 1 or unit length 3 (150 and 450 ms respectively) were included, to equate the acoustic context of events used in the analysis. Also, the first 12 sounds of each sequence (2 whole patterns) were excluded from the analysis, to allow for the build-up of expectations. Epochs with a voltage change of more than 150 microvolt in a 200 ms sliding window were rejected from further analysis. For each condition and participant, epochs were averaged to obtain ERPs and baseline corrected using the average voltage of the 50 ms window preceding each sound. Finally, ERPs were averaged over participants to obtain grand average waveforms.

Peak latencies for the P1 and N1 responses were determined independent from the statistical analysis, from the average waveform collapsed over all conditions. P1 peaked at 58 ms after tone onset. We defined P1 amplitude as the average amplitude in a 20 ms window around the peak (48-68 ms). N1 peaked at 124 ms, and was more distributed in time. Thus, we defined N1 amplitude as the average amplitude from a 40 ms window around the peak (104-144 ms). Auditory evoked potentials are known to be maximal over fronto-central electrodes (Picton et al., 1974; Ruhnau, Herrmann, Maess, & Schröger, 2011), which was also observed in the current dataset. Therefore, ERP amplitudes were computed from the average

of a cluster of 15 fronto-central electrodes: F3, F1, Fz, F2, F4, FC3, FC1, FCz, FC2, FC4, C3, C1, Cz, C2, and C4. All statistics and figures reported here are based on the average amplitude from this region of interest.

Statistical analysis. Amplitudes for P1 and N1 were entered into repeated measures ANOVAs, with four within-subject factors: Periodicity (periodic, aperiodic), Predictability (predictable, unpredictable), Position (on the beat, off the beat), and Attention (attended, unattended). For significant interactions ($p<0.05$), subsequent tests of simple effects were performed. For all simple effects, we report uncorrected p-values. Effect sizes are reported as partial eta squared. Analyses were performed in SPSS 24.

Comparing beat-based and memory-based expectations directly. While the factorial design included both Periodicity and Predictability as factors, the ANOVAs do not allow for a direct comparison of their effects. To directly compare the effects of beat-based and memory-based expectations, we performed two additional analyses. In both analyses, the effect of beat-based expectations was quantified as the difference between responses on the beat in periodic and aperiodic sequences. The effect of memory-based expectations was quantified as the difference between responses in predictable and unpredictable sequences. For the latter, we only included responses on the beat, to make sure that possible differences between beat-based and memory-based expectations could not be attributed to differences in grouping.

First, we compared the effects of beat-based and memory-based expectations in the P1 and N1 windows directly using paired-samples T-tests. To quantify the possibility that no differences were present, we performed both traditional and Bayesian T-tests using JASP (JASP Team, 2019; Wagenmakers et al., 2018). We estimated Bayes factors using a Cauchy prior distribution ($r = 0.71$), with as a null hypothesis no differences between the effects of beat-based and memory-based expectations. In addition, we performed a robustness check as

implemented in JASP, to assess whether the results would change with a different prior ($r = 1$) as proposed originally for Bayesian T-tests (Jeffreys, 1961; Wagenmakers et al., 2018).

Second, we directly compared the effects of memory-based and beat-based expectations using cluster-based permutation tests. With this approach we could examine potential differences at all timepoints and at all electrodes, while taking into account the multiple comparisons along both the spatial and time axes. As ERP components often overlap in time, their real peaks may be obscured in grand average waveforms (Luck, 2005). The use of cluster-based permutation testing allowed us to make sure we did not miss potential differences between beat-based and memory-based expectations by selectively examining peak time windows and selected clusters of electrodes. Cluster-based permutation tests were performed using the Fieldtrip toolbox (Oostenveld, Fries, Maris, & Schoffelen, 2011). All timepoints in the 150 ms time window following the onset of sound events were included. We limited the analysis to the first 150 ms, to avoid contamination of subsequent sounds, which in some but not all conditions, could occur 150 ms following an event. Clusters were formed based on adjacent time-electrode samples that survived a statistical threshold of $p<0.01$ when comparing the conditions of interest with dependent samples T-tests. Clusters were subsequently evaluated with permutation tests, using 2000 iterations.

Results

Behavioral results

Hit rates and reaction times for all conditions from both Experiment 1 and 2 are shown in Table 1 and Table 2 and Figure 2. In Experiment 2, all results from Experiment 1 were replicated with comparable effect sizes.

Memory-based expectations. In general, targets were detected more often and faster in predictable than in unpredictable sequences, as reflected in a significant main effect of Predictability for hit rates (Experiment 1: $F(1,29) = 39.4$, $p < 0.001$, $\eta_p^2 = 0.58$; Experiment 2: $F(1,30) = 46.4$, $p < 0.001$, $\eta_p^2 = 0.61$) and reaction times (Experiment 1: $F(1,28) = 19.6$, $p < 0.001$, $\eta_p^2 = 0.41$; Experiment 2: $F(1,30) = 25.0$, $p < 0.001$, $\eta_p^2 = 0.46$). The simple effect of Predictability was significant for all comparisons for hit rates (all uncorrected ps < 0.016, see Table 1) and reaction times (all uncorrected ps < 0.023, except for targets off the beat in periodic sequences in Experiment 2; $p = 0.055$). Thus, as expected, memory-based expectations led to improved detection of targets at expected time points. Figure 3 (top) shows the main effect of Predictability on target detection performance collapsed over the levels of Periodicity and Position.

		Experiment 1 (N=30)			Experiment 2 (N=31)		
		Periodic	Aperiodic	Effect of beat-based expectations	Periodic	Aperiodic	Effect of beat-based expectations
Beat	Predictable	82 (14)	81 (15)	+1	79 (17)	79 (17)	0
	Unpredictable	73 (19)	69 (23)	+4	72 (17)	67 (20)	+5**
Effect of memory-based expectations		+9**	+12**		+7**	+12**	
Offbeat	Predictable	63(20)	68 (21)	-5*	64 (21)	72 (19)	-8**
	Unpredictable	52 (22)	61 (26)	-9**	51 (20)	60 (21)	-9**
Effect of memory-based expectations		+11**	+7*		+13**	+12**	

Table 1. Hit rates for all conditions in Experiment 1 and Experiment 2. Hit rates are reported as percentage hits. Standard deviations in brackets. *Simple effect significant at p<0.05 (uncorrected). **Simple effect significant at p<0.01 (uncorrected).

Figure 2. Hit rates (top) and reaction times (bottom) show improved target detection for events at expected time points. Memory-based expectations lead to better and faster target detection in Experiment 1 (left) and Experiment 2 (right). Beat-based expectations only lead to improvements in the absence of memory-based expectations in Experiment 2, but consistently lead to deteriorated detection of targets off the beat in both experiments.

Beat-based expectations. In Experiment 2, we found a main effect of Periodicity on hit rates ($F(1,30) = 6.59$, $p = 0.016$, $\eta_p^2 = 0.18$). However, as expected, the effect of Periodicity on hit rates in both experiments depended on the position of the target. On the beat, targets were detected more often in periodic than aperiodic sequences, while off the beat, targets were detected more often in aperiodic than periodic sequences, as reflected in a significant interaction between Periodicity and Position (Experiment 1: $F(1,29) = 35.0$, $p < 0.001$, $\eta_p^2 = 0.54$; Experiment 2: $F(1,30) = 39.3$, $p < 0.001$, $\eta_p^2 = 0.57$). In line with the

results for hit rates, numerically, detection of targets off the beat, but not on the beat, was slower in periodic than aperiodic sequences. However, the interaction between Periodicity and Position did not reach significance for the reaction times (Experiment 1: $F(1,28) = 3.74$, $p = 0.063$, $\eta_p^2 = 0.12$; Experiment 2: $F(1,30) = 1.91$, $p = 0.18$, $\eta_p^2 = 0.06$). Figure 3 (bottom) shows the main effect of Periodicity on target detection performance separate for on-beat and off-beat targets, and collapsed over the levels of Predictability.

		Experiment 1 (N=29)			Experiment 2 (N=31)		
		Periodic	Aperiodic	Effect of beat-based expectations	Periodic	Aperiodic	Effect of beat-based expectations
Beat	Predictable	538 (78)	541 (78)	-3	517 (59)	509 (60)	+8
	Unpredictable	557 (74)	571 (80)	-14	542 (66)	554 (75)	-12
Effect of memory-based expectations		-19*	-30**		-25**	-45**	
Offbeat	Predictable	573 (84)	571 (81)	+2	550 (93)	531 (76)	+19
	Unpredictable	601 (92)	591 (96)	+10	569 (81)	571 (96)	-2
Effect of memory-based expectations		-28*	-20*		-19	-40**	

Table 2. Reaction Times in ms for all conditions in Experiment 1 and Experiment 2. Standard deviations in brackets. *Simple effect significant at p<0.05 (uncorrected). **Simple effect significant at p<0.01 (uncorrected).

Interaction between beat-based and memory-based expectations. The Predictability and Periodicity of the sequences also affected hit rates in interaction, though depending on the position of the target, as reflected in a significant three-way interaction between Periodicity, Position, and Predictability (Experiment 1: $F(1,29) = 6.69$, $p = 0.015$, $\eta_p^2 = 0.19$; Experiment 2: $F(1,30) = 5.59$, $p = 0.025$, $\eta_p^2 = 0.16$). Off the beat, the simple effect of Periodicity was significant for all comparisons (all uncorrected ps < 0.029). Thus, beat-based expectations led to decreased detection of targets when they were presented out of phase with the periodicity and therefore occurred at time points that were mispredicted with regards to the beat. On the beat, the simple effect of Periodicity only reached significance for

Figure 3. Temporal expectations improve target detection. Memory-based expectations (A) led to both higher hit rates and faster reaction times (main effect of Predictability). Beat-based expectations mainly affected hit rates by deteriorating detection for mispredicted events (off-beat; bottom of panel B). Thick lines represent the mean, thin lines are single participants.

unpredictable sequences, and only in Experiment 2 (uncorrected $p = 0.003$), showing improved detection through beat-based expectations only in the absence of memory-based expectations. In Experiment 2, we also found an interaction between Periodicity and Predictability for reaction times, with a stronger effect of Predictability in aperiodic than periodic sequences ($F(1,30) = 6.67$, $p = 0.015$, $\eta_p^2 = 0.18$).

In addition to effects of memory-based and beat-based expectations, we found a main effect of Position for hit rates (Experiment 1: $F(1,29) = 99.9$, $p < 0.001$, $\eta_p^2 = 0.78$;

Experiment 2: $F(1,30) = 100.3$, $p < 0.001$, $\eta_p^2 = 0.77$) and reaction times (Experiment 1: $F(1,28) = 37.1$, $p < 0.001$, $\eta_p^2 = 0.57$; Experiment 2: $F(1,30) = 14.4$, $p = 0.001$, $\eta_p^2 = 0.32$). Targets on the beat were detected more often and faster than targets off the beat, even in aperiodic sequences, most likely reflecting differences in surface grouping.

Thus, behaviorally, memory-based expectations improved target detection both in terms of response accuracy and response speed (see Figure 3). Beat-based expectations similarly lead to improved target detection for targets at expected time points (on-beat), but only when no memory-based expectations were present, and only in terms of response accuracy, not response speed. The improvements in target detection afforded by beat-based expectations were thus small and dependent on memory-based expectations. In contrast, at unexpected time points (off-beat, "mispredicted"), beat-based expectations lead to decrements in performance, both in the absence and presence of memory-based expectations. Interestingly, while we found an interaction between beat-based and memory-based expectations, it was in the opposite direction of what we expected: rather than enhancing each other, when both types of expectations were present, their effects were diminished.

EEG results

Figure 4 shows the auditory evoked potentials for all conditions. Average amplitudes as extracted from the P1 and N1 time-windows and fronto-central region of interest are depicted in Figure 5.

Memory-based expectations. As expected, both P1 and N1 responses were smaller for predictable than unpredictable events, reflected in a main effect of Predictability (P1: $F(1,30) = 10.3$, $p = 0.003$, $\eta_p^2 = 0.26$; N1: $F(1,30) = 4.52$, $p = 0.042$, $\eta_p^2 = 0.13$), showing attenuation through memory-based expectations. The effect of Predictability did not depend on task relevance (e.g., none of the interactions including Predictability and Attention reached significance: all $ps > 0.09$). Figure 6 shows a summary of the main effects of memory-based

Figure 4. Temporal expectations reduce auditory responses. Temporally expected events evoked smaller auditory P1 and N1 responses, regardless of whether they were attended (task-relevant) or not. Grand-average waveforms are shown for each condition separately. The time windows used for the ANOVA analyses of P1 and N1 are indicated with rectangles. Both memory-based and beat-based expectations attenuated P1 and N1 responses to expected events. N1 responses were additionally enhanced for events off the beat in periodic sequences, i.e., when they were unexpected with regard to the periodicity.

expectations on auditory-evoked potentials, collapsed over the different levels of Position and

Periodicity. The attenuation of the auditory P1 and N1 responses associated with memory-

based expectations suggests processes related to suppression of predicted information, rather

than memory-based expectations leading to changes in sensory gain. While statistically

independent of task relevance, numerically, for the P1 response the effects of memory-based

expectations were larger in attended than unattended conditions.

Beat-based expectations. Figures 7 and 8 show a summary of the effects of beat-

based expectations on auditory-evoked potentials, collapsed over the different levels of

Predictability and separate for responses to events on the beat (Figure 7) and off the beat

Figure 5. Temporal expectations reduce auditory responses. Amplitudes of P1 and N1 averaged over the time windows used for analysis (see Figure 4) and fronto-central region of interest are shown.

(Figure 8). Unexpectedly, for P1, the effect of Periodicity did not depend on the position of an event (no interaction between Periodicity and Position: $F(1,30) = 0.036$, $p = 0.85$, $\eta_p^2 = 0.001$). Instead, we found a main effect of Periodicity ($F(1,30) = 11.3$, $p = 0.002$, $\eta_p^2 = 0.27$), with smaller P1 responses to events in periodic than aperiodic sequences, both on-beat and off-beat. For the N1 response, we did find a significant interaction between Periodicity and Position ($F(1,30) = 12.7$, $p = 0.001$, $\eta_p^2 = 0.30$). Like for memory-based expectations, the

Figure 6. Memory-based expectations attenuated auditory responses. Auditory evoked potentials showing the main effect of Predictability on P1 and N1 responses are shown on the left. The main effect of Predictability is depicted as the difference between responses to events in predictable and unpredictable sequences, collapsed over the levels of Periodicity and Position (i.e., independent of beat-based expectations). In both attended and unattended conditions, P1 and N1 responses were larger for unpredictable than for predictable events. On the right, P1 and N1 amplitudes are shown for all participants separately, with the mean plotted on top.

Figure 7. Beat-based expectations attenuated auditory responses (on-beat). Auditory evoked potentials showing the effect of Periodicity on the beat on P1 and N1 responses are shown on the left. The effect of Periodicity is depicted as the difference between responses to on-beat events in periodic and aperiodic sequences, collapsed over the levels of Predictability (i.e., independent of memory-based expectations). In both attended and unattended conditions, P1 and N1 responses were smaller in amplitude for periodic than aperiodic events on the beat. On the right, P1 and N1 amplitudes are shown for all participants separately, with the mean plotted on top. Note that here, the confidence intervals around ERP waveforms represent the variability of the difference waves, in which most of the between-subject variance has been eliminated through subtraction.

effects of beat-based expectations were independent of task-relevance (e.g., none of the interactions including Periodicity and Attention reached significance: all $ps>0.06$).

On the beat, both P1 and N1 responses were smaller to events in periodic than aperiodic sequences (though only marginally so for the N1: $p = 0.055$), Smaller responses to events in periodic (i.e., beat-based expected) than aperiodic (no beat-based expectations present) sequences could indicate attenuation of expected events through beat-based expectations, in line with predictive processing, and similar to the effects of memory-based expectations on the P1 and N1 response. Interestingly, similar to the effects of memory-based expectations, while statistically independent of task relevance, the effects of beat-based expectations on the P1 response were numerically driven by the attended condition.

Off the beat, N1 responses were larger to events in periodic than aperiodic sequences ($p < 0.001$). Thus, N1 responses were smallest for events that were expected (i.e., on-beat, in phase with the periodicity), largest for events that were unexpected (i.e., off-beat, "mispredicted", out of phase with the periodicity), with the amplitude of the N1 to events that were neither expected nor unexpected (i.e., in aperiodic sequences where no beat-based expectations were present) in between. Attenuation of responses to expected events, and enhancement of responses to unexpected events suggests that first order predictions (cf. (Koelsch et al., 2019), rather than changes in sensory gain, underlie the effects of beat-based expectations on perception, similar to the effects of memory-based expectations. However, off the beat, we also found smaller P1 responses to events in periodic than aperiodic sequences, in line with resource withdrawal and smaller responses off the beat in the presence of an underlying periodic beat, as proposed by DAT. This divergent result for P1 responses off the beat is hard to interpret and warrants further research.

In addition to the effects of memory-based and beat-based expectations, P1 responses were larger in amplitude for attended than unattended events (main effect of Attention:

Figure 8. Beat-based expectations enhanced N1 responses for unexpected sounds (off-beat).
Auditory evoked potentials showing the effect of Periodicity off the beat on P1 and N1 responses are shown on the left. The effect of Periodicity is depicted as the difference between responses to off-beat events in periodic and aperiodic sequences, collapsed over the levels of Predictability (i.e., independent of memory-based expectations). In both attended and unattended conditions, N1 responses were larger in amplitude for periodic than aperiodic events off the beat, while P1 responses were larger for aperiodic events. On the right, P1 and N1 amplitudes are shown for all participants separately, with the mean plotted on top. Note that here, the confidence intervals around ERP waveforms represent the variability of the difference waves, in which most of the between-subject variance has been eliminated through subtraction.

$F(1,30) = 74.3$, $p < 0.001$, $\eta_p^2 = 0.71$). Finally, we found a main effect of Position on N1

responses ($F(1,30) = 10.3$, $p = 0.003$, $\eta_p^2 = 0.26$). However, the effect of position depended

on Attention (interaction between Position and Attention: $F(1,30) = 15.9$, $p < 0.001$, $\eta_p^2 = 0.35$). In the attended condition, N1 responses to events off the beat were larger than to events

on the beat ($p < 0.001$), while in the unattended condition, there was no difference between

responses on and off the beat ($p = 0.74$). Likely, these effects of Position, similar to the effects

in the behavioral experiments, were due to grouping differences.

To check whether observed results were not confounded by our choice of earlobe

reference, we repeated the analysis with mastoid reference, as is more customary in auditory

ERPs analyses. Note that these results may have been confounded with the PAMR response,

which was present to some degree in several subjects. With mastoid reference, results

generally were the same as with earlobe reference. However, with mastoid reference we

found a four-way interaction between Attention, Predictability, Periodicity and Position for

the P1 ($F(1,30) = 5.18$, $p = 0.030$, $\eta_p^2 = 0.15$). This four-way interaction did not reach

significance for the data referenced to earlobes ($F(1,30) = 2.64$, $p = 0.11$, $\eta_p^2 = 0.081$). To

pursue the interaction in the mastoid-referenced data, we split the data between attended and

unattended conditions. In the unattended condition, no significant effects were present. In the

attended condition, we found a three-way interaction between Periodicity, Predictability, and

Position ($F(1,30) = 4.50$, $p = 0.04$, $\eta_p^2 = 0.13$). The effect of Periodicity on the P1 response

was significant only for events off the beat in predictable sequences (uncorrected $p = 0.005$),

and for events on the beat in unpredictable sequences (uncorrected $p = 0.005$). While there

were thus small differences in significance between the data sets, the main findings – the

effect of Predictability for both P1 and N1 and interaction between Periodicity and Position

for N1 – were robust to changes in reference.

Comparing beat-based and memory-based expectations directly. Finally, and crucial to our main question, we compared the effects of memory-based and beat-based temporal expectations on auditory-evoked responses directly. We subtracted responses to events on the beat in unpredictable from predictable sequences to index the effects of memory-based expectations, and we subtracted responses to events on the beat in aperiodic from periodic sequences to index the effects of beat-based expectations. Subsequently, we compared the results of this subtraction procedure, both in the restricted time windows around the P1 and N1 peaks, using Bayesian and traditional T-tests, and in a window from 0 to 150 ms after the onset of sound events using cluster-based permutation testing. As the ANOVA yielded somewhat inconclusive results regarding the effects of task relevance, here we report analyses split over attended and unattended conditions, as well as analyses in which we collapsed over attention conditions.

As is visible in Figure 9, the difference waves indexing the two different types of temporal expectations examined have a very similar morphology. Indeed, traditional t-tests showed no significant differences between the effects of Predictability and Periodicity in the P1 window (attended: $t(30) = 1.24$, $p = 0.23$; unattended: $t(30) = -0.88$, $p = 0.39$; combined: $t(30) = 0.33$, $p = 0.74$), nor in the N1 window (attended: $t(30) = 0.57$, $p = 0.57$; unattended: $t(30) = -0.97$, $p = 0.34$; combined: $t(30) = -0.35$, $p = 0.73$). The Bayes factors indicated evidence in favor of the null hypothesis (no difference between conditions), for the P1 window (attended: $BF_{01} = 2.6$; unattended: $BF_{01} = 3.7$; combined: $BF_{01} = 5.0$) and the N1 window (attended: $BF_{01} = 4.5$; unattended: $BF_{01} = 3.4$; combined: $BF_{01} = 4.9$). In other words, the data were 2.6 to 5.0 times more likely to occur under the null hypothesis than in the presence of differences between the effects of Periodicity and Predictability. Bayes factors >3 are regarded as moderate evidence in favor of the null hypothesis (Wagenmakers et al., 2018). The robustness check indicated that the results did not change as a function of the prior used.

With the more traditional prior (r = 1), Bayes factors in favor of the null hypothesis ranged from 3.5 to 6.8.

Figure 9. Beat-based and memory-based expectations similarly affected auditory stimulus processing. Effects of beat-based expectations are defined as the difference between responses to events on the beat in periodic and aperiodic sequences. Effects of memory-based expectations are defined as the difference between responses to events on the beat predictable and unpredictable sequences. When comparing the effects of beat-based and memory-based expectations directly using t-tests and cluster-based permutation tests, no significant differences were found.

Cluster-based testing on the ERP waveforms showed no significant differences between the effects of Periodicity and Predictability on the auditory evoked potential, neither in attended, nor in unattended conditions, nor when we collapsed the data over attended and unattended conditions. Thus, using traditional T-tests, Bayesian T-tests, and cluster-based testing, we could not find evidence suggesting differences between the effects of memory-based and beat-based expectations on early auditory responses.

Discussion

Temporal expectations facilitate sensory processing and perception in dynamic environments, and play an important role in synchronizing our actions to regularities in the outside world, for example, when dancing to a beat in music (Honing & Bouwer, 2019; McGarry et al., 2019). Currently, little is still known about whether shared or separate mechanisms contribute to temporal expectations based on different types of structure in the environment. Specifically, it is unclear whether beat-based expectations (based on periodicity of the input) and memory-based expectations (based on learned predictions of absolute intervals) affect sensory processing and performance in similar ways, independently of each other or in interaction (Breska & Deouell, 2017b; Nobre & Rohenkohl, 2014). Moreover, to what extent these effects rely on task relevance is also still unclear.

In the current study, we show that beat-based and memory-based expectations cannot be differentiated in terms of their effects on auditory processing and/or performance. Both types of expectations lead to enhanced detection of expected events, and to attenuation of auditory responses to those events, the latter independent of task relevance. Also, beat-based and memory-based expectations interacted, with smaller behavioral effects when both were present. These findings are in line with the notion that beat-based and memory-based expectations are subserved by a shared mechanism for temporal predictive processing (Breska & Deouell, 2017b; Rimmele et al., 2018; Teki et al., 2012). Yet, memory-based expectations enhanced target detection even for events that were fully predictable based on the beat. Also, beat-based expectations lead to reduced target detection and enhanced auditory responses to events out of phase with the periodicity, even when these events were fully predictable based on memory. These findings suggest that while the effects of beat-based and memory-based expectations on sensory processing and performance are the same, the underlying computation may in fact be separate, with beat-based expectations relying (in part) on a

rhythmic processing mode characterized by withdrawal of resources off the beat (Breska &

Deouell, 2017b). Below, we discuss these findings and their theoretical implications in detail.

Behaviorally, both beat-based and memory-based expectations facilitated target

detection in a rhythmic sound sequence, as reflected by enhanced hit rates for targets with

expected timing. Although only memory-based expectations improved response speed, the

results for beat-based expectations were numerically in the same direction. Thus, unlike

previous research (Morillon et al., 2016), we did not find a qualitative difference between

beat-based and memory-based expectations on performance. However, in previous work,

beat-based expectations were elicited by isochronous sequences, which also elicit memory-

based expectations, rendering interpretation of their findings difficult.

That the effects of beat-based and memory-based expectations on auditory processing

may rely (in part) on shared mechanisms may further be supported by the observed interaction

between the two types of expectations. Beat-based facilitation of detection rates only reached

significance in the absence of memory-based expectations. Given that detection rates on

average did not exceed 82 percent, even in the fully predictable sequences, it is unlikely that

the absence of a beat-based effect here was due to a ceiling effect. Moreover, we expected

beat-based expectations to facilitate memory-based expectations, similar to the facilitation of

content predictions ("what") afforded by temporal expectations (Auksztulewicz et al., 2018;

Hoch, Tyler, & Tillmann, 2013; Schwartze, Rothermich, Schmidt-Kassow, & Kotz, 2011;

Selchenkova, Jones, & Tillmann, 2014). However, if anything, the effect of predictability on

target detection was smaller in the periodic than the aperiodic sequences. Thus, when both

types of expectations were present, their effects on auditory processing and performance were

diminished. This may suggest that beat-based and memory-based expectations also to some

extent compete for limited capacity temporal processing to form expectations, leading to

interference when both need to be engaged.

In correspondence to our behavioral findings, our ERP results also do not suggest qualitative differences between the effects of beat-based and memory-based expectations. Both P1 and N1 responses to expected events were attenuated, in line with theories about predictive processing, and explained by assuming that the brain only processes input that is not predicted (Friston, 2005; Schröger, Marzecová, et al., 2015). Effects of beat-based expectations are often explained by entrainment models and DAT, which assume increases in sensory gain at expected time points (Henry & Herrmann, 2014; Large & Jones, 1999), leading to enhanced, rather than attenuated sensory responses (Haegens & Zion Golumbic, 2018). Our results, however, do not show enhanced auditory responses due to beat-based expectations.

While the changes in sensory gain suggested by entrainment have often been equated with fluctuations in attention (Henry & Herrmann, 2014; Lakatos et al., 2008; Large, Herrera, & Velasco, 2015), other studies have also shown that entrainment leads to attenuation rather than enhancement of sensory responses (O'Connel et al., 2015; van Atteveldt et al., 2015), like the beat-based expectations in the current study. Moreover, the effects of periodicity, which guides expectations bottom-up, can be dissociated from the effects of task relevance and general top-down attention mechanisms (Kunert & Jongman, 2017). Thus, while entrainment may lead to fluctuations in neural excitability, these may be related to predictions, rather than attention, as proposed by DAT.

Several studies have found enhancement of sensory responses when manipulating temporal expectations, which may seem contradictory to the current findings. However, first, the observed enhancement may depend on the use of different manipulations of temporal expectations. For example, when temporal expectations are cue-based (Hsu et al., 2013), they may indeed be manipulations of the relevance of a time point, rather than prediction (Lange, 2013). Second, when comparing responses to sounds in phase and out of phase with some

periodicity, differences in evoked responses may be due to grouping differences between sounds on and off the beat (Schnupp, Rajendran, Harper, Garcia-Lazaro, & Lesica, 2017). Finally, the perception of hierarchical structure in music (meter) may lead to perceived illusory metrical accents (Bouwer, Burgoyne, Odijk, Honing, & Grahn, 2018; Repp, 2010), causing enhanced responses on the beat when compared to off the beat. Indeed, two studies that specifically manipulated beat-based expectations by asking participants to imagine accents on the beat found enhancement of sensory responses (Iversen, Repp, & Patel, 2009; Schaefer, Vlek, & Desain, 2010). To sum up, it may be that entrainment and beat-based expectations are not necessarily related to fluctuations in attention, and that previously reported enhancement of auditory responses is due to grouping and metrical accenting, which we controlled for in the current study by including the aperiodic sequences. Which specific task- and stimulus-related factors cause auditory responses to be enhanced, and whether attention plays a role in this at all, remains an interesting question for further research.

In addition to considering the effects of beat-based expectations at expected time points, we also looked at how beat-based expectations affected processing at time points that were unexpected, or rather, mispredicted, based on the beat (i.e., off-beat). On these positions N1 responses were larger for events in periodic sequences than for events with similar grouping structure in aperiodic sequences. Enhancement of auditory processing of events off the beat when periodicity is present is in line with these events being more unexpected than their aperiodic counterparts. Behaviorally, we found deteriorated target detection for events off the beat in periodic sequences when compared to aperiodic sequences, in line with reduced processing of events off the beat. Importantly, detection was hampered even if event timing was fully predictable based on learning the sequence of intervals.

Our findings illustrate the importance of also examining effects of beat-based expectations off the beat in distinguishing between beat-based and memory-based

expectations. They support the view that beat-based expectations may allow the brain to go into the more efficient "rhythmic mode" of processing, instead of a continuous "vigilance mode" associated with non-periodic input. Rhythmic mode may be accompanied by automatic suppression of out of phase input when entrainment occurs (Schroeder & Lakatos, 2009; Zoefel & Vanrullen, 2017). Beat-based, but not memory-based expectations have indeed been associated with withdrawal of resources from unexpected moments in time, as apparent from immediate CNV resolution after expected time points for beat-based, but not memory-based expectations (Breska & Deouell, 2017b). A rhythmic processing mode characterized by withdrawal of resources off the beat, rather than focusing of resources on the beat, could also explain why beat-based expectations only weakly affected responses on the beat, especially when memory-based expectations were also present. With the current design, we could not probe events that were mispredicted in terms of memory-based expectations. Thus, we cannot be sure that memory-based expectations do not show similar withdrawal of resources from time points where no event is expected. However, focusing on the effects of expectations at unexpected instead of expected time points may be an interesting way to compare beat-based and memory-based expectations in the future.

We did not find any statistically significant interactions between the effects of expectations and task relevance. Off the beat, the independence of the effect of beat-based expectations of task relevance may be related to the fact that off the beat, events were mispredicted, rather than unpredicted (e.g., the aperiodic sequences did not allow for the formation of beat-based expectations, making all events unpredicted based on a beat, while in the periodic sequences, events off the beat were not in line with the beat-based expectations that could be formed). The effects of mispredicted stimuli have been shown to be independent of task relevance (Hsu et al., 2018), reminiscent of the independence of the MMN, an ERP

response to mispredicted sounds, from task relevance (Näätänen, Paavilainen, Rinne, & Alho, 2007).

For the facilitating effects of both beat-based (on-beat) and memory-based expectations, no interaction with task relevance was present either. However, based on the current data, we cannot rule out that task relevance may facilitate effects of temporal expectations, as numerically, effects were larger in the attended than unattended condition. Indeed, with mastoid reference, the (four-way) interaction with Attention did reach significance for the P1 response. We can therefore not draw strong conclusions about the interaction between task relevance and temporal expectations, other than noting that task relevance does not seem to be a prerequisite for expectations to develop. Whether task relevance interacts with expectations to shape perception is currently a matter of active debate. Recently, several studies have found interactions between prediction and attention, when looking at feature predictions (Auksztulewicz & Friston, 2015; Smout, Tang, Garrido, & Mattingley, 2019), and spatial predictions (Fardo et al., 2017; Marzecová et al., 2017). However, none of these studies looked at temporal predictions, and none found interacting effects as early as the P1 and N1, as studied here. Indeed, a recent study specifically aimed at uncovering earlier effects of attention and prediction found interacting effects of attention and prediction on the P3 in the absence of effects earlier in the processing stream (Alilović, Timmermans, Reteig, van Gaal, & Slagter, 2019). Thus, the lack of interaction observed here may be due to the type of predictions we studied, and the latency of the effect. Moreover, it has been suggested that temporal expectations are fundamentally different from feature and spatial expectations (Auksztulewicz et al., 2018; Sherwell et al., 2017), which would hint that the absence of an interaction between temporal expectations and task relevance may in fact be informative of the underlying process.

In the current study, we strictly controlled the acoustic context preceding the sounds of interest (i.e., the last time point prior to the events used in the analysis that differed in terms of the occurrence of preceding sounds was at -300 ms relative to onset). Yet, we cannot rule out that the longer latency portion of responses to sounds preceding the sounds of interest bled into the ERP waveforms to some extent. In particular, this could have affected the results by affecting the baseline. In addition, if beat-based expectations elicit low frequency entrainment, differences in the phase of the underlying oscillation could lead to a baseline offset. Issues with differences in the baseline can be avoided by using stringent high-pass filtering, eliminating both the beat frequency and the longer-latency portion of the auditory ERP (cf. Chang, Gavin, & Davies, 2012), but such filters have been shown to lead to significant distortions of the P1 waveform, and can even shift low-frequency information from the N1 range into the P1 response (Liljander, Holm, Keski-Säntti, & Partanen, 2016; Tanner, Morgan-Short, & Luck, 2015). While filtering as such is not optimal to eliminate potential confounds in the baseline, here, for the attenuating effects of expectations on both P1 and N1, such a confound is unlikely. Differences in baseline activity would result in a general shift of the ERP waveform, but the attenuation for expected events was present both for P1 and N1 – components of opposite polarity. Thus, our conclusions with regards to the qualitative effects of temporal expectations, attenuation instead of enhancement, are independent of baseline shifts.

For the N1 enhancement we observed for off-beat events in periodic sequences, the effect of baseline differences is harder to rule out, as here, the direction of the effect was opposite for the P1 (attenuation, see Figure 8). This touches upon a fundamental issue when examining temporal expectations in general, and beat-based expectations in particular. Humans are most apt at perceiving a beat in auditory stimuli (Grahn, 2012), at a tempo of around 100-120 beats per minute (London, 2012). When moving away from isochronous

stimuli, this will automatically result in rapid successions of sounds, making it difficult to completely eliminate effects of previous sounds in ERPs, even in a paradigm as highly controlled as ours. Also, this prevents the analysis of ERPs after subsequent sounds are presented, in our case 150 ms after the onset of the sounds of interest, even if some effects of expectations may only be visible at longer latencies (Alilović et al., 2019). Balancing the need for ecologically valid stimuli, optimal for eliciting (beat-based) temporal expectations, with the need for strict experimental control allowing for easy assessment of separate ERP components, remains a big challenge for future research (Bouwer et al., 2014, 2016; Honing, Bouwer, & Háden, 2014).

Conclusion

To sum up, in the current study, we could not differentiate between beat-based and memory-based temporal expectations in terms of their effects on the detection of, and auditory responses to targets with predictable timing, nor in terms of how they were affected by task relevance. Also, beat-based and memory-based expectations to some extent interfered with each other. Similar effects on ERPs and behavior and the presence of interference may point at a shared underlying mechanism, which may be surprising given the evidence from clinical studies (Breska & Ivry, 2018) and research in nonhuman animals (Honing et al., 2018) that suggests that beat-based expectations are distinct from other types of temporal expectations. Indeed, we also found evidence for distinct processing of beat-based and memory-based expectations: When the timing of events was fully predictable based on memory, the presence of beat-based expectations still deteriorated target detection and enhanced sensory responses for events off the beat (at unexpected moments), and when events were fully predictable based on the beat, memory-based expectations still improved target detection.

To reconcile these findings, we propose that beat-based and memory-based expectations overlap in their effect on auditory responses and behavior, in line with both types of expectations serving the same function, but nonetheless, have partly separate underlying mechanisms to form expectations. Future research will have to focus on distinguishing between possible separate underlying mechanisms by examining the neural dynamics and neural networks involved in different types of temporal expectations. This may also elucidate at which point in the processing stream different types of temporal expectations interact, be it at early stages, during the formation of expectations, or at later stages, where expectations exert their effect on perception and behavior. The special status of beat-based expectations, which has been linked with evolutionary advantages of music (Honing et al., 2015), thus remains an open question. We contribute to this question by being the first to look at the orthogonal effects of beat-based and memory-based expectations on responses to auditory rhythm, and by showing how focusing on the effects of expectations on processing at unexpected, rather than expected time points, may provide a fruitful way to differentiate beat-based from other types of expectations in future research.

References

Abecasis, D., Brochard, R., Granot, R., & Drake, C. (2005). Differential brain response to metrical accents in isochronous auditory sequences. *Music Perception*, *22*(3), 549–562. https://doi.org/10.1525/mp.2005.22.3.549

Alilović, J., Timmermans, B., Reteig, L. C., van Gaal, S., & Slagter, H. A. (2019). No evidence that predictions and attention modulate the first feedforward sweep of cortical information processing. *Cerebral Cortex*, *29*(5), 2261–2278. https://doi.org/10.1093/cercor/bhz038

Arnal, L. H., Doelling, K. B., & Poeppel, D. (2014). Delta-beta coupled oscillations underlie temporal prediction accuracy. *Cerebral Cortex*, *25*(9), 3077–3085. https://doi.org/10.1093/cercor/bhu103

Auksztulewicz, R., & Friston, K. J. (2015). Attentional enhancement of auditory mismatch responses: a DCM/MEG study. *Cerebral Cortex*, *25*(11), 4273–4283. https://doi.org/10.1093/cercor/bhu323

Auksztulewicz, R., Schwiedrzik, C. M., Thesen, T., Doyle, W., Devinsky, O., Nobre, A. C., … Melloni, L. (2018). Not all predictions are equal: "What" and "when" predictions modulate activity in auditory cortex through different mechanisms. *The Journal of Neuroscience*, *38*(40), 8680–8693. https://doi.org/10.1523/JNEUROSCI.0369-18.2018

Bell, S. L., Smith, D. C., Allen, R., & Lutman, M. E. (2004). Recording the middle latency response of the auditory evoked potential as a measure of depth of anaesthesia. A technical note. *British Journal of Anaesthesia*, *92*(3), 442–445. https://doi.org/10.1093/bja/aeh074

Bouwer, F. L., Burgoyne, J. A., Odijk, D., Honing, H., & Grahn, J. A. (2018). What makes a rhythm complex? The influence of musical training and accent type on beat perception. *PLOS ONE*, *13*(1), e0190322. https://doi.org/10.1371/journal.pone.0190322

Bouwer, F. L., & Honing, H. (2015). Temporal attending and prediction influence the

perception of metrical rhythm: Evidence from reaction times and ERPs. *Frontiers in

Psychology*, *6*(July), 1094. https://doi.org/10.3389/fpsyg.2015.01094

Bouwer, F. L., Van Zuijen, T. L., & Honing, H. (2014). Beat processing is pre-attentive for

metrically simple rhythms with clear accents: An ERP study. *PLoS ONE*, *9*(5), e97467.

https://doi.org/10.1371/journal.pone.0097467

Bouwer, F. L., Werner, C. M., Knetemann, M., & Honing, H. (2016). Disentangling beat

perception from sequential learning and examining the influence of attention and musical

abilities on ERP responses to rhythm. *Neuropsychologia*, *85*, 80–90.

https://doi.org/10.1016/j.neuropsychologia.2016.02.018

Breska, A., & Deouell, L. Y. (2014). Automatic bias of temporal expectations following

temporally regular input independently of high-level temporal expectation. *Journal of

Cognitive Neuroscience*, *26*(7), 1555–1571. https://doi.org/10.1162/jocn_a_00564

Breska, A., & Deouell, L. Y. (2016). When synchronizing to rhythms is not a good thing:

Modulations of preparatory and post-target neural activity when shifting attention away

from on-beat times of a distracting rhythm. *The Journal of Neuroscience*, *36*(27), 7154–

7166. https://doi.org/10.1523/JNEUROSCI.4619-15.2016

Breska, A., & Deouell, L. Y. (2017a). Dance to the rhythm, cautiously: Isolating unique

indicators of oscillatory entrainment. *PLOS Biology*, *15*(9), e2003534.

https://doi.org/10.1371/journal.pbio.2003534

Breska, A., & Deouell, L. Y. (2017b). Neural mechanisms of rhythm-based temporal

prediction: Delta phase-locking reflects temporal predictability but not rhythmic

entrainment. *PLOS Biology*, *15*(2), e2001665.

https://doi.org/10.1371/journal.pbio.2001665

Breska, A., & Ivry, R. B. (2018). Double dissociation of single-interval and rhythmic

temporal prediction in cerebellar degeneration and Parkinson's disease. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(48), 12283–12288. https://doi.org/10.1073/pnas.1810596115

Brochard, R., Abecasis, D., Potter, D. D., Ragot, R., & Drake, C. (2003). The "ticktock" of our internal clock: direct brain evidence of subjective accents in isochronous sequences. *Psychological Science*, *14*(4), 362–366. https://doi.org/10.1111/1467-9280.24441

Chang, W.-P., Gavin, W. J., & Davies, P. L. (2012). Bandpass filter settings differentially affect measurement of P50 sensory gating in children and adults. *Clinical Neurophysiology*, *123*(11), 2264–2272. https://doi.org/10.1016/J.CLINPH.2012.03.019

Clark, A. (2013). Whatever next? Predictive brains, situated agents, and the future of cognitive science. *Behavioral and Brain Sciences*, *36*(3), 181–204. https://doi.org/10.1017/S0140525X12000477

Cope, T. E., Grube, M., Singh, B., Burn, D. J., & Griffiths, T. D. (2014). The basal ganglia in perceptual timing: Timing performance in Multiple System Atrophy and Huntington's disease. *Neuropsychologia*, *52*(1), 73–81. https://doi.org/10.1016/j.neuropsychologia.2013.09.039

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*, 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Escoffier, N., Herrmann, C. S., Schirmer, A., Escof, N., Herrmann, C. S., & Schirmer, A. (2015). Auditory rhythms entrain visual processes in the human brain: Evidence from evoked oscillations and event-related potentials. *NeuroImage*, *111*, 267–276. https://doi.org/10.1016/j.neuroimage.2015.02.024

Fardo, F., Auksztulewicz, R., Allen, M., Dietz, M. J., Roepstorff, A., & Friston, K. J. (2017). Expectation violation and attention to pain jointly modulate neural gain in

somatosensory cortex. *NeuroImage, 153*, 109–121.

https://doi.org/10.1016/J.NEUROIMAGE.2017.03.041

Feldman, H., & Friston, K. J. (2010). Attention, uncertainty, and free-energy. *Frontiers in Human Neuroscience, 4*(December), 215. https://doi.org/10.3389/fnhum.2010.00215

Fitzroy, A. B., & Sanders, L. D. (2015). Musical meter modulates the allocation of attention across time. *Journal of Cognitive Neuroscience, 27*(12), 2339–2351. https://doi.org/10.1162/jocn

Forth, J., Agres, K., Purver, M., & Wiggins, G. A. (2016). Entraining IDyOT: Timing in the Information Dynamics of Thinking. *Frontiers in Psychology, 7*, 1575. https://doi.org/10.3389/fpsyg.2016.01575

Friston, K. J. (2005). A theory of cortical responses. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences, 360*(1456), 815–836. https://doi.org/10.1098/rstb.2005.1622

Grahn, J. A. (2012). See what I hear? Beat perception in auditory and visual rhythms. *Experimental Brain Research, 220*(1), 51–61. https://doi.org/10.1007/s00221-012-3114-8

Grahn, J. A., & Brett, M. (2007). Rhythm and beat perception in motor areas of the brain. *Journal of Cognitive Neuroscience, 19*(5), 893–906. https://doi.org/10.1162/jocn.2007.19.5.893

Grahn, J. A., Henry, M. J., & McAuley, J. D. (2011). FMRI investigation of cross-modal interactions in beat perception: audition primes vision, but not vice versa. *NeuroImage, 54*(2), 1231–1243. https://doi.org/10.1016/j.neuroimage.2010.09.033

Grahn, J. A., & Schuit, D. (2012). Individual differences in rhythmic ability: Behavioral and neuroimaging investigations. *Psychomusicology: Music, Mind, and Brain, 22*(2), 105–121. https://doi.org/10.1037/a0031188

Haegens, S., & Zion Golumbic, E. (2018). Rhythmic facilitation of sensory processing: A

   critical review. *Neuroscience and Biobehavioral Reviews*, *86*(March), 150–165.

   https://doi.org/10.1016/j.neubiorev.2017.12.002

Henry, M. J., & Herrmann, B. (2014). Low-frequency neural oscillations support dynamic

   attending in temporal context. *Timing & Time Perception*, *2*(1), 62–86.

   https://doi.org/10.1163/22134468-00002011

Henry, M. J., & Obleser, J. (2012). Frequency modulation entrains slow neural oscillations

   and optimizes human listening behavior. *Proceedings of the National Academy of

   Sciences of the United States of America*, *109*(49), 2009–2100.

   https://doi.org/10.1073/pnas.1213390109

Hoch, L., Tyler, M. D., & Tillmann, B. (2013). Regularity of unit length boosts statistical

   learning in verbal and nonverbal artificial languages. *Psychonomic Bulletin & Review*,

   20, 142–147. https://doi.org/10.3758/s13423-012-0309-8

Honing, H., & Bouwer, F. L. (2019). Rhythm. In P. J. Rentfrow & D. Levitin (Eds.),

   Foundations in Music Psychology: Theory and Research. Cambridge, MA: MIT Press.

Honing, H., Bouwer, F. L., & Háden, G. P. (2014). Perceiving temporal regularity in music:

   The role of auditory event-related potentials (ERPs) in probing beat perception. In H.

   Merchant & V. de Lafuente (Eds.), *Neurobiology of Interval Timing* (Vol. 829, pp. 305–

   323). New York: Springer. https://doi.org/10.1007/978-1-4939-1782-2_16

Honing, H., Bouwer, F. L., Prado, L., & Merchant, H. (2018). Rhesus monkeys (Macaca

   mulatta) sense isochrony in rhythm, but not the beat: Additional support for the gradual

   audiomotor evolution hypothesis. *Frontiers in Neuroscience*, *12*, 475.

   https://doi.org/10.3389/fnins.2018.00475

Honing, H., Cate, C., Peretz, I., Trehub, S. E., ten Cate, C., Peretz, I., & Trehub, S. E. (2015).

   Without it no music: Cognition, biology and evolution of musicality. *Philosophical

Transactions of the Royal Society of London. Series B, Biological Sciences, *370*, 20140088. https://doi.org/10.1098/rstb.2014.0088

Honing, H., & Merchant, H. (2014). Differences in auditory timing between human and non-human primates. *Behavioral and Brain Sciences*, *27*(6), 557–558.

Hsu, Y.-F., Bars, S. Le, & Ha, J. A. (2015). Distinctive representation of mispredicted and unpredicted prediction errors in human electroencephalography. *The Journal of Neuroscience, 35*(43), 14653–14660. https://doi.org/10.1523/JNEUROSCI.2204-15.2015

Hsu, Y.-F., Hämäläinen, J. A., & Waszak, F. (2013). Temporal expectation and spectral expectation operate in distinct fashion on neuronal populations. *Neuropsychologia*, *51*(13), 2548–2555. https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2013.09.018

Hsu, Y.-F., Hämäläinen, J. A., & Waszak, F. (2018). The processing of mispredicted and unpredicted sensory inputs interact differently with attention. *Neuropsychologia*, *111*, 85–91. https://doi.org/10.1016/j.neuropsychologia.2018.01.034

Iversen, J. R., Patel, A. D., Nicodemus, B., & Emmorey, K. (2015). Synchronization to auditory and visual rhythms in hearing and deaf individuals. *Cognition*, *134*, 232–244. https://doi.org/10.1016/j.cognition.2014.10.018

Iversen, J. R., Repp, B. H., & Patel, A. D. (2009). Top-down control of rhythm perception modulates early auditory responses. *Annals of the New York Academy of Sciences*, *1169*, 58–73. https://doi.org/10.1111/j.1749-6632.2009.04579.x

JASP Team. (2019). JASP. Retrieved from https://jasp-stats.org

Jeffreys, H. (1961). *Theory of probability* (3rd ed.). Oxford, UK: Oxford University Press.

Jones, M. R., & Boltz, M. G. (1989). Dynamic attending and responses to time. *Psychological Review, 96*(3), 459–491. https://doi.org/10.1037/0033-295X.96.3.459

Koelsch, S., Vuust, P., & Friston, K. J. (2019). Predictive processes and the peculiar case of

music. *Trends in Cognitive Sciences*, *23*(1), 63–77.

https://doi.org/10.1016/J.TICS.2018.10.006

Kok, P., Rahnev, D. A., Jehee, J. F. M., Lau, H. C., & de Lange, F. P. (2012). Attention

reverses the effect of prediction in silencing sensory signals. *Cerebral Cortex*, *22*(9),

2197–2206. https://doi.org/10.1093/cercor/bhr310

Kunert, R., & Jongman, S. R. (2017). Entrainment to an auditory signal: Is attention

involved? *Journal of Experimental Psychology: General*, *146*(1), 77–88.

https://doi.org/10.1037/xge0000246

Lakatos, P., Karmos, G., Mehta, A. D., Ulbert, I., & Schroeder, C. E. (2008). Entrainment of

neuronal oscillations as a mechanism of attentional selection. *Science*, *320*(5872), 110–

113. https://doi.org/10.1126/science.1154735

Lakatos, P., Musacchia, G., O'Connel, M. N., Falchier, A. Y. Y., Javitt, D. C. C., Schroeder,

C. E., & O'Connel, M. N. (2013). The spectrotemporal filter mechanism of auditory

selective attention. *Neuron*, *77*(4), 750–761.

https://doi.org/10.1016/j.neuron.2012.11.034

Lange, K. (2009). Brain correlates of early auditory processing are attenuated by expectations

for time and pitch. *Brain and Cognition*, *69*(1), 127–137.

https://doi.org/10.1016/j.bandc.2008.06.004

Lange, K. (2013). The ups and downs of temporal orienting: A review of auditory temporal

orienting studies and a model associating the heterogeneous findings on the auditory N1

with opposite effects of attention and prediction. *Frontiers in Human Neuroscience*,

7(June), 263. https://doi.org/10.3389/fnhum.2013.00263

Large, E. W., Herrera, J. A., & Velasco, M. J. (2015). Neural networks for beat perception in

musical rhythm. *Frontiers in Systems Neuroscience*, *9*(November), 159.

https://doi.org/10.3389/fnsys.2015.00159

Large, E. W., & Jones, M. R. (1999). The dynamics of attending: how people track time-varying events. *Psychological Review*, *106*(1), 119–159. https://doi.org/10.1037/0033-295X.106.1.119

Lawrance, E. L. A., Harper, N. S., Cooke, J. E., & Schnupp, J. W. H. (2014). Temporal predictability enhances auditory detection. *The Journal of the Acoustical Society of America*, *135*(6), EL357–EL363. https://doi.org/10.1121/1.4879667

Liljander, S., Holm, A., Keski-Säntti, P., & Partanen, J. V. (2016). Optimal digital filters for analyzing the mid-latency auditory P50 event-related potential in patients with Alzheimer's disease. *Journal of Neuroscience Methods*, *266*, 50–67. https://doi.org/10.1016/J.JNEUMETH.2016.03.013

London, J. (2012). *Hearing in time: Psychological aspects of musical meter.* (2nd ed.). Oxford: Oxford University Press.

Luck, S. (2005). *An introduction to the event-related potential technique*. Cambridge, MA: MIT Press. Retrieved from http://mitpress.mit.edu/catalog/item/default.asp?tid=10677&ttype=2

Marzecová, A., Widmann, A., Sanmiguel, I., Kotz, S. A., & Schröger, E. (2017). Interrelation of attention and prediction in visual processing: Effects of task-relevance and stimulus probability. *Biological Psychology*, *125*, 76–90. https://doi.org/10.1016/j.biopsycho.2017.02.009

McGarry, L. M., Sternin, A., & Grahn, J. A. (2019). Music and Movement. In P. J. Rentfrow & D. J. Levitin (Eds.), *Foundations in Music Psychology: Theory and Research*. Cambridge, MA: MIT Press.

Merchant, H., Grahn, J. A., Trainor, L. J., Rohrmeier, M. A., & Fitch, W. T. (2015). Finding the beat: a neural perspective across humans and non-human primates. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *370*,

20140093. https://doi.org/10.1098/rstb.2014.0093

Merchant, H., & Honing, H. (2014). Are non-human primates capable of rhythmic

entrainment? Evidence for the gradual audiomotor evolution hypothesis. *Frontiers in*

Neuroscience, *7*(January), 274. https://doi.org/10.3389/fnins.2013.00274

Morillon, B., Schroeder, C. E., Wyart, V., & Arnal, L. H. (2016). Temporal prediction in lieu

of periodic stimulation. *The Journal of Neuroscience*, *36*(8), 2342–2347.

https://doi.org/10.1523/JNEUROSCI.0836-15.2016

Näätänen, R. N., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity

(MMN) in basic research of central auditory processing: a review. *Clinical*

Neurophysiology, *118*(12), 2544–2590. https://doi.org/10.1016/j.clinph.2007.04.026

Nobre, A. C., & Rohenkohl, G. (2014). Time for the Fourth Dimension in Attention. In A. C.

Nobre & S. Kastner (Eds.), *The Oxford Handbook of Attention* (pp. 676–721). New

York, NY: Oxford University Press.

https://doi.org/10.1093/oxfordhb/9780199675111.013.036

Nobre, A. C., & van Ede, F. (2018). Anticipated moments: temporal structure in attention.

Nature Publishing Group, *19*, 34–48. https://doi.org/10.1038/nrn.2017.141

O'Connel, M. N., Barczak, A., Ross, D., McGinnis, T., Schroeder, C. E., & Lakatos, P.

(2015). Multi-scale entrainment of coupled neuronal oscillations in primary auditory

cortex. *Frontiers in Human Neuroscience*, *9*(December), 655.

https://doi.org/10.3389/fnhum.2015.00655

Obleser, J., Henry, M. J., & Lakatos, P. (2017). What do we talk about when we talk about

rhythm? *PLOS Biology*, *15*(9), e2002794. https://doi.org/10.1371/journal.pbio.2002794

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source

software for advanced analysis of MEG, EEG, and invasive electrophysiological data.

Computational Intelligence and Neuroscience, *2011*, 156869.

https://doi.org/10.1155/2011/156869

Paris, T., Kim, J., & David, C. (2016). The processing of attended and predicted sounds in time. *Journal of Cognitive Neuroscience, 28*(1), 158–165. https://doi.org/10.1162/jocn_a_00885

Picton, T. W., Hillyard, S. A., Krausz, H. I., & Galambos, R. (1974). Human auditory evoked potentials. I: Evaluation of components. *Electroencephalography and Clinical Neurophysiology, 36*, 179–190. https://doi.org/10.1016/0013-4694(74)90155-2

Potter, D. D., Fenwick, M., Abecasis, D., & Brochard, R. (2009). Perceiving rhythm where none exists: event-related potential (ERP) correlates of subjective accenting. *Cortex, 45*(1), 103–109. https://doi.org/10.1016/j.cortex.2008.01.004

Povel, D.-J., & Essens, P. (1985). Perception of Temporal Patterns. *Music Perception, 2*(4), 411–440. https://doi.org/10.2307/40285311

Repp, B. H. (2010). Do metrical accents create illusory phenomenal accents? *Attention, Perception & Psychophysics, 72*(5), 1390–1403. https://doi.org/10.3758/APP

Rimmele, J. M., Jolsvai, H., & Sussman, E. S. (2011). Auditory target detection is affected by implicit temporal and spatial expectations. *Journal of Cognitive Neuroscience, 23*(5), 1136–1147. https://doi.org/10.1162/jocn.2010.21437

Rimmele, J. M., Morillon, B., Poeppel, D., & Arnal, L. H. (2018). Proactive sensing of periodic and aperiodic auditory patterns. *Trends in Cognitive Sciences, 22*(10), 870–882. https://doi.org/10.1016/J.TICS.2018.08.003

Rohenkohl, G., Coull, J. T., & Nobre, A. C. (2011). Behavioural dissociation between exogenous and endogenous temporal orienting of attention. *PloS One, 6*(1), e14620. https://doi.org/10.1371/journal.pone.0014620

Rohenkohl, G., Cravo, A. M., Wyart, V., & Nobre, A. C. (2012). Temporal expectation improves the quality of sensory information. *The Journal of Neuroscience, 32*(24),

8424–8428. https://doi.org/10.1523/JNEUROSCI.0804-12.2012

Ruhnau, P., Herrmann, B., Maess, B., & Schröger, E. (2011). Maturation of obligatory auditory responses and their neural sources: evidence from EEG and MEG. *NeuroImage*, *58*(2), 630–639. https://doi.org/10.1016/j.neuroimage.2011.06.050

Sanabria, D., & Correa, Á. (2013). Electrophysiological evidence of temporal preparation driven by rhythms in audition. *Biological Psychology*, *92*(2), 98–105. https://doi.org/10.1016/j.biopsycho.2012.11.012

Schaefer, R. S., Vlek, R. J., & Desain, P. (2010). Decomposing rhythm processing: Electroencephalography of perceived and self-imposed rhythmic patterns. *Psychological Research*, 95–106. https://doi.org/10.1007/s00426-010-0293-4

Schnupp, J. W. H., Rajendran, V. G., Harper, N. S., Garcia-Lazaro, J. A., & Lesica, N. A. (2017). Midbrain adaptation may set the stage for the perception of musical beat. *Proceedings of the Royal Society B*, *284*, 20171455. https://doi.org/10.1098/rspb.2017.1455

Schroeder, C. E., & Lakatos, P. (2009). Low-frequency neuronal oscillations as instruments of sensory selection. *Trends in Neurosciences*, *32*(1), 9–18. https://doi.org/10.1016/j.tins.2008.09.012

Schröger, E., Kotz, S. A., & SanMiguel, I. (2015). Bridging prediction and attention in current research on perception and action. *Brain Research*, *1626*, 1–13. https://doi.org/10.1016/j.brainres.2015.08.037

Schröger, E., Marzecová, A., & Sanmiguel, I. (2015). Attention and prediction in human audition: A lesson from cognitive psychophysiology. *The European Journal of Neuroscience*, *41*(5), 641–664. https://doi.org/10.1111/ejn.12816

Schultz, B. G., Stevens, C. J., Keller, P. E., & Tillmann, B. (2013). The implicit learning of metrical and nonmetrical temporal patterns. *Quarterly Journal of Experimental*

Psychology (2006), *66*(2), 360–380. https://doi.org/10.1080/17470218.2012.712146

Schwartze, M., Farrugia, N., & Kotz, S. A. (2013). Dissociation of formal and temporal

predictability in early auditory evoked potentials. *Neuropsychologia*, *51*(2), 320–325.

https://doi.org/10.1016/j.neuropsychologia.2012.09.037

Schwartze, M., & Kotz, S. A. (2013). A dual-pathway neural architecture for specific

temporal prediction. *Neuroscience and Biobehavioral Reviews*, *37*(10), 2587–2596.

https://doi.org/10.1016/j.neubiorev.2013.08.005

Schwartze, M., Rothermich, K., Schmidt-Kassow, M., & Kotz, S. A. (2011). Temporal

regularity effects on pre-attentive and attentive processing of deviance. *Biological*

Psychology, *87*(1), 146–151. https://doi.org/10.1016/j.biopsycho.2011.02.021

Selchenkova, T., Jones, M. R., & Tillmann, B. (2014). The influence of temporal regularities

on the implicit learning of pitch structures. *The Quarterly Journal of Experimental*

Psychology, *67*(12), 2360–2380. https://doi.org/10.1080/17470218.2014.929155

Sherwell, C., Garrido, M. I., & Cunnington, R. (2017). Timing in predictive coding: The roles

of task relevance and global probability. *Journal of Cognitive Neuroscience*, *29*(5), 780–

792. https://doi.org/10.1162/jocn_a_01085

Smout, C. A., Tang, M. F., Garrido, M. I., & Mattingley, J. B. (2019). Attention promotes the

neural encoding of prediction errors. *PLOS Biology*, *17*(2), e2006812.

https://doi.org/10.1371/journal.pbio.2006812

Tanner, D., Morgan-Short, K., & Luck, S. J. (2015). How inappropriate high-pass filters can

produce artifactual effects and incorrect conclusions in ERP studies of language and

cognition. *Psychophysiology*, *52*(8), 997–1009. https://doi.org/10.1111/psyp.12437

Teki, S., Grube, M., & Griffiths, T. D. (2012). A unified model of time perception accounts

for duration-based and beat-based timing mechanisms. *Frontiers in Integrative*

Neuroscience, *5*(January), 1–7. https://doi.org/10.3389/fnint.2011.00090

Teki, S., Grube, M., Kumar, S., & Griffiths, T. D. (2011). Distinct neural substrates of

    duration-based and beat-based auditory timing. *The Journal of Neuroscience*, *31*(10),

    3805–3812. https://doi.org/10.1523/JNEUROSCI.5561-10.2011

ten Oever, S., Schroeder, C. E., Poeppel, D., van Atteveldt, N., & Zion-Golumbic, E. (2014).

    Rhythmicity and cross-modal temporal cues facilitate detection. *Neuropsychologia*, *63*,

    43–50. https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2014.08.008

Tierney, A. T., & Kraus, N. (2013). Neural responses to sounds presented on and off the beat

    of ecologically valid music. *Frontiers in Systems Neuroscience*, *7*(May), 14.

    https://doi.org/10.3389/fnsys.2013.00014

Todorovic, A., Schoffelen, J., Ede, F. Van, Maris, E., & de Lange, F. P. (2015). Temporal

    expectation and attention jointly modulate auditory oscillatory activity in the beta band.

    PloS One, *10*(3), e0120288. https://doi.org/10.1371/journal.pone.0120288

van Atteveldt, N., Musacchia, G., Zion-Golumbic, E., Sehatpour, P., Javitt, D. C., &

    Schroeder, C. E. (2015). Complementary fMRI and EEG evidence for more efficient

    neural processing of rhythmic vs. unpredictably timed sounds. *Frontiers in Psychology*,

    6(OCT), 1–11. https://doi.org/10.3389/fpsyg.2015.01663

van der Weij, B., Pearce, M. T., & Honing, H. (2017). A probabilistic model of meter

    perception: simulating enculturation. *Frontiers in Psychology*, *8*(May), 824.

    https://doi.org/10.3389/fpsyg.2017.00824

Wagenmakers, E.-J., Love, J., Marsman, M., Jamil, T., Ly, A., Verhagen, J., … Morey, R. D.

    (2018). Bayesian inference for psychology. Part II: Example applications with JASP.

    Psychonomic Bulletin & Review, *25*(1), 58–76. https://doi.org/10.3758/s13423-017-

    1323-7

Zarco, W., Merchant, H., Prado, L., & Mendez, J. C. (2009). Subsecond timing in primates:

    comparison of interval production between human subjects and rhesus monkeys. *Journal*

of Neurophysiology, *102*(6), 3191–3202. https://doi.org/10.1152/jn.00066.2009

Zoefel, B., & Vanrullen, R. (2017). Oscillatory mechanisms of stimulus processing and

selection in the visual and auditory systems: State-of-the-art, speculations and

suggestions. *Frontiers in Neuroscience*, *11*(May), 296.

https://doi.org/10.3389/fnins.2017.00296