1

Butterfly abundance declines over 20 years of systematic monitoring in Ohio, USA

Tyson Wepprich[1]*, Jeffrey R. Adrion[2], Leslie Ries[3], Jerome Wiedmann[4], Nick M. Haddad[5]

1. Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon,

United States of America

2. Institute of Ecology and Evolution, University of Oregon, Eugene, Oregon, United States

of America

3. Department of Biology, Georgetown University, Washington, D.C., United States of

America

4. The Ohio Lepidopterists, Columbus, Ohio, United States of America

5. Department of Integrative Biology and W.K. Kellogg Biological Station, Michigan State

University, Hickory Corners, Michigan, United States of America

* Corresponding author

Email: tyson.wepprich@oregonstate.edu (TW)

Abstract

Severe insect declines make headlines, but are rarely based on systematic monitoring outside of Europe. We estimate the rate of change in total butterfly abundance and the population trends for 81 species using 21 years of systematic monitoring in Ohio, USA. Total abundance is declining at 2% per year, resulting in a cumulative 33% reduction in butterfly abundance. Three times as many species have negative population trends compared to positive trends. The rate of total decline and the proportion of species in decline mirror those documented in long-term European monitoring. Multiple environmental changes such as climate change, habitat degradation, and agricultural practices may contribute to these declines in Ohio and shift the makeup of the butterfly community by benefiting some species over others. Our analysis of life-history traits associated with population trends shows an impact of climate change, as species with northern distributions and fewer annual generations declined more rapidly. However, even common and invasive species associated with human-dominated landscapes are declining, suggesting widespread environmental causes for these trends. Declines in common species, although they may not be close to extinction, will have an outsized impact on the ecosystem services provided by insects. These results from the most extensive, systematic insect monitoring programs in North America demonstrate an ongoing defaunation in butterflies that on an annual scale might be imperceptible, but cumulatively has reduced butterfly numbers by a third over 20 years.

Introduction

Defaunation, or the drastic loss of animal species and declines in abundance, threatens to destabilize ecosystem functioning globally (1). In comparison to studies of vertebrate

populations, monitoring of changes in insect diversity is more difficult and far less prevalent

(2,3). Despite this, a global analysis of long-term population trends across 452 species estimated

that insect abundance had declined 45% over 40 years (1). Recently, more extreme declines in

insect biomass have been observed upon resampling after 2-4 decades (4,5). Losses of total

biomass or total abundance across all species may be more consequential than local declines in

species diversity, as common insect species contribute the most to ecosystem functions, such as

pollination (6). However, our knowledge of insect declines is skewed towards European

monitoring programs, including in global analyses (1). In this study, we analyze long-term,

region-wide trends in abundance across a diversity of species for an entire insect group in North

America to examine the scope of insect defaunation.

The best source of data to assess insect defaunation comes from large-scale, systematic

monitoring programs of multiple species (3). Through these efforts, trained volunteers or citizen

scientists have contributed much of the evidence for biotic responses to anthropogenic climate

warming through changes in insect phenology and range distributions (7,8). Unlike citizen

science reporting of opportunistic observations or species checklists, many insect monitoring

programs use a systematic protocol developed specifically for volunteers to track butterfly

abundances through time, both within and between seasons, and over large spatial scales (9).

Pollard-based monitoring programs, modeled after the first nationwide Butterfly Monitoring

Scheme launched in the United Kingdom in 1977 (UKBMS), use weekly standardized counts on

fixed transects (10). Their widespread adoption enables regional comparisons of insect responses

to environmental change or defaunation (11,12). We compare our analysis with exemplary long-

term monitoring schemes from Europe to test if the rate of insect declines generalizes across

continents.

The best source of abundance data for assessment of chronic insect decline, and the most prominent source of data in (1), is within the butterflies. Due to the relative ease and popularity of monitoring butterflies, environmental assessments use them as an indicator taxa for the general trajectory of biodiversity, assuming that they experience comparable pressures from land-use change, climate change, and habitat degradation as other insect taxa (13–15). Intensive long-term monitoring of individual butterfly species has provided rigorous, quantitative estimates of declines. Most prominently, the Eastern North American Monarch has declined by over 85% (16) and the Western North American Monarch by over 95% (17) over the past two decades. Severe declines have also been observed in some of the rarest butterflies (18,19). These data from individual species of conservation concern may not represent a broader trend across butterflies, which is what we aim to document in this study.

Volunteers, organized and trained by The Ohio Lepidopterists, have assembled the most extensive dataset of systematic butterfly counts that stands alone in North America in terms of the spatial extent and sampling frequency of Pollard walks (9). Three other monitoring programs in the United States have documented long-term, multi-species population trends. In Massachusetts, based on species lists from field trips, climate-driven community shifts explain how the relative likelihood of species observations change over 18 years (20). Art Shapiro and colleagues have made biweekly presence/absence observations on 11 fixed transects along an elevational gradient in California over 40 years to document species richness changes in response to climate and land-use, increasing abundance at a high elevation site, and impacts of agricultural practices on abundance at low elevation sites (21,22). Several teams have monitored declines in specialist butterflies restricted to native prairie patches in the Midwestern states with transect or timed survey methods over 26 years (23,24). The number of Pollard-based monitoring

programs has increased sharply since 2010 in the United States (9), which could potentially track

to test how widespread and consistent butterfly trends are.

Here, we used 21 years of weekly butterfly surveys across 104 sites to assess abundance

trends for butterflies in Ohio. We estimate population trends for 81 species and test for their

association with life-history traits and phylogenetic relatedness. We review findings from

European butterfly monitoring schemes for quantitative comparison with the rate of abundance

changes in Ohio. This analysis provides evidence of widespread insect defaunation and species'

declines from the most extensive, systematic monitoring program in North America.

Materials and methods

Study sites

We studied butterfly population trends across the state of Ohio in the Midwestern USA.

Over its 116,100 km^2 land area, Ohio has a mosaic of habitat types due to its partially glaciated

history and its place at the confluence of Midwestern prairies, the Appalachian Mountains, and

the boreal forest (25). Only remnants of wetland and prairie habitat remain in the state due to

human modification of the landscape. Some rare butterflies have declined due to forest

succession following suppression of disturbances (26). Agriculture and pastures (50%), forest

(30%), and urban development (10%) are the predominant land-use/land cover classes (27).

Monitoring sites have a Northeast to Southwest gradient in their mean annual

temperatures (mean 18.8°C, range from 14.0°C to 23.6°C) from interpolated daily temperatures

from Daymet over 1996-2016 (Thornton et al. 1997). Mean annual temperatures at these sites

grew at a linear trend of 0.3°C per decade and growing season length has increased by 60 degree-

days (base 5°C) per decade from 1980-2016. Monitoring sites span the state but are concentrated

near cities (Fig 1). On average, within a radius of 2 kilometers, monitoring sites have 24%

cropland and pasture, 34% forest, and 30% urban land-use based on the National Land Cover

Dataset (29). Although not considered in this study, impervious surfaces from urban

development influence temperature-dependent butterfly phenology in Ohio through the urban

heat island effect, which may not be captured in these gridded temperature interpolations (30).

Fig 1: Transect locations monitored by volunteers with the Ohio Lepidopterists. Of the 147

sites, this analysis used the 104 sites monitored for three or more years.

Monitoring surveys

Trained volunteers contributed 24,405 butterfly surveys from 1996 to 2016 as part of the

Ohio Lepidopterists Long-term Monitoring of Butterflies program. Volunteers survey on fixed

paths at approximately weekly intervals during the entire growing season from April through

October (median 23 of 30 weeks surveyed per year per site) and count every species within an

approximate 5-meter buffer around the observer (10). Surveys are constrained to times of good

weather to increase the detectability of butterflies and last a mean 85 minutes in duration. The

annual number of monitored sites ranged from 13 in 1996 to a maximum of 80 in 2012. We

limited our analysis of abundance trends to the 104 sites with three or more years of monitoring

data and 10 or more surveys per year at each site (Fig 1). We included observations of all sites

with at least 5 surveys per year in phenology models that we used to interpolate missing counts

before estimating abundance (31).

All species (102) with population indices estimated by phenology models contributed to

the total abundance analysis. We limited species-specific analysis to 81 with sufficient

population indices for estimating trends (present at five or more sites and for 10 or more years).

Species naming conventions in the monitoring program follow those used in Opler and Krizek

(1984) and Iftner et al. (1992) except for combining all observations of *Celastrina ladon* (Spring

Azure) and *Celastrina neglecta* (Summer Azure) as an unresolved species complex.

Population indices

We estimated population indices for each site x year x species by adapting methods

established for the UKBMS that account for missing surveys and butterfly phenology over the

season (31,33). We used generalized additive models for each species to estimate variation in

counts in order to interpolate missing surveys with model predictions (31,34). To account for

seasonal, spatial, and interannual variation in species phenology, we extended the regional

generalized additive model approach (12, Supplement 1) by including spatially-explicit site

locations and converted calendar dates of observations to degree-days (35), which can improve

butterfly phenology predictions. We calculated the population index by integrating over the

weekly counts and missing survey interpolations using the trapezoid method (31).

Controlling for confounding factors

We accounted for differences in sampling across sites and years so that our modeled

trends would capture changes in abundance rather than changes in detection probability (36).

True abundance is confounded with detection probability when using counts from Pollard walks

(37). Butterfly monitoring protocols that account for detection probability like distance sampling

are commonly used for single-species studies (38), but untenable for scaling up to a statewide

program. Most analyses of Pollard walks assume no systematic change in detectability (but see

(39)) because counts correlate closely with true abundance estimates from distance sampling

(40,41). We used two covariates to account for variation in sampling and its influence on

population indices (20,36,42). We tracked the number of species reported in each survey, or list-

length, which is a synthetic measure of factors influencing detectability such as weather

conditions, site quality, and observer effort (20,43,44). We treated the total duration of surveys in

minutes as an offset in the models of population trends, which converts the population indices to

a rate of butterflies observed per minute. Because we interpolated missing surveys for the

population indices, we projected what the total duration would be if all 30 weeks had been

surveyed at the mean duration reported for that site x year.

Sampling across the state is nonrandom because participants choose transect locations, a

common practice in volunteer-based monitoring programs. Since sites generally cluster near

human population centers with a greater proportion of developed land-use and a lesser

proportion of agriculture, we assumed that population trends at the 104 sites across the state

sufficiently capture the broader statewide trends (36). Comparisons between the UKBMS

volunteer-placed transects and a broader survey with stratified, random sampling show

congruence between species trends estimated from each monitoring strategy (45).

Population Trends

We used generalized linear mixed models to estimate temporal trends in relative

abundance for 81 species from their population indices (46). We modeled population indices at

each site and year as an over-dispersed Poisson random variable with covariates on the log-link

scale.

$$\log(PopulationIndex) = \beta_0 + \beta_1 \times year + \beta_2 \times listlength + \log(duration) + siteID \\ + yearID + siteyearID$$

We included the year and mean list length for each population index as covariates, which

were centered to aid in model fitting and interpretation (47). We used the coefficient for year (β_2)

as the annual trend in population indices as our main result. We controlled for changes in

sampling by using the total duration of surveys as a model offset, converting the dependent

variable to a rate of butterflies counted per minute. Random effects of individual sites and years

account for spatial and temporal variation in population counts deviating from the statewide

trend. We accounted for over-dispersion in the Poisson-distributed counts with the random effect

siteyearID for each unique observation (48). We modeled trends in total abundance using the

same modeling approach, but summed across 102 species' population indices for each site x year

observation. We interpreted trends as an annual rate by taking the geometric mean rate of change

between the predicted abundance between two points in time after setting the list-length

covariate to its mean and excluding the random effects (46). For comparisons with other

monitoring programs, we used a *p*-value threshold of 0.05 to classify trends as positive, negative,

or stable.

Our approach is similar to that used by the UKBMS and other European monitoring

programs which use generalized linear models in TRIM software (49). One key difference is that

our site and annual fluctuations from the temporal trend were derived from random effects rather

than fixed effects, which reduces spurious detection of trends (42). Another key difference is that

TRIM does not allow for continuous covariates, which we used to account for sampling variation

instead of assuming no confounding pattern in sampling effort. To validate that our modeling

choices did not unreasonably influence the results, we used three alternative approaches: (1) a

Poisson-based generalized linear model (equation 1 without the random effect *siteyearID*); (2) a

nonlinear generalized additive mixed model with a smoothing spline replacing the linear

temporal trend (42); and (3) a TRIM model with over-dispersion and serial temporal correlation

but no sampling covariates or offsets (49). We compared similarity in the total abundance trends,

the correlation of species' trends between model alternatives, and the classification of species'

trends.

Comparison with other studies

We compare our findings to three European long-term, regional butterfly monitoring

programs with systematic Pollard walks that publish regular updates on total abundance and

species' trends (39,50,51). Although all programs analyzed counts with Poisson regression, we

had to standardize them differently depending on the data available and their modeling

approaches. The UKBMS reports total abundance indicators as the geometric mean of species

trends from two groups: specialist and countryside species (50). We used the reported smoothed

annual index values for these indicators because the first year of monitoring is an outlier that

exaggerates declines (UK Biodiversity Indicators 2018, http://jncc.defra.gov.uk/page-4236). We

used the Dutch Butterfly Monitoring Scheme's reported cumulative annual trend in total

butterflies counted across all transects after correction for missing surveys (51). For the Catalan

Butterfly Monitoring Scheme, we extracted annual population indices from the 2015-2016

annual report (52) with WebPlotDigitizer 4.1 (53) and performed a Poisson regression over time

with annual random effects to obtain a comparable abundance trend. We converted total

abundance trends into annual percent rates for comparison. We tallied the increases and

decreases in species' trends for each region reported by the monitoring program, without

accounting for differences in their statistical approaches.

Species' traits

To explore potential mechanisms that might explain species-level variation in abundance

trends, we modeled the estimates of species' temporal trends (β_1) as a response to life history

traits (20,30). Of the 81 species, 14 were classified as migratory species and 4 (*Colias*

eurytheme, *Lycaena phlaeus, Pieris rapae*, and *Thymelicus lineola*) were considered naturalized

species that were introduced to the state 90 or more years ago (25,32). We analyzed traits models

both across all species and after excluding migratory species, which would have population

trends driven by factors outside of Ohio. We collected traits that relate to insect responses to

climate change and habitat change, as these are two primary drivers of butterfly community

changes (7,20,21).

We tested if butterflies with traits making them more adaptive to a warming climate have

more positive population trends. Voltinism, or the number of generations per year, increases in

warmer years and warmer regions within many species in Ohio (54), compared with obligate

univoltine species that do not adjust their lifecycle based on changing growing season length. We

assigned voltinism observed in Ohio (1-4 generations per year) to an integer scale based on

visualization of phenology models and (25). We compared species with different range

distributions, assuming that species distributed in warmer, Southern regions would be more

3likely to increase in Ohio as the climate warms. We assigned species' ranges as Southern, core,

or Northern by range maps and county records (25,32). The life stage in which species

overwinter contributes to its ability to respond to warming with shifts in phenology (20,55).

We would expect more generalist species, in host plant requirements and habitat

preferences, to have more positive population trends in a landscape heavily modified by human

use (21,50). For host plant requirements, we gathered two traits from the literature that describe

host plant category (forb, graminoid, or woody) and whether the butterfly's host plant

requirements span multiple plant families or are limited to one plant family or genus (25). Mean

wing size from (32) was used as a surrogate of dispersal ability between habitats, which is

expected to increase ability to access resources in a fragmented landscape. Three of the authors

assigned species as wetland-dependent or human-disturbance tolerant species, which we

aggregated into two binary variables to test if specialist or generalist habitat preferences correlate

with abundance trends.

We used univariate linear models for each life history trait both for all 81 species and

with the 14 migratory species excluded. To account for the phylogenetic relatedness and the non-

independence across species, we compared phylogenetic generalized least squares models that

estimated branch length transformations with Pagel's lambda by maximum likelihood (56). The

phylogenetic models excluded three species without gene sequences available.

Phylogenetic tree

We obtained coding sequences for the most widely used DNA barcoding locus, the

mitochondrial cytochrome c oxidase subunit I gene COI-5P, from GenBank (57). For species not

found in GenBank, we obtained coding sequences from The Barcode of Life Data System (58).

When possible, we obtained sequences from multiple sampling locations in North America.

Owing to the relatively small size of our multiple-species alignment—i.e. a single

mtDNA locus, 651 base pairs in length—we decided to take both a constrained and

unconstrained maximum likelihood approach to estimate the genealogical relationships of our

samples. Some of the species from our analysis, though not all, were recently used in a more

comprehensive phylogenetic analysis of butterflies (59), thus prompting us to constrain the

phylogenetic backbone of our tree using family-level relationships. We report details of our

workflow in Supplement 1.

Statistical analysis

We used R 3.5.2 for analysis (60) and share the data and our code on Dryad. We fit

generalized additive models with the *mgcv* package (34), generalized linear mixed models with

the *lme4* package (Bates et al. 2015), and phylogenetic generalized least squares models with the

ape and *caper* packages (62,63). Confidence intervals for the temporal trends were estimated

with bootstrapped model fits with the *merTools* package (64). We estimated the goodness of fit

with R^2 developed for generalized linear mixed models that give marginal and conditional R^2

values for the fixed effects and the fixed + random effects, respectively (65,66).

Results

The statewide relative abundance across all species declined at an annual rate of 2.0% (β_1

= -0.020, std. err. 0.005, $p < 0.001$), accumulating a 33% decline over 1996-2016 (Table 1, Fig

2). Among population trends, more than three times as many species are declining than

increasing in abundance at our threshold of $p < 0.05$ (32 versus 9, respectively) (Table 2, Fig B-

D in S1 Appendix). Positive and negative species trends are distributed across the phylogenetic

tree (Fig A in S1 Appendix).

Table 1: Generalized linear mixed model of total abundance across all species. The natural

logarithm of the total survey duration across the monitoring season was an offset in the model.

The model's marginal R^2 was 0.20 for its fixed effects and its conditional R^2 was 0.61 when

including variation in sites, years, over-dispersion with random effects parameters.

Fixed effects	B	std.error	z statistic	p.value
Intercept	1.33	0.0506	26.4	<0.001
Year (numeric)	-0.0203	0.00496	-4.11	<0.001
List-length	0.104	0.00587	17.7	<0.001
Random effects	std. dev.	# groups		
Site x year	0.278	1005		
Site	0.417	104		
Year (factor)	0.121	21		

Table 2: Species' abundance trends over time. Trends are the coefficient of year in our regression models with the accompanying standard error and *p*-value for the coefficient (equation 1). Included are the number of population indices calculated for each species for use in abundance model (Site x year) and total number of butterflies recorded for all years. Bold font indicates trends classified as increasing or decreasing ($p < 0.05$).

Species		Sample size				GLMM temporal trend		
Common	Latin	Total # counted	Sites	Years	Site/ year	Trend coef.	Std. error	P
Aphrodite Fritillary	*Speyeria aphrodite*	477	9	16	131	*-0.233*	*0.060*	*<0.001*
Baltimore	*Euphydryas phaeton*	818	7	17	83	*-0.224*	*0.071*	*0.002*
American Copper	*Lycaena phlaeas*	10,255	31	21	359	*-0.193*	*0.024*	*<0.001*
Hoary Edge Skipper	*Achalarus lyciades*	291	7	19	88	*-0.178*	*0.061*	*0.003*
Milbert's Tortoise Shell	*Nymphalis milberti*	140	8	16	101	*-0.174*	*0.065*	*0.008*
European Skipper	*Thymelicus lineola*	46,549	57	21	609	*-0.173*	*0.021*	*<0.001*
Southern Cloudywing	*Thorybes bathyllus*	667	15	20	194	*-0.129*	*0.037*	*<0.001*
Falcate Orangetip	*Anthocharis midea*	756	8	18	103	*-0.123*	*0.040*	*0.002*
Dreamy Duskywing	*Erynnis icelus*	879	18	21	260	*-0.120*	*0.024*	*<0.001*
Swarthy Skipper	*Nastra lherminier*	448	7	17	78	*-0.114*	*0.041*	*0.006*
Tawny Emperor	*Asterocampa clyton*	937	27	19	308	*-0.114*	*0.036*	*0.002*
Leonard's Skipper	*Hesperia leonardus*	1,348	11	20	144	*-0.110*	*0.025*	*<0.001*
White M Hairstreak	*Parrhasius m-album*	95	7	15	110	-0.105	0.081	0.195
Northern Cloudywing	*Thorybes pylades*	547	16	20	210	*-0.095*	*0.033*	*0.004*
Coral Hairstreak	*Satyrium titus*	607	15	21	217	*-0.094*	*0.025*	*<0.001*
Juvenal's Duskywing	*Erynnis juvenalis*	3,838	38	21	487	*-0.083*	*0.020*	*<0.001*
Common Wood Nymph	*Cercyonis pegala*	21,603	77	21	788	*-0.073*	*0.013*	*<0.001*
Common Sooty Wing	*Pholisora catullus*	1,142	34	20	398	*-0.072*	*0.015*	*<0.001*
Sleepy Duskywing	*Erynnis brizo*	811	13	18	156	*-0.071*	*0.032*	*0.027*
Monarch	*Danaus plexippus*	46,070	104	21	1,005	*-0.070*	*0.023*	*0.002*
Red-spotted Purple	*Limenitis arthemis*	6,226	87	21	913	*-0.064*	*0.019*	*<0.001*
Bronze Copper	*Lycaena hyllus*	656	23	21	254	-0.063	0.039	0.103
Northern Broken-Dash	*Wallengrenia egeremet*	5,959	49	21	528	*-0.062*	*0.018*	*<0.001*
Tawny-edged Skipper	*Polites themistocles*	2,322	48	21	541	*-0.058*	*0.016*	*<0.001*
West Virginia White	*Pieris virginiensis*	214	5	16	63	-0.058	0.059	0.329
Fiery Skipper	*Hylephila phyleus*	3,917	57	19	646	-0.057	0.061	0.351
Meadow Fritillary	*Boloria bellona*	5,447	55	21	598	*-0.056*	*0.027*	*0.040*

Common Name	Scientific Name							
Orange Sulphur	*Colias eurytheme*	62,160	101	21	996	*-0.055*	*0.021*	*0.008*
Long Dash	*Polites mystic*	1,317	21	21	219	*-0.047*	*0.020*	*0.022*
American Lady	*Vanessa virginiensis*	2,029	54	21	637	-0.045	0.033	0.179
Black Swallowtail	*Papilio polyxenes*	12,410	92	21	941	*-0.044*	*0.015*	*0.004*
Gray Hairstreak	*Strymon melinus*	2,418	49	19	587	-0.044	0.026	0.089
Painted Lady	*Vanessa cardui*	5,564	80	21	873	-0.042	0.054	0.440
Great Spangled Fritillary	*Speyeria cybele*	33,573	90	21	904	*-0.041*	*0.020*	*0.047*
Hobomok Skipper	*Poanes hobomok*	6,863	51	21	576	*-0.040*	*0.014*	*0.005*
Viceroy	*Limenitis archippus*	16,079	85	21	896	*-0.039*	*0.016*	*0.014*
Cabbage White	*Pieris rapae*	304,105	104	21	1,005	*-0.038*	*0.010*	*<0.001*
Hackberry Emperor	*Asterocampa celtis*	9,992	42	20	467	*-0.037*	*0.017*	*0.033*
Striped Hairstreak	*Satyrium liparops*	155	14	18	211	-0.028	0.067	0.682
Variegated Fritillary	*Euptoieta claudia*	956	17	19	204	-0.027	0.052	0.603
Little Wood Satyr	*Megisto cymela*	76,612	87	21	878	*-0.026*	*0.009*	*0.005*
American Snout Butterfly	*Libytheana carinenta*	1,007	36	18	418	-0.025	0.050	0.612
Hickory Hairstreak	*Satyrium caryaevorum*	196	12	20	170	-0.023	0.053	0.656
Mourning Cloak	*Nymphalis antiopa*	3,214	85	21	905	-0.021	0.018	0.256
Clouded Sulphur	*Colias philodice*	49,267	102	21	998	-0.014	0.014	0.286
Spicebush Swallowtail	*Papilio troilus*	25,322	82	21	858	-0.014	0.014	0.324
Dun Skipper	*Euphyes vestris*	1,684	49	21	585	-0.014	0.012	0.224
Question Mark	*Polygonia interrogationis*	6,564	88	21	915	-0.012	0.025	0.640
Delaware Skipper	*Atrytone logan*	1,086	30	21	313	-0.011	0.029	0.697
Horace's Duskywing	*Erynnis horatius*	2,885	31	21	376	-0.011	0.023	0.633
Eastern Tiger Swallowtail	*Papilio glaucus*	29,299	101	21	996	-0.010	0.015	0.483
Pearl Crescent	*Phyciodes tharos*	180,631	104	21	1,005	-0.010	0.014	0.461
Little Yellow	*Eurema lisa*	1,681	24	18	287	-0.008	0.073	0.917
Eastern Comma	*Polygonia comma*	6,222	92	21	944	-0.007	0.011	0.561
Giant Swallowtail	*Papilio cresphontes*	1,109	28	21	322	0.002	0.019	0.912
Banded Hairstreak	*Satyrium calanus*	1,107	36	21	468	0.004	0.031	0.896
Silver-spotted Skipper	*Epargyreus clarus*	54,462	102	21	996	0.005	0.012	0.672
Red Admiral	*Vanessa atalanta*	28,637	97	21	969	0.008	0.044	0.865
Red-banded Hairstreak	*Calycopis cecrops*	795	7	17	91	0.009	0.057	0.879
Crossline Skipper	*Polites origenes*	1,087	27	21	347	0.009	0.020	0.636
Sachem	*Atalopedes campestris*	1,445	19	18	231	0.013	0.058	0.823
Peck's Skipper	*Polites peckius*	23,702	90	21	905	0.014	0.014	0.306
Northern Eyed Brown	*Satyrodes eurydice*	1,342	13	21	174	0.016	0.035	0.651
Eastern Tailed Blue	*Everes comyntas*	56,137	99	21	974	0.016	0.010	0.113
Henry's Elfin	*Callophrys henrici*	330	7	17	76	0.017	0.055	0.752
Little Glassy Wing	*Pompeius verna*	8,658	56	21	632	0.019	0.019	0.307
Silvery Checkerspot	*Chlosyne nycteis*	2,049	20	19	224	0.039	0.022	0.074

Spring/Summer Azure	*Celastrina ladon/neglecta*	63,947	103	21	1,002	*0.047*	*0.021*	*0.022*
Common Buckeye	*Junonia coenia*	15,771	73	19	834	0.050	0.067	0.459
Pipevine Swallowtail	*Battus philenor*	703	23	18	279	0.053	0.033	0.110
Least Skipper	*Ancyloxypha numitor*	27,506	84	21	844	*0.053*	*0.015*	*<0.001*
Appalachian Eyed Brown	*Satyrodes appalachia*	2,118	12	18	118	0.060	0.045	0.181
Zabulon Skipper	*Poanes zabulon*	10,960	71	21	747	*0.061*	*0.022*	*0.004*
Northern Pearly-Eye	*Enodia anthedon*	2,785	37	21	434	*0.071*	*0.020*	*<0.001*
Zebra Swallowtail	*Eurytides marcellus*	1,349	18	18	224	*0.075*	*0.030*	*0.011*
Cloudless Sulphur	*Phoebis sennae*	1,840	27	19	355	0.088	0.057	0.121
Common Checkered-Skipper	*Pyrgus communis*	3,089	33	18	357	*0.092*	*0.046*	*0.046*
Wild Indigo Duskywing	*Erynnis baptisiae*	15,209	51	19	570	*0.106*	*0.020*	*<0.001*
Harvester	*Feniseca tarquinius*	341	11	20	143	*0.122*	*0.061*	*0.046*
Sleepy Orange	*Eurema nicippe*	2,028	6	17	63	0.146	0.134	0.276
Gemmed Satyr	*Cyllopsis gemma*	1,059	6	16	81	*0.228*	*0.052*	*<0.001*

Fig 2: The relative abundance of counted butterflies in Ohio declined by 33% over 1996-

2016. Plotted are model predictions for each year based on the fixed effects of year (solid line)

and annual random effects (dots) to show annual variation about the trend line. Shading shows

95% confidence interval based on bootstrapped model fits for the temporal trend.

Both in the total trend in abundance and in the proportion of species with declines, these

results are comparable to our review of three European butterfly monitoring schemes (Table 3).

Although the longer-running programs show larger cumulative declines, the annual rate of

change ranges from -2.0% to -2.6% for Ohio, Catalonia, and the Netherlands, respectively, with

the United Kingdom total trends split between generalist species (-0.8%) and specialist species (-

2.4%). Across monitoring programs, declining species outnumber increasing species by a factor

of two to three (Table 3).

Table 3: Comparison of estimated annualized and cumulative rates of change in total abundance and species trends in regional

butterfly monitoring programs in Europe compared to this study. Number of sites represents those reported to be analyzed in total, but

may no longer be active. Number of butterflies counted per year is based on the most recent years of monitoring described in the

references.

Region (km^2)	Years	Sites	Counted/year (x 1000)	Annualized trend in total abundance (cumulative)	Species' trends			Reference
					Positive	Negative	Stable/ not signif.	
United Kingdom (242,500)	41 (1976-2017)	3,164	1,700	-0.8% (-28%) countryside -2.4% (-63%) specialist	11	22	24	(50)
Netherlands (42,508)	25 (1992-2017)	600	250	-2.0% (-40%)	11	23	13	(51)
Catalonia, Spain (32,108)	22 (1994-2016)	116	122	-2.6% (-44%)	15	46	5	(39,52)
Ohio, USA (116,100)	**20 (1996-2016)**	**104**	**80**	**-2.0% (-33%)**	**9**	**32**	**40**	**this study**

In general, traits associated with species' responses to climate were more important,

based on the predictive ability (adjusted R^2) of univariate models, than traits associated with

habitat and host plant restrictions (Fig 3, Tables A and B in S1 Appendix). Phylogenetic signal

was included in most traits, so we focus on the phylogenetic generalized least squares results.

Migratory species to Ohio had stable population trends on average compared to resident species

and the four naturalized species. The Monarch (*Danaus plexippus*) was the only migratory

species in decline. Multivoltine species with more annual generations had more positive

population trends. Species with more northern geographic ranges were associated with more

negative population trends. Species eating forb host plants had negative trends on average, but

there was no effect of host plant specialization on population trends.

Fig 3: Species' traits are associated with variation in the statewide trends in abundance. We

plot each trend compared to the six most important traits for the 78 species included in the

phylogenetic GLS models with full results in Table A in S1 Appendix. Squares represent the

regression coefficients with 95% confidence intervals shown in lines. Dots for each species are

jittered for visualization.

Our choice of modeling approach did not change the overall evidence of defaunation.

Generalized linear mixed models with Poisson-distributed errors and generalized additive mixed

models estimated declines in total abundance similar in magnitude at -1.83% and -2.13% annual

rates, respectively. The annual trend estimate from TRIM, without sampling covariates, was half

the magnitude at -0.96%. Species' trends had high correlations between pairwise comparisons,

but TRIM models had notably more positive trends compared to the other three approaches

(Table C in S1 Appendix).

Discussion

We show that the total butterfly abundance has declined by 33% over 20 years in Ohio.

This rate is faster than the global abundance trend estimated for Lepidoptera (35% over 40 years)

and corresponds more closely to the steeper declines (45% over 40 years) estimated for all

insects (1). The Ohio butterfly monitoring program, judged by the weekly frequency, 20-year

time period, and statewide spatial extent of its surveys, is the most extensive systematic insect

survey in North America and comparable to three exemplary European butterfly monitoring

schemes. The annualized 2% rate of decline in this study aligns closely with trends from

European butterfly monitoring, confirming the decline of the most closely monitored group of

insects in both Europe and North America (Table 3). With less known about other insect taxa,

butterflies provide a necessary, if imperfect, surrogate to understand the trajectory and potential

mechanisms behind broader insect trends (13). Extensive in both time and space, the decline in

butterfly abundance is the current best estimate for the rate of insect defaunation in North

America.

The proportion of butterfly species with population declines compared to population

increases is similar between Ohio (negative trends three times more numerous) and European

studies (negative trends 2-3 times more numerous) (Table 3). In other taxa, moths in the United

Kingdom show a similar proportion of species declines (67). Long-term monitoring in protected

areas, although less extensive in space, shows more positive species trends for moths in Finland

(at 67.7° latitude) and across pollinators in Spain (at 850-1750 m. elevations) (68,69). These

counterexamples show how insect communities may shift at high-latitude or high-elevation sites

with anthropogenic climate warming (21) or may persist in more remote areas. However,

butterfly monitoring in populated areas show a consistency in observed declines (Table 3) that

we argue would generalize to other landscapes dominated by human use.

We demonstrate declines in species that are generalist, widespread, and not considered

vulnerable to extinction (25,70). The four species introduced to Ohio (*Pieris rapae*, *Lycaena*

phlaeus, and *Thymelicus lineola* from Europe and *Colias eurytheme* from the western USA) are

declining more rapidly than native species (Fig 3). Although few may share concern for the most

widespread, invasive butterfly in the world's agricultural and urban settings (71), declines in

Pieris rapae could be indicative of persistent environmental stressors that would affect other

species as well. We would expect negative environmental changes to disproportionately affect

rare species prone to the demographic dangers of small populations or specialist species that rely

on a narrow range of resources or habitat (UKBMS in Table 3, Swengel et al. 2011). This pattern

of species declines would lead to biotic homogenization as rarer species are lost and common,

disturbance-tolerant species remain (72,73). However, our study adds another example of

declines in common butterfly species thought to be well-suited to human-modified habitat

(11,21,74).

The Eastern North American migratory Monarch (*Danaus plexippus*) abundance in Ohio

is declining by 7% per year. The Monarch is the only declining migratory species out of 14 in

our analysis. Despite disagreements about whether summer abundance trends have tracked

winter colony declines (75,76), our study shows that the long-term trends correspond. However,

our study's first two years have very high Monarch population indices which could be outliers

(Fig B in S1 Appendix) following the two largest recorded winter population counts (16,77).

With these two years removed, the statewide Monarch trend is a 4% decline per year, showing

that the magnitude of summer abundance trends are sensitive to the years of inclusion. Our

results align with a study using Illinois systematic monitoring data that shows a summer

abundance decline for monarchs over two decades, but at different rates across decades (78). A

more recent study showed no decline during the summer during 2004-2016 using a population

index from NABA counts (77). The trend we document comes from the sum of summer breeding

and fall migratory butterflies returning to Mexico; estimates of abundance for these separate

generations may be required to model how different stages of the lifecycle contribute to the long-

term decline in the winter colonies (77).

Even with systematic monitoring, accurate estimates of insect abundance are missing

from many species—a fifth of regularly observed species in did not meet our minimum data

requirements to for us to estimate trends. None of these species are considered to be of

conservation concern, but this also means that we would be limited in our ability to even

determine if their populations have reached threatened status. Targeted surveys of selected

species, non-adult life stages, or habitats can expand the monitoring to data-deficient species

commonly excluded by protocols designed to monitor many species efficiently (50) and can be

used to estimate demographic responses to environmental drivers not apparent from adult

butterfly counts (79). Additional life-history knowledge about species in our study could inform

how worried we should be about extreme population declines, like the Baltimore (*Euphydryas*

phaeton). We noticed a multiyear population cycle, not captured by log-linear trends or

generalized additive models, which reached a nadir at the end of our dataset and exaggerated the

Baltimore's decline (Fig B in S1 Appendix). In other cases, density-dependent population

regulation may help inform whether species are resilient to temporary declines and guide

predictions for future population trajectories (80).

Insect declines have multifaceted causes, and the relative impact of these causes is still

unknown (81). Although analysis of the causes of site differences in abundance or species trends

is beyond the scope of this study, we discuss three environmental drivers commonly associated

with global insect declines: climate change, habitat loss and fragmentation, and agricultural

intensification (81,82). If species' traits are associated with population trends, then their

relationships may suggest environmental changes driving population responses in species sharing

these traits (46,81,83). In this study, life-history traits were weakly predictive of population

trends, but their associations provide hypotheses that could be tested further (46).

Habitat loss and fragmentation

In Ohio, habitat loss and fragmentation plateaued well before butterfly monitoring

started, with human population growth slowing by 1970. In common with other Midwestern

states, Ohio had already lost tallgrass prairie species, such as the Regal Fritillary (*Speyeria*

idalia), due to this habitat conversion to agriculture (25,26). Land-use has changed slowly over

the course of the monitoring program; less than 10% of monitoring sites have had more than

2.5% change in the surrounding (2.5-km radius) developed, agriculture, or forest land cover from

2001-2011. The persistence of butterfly populations in a landscape of habitat fragments are

mediated by species' traits that permit them to either move between more isolated resources or

persist in smaller, localized populations (82,84). Wing size is one life history trait associated

with dispersal ability, but it had no association with species' population trends (Tables A and B

in S1 Appendix). However, defining habitat patches by land-use classes overlooks how mobile

insect populations are bound by resources, varying across the lifecycle, rather than area (85,86).

Although there has been little wholesale habitat conversion around our study transects,

degradation of the remaining habitat could be a cause of the general decline in butterfly

abundance.

Climate change

Species trends are associated with two life-history traits, voltinism and range distribution,

which suggest that the butterfly community is changes with the warming climate. Species that

complete more annual generations, or multivoltine species, had more positive abundance trends.

This aligns with obligate univoltine species becoming less common in Massachusetts (20), but is

the opposite of the findings in Spain where multivoltine species are in steeper declines with

exposure to increasingly dry summers (39). Multivoltine species may be more adaptive to annual

and spatial variation in seasonal temperatures as many have plasticity in the voltinism observed

within Ohio (25). For most of the species with flexible voltinism in Ohio, adding an extra

generation in warmer summers increases their annual population growth rates (54). Northern-

distributed species have more negative population trends compared to widely-distributed or

southern species. This corresponds with findings from Massachusetts and Europe that warm-

adapted species are replacing cool-adapted species as range distributions shift (20,87). Even

though these two traits should be increasing butterfly abundance for some species as the climate

warms, it has not been enough to prevent the overall decline in butterfly abundance.

Agricultural intensification

Cropland and pasture make up half of Ohio's land area, so we would expect agricultural

practices to affect statewide insect abundance. One assessment of pollinator habitat suitability

based on land-use, acres in conservation reserve programs, and crop type estimated an increase

in resources in Ohio from 1982 through 2002, followed by a stable trend (88). However,

agricultural practices can decrease insect abundance with systemic insecticides, herbicide use on

host plants or nectar resources, and nitrogen fertilization that alters the composition of

surrounding plant communities.

In Ohio, the use of neonicotinoids rapidly increased after 2004 when they became widely

used on corn and soybeans (89,90). The mechanistic link between neonicotinoid insecticides and

insect declines is established and observational studies have shown widespread impacts of their

use (91–93). Even though seed-coatings with neonicotinoids reduce broadcast spraying, the

mechanical planting of these seeds exposes widespread areas around farms to contaminated dust

that is incorporated into non-target plants and insects (94,95). In the United Kingdom and

California, neonicotinoids are associated with butterfly declines (22,96) and hinder butterfly

larval development on host plants (97). We did not design this study to test whether

neonicotinoids affect butterfly abundance in Ohio. However, the observed declines across

common, invasive species, which would typically be predicted to exploit an agricultural or

human-altered landscape, would be consistent with widespread exposure to insecticides.

Species that eat forbs as larvae have negative population declines (Fig 3). Both herbicide

use and nitrogen deposition may alter plant communities to favor grasses over forbs (98).

Milkweed losses contribute to declines in Monarch butterfly abundance, as they lose host333

plants as a result of herbicide use (78,79). In Ohio, glyphosate use has increased linearly, and is

now applied at 6 times the rate it was in 1996 (89,90). Nitrogen increases have been linked to

declines in grassland species adapted to low-nitrogen environments (99–101) and to higher

mortality during larval development on enriched host plants (102).

Conclusions

Systematic, long-term surveys of butterflies provide the most rigorous evidence for the

rate of insect declines. This study demonstrates that defaunation is happening in North America

similarly to Europe. In landscapes comprising natural areas amid heavy human land-use,

butterfly total abundance is declining at 2% per year and 2-3 times more species have population

trends declining rather than increasing. Additional Pollard-based monitoring programs in North

America, listed in (9), will enable tracking insect trends over larger spatial extents as with efforts

to integrate data across European monitoring schemes (11). The rates for other insect groups may

deviate from this baseline and were previously estimated to be declining more rapidly than

Lepidoptera (1). Expanded monitoring and support for taxonomists are imperative for other taxa

and under sampled regions, like the Tropics where most insect diversity resides. Besides the

evaluation if butterfly trends generalize to other insects, the most urgent need for science and

conservation is understanding the causes of decline and testing mitigation actions. As butterflies

are the best-monitored insect taxa, they are the best indicator of the baseline threat to the 5.5

million insects, the most diverse group of animals on earth.

Acknowledgments

We thank the volunteers and directors who contribute their time and expertise to the Ohio

butterfly monitoring program. The Ohio Department of Natural Resources provides support, the

Cleveland Museum of Natural History provides data entry and archiving, and the Ohio

Lepidopterists provides training and coordination for the Ohio butterfly monitoring program. We

thank Marjorie Weber for advice on our phylogenetic analysis. The Department of the Interior

Southeast Climate Adaptation Science Center and North Carolina State University supported TW

during earlier work with Ohio butterflies that grew into this analysis.

References

1.   Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJB, Collen B. Defaunation in the

Anthropocene. Science. 2014 Jul 25;345(6195):401–6.

2.   May RM. How many species are there on earth? Science. 1988;241(4872):1441–1449.

3.   Conrad KF, Fox R, Woiwod IP. Monitoring biodiversity: measuring long-term changes in

insect abundance. Insect Conserv Biol. 2007;203–225.

4.   Hallmann CA, Sorg M, Jongejans E, Siepel H, Hofland N, Schwan H, et al. More than 75

percent decline over 27 years in total flying insect biomass in protected areas. Lamb EG,

editor. PLOS ONE. 2017 Oct 18;12(10):e0185809.

5.   Lister BC, Garcia A. Climate-driven declines in arthropod abundance restructure a

rainforest food web. Proc Natl Acad Sci. 2018 Oct 30;115(44):E10397–406.

6.   Winfree R, W. Fox J, Williams NM, Reilly JR, Cariveau DP. Abundance of common

species, not species richness, drives delivery of a real-world ecosystem service. Shea K,

editor. Ecol Lett. 2015 Jul;18(7):626–35.

7.   Warren MS, Hill JK, Thomas JA, Asher J, Fox R, Huntley B, et al. Rapid responses of

British butterflies to opposing forces of climate and habitat change. 2001;414:5.

8.   Parmesan C. Ecological and Evolutionary Responses to Recent Climate Change. Annu

Rev Ecol Evol Syst. 2006 Dec;37(1):637–69.

9.   Taron D, Ries L. Butterfly Monitoring for Conservation. In: Daniels JC, editor. Butterfly

Conservation in North America [Internet]. Dordrecht: Springer Netherlands; 2015 [cited

2018 Dec 7]. p. 35–57. Available from: http://link.springer.com/10.1007/978-94-017-

9852-5_3

10.   Pollard E, Yates TJ. Monitoring butterflies for ecology and conservation: the British

520   butterfly monitoring scheme. Springer Science & Business Media; 1994.

521   11.   Van Swaay C, Van Strien A, Aghababyan K, Astrom S, Botham M, Brereton T, et al. The

522   European Butterfly Indicator for Grassland species: 1990-2013. 2015;

523   12.   Schmucki R, Pe'er G, Roy DB, Stefanescu C, Van Swaay CAM, Oliver TH, et al. A

524   regionally informed abundance index for supporting integrative analyses across butterfly

525   monitoring schemes. Müller J, editor. J Appl Ecol. 2016 Apr;53(2):501–10.

526   13.   Thomas JA. Monitoring change in the abundance and distribution of insects using

527   butterflies and other indicator groups. Philos Trans R Soc B Biol Sci. 2005 Feb

528   28;360(1454):339–57.

529   14.   Brereton T, Roy DB, Middlebrook I, Botham M, Warren M. The development of butterfly

530   indicators in the United Kingdom and assessments in 2010. J Insect Conserv. 2011

531   Apr;15(1–2):139–51.

532   15.   Dennis EB, Morgan BJT, Roy DB, Brereton TM. Urban indicators for UK butterflies.

533   Ecol Indic. 2017 May;76:184–93.

534   16.   Agrawal AA, Inamine H. Mechanisms behind the monarch's decline. Science.

535   2018;360(6395):1294–1296.

536   17.   Schultz CB, Brown LM, Pelton E, Crone EE. Citizen science monitoring demonstrates

537   dramatic declines of monarch butterflies in western North America. Biol Conserv.

538   2017;214:343–346.

539   18.   Belitz MW, Hendrick LK, Monfils MJ, Cuthrell DL, Marshall CJ, Kawahara AY, et al.

540   Aggregated occurrence records of the federally endangered Poweshiek skipperling

541   (Oarisma poweshiek). Biodivers Data J. 2018;(6).

19. Haddad NM. Resurrection and resilience of the rarest butterflies. PLOS Biol. 2018 Feb

6;16(2):e2003488.

20. Breed GA, Stichter S, Crone EE. Climate-driven changes in northeastern US butterfly

communities. Nat Clim Change. 2013 Feb;3(2):142–5.

21. Forister ML, McCall AC, Sanders NJ, Fordyce JA, Thorne JH, O'Brien J, et al.

Compounded effects of climate change and habitat alteration shift patterns of butterfly

diversity. Proc Natl Acad Sci. 2010 Feb 2;107(5):2088–92.

22. Forister ML, Cousens B, Harrison JG, Anderson K, Thorne JH, Waetjen D, et al.

Increasing neonicotinoid use and the declining butterfly fauna of lowland California. Biol

Lett. 2016 Aug;12(8):20160475.

23. Schlicht D, Swengel A, Swengel S. Meta-analysis of survey data to assess trends of prairie

butterflies in Minnesota, USA during 1979–2005. J Insect Conserv. 2009 Aug;13(4):429–

47.

24. Swengel SR, Schlicht D, Olsen F, Swengel AB. Declines of prairie butterflies in the

midwestern USA. J Insect Conserv. 2011 Apr;15(1–2):327–39.

25. Iftner DC, Shuey JA, Calhoun JV. Butterflies and skippers of Ohio. College of Biological

Sciences, Ohio State University; 1992.

26. Shuey JA, Calhoun JV, Iftner DC. Butterflies that are endangered, threatened, and of

special concern in Ohio. 1987;

27. Bigelow D, Borchers A. Major uses of land in the United States, 2012. 2017.

28. Thornton PE, Running SW, White MA. Generating surfaces of daily meteorological

variables over large regions of complex terrain. J Hydrol. 1997 Mar;190(3–4):214–51.

29.   Homer C, Dewitz J, Yang L, Jin S, Danielson P, Xian G, et al. Completion of the 2011

565   National Land Cover Database for the conterminous United States–representing a decade

566   of land cover change information. Photogramm Eng Remote Sens. 2015;81(5):345–354.

567   30.   Diamond SE, Cayton H, Wepprich T, Jenkins CN, Dunn RR, Haddad NM, et al.

568   Unexpected phenological responses of butterflies to the interaction of urbanization and

569   geographic temperature. Ecology. 2014 Sep;95(9):2613–21.

570   31.   Dennis EB, Freeman SN, Brereton T, Roy DB. Indexing butterfly abundance whilst

571   accounting for missing counts and variability in seasonal pattern. O'Hara RB, editor.

572   Methods Ecol Evol. 2013 Jul;4(7):637–45.

573   32.   Opler PA, Krizek GO. Butterflies east of the Great Plains: an illustrated natural history.

574   Johns Hopkins Univ Pr; 1984.

575   33.   Dennis EB, Morgan BJT, Freeman SN, Brereton TM, Roy DB. A generalized abundance

576   index for seasonal invertebrates: A Generalized Abundance Index for Seasonal

577   Invertebrates. Biometrics. 2016 Dec;72(4):1305–14.

578   34.   Wood SN. Generalized Additive Models: An Introduction with R. 2nd ed. Chapman and

579   Hall/CRC; 2017.

580   35.   Cordano EE& E. Interpol.T: Hourly interpolation of multiple temperature daily series

581   [Internet]. 2013. Available from: https://CRAN.R-project.org/package=Interpol.T

582   36.   Link WA, Sauer JR. Estimating Population Change from Count Data: Application to the

583   North American Breeding Bird Survey. Ecol Appl. 1998 May;8(2):258.

584   37.   Pellet J, Bried JT, Parietti D, Gander A, Heer PO, Cherix D, et al. Monitoring Butterfly

585   Abundance: Beyond Pollard Walks. Schweiger O, editor. PLoS ONE. 2012 Jul

586   30;7(7):e41396.

587   38.   Henry EH, Anderson CT. Abundance estimates to inform butterfly management: double-

588   observer versus distance sampling. J Insect Conserv. 2016 Jun;20(3):505–14.

589   39.   Melero Y, Stefanescu C, Pino J. General declines in Mediterranean butterflies over the last

590   two decades are modulated by species traits. Biol Conserv. 2016 Sep;201:336–42.

591   40.   Haddad NM, Hudgens B, Damiani C, Gross K, Kuefler D, Pollock K. Determining

592   Optimal Population Monitoring for Rare Butterflies: *Monitoring Rare Butterflies*. Conserv

593   Biol. 2008 Aug;22(4):929–40.

594   41.   Isaac NJB, Cruickshanks KL, Weddle AM, Marcus Rowcliffe J, Brereton TM, Dennis

595   RLH, et al. Distance sampling and the challenge of monitoring butterfly populations:

596   Distance sampling and monitoring butterflies. Methods Ecol Evol. 2011 Dec;2(6):585–94.

597   42.   Knape J. Decomposing trends in Swedish bird populations using generalized additive

598   mixed models. Siriwardena G, editor. J Appl Ecol. 2016 Dec;53(6):1852–61.

599   43.   Szabo JK, Vesk PA, Baxter PWJ, Possingham HP. Regional avian species declines

600   estimated from volunteer-collected long-term data using List Length Analysis. Ecol Appl.

601   2010;20(8):2157–69.

602   44.   van Strien AJ, van Swaay CAM, Termaat T. Opportunistic citizen science data of animal

603   species produce reliable estimates of distribution trends if analysed with occupancy

604   models. Devictor V, editor. J Appl Ecol. 2013 Dec;50(6):1450–8.

605   45.   Roy DB, Ploquin EF, Randle Z, Risely K, Botham MS, Middlebrook I, et al. Comparison

606   of trends in butterfly populations between monitoring schemes. J Insect Conserv. 2015

607   Apr;19(2):313–24.

608   46.   Soykan CU, Sauer J, Schuetz JG, LeBaron GS, Dale K, Langham GM. Population trends

609   for North American winter birds based on hierarchical models. Ecosphere.

610   2016;7(5):e01351.

611   47.   Kraemer HC, Blasey CM. Centring in regression analyses: a strategy to prevent errors in

612   statistical inference. Int J Methods Psychiatr Res. 2004;13(3):141–51.

613   48.   Harrison XA. Using observation-level random effects to model overdispersion in count

614   data in ecology and evolution. PeerJ. 2014 Oct 9;2:e616.

615   49.   Pannekoek J, Van Strien A. Trim 3 Manual (TRends & Indices for Monitoring data)–

616   Statistics Netherlands. Voorburg; 2001.

617   50.   Brereton T, Botham M, Middlebrook I, Randle Z, Noble D, Harris S, et al. United

618   Kingdom Butterfly Monitoring Scheme report for 2017. Centre for Ecology & Hydrology

619   & Butterfly Conservation; 2018.

620   51.   Van Swaay CAM, Bos G, Van Grunsven RHA, Kok J, Huskens K, Van Deijk JR, et al.

621   Vlinders en libellen geteld: Jaarverslag 2017 [Internet]. De Vlinderstichting, Wageningen;

622   2018 [cited 2019 Feb 14]. Report No.: Rapport VS2018.006. Available from:

623   https://assets.vlinderstichting.nl/docs/6d51f174-b497-4777-b84a-362e344c3528.pdf

624   52.   Stefanescu C. Resum de les temporades 2015 i 2016. Cynthia Butlletí Butterfly Monit

625   Scheme Catalunya [Internet]. 2018 [cited 2019 Feb 14];14. Available from:

626   http://www.catalanbms.org/es/cynthia/

627   53.   Rohatgi A. WebPlotDigitizer. 2011.

628   54.   Wepprich TM. Effects of Climatic Variability on a Statewide Butterfly Community. North

629   Carolina State University; 2017.

630   55.   Diamond SE, Frame AM, Martin RA, Buckley LB. Species' traits predict phenological

631   responses to climate change in butterflies. 2011;92(5):8.

632   56.   Pagel M. Inferring the historical patterns of biological evolution. Nature. 1999

633   Oct;401(6756):877–84.

634   57.   Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. GenBank. Nucleic Acids

635   Res. 2016 Jan 4;44(Database issue):D67–72.

636   58.   Ratnasingham S, Hebert PD. BOLD: The Barcode of Life Data System (http://www.

637   barcodinglife. org). Mol Ecol Notes. 2007;7(3):355–364.

638   59.   Espeland M, Breinholt J, Willmott KR, Warren AD, Vila R, Toussaint EFA, et al. A

639   Comprehensive and Dated Phylogenomic Analysis of Butterflies. Curr Biol. 2018

640   Mar;28(5):770-778.e5.

641   60.   R Core Team. R: A Language and Environment for Statistical Computing [Internet].

642   Vienna, Austria: R Foundation for Statistical Computing; 2018. Available from:

643   https://www.R-project.org/

644   61.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using

645   **lme4**. J Stat Softw [Internet]. 2015 [cited 2018 Dec 4];67(1). Available from:

646   http://www.jstatsoft.org/v67/i01/

647   62.   Orme D, Freckleton R, Thomas G, Petzoldt T. The caper package: comparative analysis of

648   phylogenetics and evolution in R. R Package Version. 2013;5(2).

649   63.   Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary

650   analyses in R. Bioinformatics. 2018;xx:xxx–xxx.

651   64.   Knowles JE, Frederick C. merTools: Tools for Analyzing Mixed Effect Regression

652   Models [Internet]. 2018. Available from: https://CRAN.R-project.org/package=merTools

653   65.   Nakagawa S, Schielzeth H. A general and simple method for obtaining R^2 from

654   generalized linear mixed-effects models. O'Hara RB, editor. Methods Ecol Evol. 2013

655   Feb;4(2):133–42.

656   66.   Barton K. MuMIn: Multi-Model Inference [Internet]. 2018. Available from:

657   https://CRAN.R-project.org/package=MuMIn

658   67.   Conrad KF, Woiwod IP, Parsons M, Fox R, Warren MS. Long-term population trends in

659   widespread British moths. J Insect Conserv. 2004;8(2–3):119–136.

660   68.   Hunter MD, Kozlov MV, Itämies J, Pulliainen E, Bäck J, Kyrö E-M, et al. Current

661   temporal trends in moth abundance are counter to predicted effects of climate change in an

662   assemblage of subarctic forest moths. Glob Change Biol. 2014 Jun;20(6):1723–37.

663   69.   Herrera CM. Complex long-term dynamics of pollinator abundance in undisturbed

664   Mediterranean montane habitats over two decades. Ecol Monogr. 2019 Feb

665   1;89(1):e01338.

666   70.   Cech R, Tudor G, others. Butterflies of the east coast. Princeton University Press; 2005.

667   71.   Ryan SF, Lombaert E, Espeset A, Vila R, Talavera G, Dincă V, et al. Global invasion

668   history of the world's most abundant pest butterfly: a citizen science population genomics

669   study. bioRxiv. 2018 Dec 26;506162.

670   72.   McKinney ML, Lockwood JL. Biotic homogenization: a few winners replacing many

671   losers in the next mass extinction. Trends Ecol Evol. 1999;14(11):450–453.

672   73.   Clavel J, Julliard R, Devictor V. Worldwide decline of specialist species: toward a global

673   functional homogenization? Front Ecol Environ. 2011;9(4):222–8.

74.   Van Dyck H, Van Strien AJ, Maes D, Van Swaay CAM. Declines in Common,

Widespread Butterflies in a Landscape under Intense Human Use. Conserv Biol. 2009

Aug;23(4):957–65.

75.   Ries L, Oberhauser K, Taron D, Battin J, Rendon-Salinas E, Altizer S, et al. Connecting

eastern monarch population dynamics across their migratory cycle. Monarchs Chang

World Biol Conserv Iconic Insect Cornell Univ Press Ithaca NY. 2015;268–281.

76.   Pleasants JM, Williams EH, Brower LP, Oberhauser KS, Taylor OR. Conclusion of No

Decline in Summer Monarch Population Not Supported. Ann Entomol Soc Am. 2016 Mar

1;109(2):169–71.

77.   Saunders SP, Ries L, Neupane N, Ramírez MI, García-Serrano E, Rendón-Salinas E, et al.

Multiscale seasonal factors drive the size of winter monarch colonies. Proc Natl Acad Sci.

2019 Mar 18;201805114.

78.   Saunders SP, Ries L, Oberhauser KS, Thogmartin WE, Zipkin EF. Local and cross-

seasonal associations of climate and land use with abundance of monarch butterflies

Danaus plexippus. Ecography. 2018 Feb;41(2):278–90.

79.   Pleasants JM, Oberhauser KS. Milkweed loss in agricultural fields because of herbicide

use: effect on the monarch butterfly population: *Herbicide use and monarch butterflies*.

Insect Conserv Divers. 2013 Mar;6(2):135–44.

80.   Oliver TH, Roy DB. The pitfalls of ecological forecasting. Biol J Linn Soc. 2015

Jul;115(3):767–78.

81.   Fox R. The decline of moths in Great Britain: a review of possible causes: *The decline of*

moths in Great Britain. Insect Conserv Divers. 2013 Jan;6(1):5–19.

82.   Thomas JA. Butterfly communities under threat. Science. 2016 Jul 15;353(6296):216–8.

83.   Wong MKL, Guénard B, Lewis OT. Trait-based ecology of terrestrial arthropods. Biol

Rev [Internet]. 2018 [cited 2018 Dec 14];0(0). Available from:

https://onlinelibrary.wiley.com/doi/abs/10.1111/brv.12488

84.   Habel JC, Schmitt T. Vanishing of the common species: Empty habitats and the role of

genetic diversity. Biol Conserv. 2018 Feb;218:211–6.

85.   Dennis RLH, Shreeve TG, Dyck HV. Towards a Functional Resource-Based Concept for

Habitat: A Butterfly Biology Viewpoint. Oikos. 2003;102(2):417–26.

86.   Curtis RJ, Brereton TM, Dennis RLH, Carbone C, Isaac NJB. Butterfly abundance is

determined by food availability and is mediated by species traits. Diamond S, editor. J

Appl Ecol. 2015 Dec;52(6):1676–84.

87.   Parmesan C, Ryrholm N, Stefanescu C, Hill JK, Thomas CD, Descimon H, et al. Poleward

shifts in geographical ranges of butterfly species associated with regional warming.

Nature. 1999 Jun;399(6736):579–83.

88.   Hellerstein D. Land Use, Land Cover, and Pollinator Health: A Review and Trend

Analysis. :47.

89.   Baker NT, Stone WW. Estimated annual agricultural pesticide use for counties of the

conterminous United States, 2008–12: U.S. Geological Survey Data Series 907 [Internet].

US Geological Survey; p. 9. (U.S. Geological Survey Data Series 907). Available from:

https://dx.doi.org/10.3133/ds907.

90.   Stone WW. Estimated annual agricultural pesticide use for counties of the conterminous

United States, 1992–2009. US Geological Survey; 2013. (U.S. Geological Survey Data

Series 752).

91.　Goulson D. REVIEW: An overview of the environmental risks posed by neonicotinoid

insecticides. Kleijn D, editor. J Appl Ecol. 2013 Aug;50(4):977–87.

92.　Hallmann CA, Foppen RPB, van Turnhout CAM, de Kroon H, Jongejans E. Declines in

insectivorous birds are associated with high neonicotinoid concentrations. Nature. 2014

Jul;511(7509):341–3.

93.　Pisa L, Goulson D, Yang E-C, Gibbons D, Sánchez-Bayo F, Mitchell E, et al. An update

of the Worldwide Integrated Assessment (WIA) on systemic insecticides. Part 2: impacts

on organisms and ecosystems. Environ Sci Pollut Res [Internet]. 2017 Nov 9 [cited 2019

Feb 1]; Available from: https://doi.org/10.1007/s11356-017-0341-3

94.　Douglas MR, Tooker JF. Large-Scale Deployment of Seed Treatments Has Driven Rapid

Increase in Use of Neonicotinoid Insecticides and Preemptive Pest Management in U.S.

Field Crops. Environ Sci Technol. 2015 Apr 21;49(8):5088–97.

95.　Krupke CH, Holland JD, Long EY, Eitzer BD. Planting of neonicotinoid-treated maize

poses risks for honey bees and other non-target organisms over a wide area without

consistent crop yield benefit. Diamond S, editor. J Appl Ecol. 2017 Oct;54(5):1449–58.

96.　Gilburn AS, Bunnefeld N, Wilson JM, Botham MS, Brereton TM, Fox R, et al. Are

neonicotinoid insecticides driving declines of widespread butterflies? PeerJ. 2015 Nov

24;3:e1402.

97.　Basley K, Goulson D. Effects of Field-Relevant Concentrations of Clothianidin on Larval

Development of the Butterfly Polyommatus icarus (Lepidoptera, Lycaenidae). Environ Sci

Technol. 2018 Apr 3;52(7):3990–6.

98. Kleijn D, Snoeijing GIJ. Field Boundary Vegetation and the Effects of Agrochemical

Drift: Botanical Change Caused by Low Levels of Herbicide and Fertilizer. J Appl Ecol.

1997;34(6):1413–25.

99. Weiss SB. Cars, cows, and checkerspot butterflies: nitrogen deposition and management

of nutrient-poor grasslands for a threatened species. Conserv Biol. 1999;13(6):1476–1486.

100. Öckinger E, Hammarstedt O, Nilsson SG, Smith HG. The relationship between local

extinctions of grassland butterflies and increased soil nitrogen levels. Biol Conserv. 2006

Apr 1;128(4):564–73.

101. WallisDeVries MF, van Swaay CAM. A nitrogen index to track changes in butterfly

species assemblages under nitrogen deposition. Biol Conserv. 2017 Aug;212:448–53.

102. Kurze S, Heinken T, Fartmann T. Nitrogen enrichment in host plants increases the

mortality of common Lepidoptera species. Oecologia. 2018 Dec;188(4):1227–37.

S1 Appendix. Supplementary methods and results. Includes detailed methods for phenology

models and phylogenetic trees, one figure of species trends plotting on a cladogram, three figures

showing population trends and annual variation for 81 species, two tables of model results from

the trait analysis, and a table comparing our trend estimates with three other approaches.

Figure

Figure

Figure