**Predicting the conservation status of Europe's Data Deficient sharks and rays**

Predicting status of Data Deficient sharks

Rachel H.L. Walls[1]*, Nicholas K. Dulvy[1]

[1]Earth to Ocean Research Group, Department of Biological Sciences, Simon Fraser University,

Burnaby, BC, Canada

*Corresponding author.

*Contact*: +1(778)251-8227, rwalls@sfu.ca (Rachel HL Walls)

**Keywords**: Aichi target, Chondrichthyes, Elasmobranchii, depth refuge, General Fisheries

Commission for the Mediterranean, Red List Index

**ABSTRACT**

Shark and ray biodiversity is threatened primarily by overfishing and the globalisation of trade,

and Europe has been one of the most documented heavily fished regions for a relatively long

time. Yet, we have little idea of the conservation status of the hundreds of Data Deficient shark

and ray species. It is important to derive some insight into the status of these species, both to

understand global extinction rates and also to ensure that any threatened Data Deficient species

are not overlooked in conservation planning. Here, we developed a biological and ecological trait

model to predict the categorical conservation status of 26 Northeast Atlantic and 15

Mediterranean Sea Data Deficient sharks and rays. We first developed an explanatory model

based on all species *evaluated* on the International Union for Conservation of Nature (IUCN)

Red List of Threatened Species[TM], using maximum body size, median depth (as a proxy for

fisheries exposure), and reproductive mode, and then *predicted* the status of all Data Deficient

species. Almost half of Northeast Atlantic (46%, *n*=12 of 26), and two-thirds of Mediterranean

(67%, *n*=10 of 15) Data Deficient species are predicted to be in one of the three IUCN threatened

categories. Northeast Atlantic Data Deficient species are *predicted* to be 1.2 times more

threatened than *evaluated* species (38%, *n*=36 of 94), whereas threat levels in the Mediterranean

Sea are relative for each (66%, *n*=38 of 58). This case study is intended for extrapolation to the

global shark and ray dataset upon completion of the global IUCN Red List assessment. Trait-

based, categorical prediction of conservation status is a cost-effective approach towards

incorporating Data Deficient species into (i) estimates of lineage-wide extinction rates, (ii)

revised protected species lists, and (iii) Red List Indices, thus preventing poorly known species

from reaching extinction unnoticed.

## 1 INTRODUCTION

Despite a broadening of coverage of species and more intensive Red List assessment by the International Union for Conservation of Nature (IUCN) in the past decade, over one-sixth or around 13,465 species have been found to be Data Deficient (Bland et al., 2017). Data-deficiency is most prevalent in reptiles and amphibians, marine and freshwater organisms, invertebrates, and plants (Bland et al., 2012, 2014; Böhm et al., 2013; Callmander et al., 2005; Collen et al., 2012; Hoffmann et al., 2010). The IUCN classification means that there are insufficient data to make a more refined determination, hence Data Deficient species could range from actually being Least Concern or they could be threatened or even Extinct. Data-deficiency creates uncertainty in estimates of extinction rates, which is a key challenge to track progress towards the Convention on Biological Diversity's (CBD) Aichi Target 12: to halt the loss of biodiversity by 2020 (CBD & UNEP, 2011). Clearly, a complete understanding of which species are threatened (Vulnerable, Endangered, or Critically Endangered) is an essential first step toward tracking and improving species' status (Bland et al., 2014, 2015).

Data Deficient species are typically overlooked in conservation planning (Bland et al., 2014), with the implicit assumption that the biology and threatening processes of both Data Deficient and data-sufficient species are similar. To provide a first-approximation of the extinction rate of any taxon, the IUCN assumes Data Deficient species are equally as threatened as the data-sufficient species within a taxonomic group (Hoffmann et al., 2010). However, there are numerous reasons why the trait distribution and exposure to threatening processes might be different. For example, most recently discovered sharks have been found in the deep sea (Randhawa et al., 2015) and are relatively small-bodied, beyond the reach of most fisheries,

hence those Data Deficient deepwater species may actually be Least Concern because they have

refuge from the main threatening process of overfishing. Conversely, many recently resolved

species complexes, such as devil rays, eagle rays, and skates may be highly exposed to fisheries

and hence the newly described 'Data Deficient' species might already be highly threatened

(Iglésias et al., 2010; White & Last, 2012).

There is a vast body of work on the correlates of population trajectories and extinction risk

(Cardillo et al., 2005; McKinney, 1997; Owens & Bennett, 2000). Broadly, large body size,

small geographic range, and ecological specialisation are the biological traits most often related

to extinction risk, depending on their interaction with the appropriate threatening process (Owens

& Bennett, 2000; Reynolds et al., 2005b). Only recently has this knowledge been used to predict

extinction risk of Data Deficient species (Bland et al., 2015; Butchart & Bird, 2010; Dulvy et al.,

2014; Jetz & Freckleton, 2015). Trait-based predictions of IUCN conservation status use

biological and ecological trait data to predict the most likely categorisations for Data Deficient

species based on assessed species. The simplest approach is to make the binary prediction

whether a Data Deficient species is Least Concern or threatened. This approach has been used

with a high degree of accuracy for mammals, birds, sharks and rays (Bland et al., 2015; Butchart

& Bird, 2010; Dulvy et al., 2014; Jetz & Freckleton, 2015). The most significant advance has

been the development of ordinal (or categorical) regression which enables prediction of the

actual IUCN Red List category, based on relevant biological and ecological traits (Luiz et al.,

2016). A total of 50 of 163 groupers (family Epinephelinae) were Data Deficient, yet trait-based

ordinal regression revealed a total of three species predicted to be Critically Endangered, five to

be Endangered, and 12 to be Vulnerable (Luiz et al., 2016).

Sharks and rays represent the oldest evolutionary radiation of vertebrate Classes (Stein et al.,

2018), with an incredibly broad range of life-histories, spanning all ocean basins, and down to

great depths (Cortés, 2000; Dulvy et al., 2014; Dulvy & Forrest, 2010). This makes them ideal

for trait-based predictive modelling, while their high levels of population-relevant data-

deficiency present the opportunity to test categorical predictions on a highly Data Deficient

group for the first time. Europe represents the first region to be reassessed as part of an ongoing

global IUCN Red List reassessment of sharks and rays, as well as being one of the most

relatively data-sufficient regions for the Class (Dulvy et al., 2016; Fernandes et al., 2017; Nieto

et al., 2015).

Here, we use Europe's sharks and rays to consider three questions: (1) which biological and

ecological traits are driving extinction risk; (2) how does the proportion of *evaluated*-threatened

species compare with *predicted*-to-be-threatened Data Deficient species; and (3) which are the

most threatened Data Deficient sharks and rays? We used cumulative link mixed-effects

modeling (CLMM) to evaluate the relationship between species' trait data and conservation

status, and eventually predict the conservation status of Europe's Data Deficient sharks and rays.

This CLMM approach maintains the hierarchy of the IUCN categories while preventing the loss

of information inevitable from lumping categories together as threatened and non-threatened

(Luiz et al., 2016).  Model performance was evaluated using the Akaike Information Criterion

(AIC) with small sample size correction ($AIC_c$).

## 2 METHODS

First, we describe the IUCN Red List conservation assessment of European sharks and rays.

Second, we describe the development of an explanatory trait-based model to explain

conservation status. Third, we describe the prediction and cross-validation of the conservation

status of Europe's Data Deficient sharks and rays.

### 2.1 IUCN Red List assessment

The European Red List assessments spanned the Northeast Atlantic Ocean and the

Mediterranean and Black Seas, including the territorial waters and Exclusive Economic Zones of

all European countries in the Northeast and Eastern Central Atlantic Ocean, and the offshore

Macronesian island territories belonging to Portugal and Spain (Dulvy et al., 2016; Fernandes et

al., 2017; Nieto et al., 2015).

In total, 131 species were assessed at the regional level for Europe using the 2001 IUCN Red

List Categories and Criteria, version 3.1 (IUCN, 2012b). We convened 54 experts, composed

mainly of members of the IUCN Shark Specialist Group, and completed the 131 European

assessments over 21 months, from 2013–15. This culminated in a one-week workshop, attended

by fifteen IUCN Shark Specialist Group members, to finalise and review all assessments. The

assessed species included 50 skates and rays (Order Rajiformes), 72 sharks (Order

Carcharhiniformes, Hexanchiformes, Lamniformes, Squaliformes, Squatiniformes), and nine

chimaeras (Order Chimaeriformes). Only breeding residents of Europe were included in the

assessments, including 'visitor' species defined by the IUCN as "a taxon that does not reproduce

within a region but regularly occurs within its boundaries either now or during some period of

the last century" (IUCN, 2012a). The only visitors in Europe are currently the Smalltooth

Sawfish (*Pristis pristis*, Linnaeus 1758)*,* and Largetooth Sawfish (*Pristis pectinata*, Latham

1794). Vagrant species were not included in assessments, which by IUCN definition are "a taxon

that is currently found only occasionally within the boundaries of a region". Vagrant species

previously listed in Europe were listed as Not Applicable, and discounted from the following

analyses (e.g., the Nurse Shark, *Ginglymostoma cirratum*, Bonnaterre 1788; Nieto et al., 2015).

The IUCN Red List categories considered in this assessment are Least Concern, Near

Threatened, Vulnerable, Endangered, Critically Endangered, and Data Deficient, as there are no

sharks or rays known to be Regionally Extinct from the entire European region at present. This

ordering represents lowest to highest extinction risk, with the exception of Data Deficient, which

could include species that are both low and high risk.

**2.2 Developing an explanatory trait-based model for conservation status**

We considered three biological and ecological traits: maximum body size, median depth, and

reproductive mode (Dulvy & Forrest, 2010; Dulvy & Reynolds, 2002; Field et al., 2009; Rigby

& Simpfendorfer, 2015). Large maximum body size has been related to a greater likelihood of

decline and extinction risk due to higher catchability and slower population growth rates in

fishes, and other vertebrates (Dillingham et al., 2016; Field et al., 2009; Pardo et al., 2016).

Deeper depth ranges are associated with refuge from fishing activity, and hence, lower extinction

risk (Dulvy et al., 2014; Luiz et al., 2016). Overfishing is the greatest threat to sharks and rays

and occurs predominantly down to 400 m deep and exceptionally down to greater depths (Bailey

et al., 2009). Egg-laying (oviparous) species tend to be more fecund than live-bearing

(viviparous) species and hence may have greater maximum population growth rates, greater

variance in reproductive output, and hence scope for density-dependent compensation and lower

sensitivity to fishing mortality for adults (Dulvy & Forrest, 2010; Forrest et al., 2008).

There are inherent differences in biogeography, fisheries, and fisheries management between

Europe's major sub-regions, the Northeast Atlantic Ocean and Mediterranean Sea, which

warranted building models separately for each. There are 120 Northeast Atlantic sharks and rays

and 73 Mediterranean species, so lumping the two together as a Europe-wide status created a

bias towards Northeast Atlantic status. The IUCN categories were scored as Least Concern = 0,

Near Threatened = 1, Vulnerable = 2, Endangered = 3, and Critically Endangered = 4 (Butchart

et al., 2007). For each sub-regional model, IUCN category was the response variable and

maximum body size (cm, total length), median depth (m), reproductive mode (scored oviparous

= 1 or viviparous = 0). Median depth was used as a proxy for minimum depth and depth range to

account for exclusively shallow or deep species' distributions, while also avoiding having two

highly correlated fixed effects within a model. We also considered the interaction between size

and depth as a fixed effect. The interaction between size and depth is important because large-

bodied species are only associated with higher extinction risk if they exist within the reach of

fisheries (Dulvy et al., 2014). Size and depth were centred and scaled by two standard deviations.

Family was included as a random effect to account for phylogenetic covariation.

**2.3 Predicting conservation status**

Predictive accuracy of the explanatory model was evaluated using Area Under the Curve (AUC)

from Receiver Operating Characteristic curves (Sing et al., 2005). The AUC measure only works

for binary classification, so to test the predictive accuracy of each of the five categories

individually we scored each of the five IUCN categories separately as one, against all four other

categories scored as zero. We also grouped the threatened categories (Critically Endangered,

Endangered, Vulnerable) with a score of one, and non-threatened (Near Threatened, Least

Concern) scored as zero to determine the model accuracy for predicting threatened versus non-

threatened species. Predictive power was tested using data-sufficient species by dropping species

from the model to predict the conservation status and cross-validate against each known,

assessed conservation status. Test sets were run, comprising all data-sufficient species with all

species dropped one at a time. The model for each sub-region that was able to predict the correct

IUCN status with the highest AUC predictive accuracy measure was then used to predict the

categories of the actual Data Deficient species. The highest overall accuracy for a model was

determined by calculating the mean across all five AUC values for each IUCN category. Finally,

the IUCN categorisation for each Data Deficient species was classified using a 50% cut-off

point. All analyses were conducted in R version 3.5.2 (R Core Team, 2018), models were fit

using the clmm2 function from the ordinal package (Christensen, 2019), and performance was

evaluated with the ROCR package, version 1.0-7 (Sing et al., 2005).

**3 RESULTS**

**3.1 IUCN regional European Red List assessment**

One-fifth of the 120 Northeast Atlantic (22%, *n*=26) and 73 Mediterranean Sea (21%, *n*=15)

shark and ray species assessed in 2015 are listed as Data Deficient (Figure 1, Table 1). Most

species are assessed as Least Concern (38%) in the Northeast Atlantic, whereas the majority of

species are Critically Endangered (27%) in the Mediterranean Sea (Figure 1). Specifically, of the

data-sufficient Northeast Atlantic assessments, 38% (*n*=46) of species are Least Concern, 10%

(12) Near Threatened, 8% (9) Vulnerable, 13% (15) Endangered, and 10% (12) Critically

Endangered (Figure 1, Table 1). In the Mediterranean Sea, only 16% (12) species are Least

Concern, 11% (8) Near Threatened, 10% (7) Vulnerable, 15% (11) Endangered, and 27% (20)

Critically Endangered (Figure 1, Table 1). Sharks and rays are more threatened in Europe than

the global average (17.4%, *n*=181; Table 1). Specifically, nearly one-third (30%, *n*=36) are

threatened in the Northeast Atlantic and over half (52%, *n*=38) are threatened in the

Mediterranean Sea. Rays are approximately as threatened as sharks in both regions, in the

Northeast Atlantic 32% (*n*=14) of rays are threatened versus 33% (*n*=22) of sharks, and in the

Mediterranean Sea 50% (*n*=16) of rays are threatened versus 55% (*n*=22) of sharks (Table 1).

**3.2 Biological and ecological predictors of conservation status**

Large-bodied sharks and rays are more likely to be in higher categories of threat across Europe,

particularly in the Mediterranean Sea where threat levels are generally higher (Figure 2a,b).

When considering maximum body size in the Northeast Atlantic only, for every one unit increase

in maximum body size (i.e. cm total length), the odds of a species being in an IUCN category of

equal or higher threat increase by 0.98 (Figure 3a, Table S1). Similarly, in the Mediterranean

Sea, for every one unit increase in maximum body size, the odds of a species being in an IUCN

category of equal or higher threat increase by 0.94 (Figure 3a, Table S1). All other things being

equal, a shark or ray of three metres total length in the Northeast Atlantic has a 71.7%

probability of being in a threatened category (e.g. the Sandbar Shark, *Carcharhinus plumbeus*,

Nardo 1827) compared to a 1.5 m species, which has a 39.4% probability of the same (e.g. the

Angular Roughshark, *Oxynotus centrina*, Linnaeus 1758; Figure 2a). Whereas, in the

Mediterranean Sea the Sandbar Shark is 84.5% likely to be in a threatened category and the

Angular Roughshark is 62.1% likely to be threatened in this sub-region (Figure 2b). Hence, the

conservation status of a 1.5 m shark or ray in the Mediterranean Sea is closer to that of a three

metre species in the Northeast Atlantic, showing much less difference in likely conservation

status between similar sized species in the Mediterranean Sea.

Sharks and rays with greater depth distributions are more likely to be in lower categories of

threat in the Northeast Atlantic (Figure 2c), but this pattern is muted in the Mediterranean Sea

because threat levels are generally high for species across all depth distributions (Figure 2d).

When considering median depth in the Northeast Atlantic only, for every one unit increase in

median depth (i.e. metres), the odds of a species being in an IUCN category of higher threat

decrease by 0.04 (Figure 3c, Table S1). In the Mediterranean Sea, for every one unit increase in

median depth, the odds of a species being in a higher category of threat decrease by 0.24 (Figure

3c, Table S1). A similar-sized shark or ray with a median depth of 200 m has a 60% chance of

being threatened in the Northeast Atlantic (e.g. the Nursehound, *Scyliorhinus stellaris*, Linnaeus

1758), compared with a species with a median depth of 1,000 m (e.g. the Blackmouth Catshark,

*Galeus melastomus*, Rafinesque 1810), which has a 21.5% chance of being threatened in the

same sub-region (Figure 2c). In the Mediterranean Sea the difference in risk is muted because of

the greater reach of fisheries there: the Nursehound is 71.1% likely to be threatened, while the

Blackmouth Catshark is 40.6% likely to be threatened in this sub-region (Figure 2d). Again,

there is less differentiation between shallow and deepwater conservation status for

Mediterranean species than Northeast Atlantic, and a higher likelihood of being threatened

overall.

When maximum size, median depth, and reproductive mode are all considered, the odds of an

egg-laying (oviparous) species being in a higher threat category decrease by 0.14 in the

Northeast Atlantic (Figure 3b, Table S1). This effect was not significant in the Mediterranean

Sea, again because the trait sensitivity is overridden or muted by the higher degree of exposure to

fishing (Figure 3b, Table S1).

The most at-risk shark and ray species across Europe are therefore larger-bodied species

restricted to the most heavily fished 0–400 m depth zone. The interaction between size and depth

is such that for every unit increase in both size (cm) and depth (m), the odds of a shark or ray

being in a higher category of threat decrease by 0.02 in the Northeast Atlantic, and by 0.05 in the

Mediterranean Sea (Figure 3d, Table S1).

**3.3 Predicted versus evaluated conservation status**

The model with highest predictive accuracy (AUC) for both European sub-regions includes body

size, reproductive mode, and the interaction between size and depth (Table 2, Figure 3). For the

Northeast Atlantic, more than eight times out of ten, the top model predicts the correct category

for the Critically Endangered, Endangered, Least Concern, or grouped threatened categories

(Table S2). The Vulnerable category is predicted correctly more than six times out of ten, and the

Near Threatened less than four times out of ten. The top model for the Mediterranean Sea

predicts the Least Concern category correctly more than eight times out of then, and the

Critically Endangered and Near Threatened categories more than seven times out of ten. Both

sub-regional top models are weaker at predicting mid-range categories, with the Endangered

category predicted correctly less than five times and the Vulnerable category less than six times

out of ten (Table S2).

Almost half of Northeast Atlantic, and two-thirds of Mediterranean Data Deficient sharks and

rays are predicted-to-be-threatened with an elevated risk of extinction (Figure 4c). This

percentage of *predicted* threatened sharks and rays is greater than *evaluated* threat levels in the

Northeast Atlantic (46% *predicted* versus 38% *evaluated* threatened), and similar to *evaluated*

threatened in the Mediterranean Sea (67% *predicted* versus 66% *evaluated* threatened). The 12

Northeast Atlantic species predicted-to-be-threatened comprise 11 sharks and one ray (Figure

4a). All 12 Northeast Atlantic predicted-to-be-threatened species range from 89–640 cm total

length, with depth ranges overlapping with fishing activity, and are viviparous. The ten

predicted-to-be-threatened Mediterranean species comprise nine sharks and one ray (Figure 4b,

Table S3). All nine species range from 114–427 cm total length, overlap significantly with the

heavily fished depth zone, and are viviparous.

The distribution of both *evaluated* and *predicted* Northeast Atlantic listings in each case shows a

median categorisation of Near Threatened, whereas in the Mediterranean Sea the median

categorisation for both is Endangered (Figure S1). Overall, Northeast Atlantic and Mediterranean

Sea listings have opposing distributions, with the majority of Northeast Atlantic species non-

threatened and the majority of Mediterranean species threatened (Figure S1).

**3.4 Europe's most threatened Data Deficient sharks and rays**

The species predicted to have the most elevated extinction risk (i.e. Critically Endangered)

across Europe are all viviparous, large-bodied (349–640 cm total length) sharks whose median

depths range from 40.5–350 m, hence overlapping greatly with the heavily fished zone (0–400 m

depth; Table S3). In the Northeast Atlantic, the Great White Shark (*Carcharodon carcharias*,

Linnaeus 1758) and Great Hammerhead Shark (*Sphyrna mokarran*, Rüppell 1837) are predicted

to be Critically Endangered (Figure 4a; Table S3). While in the Mediterranean Sea, the Dusky

Shark (*Carcharhinus obscurus*, Lesueur 1818), Copper Shark (*Carcharhinus brachyurus*,

Günther 1870), and Longfin Mako (*Isurus paucus*, Guitart 1966) are predicted to be Critically

Endangered (Figure 4b; Table S3). With this categorical regression approach, we identify a total

of 14 Critically Endangered species in the Northeast Atlantic (approximately one third of

threatened) and 23 Critically Endangered species in the Mediterranean Sea (approximately one

half of threatened; Figure 5). If conservation efforts were focused on all imperilled species, i.e.

the combined *evaluated*-threatened and *predicted*-to-be-threatened species, there would be a

target list of 48 species to protect in the Northeast Atlantic and 48 species in the Mediterranean

Sea (Table 3, Figure 5).

**4 DISCUSSION**

**4.1 Regional versus global IUCN Red List Status**

Here, we show that sharks and rays are proportionally more threatened in the two main sub-

regions of Europe than the global reported threat rate, particularly when we account for the

predicted risk status of Data Deficient species. Overall, we estimate that there are 40% (48)

imperilled species (*evaluated* threatened and *predicted*-to-be-threatened) in the Northeast

Atlantic and 67% (48) in the Mediterranean Sea. Compared with other vertebrate groups, these

threat levels not only exceed those of global sharks and rays (23.9%, *n*=249 of 1,041, (Dulvy et

al., 2014), but also that of amphibians: the most imperilled assessed group to date (41%, *n*=2,561

of 6,284, Hoffmann et al., 2010). Furthermore, whereas almost half of global sharks and rays are

Data Deficient (46.8%, *n*=487), approximately one-fifth of Europe's species are Data Deficient,

which is also closer to global amphibian data deficiency proportionally (26%, *n*=1,597;

Hoffmann et al., 2010). The high levels of threat and relatively low levels of data deficiency in

Europe result from the region's comparably long-standing history of fishing and data collection

compared with the rest of the world (Barrett et al., 2004; Hoffmann, 1996). We next consider: (1)

the biological and ecological traits driving these regional threat levels, (2) the differences

between *evaluated* and *predicted* conservation status, and (3) how categorically predicting such

could help narrow the focus of conservation efforts overall.

**4.2 Biological and ecological predictors of conservation status**

Sharks and rays with both larger maximum body size and shallower depth distribution are more

likely to face an elevated risk of extinction than smaller (faster-growing) species that live

predominantly in deeper water. Fishing is the greatest threat to sharks and rays (McClenachan, et

al., 2012), and it is greatest from 0–400 m depth, but in European waters lower levels of fishing

activity occur down to at least 1,000 m (Amoroso et al., 2018; Morato et al., 2006). In waters

deeper than the reach of fisheries, a species can be very large-bodied and not threatened at all,

because body size has little influence over conservation status unless it is combined with a major

threat (Fernandes et al., 2017; Owens & Bennett, 2000; Reynolds et al., 2005a). For example, the

Goblin Shark (*Mitsukurina owstoni*, Jordan 1898) reaches 617 cm total length with a depth range

of 40–1,569 m. This deepwater shark is listed as Least Concern in the Northeast Atlantic as the

majority of its depth range offers refuge from fishing activity. By contrast, the Common

Thresher Shark (*Alopias vulpinus*, Bonnaterre 1788) reaches 573 cm total length, has a depth

range of 0–366 m (i.e. entirely overlapping with the heavily fished zone), and is listed as

Endangered in the same sub-region. Large body size has been associated with increased

probability of extinction for numerous taxonomic groups (e.g. Cardillo et al., 2011; Comeros-

Raynal et al., 2016; Field et al., 2009). Large body size is known to be correlated to a slow

speed-of-life, but also as an impediment to evading capture in fishing gear. For shark and ray

conservation, accounting for this relationship between size and susceptibility to capture is

complicated by the issue of bycatch. More sharks and rays are threatened by incidental catch

than by actual target fisheries (Dulvy et al., 2014), where their higher intrinsic sensitivity is not

accounted for by fisheries management regimes that are focused on faster growing, less sensitive

teleost (bony fish) species. Across Europe, the predominant fishing techniques are highly

unselective, such as multi-species trawling (Smith & Garcia, 2014). The incentive for fishers to

increase selectivity to benefit non-target species is low when this action would undoubtedly

coincide with reduced target catch. The consequent unselective fishing of non-target species is a

major driver of the high threat levels among Europe's sharks and rays and could lead to

overlooked local extinctions. This predicament alone presents incentive to better understand the

status of Data Deficient species, particularly in heavily fished waters such as Europe.

The conservation status of sharks and rays in the Mediterranean Sea appears much worse than

the Northeast Atlantic, which can partly be explained by the lack of depth refuge for sharks and

rays from heavy fishing activity in this sub-region. The Mediterranean Sea has a longer history

of fishing than the Northeast Atlantic, and nowadays, that fishing is not managed as efficiently as

it is in the Northeast Atlantic (Fernandes et al., 2017; Smith & Garcia, 2014). A semi-enclosed

sea equates to many more sites for landing catches, none of which are being consistently

monitored. Further exacerbating this lack of monitoring, the Mediterranean fishery principally

comprises higher numbers of smaller artisanal vessels, compared with fewer, more readily

trackable commercial vessels in the Northeast Atlantic (Smith & Garcia, 2014). Semi-enclosed

seas are also more susceptible to other major threatening events than open oceans, such as ocean

acidification, rising temperatures, and coastal pollution and development (Caddy, 2000). Despite

these logical contributors to the higher threat levels seen among Mediterranean sharks and rays,

the difference in conservation status between both major European sub-regions can largely be

attributed to the differing taxonomic and hence trait composition (120 Northeast Atlantic and 73

Mediterranean Sea sharks and rays). There are 35 deepwater shark and ray species that exist in

the Northeast Atlantic exclusively outside the reach of fisheries, and are all therefore listed as

Least Concern, which do not occur in the Mediterranean Sea. If those species are removed from

the Northeast Atlantic species list, we see the same number of imperilled species in each sub-

region, and a much more similar overall proportion of threat (Table S4). Meanwhile, the median

depths of all 73 Mediterranean Sea species overlap to some degree (if not entirely) with the

heavily fished 0–400 m depth zone. This explains why median depth, and the interaction

between maximum body size and median depth, are weaker explanatory variables in this sub-

region: depth refuge from fishing activity simply does not exist for sharks and rays in the

Mediterranean Sea.

Perhaps the prevailing threat resultant from lacking refuge in the Mediterranean Sea also

explains to some extent why oviparous species are not significantly lower-risk in this sub-region

than viviparous species. Oviparity is characteristically associated with faster population growth

rates than viviparity (Field et al., 2009). All of the most threatened sharks and rays in Europe are

viviparous, likely because they are less able to withstand fishing pressure as effectively as

typically faster-growing, egg-laying species.

**4.3 *Predicted* versus *evaluated* conservation status**

To provide taxon-wide estimates of extinction risk in the face of uncertainty, the IUCN assumes

that the fraction of threatened Data Deficient species is the same as the proportion of *evaluated-*

threatened species. While pragmatic, this is an assumption to be tested. A recent estimate of

which Data Deficient sharks and rays might be classified as Least Concern or threatened

revealed 14% of Data Deficient species were *predicted*-to-be-threatened ($n$=68 of 487), and

overall 17.8% were *evaluated*-threatened (Dulvy et al., 2014). Taken together, there is an overall

estimated global threat level of 23.9% imperilled sharks and rays (Dulvy et al., 2014). By

comparison, this is much lower than the IUCN equal ratio approach, which hence yields an

inflated estimate of 33% of sharks and rays threatened (Hoffmann et al., 2010).

This 1:1 ratio of *predicted*-to-*evaluated* threatened species proportions holds true for global birds

(Class Aves), in which knowledge is significantly greater than for other taxa and hence there are

few Data Deficient species (0.6%, $n$=63 of 10 500; Butchart & Bird, 2010). This 1:1 ratio

approach, however, yields a 50% underestimation of globally Data Deficient threatened

mammals, where one-third of *evaluated* species are threatened, whereas two-thirds of Data

Deficient species are *predicted*-to-be-threatened (1:2, Jetz & Freckleton, 2015). In the case of

mammals, reliance on this ratio could have devastating implications for species extinction rates

by overlooking 50% of the recently predicted-to-be-threatened Data Deficient species.

Conversely, using the 1:1 ratio of *evaluated* to *predicted* threatened status would overestimate

globally Data Deficient threatened groupers, for which the proportion of *evaluated* threat is

actually three times higher than that of *predicted* threat (3:1, Luiz et al., 2016).

Northeast Atlantic sharks and rays have a similar threat distribution, and hence, negative

conservation implications to global mammals if the IUCN's 1:1 ratio were to be relied upon

(1:1.2 *evaluated* to *predicted* threat). Global sharks and rays have the opposite pattern, whereby

conservation resources might be wasted on the protection of Data Deficient species according to

this ratio. Yet, despite the high levels of data deficiency among Mediterranean sharks and rays

compared with global birds, the present study shows this 1:1 ratio to be appropriate for this sub-

regional taxonomic group also. The inconsistency in these risk ratio patterns across taxonomic

groups, and geographic regions within taxonomic groups, highlights the need for taxon-specific

predictions of threat among Data Deficient listings.

**4.4 Updating protected species lists in Europe**

There are a number of lists that flag species for protection, but many of these are now out-of-

date. Ideally, all of the imperilled sharks and rays in Europe would be listed in the appendices of

the appropriate conservation-focused conventions in the region, and their exploitation monitored

and managed accordingly. In reality, only eight of the 48 imperilled species identified here are

listed on the Oslo-Paris convention in the Northeast Atlantic (Table 3). While of the 48

imperilled species identified here in the Mediterranean Sea, only three (and five) are listed on

Appendix II (and Appendix III) of the Berne Convention, nine on Appendix III of the Barcelona

Convention, and 23 on the General Fisheries Commission for the Mediterranean priority species

list (Table 3). The Great Hammerhead Shark is currently one of the 24 species included on the

GFCM priority species list for the Mediterranean Sea, but since the 2015 European Red List

reassessment this species is considered a Vagrant in this basin. We therefore consider only 23

species on the GFCM priority species list in this study (Table 3). Clearly, there is significant

scope to update these lists to ensure protection of all imperilled species in both sub-regions.

**4.5 Incorporating predictions into a Red List Index for 2020 target tracking**

Categorical predictions of IUCN status enable the inclusion of Data Deficient species in

aggregate species conservation status analyses, from which they are currently excluded for all

taxonomic groups. The Convention on Biological Diversity's Aichi Targets are monitored using

indicators, such as the IUCN's Red List Index, which is an indicator of the change in aggregate

extinction risk over time (Brooks et al., 2015). Predicting IUCN status for Data Deficient species

enables their addition to such indices, which would in turn give conservation planners a more

holistic idea of conservation status. With sufficient model accuracy, it is likely more informative

to include these predictions in such indices than to exclude them altogether. Upon completion of

the ongoing global reassessment of sharks and rays, this methodology can be extrapolated to the

global dataset for inclusion in the global Red List Index. This approach would prove even more

accurate for highly data-sufficient groups such as birds. Resource limitations have hindered

scientists and conservationists from focusing on Data Deficient species historically, but

categorical predictions of conservation status are a cost-effective solution to this shortcoming

(Bland et al., 2015), at least until data availability and resources allow for fully comprehensive

IUCN assessment of these species. This case study, and the extrapolation to the highly Data

Deficient global shark and ray dataset, will ideally be the first step towards applying this

predictive approach to some more Data Deficient groups, such as plants and invertebrates.

**ACKNOWLEDGEMENTS**

We thank the IUCN Species Survival Commission Shark Specialist Group members, staff, and

volunteers, and the IUCN head office for their logistical support throughout the Red List

assessment process. In particular, we thank Alvaro Abella, David Allen, Michel Bariche, Tom

Blasdale, Mohamed N Bradaï, Elena Buscher, Maurice Clarke, Simona Clo, Andrey Dolgov,

Manuel Dureuil, Jim Ellis, Edward Farrell, Francesco Ferretti, Sonia Fordham, Sarah Fowler,

Karen Frazer, Mariana García, Javier Guallart, Lucy Harrison, Ali Hood, Samuel Iglésias,

Armelle Jung, James Kemp, Pete Kyne, Julia Lawson, Sophy McCully, Ana Nieto, Giuseppe

Notarbartolo di Sciara, Caroline Pollock, Gina Ralph, Fabrizio Serena, Bernard Seret, Alen

Soldo, Matthias Stehmann, Heike Zidowitz. We thank Tobias Sing, Dan Greenberg, Nathan

Pacoureau, Chris Brown, and Rylee Murray for statistical advice, and members of the Dulvy

Lab, Earth$_2$Ocean Lab and Stats Beerz for comment on drafts and statistical advice.

**Funding**: The European Red List of marine fishes was a project funded by the European

Commission (Directorate General for the Environment under Service Contract No.

070307/2011/607526/SER/B.3). This project was funded by the Shark Conservation Fund as part

of the Global Shark Trends Project. Rachel Walls and Nicholas Dulvy were supported by a

Natural Science and Engineering Research Council Discovery and Accelerator Award and

Nicholas Dulvy was supported by the Canada Research Chairs Program. Rachel Walls was also

supported by Simon Fraser University with three Graduate Fellowship awards.

**REFERENCES**

Amoroso, R. O., Pitcher, C. R., Rijnsdorp, A. D., McConnaughey, R. A., Parma, A. M.,

Suuronen, P., … Jennings, S. (2018). Bottom trawl fishing footprints on the world's

continental shelves. *Proceedings of the National Academy of Sciences*, *115*(43), E10275–

E10282. http://doi.org/10.1073/pnas.1802379115

Bailey, D. M., Collins, M. A., Gordon, J. D. M., Zuur, A. F., & Priede, I. G. (2009). Long-term

changes in deep-water fish populations in the northeast Atlantic: A deeper reaching effect of

fisheries? *Proceedings of the Royal Society B: Biological Sciences*, *276*(1664), 1965–1969.

http://doi.org/10.1098/rspb.2009.0098

Barrett, J. H., Locker, A. M., & Roberts, C. M. (2004). The origins of intensive marine fishing in

medieval Europe: the English evidence. *Proceedings of the Royal Society B: Biological*

*Sciences*, *271*, 2417–2421. http://doi.org/10.1098/rspb.2004.2885

Bland, L. M., Bielby, J., Kearney, S., Orme, C. D. L., Watson, J. E. M., & Collen, B. (2017).

Toward reassessing data-deficient species. *Conservation Biology*, *31*(3), 531–539.

http://doi.org/10.1111/cobi.12850

Bland, L. M., & Böhm, M. (2016). Overcoming data deficiency in reptiles. *Biological*

*Conservation*, *204*, 16–22. http://doi.org/10.1016/j.biocon.2016.05.018

Bland, L. M., Collen, B., Orme, C. D. L., & Bielby, J. (2012). Data uncertainty and the

selectivity of extinction risk in freshwater invertebrates. *Diversity and Distributions*, *18*(12),

1211–1220. http://doi.org/10.1111/j.1472-4642.2012.00914.x

Bland, L. M., Collen, B., Orme, C. D. L., & Bielby, J. (2014). Predicting the conservation status

of data-deficient species. *Conservation Biology*, *29*(1), 250–259.

http://doi.org/10.1111/cobi.12372

Bland, L. M., Collen, B., Orme, C. D. L., & Bielby, J. (2015). Predicting the conservation status

of data-deficient species. *Conservation Biology*, *29*(1), 250–259.

http://doi.org/10.1111/cobi.12372

Bland, L. M., Orme, C. D. L., Bielby, J., Collen, B., Nicholson, E., & Mccarthy, M. A. (2015).

Cost-effective assessment of extinction risk with limited information. *Journal of Applied*

*Ecology*, *52*(4), 861–870. http://doi.org/10.1111/1365-2664.12459

Böhm, M., Collen, B., Baillie, J. E. M., Bowles, P., Chanson, J., Cox, N., … Zug, G. (2013). The

conservation status of the world's reptiles. *Biological Conservation*, *157*, 372–385.

http://doi.org/10.1016/j.biocon.2012.07.015

Brooks, T. M., Butchart, S. H. M., Cox, N. A., Heath, M., Hilton-Taylor, C., Hoffmann, M., …

Smart, J. (2015). Harnessing biodiversity and conservation knowledge products to track the

Aichi Targets and Sustainable Development Goals. *Biodiversity*, *16*(2–3), 157–174.

http://doi.org/10.1080/14888386.2015.1075903

Butchart, S. H. M., Akçakaya, H. R., Chanson, J., Baillie, J. E. M., Collen, B., Quader, S., …

Hilton-Taylor, C. (2007). Improvements to the Red List Index. *PLoS ONE*, *2*(1).

http://doi.org/10.1371/journal.pone.0000140

Butchart, S. H. M., & Bird, J. P. (2010). Data Deficient birds on the IUCN Red List: What don't

we know and why does it matter? *Biological Conservation*, *143*(1), 239–247.

http://doi.org/10.1016/j.biocon.2009.10.008

Caddy, J. (2000). Marine catchment basin effects versus impacts of fisheries on semi-enclosed

seas. *ICES Journal of Marine Science*, *57*(3), 628–640.

http://doi.org/10.1006/jmsc.2000.0739

Callmander, M. W., Schatz, G. E., & Lowry, P. P. (2005). IUCN Red List assessment and the

Global Strategy for Plant Conservation: Taxonomists must act now. *Taxon*, *54*(4), 1047–

1050. http://doi.org/10.2307/25065491

Cardillo, M., Mace, G., Jones, K. E., Bielby, J., Bininda-Emonds, O. R. P., Sechrest, W., …

Purvis, A. (2005). Multiple causes of high extinction risk in large mammal species. *Science*,

*309*(5738), 1239–1241.

Cardillo, M., Mace, G. M., Jones, K. E., Bielby, J., Bininda-emonds, O. R. P., & Sechrest, W.

(2011). Multiple Causes of High Extinction Risk in Large Mammal Species. *Science*,

*1239*(2005), 1239–1241. http://doi.org/10.1126/science.1116030

CBD, & UNEP. (2011). The Strategic Plan for Biodiversity 2011–2020 and the Aichi Targets

"Living in Harmony with Nature." *A Ten-Year Framework for Action by All Countries and*

*Stakeholders to Save Biodiversity and Enhance Its Benefits for People. Secretariat of the*

*Convention*, 2. Retrieved from www.cbd.int

Christensen, R. H. B. (2019). Package 'ordinal.' Retrieved from https://cran.r-

project.org/web/packages/ordinal/ordinal.pdf

Collen, B., Böhm, M., Kemp, R., & Baillie, J. E. M. (2012). *Spineless: status and trends of the*

*world's invertebrates*. London, United Kingdom.

Comeros-Raynal, M. T., Polidoro, B. A., Broatch, J., Mann, B. Q., Gorman, C., Buxton, C. D.,

… Carpenter, K. E. (2016). Key predictors of extinction risk in sea breams and porgies

(Family: Sparidae). *Biological Conservation*, *202*, 88–98.

http://doi.org/10.1016/j.biocon.2016.08.027

Cortés, E. (2000). Life history patterns and correlations in sharks. *Reviews in Fisheries Science*,

*8*(4), 299–344.

Dillingham, P. W., Moore, J. E., Fletcher, D., Cortés, E., Alexandra Curtis, K., James, K. C., &

Lewison, R. L. (2016). Improved estimation of intrinsic growth rmax for long-lived species:

Integrating matrix models and allometry. *Ecological Applications*, *26*(1), 322–333.

http://doi.org/10.1890/14-1990

Dulvy, N., Allen, D., Ralph, G., & Walls, R. (2016). The conservation status of Sharks, Rays and

in the Mediterranean Sea., 6+6 pp.

Dulvy, N., & Forrest, R. (2010). Life histories, population dynamics, and extinction risks in

chondrichthyans. In J. Carrier, J. Musick, & M. Heithaus (Eds.), *Sharks and Their Relatives*

*II: Biodiversity, Adaptive Physiology, and Conservation* (pp. 635–676). CRC Press: Boca

Raton.

Dulvy, N. K., & Forrest, R. E. (2010). Biodiversity, adapative physiology, and conservation. In J.

C. Carrier, J. A. Musick, & M. R. Heithaus (Eds.), *Sharks and their relatives II* (pp. 640–

670). Boca Raton, FL: CRC Press.

Dulvy, N. K., Fowler, S. L., Musick, J. A., Cavanagh, R. D., Kyne, P. M., Harrison, L. R., …

White, W. T. (2014). Extinction risk and conservation of the world's sharks and rays. *ELife*,

*3*. http://doi.org/10.7554/eLife.00590

Dulvy, N. K., & Reynolds, J. D. (2002). Predicting extinction vulnerability in skates.

*Conservation Biology*, *16*(2), 440–450. http://doi.org/10.1046/j.1523-1739.2002.00416.x

Fernandes, P. G., Ralph, G. M., Nieto, A., García Criado, M., Vasilakopoulos, P., Maravelias, C.

D., … Carpenter, K. E. (2017). Coherent assessments of Europe's marine fishes show

regional divergence and megafauna loss. *Nature Ecology & Evolution*, *1*(May), 0170.

http://doi.org/10.1038/s41559-017-0170

Field IC, Meekan M, Buckworth RC, & Bradshaw CJA. (2009). *Susceptibility of Sharks, Rays*

*and Chimeras to Global Extinction. Advances in Marine Biology* (1st ed., Vol. 56). Elsevier

Ltd. http://doi.org/10.1016/S0065-2881(09)56004-X

Forrest, R. E., Martell, S. J. D., Melnychuk, M. C., & Walters, C. J. (2008). An age-structured

model with leading management parameters, incorporating age-specific selectivity and

maturity. *Canadian Journal of Fisheries and Aquatic Sciences*, *65*(2), 286–296.

http://doi.org/10.1139/f07-156

Hoffmann, M., Hilton-taylor, C., Angulo, A., Böhm, M., Brooks, T. M., Butchart, S. H. M., …

Russell, A. (2010). The Impact of Conservation on the Status of the World's Vertebrates.

*Science*, *330*(6010), 1503–1509. http://doi.org/10.1126/science.1194442

Hoffmann, R. C. (1996). Economic Development and Aquatic Ecosystems in Medieval Europe.

*American Historical Review*, *101*, 631–669. http://doi.org/10.4324/9781315252094-5

Iglésias, S. P., Toulhoat, L., & Sellos, D. Y. (2010). Taxonomic confusion and market

mislabelling of threatened skates: important consequences for their conservation status.

*Aquatic Conservation: Marine and Freshwater Ecosystems*, *20*(3), 319–333.

http://doi.org/10.1002/aqc.1083

IUCN. (2012a). *Guidelines for application of IUCN Red List criteria at regional and national*

*levels: Version 4.0* (4th ed.). Gland, Switzerland and Cambridge, UK: IUCN, Gland,

Switzerland. Retrieved from www.iucn.org/publications

IUCN. (2012b). *IUCN Red List Categories and Criteria: version 3.1*. (IUCN, Ed.) (Second).

Gland, Switzerland and Cambridge, UK: IUCN, Gland, Switzerland.

http://doi.org/https://www.iucnredlist.org/static/categories_criteria_3_1

Jetz, W., & Freckleton, R. P. (2015). Towards a general framework for predicting threat status of

data-deficient species from phylogenetic, spatial and environmental information.

*Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*,

*370*(1662), 20140016. http://doi.org/10.1098/rstb.2014.0016

Luiz, O. J., Woods, R. M., Madin, E. M. P., & Madin, J. S. (2016). Predicting IUCN Extinction

Risk Categories for the World's Data Deficient Groupers (Teleostei: Epinephelidae).

*Conservation Letters*, *9*(5), 342–350. http://doi.org/10.1111/conl.12230

McClenachan, L., Cooper, A. B., Carpenter, K. E., & Dulvy, N. K. (2012). Extinction risk and

bottlenecks in the conservation of charismatic marine species. *Conservation Letters*, *5*(1),

73–80. http://doi.org/10.1111/j.1755-263X.2011.00206.x

McKinney, M. L. (1997). Extinction vulnerability and selectivity: Combining ecological and

paleontological views. *Annual Review of Ecology and Systematics*, *28*(1), 495–516.

http://doi.org/10.1146/annurev.ecolsys.28.1.495

Morato, T., Watson, R., Pitcher, T. J., & Pauly, D. (2006). Fishing down the deep, 23–33.

Nieto,  a., Ralph, G. M., Comeros-Raynal, M. T., Kemp, J., García Criado, M., Allen, D. J., …

Williams, J. T. (2015). European Red List of Marine Fishes, 88.

http://doi.org/10.2779/082723

Owens, I. P. F., & Bennett, P. M. (2000). Ecological basis of extinction risk in birds : Habitat

loss versus human persecution and introduced predators. *PNAS*, *97*(22), 12144–12148.

Pardo, S. A., Kindsvater, H. K., Cuevas-Zimbrón, E., Sosa-Nishizaki, O., Pérez-Jiménez, J. C.,

& Dulvy, N. K. (2016). Growth, productivity, and relative extinction risk of a data-sparse

devil ray. *Scientific Reports*, *6*, 1–10. http://doi.org/10.1038/srep33745

Randhawa, H. S., Poulin, R., & Krkošek, M. (2015). Increasing rate of species discovery in

sharks coincides with sharp population declines: Implications for biodiversity. *Ecography*,

*38*(1), 96–107. http://doi.org/10.1111/ecog.00793

Reynolds, J. D., Webb, T. J., & Hawkins, L. A. (2005a). Life history and ecological correlates of

extinction risk in European freshwater fishes. *Canadian Journal of Fisheries and Aquatic*

*Sciences*, *62*(4), 854–862. http://doi.org/10.1139/f05-066

Reynolds, J. D., Webb, T. J., & Hawkins, L. A. (2005b). Life history and ecological correlates of

extinction risk in European freshwater fishes 1, *862*(August 2003), 854–862.

http://doi.org/10.1139/F05-066

Rigby, C., & Simpfendorfer, C. A. (2015). Patterns in life history traits of deep-water

chondrichthyans. *Deep-Sea Research Part II: Topical Studies in Oceanography*, *115*, 30–

40. http://doi.org/10.1016/j.dsr2.2013.09.004

Sing, T., Sander, O., Beerenwinkel, N., & Lengauer, T. (2005). ROCR: Visualizing classifier

performance in R. *Bioinformatics*, *21*(20), 3940–3941.

http://doi.org/10.1093/bioinformatics/bti623

Smith, A. D. M., & Garcia, S. M. (2014). Fishery management: Contrasts in the mediterranean

and the atlantic. *Current Biology*, *24*(17), R810–R812.

http://doi.org/10.1016/j.cub.2014.07.031

Stein, R. W., Mull, C. G., Kuhn, T. S., Aschliman, N. C., Davidson, L. N. K., Joy, J. B., …

Mooers, A. O. (2018). Global priorities for conserving the evolutionary history of sharks,

rays and chimaeras. *Nature Ecology and Evolution*, *2*(February), 1–11.

http://doi.org/10.1038/s41559-017-0448-4

White, W. T., & Last, P. R. (2012). A review of the taxonomy of chondrichthyan fishes: a

modern perspective. *Journal of Fish Biology*, *80*(5), 901–917. http://doi.org/10.1111/j.1095-

8649.2011.03192.x

 **TABLES LEGENDS**

 **Table 1**

**Global and European IUCN Red Listings of sharks and rays.** Observed number and (percent) of global (2014), Northeast Atlantic (2015), and Mediterranean Sea (2015) sharks, rays (i.e. all rays and skates), and chimaeras in each IUCN Red List category. CR: Critically Endangered, EN: Endangered, VU: Vulnerable, thr: threatened (CR+EN+VU), NT: Near Threatened, LC: Least Concern, DD: Data Deficient (*Dulvy et al., 2014).

**Table 2**

**Summary of top Cumulative Link Mixed-effects Models for predicting IUCN status of Northeast Atlantic and Mediterranean sharks and rays.** Models included all evaluated species (*n*=94 Northeast Atlantic and *n*=58 Mediterranean Sea). Top predictive models for both sub-regions with ΔAIC <2 included maximum size (cm), reproductive mode (oviparous=1, viviparous=0), and the interaction between maximum size and median depth (m). Maximum size and median depth were centred and standardised by two standard deviations. Each species was dropped one-at-a-time from the model and the IUCN status predicted. Comparison between evaluated and predicted statuses determined the predictive accuracy of each model. Model accuracy was measured as the Area Under the Curve (AUC) from the ROCR package in R version 3.5.2 (Sing et al., 2005) by scoring each category as one and all four other categories as zero to determine the predictive accuracy of all five separately (Critically Endangered, Endangered, Vulnerable, Near Threatened, Least Concern). To determine the top predictive model overall, the mean of all five category AUC values was calculated. Left to right: Loglik = log likelihood, $AIC_c$ = AIC corrected for small sample size, ΔAIC = delta AIC, AIC wt = AIC

weight, mean AUC = Area Under Curve averaged across the five AUC values for each IUCN

category (see Table S2 for complete list of AUC values).

**Table 3**

**Current consideration of all imperilled sharks and rays in European waters by regional**

**conventions and priority species lists.** Relevant listings of all imperilled (i.e. *evaluated-*

threatened and *predicted*-to-be-threatened) sharks and rays in Europe on regional and global

protection-focused conventions, by major sub-region (Northeast Atlantic then Mediterranean

Sea). Blanks indicate no listing, while hyphens indicate inapplicability of a convention to a

species within a certain sub-region. Where a convention has multiple appendices, the applicable

appendix number is indicated (e.g. A2, A3) instead of a tick mark. Species are listed

taxonomically within each threatened IUCN category – Critically Endangered (CR), Endangered

(EN), and Vulnerable (VU) – in descending order of threat. Conventions left to right: Oslo-Paris

Convention (OSPAR; applicable to Northeast Atlantic Ocean only); Berne Convention

(applicable to Mediterranean Sea only); Barcelona Convention (Mediterranean Sea only); and

the General Fisheries Commission for the Mediterranean (GFCM) priority species list

(Mediterranean Sea only). The Great Hammerhead Shark (*Sphyrna mokarran*) is currently one of

24 species included on the GFCM priority species list, but has not been included in the

Mediterranean section of this table as it is now considered a Vagrant species in the

Mediterranean Sea, as per IUCN definition (IUCN, 2012a).

**FIGURE LEGENDS**

**Figure 1**

**Percent *evaluated* and *predicted* IUCN categorisations of Europe's sharks and rays.** Dark bars represent the percentage of species officially evaluated on the IUCN Red List, while light bars represent the percentage of Data Deficient species predicted to be under each category as per the results of the present study. Of the 120 species in the Northeast Atlantic, 94 were *evaluated* and 26 were Data Deficient and *predicted* for. In the Mediterranean Sea, 58 of 73 species were *evaluated* and 15 were Data Deficient and *predicted* for. The IUCN categories from highest to lowest threat are: CR = Critically Endangered, EN = Endangered, VU = Vulnerable, NT = Near Threatened, and LC = Least Concern.

**Figure 2**

**The effects of size and depth on shark and ray conservation status in Europe.** Histograms of the probability of an evaluated shark or ray being listed as either Critically Endangered (CR), Endangered (EN), Vulnerable (VU), Near Threatened (NT), or Least Concern (LC) based on single-trait Cumulative Link Mixed-effects Model outputs for maximum body size (cm; panels a and b) and median depth (m; panels c and d). Data include all evaluated species (*n*= 94 Northeast Atlantic, panels a and c; and *n*=58 Mediterranean Sea, panels b and d) and exclude all Data Deficient species. Dark grey vertical bars indicate large (300 cm total length, a,b) or shallow (200 m median depth, c,d) species; light grey bars represent small (150 cm total length, a,b) or deep (1,000 m median depth, c,d) species. Brackets beside bars indicate the probability of each species being categorised as threatened (CR, EN, or VU) on the IUCN Red List.

**Figure 3**

**Effects of biological and ecological traits on Europe's shark and ray conservation status.**

Standardized effect sizes with 95% confidence intervals. Cumulative link mixed effect models

with maximum body size (a), reproductive mode (b), median depth (c), and the interaction

between size and depth (d) as fixed effects and taxonomic Family as a random effect to account

for phylogenetic non-independence. Circular and triangular points represent the best explanatory

and predictive model for the Northeast Atlantic and Mediterranean Sea, respectively, which in

both cases included maximum body size, reproductive mode, and the interaction between

maximum size and median depth. Data for maximum size and median depth were centred and

standardised by two standard deviations, while reproductive mode is a binary trait where

oviparous species = 1 and viviparous species = 0.

**Figure 4**

*Predicted* **and** *evaluated* **conservation status of Europe's sharks and rays.** Top Cumulative

Link Mixed-effectd Models including maximum body size, reproductive mode, and the

interaction between size and median depth as fixed effects (for both sub-regions) and taxonomic

Family as a random effect to account for phylogenetic non-independence. Panel a shows the

probability of all 26 Data Deficient Northeast Atlantic species, and panel b of all 15

Mediterranean Sea species, being in each IUCN Red List category based on these top

explanatory models. The vertical line cutting down panels a and b represents the 50% cut-off

classification used to assign the final IUCN categorisations (according to the category bar the

line crosses). Data for size and depth were centred and standardised by two standard deviations,

while reproductive mode is a binary trait where oviparous species = 1 and viviparous species = 0.

Panel c shows the proportion of sharks and rays in the Northeast Atlantic (left) and

Mediterranean Sea (right) both *evaluated* and *predicted* to be in each IUCN category. Percentage

values within each yellow bar indicate the total percentage of *evaluated* threatened and

*predicted*-to-be-threatened species in each set, while numbers within brackets below each bar

indicate the total number of species included in each set

**Figure 5**

**Informing shark and ray conservation efforts in Europe with categorical predictions.** Solid

grey bars represent species of all IUCN categories excluding those officially evaluated by the

IUCN as Critically Endangered (CR), which are represented by solid red blocks. There are 120

shark and ray species in the Northeast Atlantic (left) and 73 in the Mediterranean Sea (right).

Horizontal red lines indicate the addition of all Data Deficient species *predicted* to be Critically

Endangered, to the *evaluated* block. Orange lines indicate all *evaluated* and *predicted*-to-be-

Endangered and Critically Endangered species, while yellow lines show all imperilled (i.e.

*evaluated* and *predicted*-to-be-threatened) species (Vulnerable, Endangered, and Critically

Endangered). Numbers beside bars indicate total number of species within each relevant

grouping.

**Table 1**

**Global and European IUCN Red Listings of sharks and rays.** Observed number and (percent) of global (2014), Northeast Atlantic

(2015), and Mediterranean Sea (2015) sharks, rays (i.e. all rays and skates), and chimaeras in each IUCN Red List category. CR:

Critically Endangered, EN: Endangered, VU: Vulnerable, thr: threatened (CR+EN+VU), NT: Near Threatened, LC: Least Concern,

DD: Data Deficient (*Dulvy et al., 2014).

| Geographic scope | Total species | Total thr species (%) | Total CR (%) | Total EN (%) | Total VU (%) | Total NT (%) | Total LC (%) | Total DD (%) |
|---|---|---|---|---|---|---|---|---|
| All global* | 1,041 | 181(17) | 25(2) | 43(4) | 113(11) | 132(13) | 241(23) | 487(47) |
| All Northeast Atlantic | 120 | 38(32) | 12(10) | 15(13) | 11(9) | 12(10) | 48(40) | 22(18) |
| All Mediterranean Sea | 73 | 39(53) | 20(27) | 11(15) | 8(11) | 9(12) | 12(17) | 13(18) |
| Rays (Northeast Atlantic) | 44 | 14(32) | 6(14) | 3(7) | 5(11) | 6(14) | 22(50) | 2(4) |
| Sharks (Northeast Atlantic) | 67 | 22(33) | 6(9) | 12(18) | 4(6) | 5(7) | 16(24) | 24(36) |
| Chimaeras (Northeast Atlantic) | 9 | - | - | - | - | 1(10) | 8(90) | - |
| Rays (Mediterranean) | 32 | 16(50) | 8(25) | 5(16) | 3(9) | 5(16) | 7(22) | 4(12) |
| Sharks (Mediterranean) | 40 | 22(55) | 12(30) | 6(15) | 4(10) | 2(5) | 5(12.5) | 11(27.5) |
| Chimaeras (Mediterranean) | 1 | - | - | - | - | 1(100) | - | - |

**Table 2**

**Summary of top Cumulative Link Mixed-effects Models for predicting IUCN status of Northeast Atlantic and Mediterranean**

**sharks and rays.** Models included all evaluated species (*n*=94 Northeast Atlantic and *n*=58 Mediterranean Sea). Top predictive

models for both sub-regions with ΔAIC <2 included maximum size (cm), reproductive mode (oviparous=1, viviparous=0), and the

interaction between maximum size and median depth (m). Maximum size and median depth were centred and standardised by two

standard deviations. Each species was dropped one-at-a-time from the model and the IUCN status predicted. Comparison between

evaluated and predicted statuses determined the predictive accuracy of each model. Model accuracy was measured as the Area Under

the Curve (AUC) from the ROCR package in R version 3.5.2 (Sing et al., 2005) by scoring each category as one and all four other

categories as zero to determine the predictive accuracy of all five separately (Critically Endangered, Endangered, Vulnerable, Near

Threatened, Least Concern). To determine the top predictive model overall, the mean of all five category AUC values was calculated.

Left to right: Loglik = log likelihood, AIC$_c$ = AIC corrected for small sample size, ΔAIC = delta AIC, AIC wt = AIC weight, mean

AUC = Area Under Curve averaged across the five AUC values for each IUCN category (see Table S2 for complete list of AUC

values).

| Region | Model hypothesis | Loglik | AIC$_c$ | ΔAIC | AIC wt | Mean AUC |
|---|---|---|---|---|---|---|
| **Northeast Atlantic** | IUCN status ~ Max size + Reproduction + Max size*Med depth | -91.553 | 201.106 | 5.684e-14 | 0.122 | 0.711 |
| **Mediterranean** | IUCN status ~ | -71.528 | 161.055 | 1.833 | 0.066 | 0.657 |

| Sea | Max size + Reproduction + Max size*Med depth |
|---|---|

**Table 3**

**Current consideration of all imperilled sharks and rays in European waters by regional conventions and priority species lists.**

Relevant listings of all imperilled (i.e. *evaluated*-threatened and *predicted*-to-be-threatened) sharks and rays in Europe on regional and

global protection-focused conventions, by major sub-region (Northeast Atlantic then Mediterranean Sea). Blanks indicate no listing,

while hyphens indicate inapplicability of a convention to a species within a certain sub-region. Where a convention has multiple

appendices, the applicable appendix number is indicated (e.g. A2, A3) instead of a tick mark. Species are listed taxonomically within

each threatened IUCN category – Critically Endangered (CR), Endangered (EN), and Vulnerable (VU) – in descending order of threat.

Conventions left to right: Oslo-Paris Convention (OSPAR; applicable to Northeast Atlantic Ocean only); Berne Convention

(applicable to Mediterranean Sea only); Barcelona Convention (Mediterranean Sea only); and the General Fisheries Commission for

the Mediterranean (GFCM) priority species list (Mediterranean Sea only). The Great Hammerhead Shark (*Sphyrna mokarran*) is

currently one of 24 species included on the GFCM priority species list, but has not been included in the Mediterranean section of this

table as it is now considered a Vagrant species in the Mediterranean Sea, as per IUCN definition (IUCN, 2012a).

| Species | Evaluated IUCN status | Predicted IUCN status | OSPAR (NEA) | Berne Convention (Med) | Barcelona Convention (Med) | GFCM priority species (Med) |
|---|---|---|---|---|---|---|
| *Squatina squatina* | Critically Endangered | - | ⍰ | - | - | - |
| *Squatina aculeata* | Critically Endangered | - | | - | - | - |
| *Squatina oculata* | Critically Endangered | - | | - | - | - |
| *Centrophorus granulosus* | Critically Endangered | - | ⍰ | - | - | - |

| Species | Evaluated IUCN status | Predicted IUCN status | OSPAR (NEA) | Bern Convention (Med) | Barcelona Convention (Med) | GFCM priority species (Med) |
|---|---|---|---|---|---|---|
| *Odontaspis ferox* | Critically Endangered | - | | - | - | - |
| *Lamna nasus* | Critically Endangered | - | ▯ | - | - | - |
| *Carcharodon carcharias* | Data Deficient | Critically Endangered | - | - | - | - |
| *Sphyrna mokarran* | Data Deficient | Critically Endangered | - | - | - | - |
| *Rostroraja alba* | Critically Endangered | - | ▯ | - | - | - |
| *Dipturus batis* | Critically Endangered | - | ▯ | - | - | - |
| *Pristis pristis* | Critically Endangered | - | - | - | - | - |
| *Pristis pectinata* | Critically Endangered | - | - | - | - | - |
| *Gymnura altavela* | Critically Endangered | - | - | - | - | - |
| *Pteromylaeus bovinus* | Critically Endangered | - | - | - | - | - |
| *Echinorhinus brucus* | Endangered | - | | - | - | - |
| *Centrophorus lusitanicus* | Endangered | - | | - | - | - |
| *Centrophorus squamosus* | Endangered | - | ▯ | - | - | - |
| *Deania calcea* | Endangered | - | | - | - | - |
| *Centroscymnus coelolepis* | Endangered | - | | - | - | - |
| *Dalatias licha* | Endangered | - | | - | - | - |
| *Squalus acanthias* | Endangered | - | ▯ | - | - | - |
| *Alopias vulpinus* | Endangered | - | | - | - | - |
| *Alopias superciliosus* | Endangered | - | | - | - | - |
| *Cetorhinus maximus* | Endangered | - | ▯ | - | - | - |
| *Isurus oxyrinchus* | Data Deficient | Endangered | | - | - | - |
| *Isurus paucus* | Data Deficient | Endangered | | - | - | - |
| *Galeocerdo cuvier* | Data Deficient | Endangered | | - | - | - |
| *Sphyrna zygaena* | Data Deficient | Endangered | | - | - | - |
| *Sphyrna lewini* | Data Deficient | Endangered | | - | - | - |
| *Carcharhinus plumbeus* | Endangered | - | | - | - | - |
| *Carcharhinus longimanus* | Endangered | - | | - | - | - |
| *Carcharhinus obscurus* | Data Deficient | Endangered | | - | - | - |

| Species | Evaluated IUCN status | Predicted IUCN status | OSPAR (NEA) | Bern Convention (Med) | Barcelona Convention (Med) | GFCM priority species (Med) |
|---|---|---|---|---|---|---|
| *Carcharhinus falciformis* | Data Deficient | Endangered | - | - | - | |
| *Glaucostegus cemiculus* | Endangered | - | - | - | - | |
| *Rhinobatos rhinobatos* | Endangered | - | - | - | - | |

| Species | Evaluated IUCN status | Predicted IUCN status | OSPAR (NEA) | Bern Convention (Med) | Barcelona Convention (Med) | GFCM priority species (Med) |
|---|---|---|---|---|---|---|
| *Mobula mobular* | Endangered | - | | - | - | - |
| *Hexanchus nakamurai* | Data Deficient | Vulnerable | | - | - | - |
| *Centrophorus uyato* | Vulnerable | - | | - | - | - |
| *Oxynotus centrina* | Vulnerable | - | | - | - | - |
| *Pseudotriakis microdon* | Data Deficient | Vulnerable | | - | - | - |
| *Galeorhinus galeus* | Vulnerable | - | | - | - | - |
| *Mustelus mustelus* | Vulnerable | - | | - | - | - |
| *Leucoraja circularis* | Vulnerable | - | | - | - | - |
| *Leucoraja fullonica* | Vulnerable | - | | - | - | - |
| *Rhinoptera marginata* | Data Deficient | Vulnerable | | - | - | - |
| *Myliobatis aquila* | Vulnerable | - | | - | - | - |
| *Dasyatis pastinaca* | Vulnerable | - | | - | - | - |
| *Dasyatis centroura* | Vulnerable | - | | - | - | - |
| *Squatina squatina* | Critically Endangered | - | - | A3 | | ⬚ |
| *Squatina aculeata* | Critically Endangered | - | - | | | ⬚ |
| *Squatina oculata* | Critically Endangered | - | - | | | ⬚ |
| *Centrophorus granulosus* | Critically Endangered | - | - | | A3 | |
| *Oxynotus centrina* | Critically Endangered | - | - | | | ⬚ |
| *Odontaspis ferox* | Critically Endangered | - | - | | | ⬚ |
| *Carcharias taurus* | Critically Endangered | - | - | | | ⬚ |
| *Lamna nasus* | Critically Endangered | - | - | A3 | | ⬚ |
| *Carcharodon carcharias* | Critically Endangered | - | - | A2 | | ⬚ |
| *Isurus oxyrinchus* | Critically Endangered | - | - | A3 | | ⬚ |
| *Isurus paucus* | Data Deficient | Critically Endangered | - | | | |
| *Sphyrna zygaena* | Critically Endangered | - | - | | | ⬚ |

| Species | Evaluated IUCN status | Predicted IUCN status | OSPAR (NEA) | Bern Convention (Med) | Barcelona Convention (Med) | GFCM priority species (Med) |
|---|---|---|---|---|---|---|
| *Carcharhinus brachyurus* | Data Deficient | Critically Endangered | - | | | |
| *Carcharhinus obscurus* | Data Deficient | Critically Endangered | - | | | |
| *Prionace glauca* | Critically Endangered | - | - | A3 | A3 | |
| *Leucoraja circularis* | Critically Endangered | - | - | | | ☐ |
| *Leucoraja melitensis* | Critically Endangered | - | - | | | ☐ |
| *Leucoraja fullonica* | Critically Endangered | - | - | | | |
| *Dipturus batis* | Critically Endangered | - | - | | | ☐ |
| *Pristis pristis* | Critically Endangered | - | - | | | ☐ |
| *Pristis pectinata* | Critically Endangered | - | - | | | ☐ |
| *Gymnura altavela* | Critically Endangered | - | - | | | ☐ |
| *Pteromylaeus bovinus* | Critically Endangered | - | - | | | |
| *Hexanchus nakamurai* | Data Deficient | Endangered | - | | | |
| *Echinorhinus brucus* | Endangered | - | - | | | |
| *Somniosus rostratus* | Data Deficient | Endangered | - | | | |
| *Squalus acanthias* | Endangered | - | - | | A3 | ☐ |
| *Alopias vulpinus* | Endangered | - | - | | A3 | |
| *Alopias superciliosus* | Endangered | - | - | | | |
| *Cetorhinus maximus* | Endangered | - | - | A2 | | ☐ |
| *Carcharhinus altimus* | Data Deficient | Endangered | - | | | |
| *Carcharhinus plumbeus* | Endangered | - | - | | A3 | |
| *Carcharhinus limbatus* | Data Deficient | Endangered | - | | | |
| *Rostroraja alba* | Endangered | - | - | A3 | | ☐ |
| *Raja radula* | Endangered | - | - | | | |
| *Glaucostegus cemiculus* | Endangered | - | - | | | ☐ |
| *Rhinobatos rhinobatos* | Endangered | - | - | | | ☐ |
| *Rhinoptera marginata* | Data Deficient | Endangered | - | | | |
| *Mobula mobular* | Endangered | - | - | A2 | | ☐ |
| *Heptranchias perlo* | Data Deficient | Vulnerable | - | | A3 | |

| Species | Evaluated IUCN status | Predicted IUCN status | OSPAR (NEA) | Bern Convention (Med) | Barcelona Convention (Med) | GFCM priority species (Med) |
|---|---|---|---|---|---|---|
| *Dalatias licha* | Vulnerable | | - | - | | |
| *Galeorhinus galeus* | Vulnerable | | - | - | - | ⬚ |
| *Mustelus asterias* | Vulnerable | - | - | | A3 | |
| *Mustelus punctulatus* | Data Deficient | Vulnerable | - | | A3 | |
| *Mustelus mustelus* | Vulnerable | - | - | | A3 | |
| *Myliobatis aquila* | Vulnerable | - | - | | | |
| *Dasyatis pastinaca* | Vulnerable | - | - | | | |
| *Dasyatis centroura* | Vulnerable | - | - | | | |

**FIGURES**

Figure 1

**Percent *evaluated* and *predicted* IUCN categorisations of Europe's sharks and rays.** Dark

bars represent the percentage of species officially evaluated on the IUCN Red List, while light

bars represent the percentage of Data Deficient species predicted to be under each category as

per the results of the present study. Of the 120 species in the Northeast Atlantic, 94 were

*evaluated* and 26 were Data Deficient and *predicted* for. In the Mediterranean Sea, 58 of 73

species were *evaluated* and 15 were Data Deficient and *predicted* for. The IUCN categories from

highest to lowest threat are: CR = Critically Endangered, EN = Endangered, VU = Vulnerable,

NT = Near Threatened, and LC = Least Concern.

**Figure 2**

**The effects of size and depth on shark and ray conservation status in Europe.** Histograms of

the probability of an evaluated shark or ray being listed as either Critically Endangered (CR),

Endangered (EN), Vulnerable (VU), Near Threatened (NT), or Least Concern (LC) based on

single-trait Cumulative Link Mixed-effects Model outputs for maximum body size (cm; panels a

and b) and median depth (m; panels c and d). Data include all evaluated species (*n*= 94 Northeast

Atlantic, panels a and c; and *n*=58 Mediterranean Sea, panels b and d) and exclude all Data

Deficient species. Dark grey vertical bars indicate large (300 cm total length, a,b) or shallow

(200 m median depth, c,d) species; light grey bars represent small (150 cm total length, a,b) or

deep (1,000 m median depth, c,d) species. Brackets beside bars indicate the probability of each

species being categorised as threatened (CR, EN, or VU) on the IUCN Red List.

**Figure 3**

**Effects of biological and ecological traits on Europe's shark and ray conservation status.**

Standardized effect sizes with 95% confidence intervals. Cumulative link mixed effect models

with maximum body size (a), reproductive mode (b), median depth (c), and the interaction

between size and depth (d) as fixed effects and taxonomic Family as a random effect to account

for phylogenetic non-independence. Circular and triangular points represent the best explanatory

and predictive model for the Northeast Atlantic and Mediterranean Sea, respectively, which in

both cases included maximum body size, reproductive mode, and the interaction between

maximum size and median depth. Data for maximum size and median depth were centred and

standardised by two standard deviations, while reproductive mode is a binary trait where

oviparous species = 1 and viviparous species = 0.

**Figure 4**

***Predicted* and *evaluated* conservation status of Europe's sharks and rays.** Top Cumulative Link Mixed-effectd Models including

maximum body size, reproductive mode, and the interaction between size and median depth as fixed effects (for both sub-regions) and

taxonomic Family as a random effect to account for phylogenetic non-independence. Panel a shows the probability of all 26 Data

Deficient Northeast Atlantic species, and panel b of all 15 Mediterranean Sea species, being in each IUCN Red List category based on

these top explanatory models. The vertical line cutting down panels a and b represents the 50% cut-off classification used to assign the

final IUCN categorisations (according to the category bar the line crosses). Data for size and depth were centred and standardised by

two standard deviations, while reproductive mode is a binary trait where oviparous species = 1 and viviparous species = 0. Panel c

shows the proportion of sharks and rays in the Northeast Atlantic (left) and Mediterranean Sea (right) both *evaluated* and *predicted* to

be in each IUCN category. Percentage values within each yellow bar indicate the total percentage of *evaluated* threatened and

*predicted*-to-be-threatened species in each set, while numbers within brackets below each bar indicate the total number of species

included in each set.

**Figure 5**

**Informing shark and ray conservation efforts in Europe with categorical predictions.** Solid

grey bars represent species of all IUCN categories excluding those officially evaluated by the

IUCN as Critically Endangered (CR), which are represented by solid red blocks. There are 120

shark and ray species in the Northeast Atlantic (left) and 73 in the Mediterranean Sea (right).

Horizontal red lines indicate the addition of all Data Deficient species *predicted* to be Critically

Endangered, to the *evaluated* block. Orange lines indicate all *evaluated* and *predicted*-to-be-

Endangered and Critically Endangered species, while yellow lines show all imperilled (i.e.

*evaluated* and *predicted*-to-be-threatened) species (Vulnerable, Endangered, and Critically

Endangered). Numbers beside bars indicate total number of species within each relevant

grouping.

Angular Roughshark
150 cm total length
39.4%

Sandbar Shark
300 cm total length
71.7%

Blackmouth Catshark
1,000 m median depth
21.5%

Nursehound
200 m median depth
60%

**CR**

**EN**

**VU**

**NT**

**LC**

**Northeast Atlantic**

Angular Roughshark
62.1%

Sandbar Shark
84.5%

Blackmouth Catshark
40.6%

Nursehound
71.1%

**Mediterranean Sea**

Maximum size (cm)

Median depth (m)

Probability