1

State-dependent differences in the frequency of TMS-evoked potentials

Candice T. Stanfield[1]*, Martin Wiener[1]

[1] Department of Psychology, George Mason University, Fairfax, Virginia, United States of

America

* Corresponding author

E-mail: cstanfie@gmu.edu (CTS)

Abstract

Previous evidence suggests different cortical areas naturally oscillate at distinct frequencies, reflecting tuning properties of each region. The concurrent use of transcranial magnetic stimulation (TMS) and electroencephalography (EEG) has been used to perturb cortical regions, resulting in an observed post-stimulation response that is maximal at the natural frequency of that region. However, little is known about the spatial extent of TMS-induced activation differences in cortical regions when comparing resting state (passive) versus active task performance. Here, we employed TMS-EEG to directly perturb three cortical areas in the right hemisphere while measuring the resultant changes in maximal evoked frequency in healthy human subjects during a resting state (N=12) and during an active sensorimotor task (N=12). Our results revealed that the brain engages a higher dominant frequency mode when actively engaged in a task, such that the frequency evoked during a task is consistently higher across cortical regions, regardless of the region stimulated. These findings suggest that a distinct characteristic of active performance versus resting state is a higher state of natural cortical frequencies.

Introduction

The influence of task-evoked activation and behavior on the modification of spontaneously occurring patterns of neural activity remains a fundamental question in neuroscience. For decades, non-invasive brain stimulation techniques, such as transcranial magnetic stimulation (TMS), have been used to modulate neural activity in humans and other mammals. Furthermore, in numerous reports, concurrent TMS and electroencephalography

(EEG) has been employed to examine cortical reactivity and connectivity. A variety of research using TMS, and some using concurrent TMS-EEG, demonstrates TMS-evoked behavioral and neural effects that are dependent on whether the subject is engaged in a task or not [1-11], as well as differences in neural activation during wakeful versus sleeping states [12].

In neural stimulation research, more is known about the influence exogenous factors have on the brain's electrical response to TMS (frequency and intensity of stimulation; positioning and orientation of the stimulation coil), as opposed to endogenous factors (e.g., global brain state). However, over the past decade, there has been an emergence of research using concurrent TMS-EEG to investigate the influence of endogenous factors on neural response. One such study observed an increase in amplitude and spatial spread during the performance of a short-term memory task [1]. Moreover, the observed task-related excitability increased as a result of stimulation to the cortical area, including spread of TMS-evoked currents to functionally connected areas. Globally, the dominant frequency recorded at the scalp matched that of the stimulated area. Yet, local cortical areas oscillated at a rate closer to its own natural frequency, even when not directly stimulated. Lower-frequency oscillatory peaks were observed in the frontal and parietal cortex (7 Hz in the theta band, and 10 Hz in the alpha band, respectively), reflecting synchronization of local cortical oscillations to parallel networks engaged in task performance. Similar results were reported in a study that provided the first direct evidence for causal entrainment of brain oscillations by short rhythmic TMS bursts while recording resultant EEG responses [2]. The TMS entrainment evoked spatially specific and frequency-specific oscillatory signatures that mimic naturally occurring task-related modulations that are of functional significance. Overall, these task-dependent changes exemplify the importance of further investigation into the influence of endogenous factors, such as global brain state.

Task-dependent changes have also been observed at the single-cell level, with concurrent

single-pulse TMS administered to awake rhesus monkeys [3]. During performance of a visually-

guided grasping task, action potentials in individual neurons within the parietal area PFG were

recorded extracellularly, while either low intensity (60% of the resting Motor Threshold; rMT) or

high intensity (120% of the rMT) stimulation was being administered. Unlike in previous

observations of anesthetized animals, single-pulse stimulation induced a highly localized and

transient excitation followed by reduced activity, corresponding with a significantly longer

grasping time. Thus, the stimulation interfered with task-related activity in parietal neurons,

while simultaneously causing behavioral effects. Additionally, the stimulation induced a highly

localized and short-lived excitation of single neurons in the parietal cortex; however, the TMS-

induced activity and task-related activity did not linearly summate in the PFG neurons. As such,

the spatial spread of TMS-induced spiking activity appeared dissociable from TMS-induced

oscillatory activity, which tends to spread more remotely.

Evidence showing the influence the state of activation has on neural response exists in

humans, as well. TMS-evoked behavioral effects were observed during visual task performance,

while single-pulse TMS was administered over three posterior occipital stimulation sites: left

lateral posterior occipital, right posterior occipital, and medial posterior occipital [5]. Results

showed modulation of activity in the visual cortex through the adaptation of the colored stimuli.

These findings could argue that it is possible for TMS to elicit a state-dependent effect of color;

thus, perceptually facilitating attributes encoded by the least active neural population. To test this

conclusion, an ancillary experiment administered triple-pulse TMS during a visual stimulus

orientation and color task. Here, evidence showed state dependency generalized to the adaptation

of the two attributes; however, adaptation effects were found to either interfere with or improve

the adapting attribute (color), and appeared dependent on the relationship between the adapting

stimulus and the test stimulus. Overall, the authors concluded that behavioral and perceptual

effects of TMS were dependent upon the level of adaptation that occurred within the neural

population during stimulation.

Further exploration into the state-dependent phenomenon has occurred in investigating

functional differences between the wakeful and sleeping brain in humans [12]. During

wakefulness, TMS induced a sustained response, initially with high-frequency oscillatory

activity (20-35 Hz) occurring in the first 100 ms, followed by slower oscillations (8-12 Hz)

persisting until 300 ms. Maximal TMS-evoked current over the rostral premotor cortex during

wakefulness resulted in spatially and temporally differentiated patterns of activation that

propagated along its anatomical connections. Furthermore, during transition into stage 1 sleep,

the TMS-evoked response grew stronger at early latencies but became shorter in duration. TMS-

evoked response during stage 1 sleep initially propagated from the right premotor cortex to the

homotopic contralateral site, but was not sustained nor reached prefrontal for parietal areas.

Conversely, maximal TMS-evoked current density remained restricted to the stimulation area

during non-rapid eye movement (NREM; sleep stages 2 and 3) sleep. Overall, results suggest

that cortical effective connectivity diminishes as we fade out of consciousness.

In addition to TMS-evoked activation studies, a large body of research has focused on

oscillatory signatures arising from macro- and micro-scale neural recordings. Among these

studies has arisen the concept of natural frequencies of human cortical modules, suggesting that

distinct regions of the cortex may naturally oscillate at distinct frequencies [13]. Of great interest

is an expansion of the natural frequencies concept, reporting evoked dominant oscillations in

specific cortical regions, with posterior regions naturally resonating at lower frequencies (~10Hz,

alpha) and anterior regions at higher frequencies (~40Hz, gamma) [14–15].

While the above results suggest state-dependent differences in cortical oscillations and

neuronal connectivity, that may be examined using TMS-EEG, the difference in evoked

oscillatory activity between resting and active task states has yet to be investigated. To further

explore this, in the present study we tested two groups of subjects (N=12 each) while

simultaneously applying single-pulse TMS and recording resultant EEG responses. One group

was tested while subjects were at rest, similar to the previous reports, while a second group was

tested while actively engaged in a simple sensorimotor task. We repeated the methods and

analysis regimes of the prior studies, wherein TMS was administered and frequency spectra data

were obtained to measure the maximum evoked frequency at each stimulation site [14].

Materials and methods

Subjects. Twenty-four right-handed subjects (12 females, age 19–36 years) with normal

or corrected-to-normal visual acuity participated in this study, and were randomized into one of

two groups: twelve subjects participated in the passive experiment (7 females, mean age 22.5

years), while the other twelve subjects participated in the active experiment (5 females, mean age

25.2 years). Following safety and ethical guidelines [16], all subjects were eligible to receive

TMS. All subjects provided informed consent and all protocols were approved by the George

Mason University Institutional Review Board.

TMS. A focal figure-of-eight coil with 70mm wing diameter driven by a Magstim Rapid²

biphasic stimulator (Magstim Inc., Wales, UK) was used to non-invasively stimulate the

subjects' cortex. On the right hemisphere of the scalp, three cortical sites were selected over the

EEG electrodes P4 (occipital), C4 (parietal), and F4 (frontal). These sites were chosen as

homologous regions to those stimulated previously in a study demonstrating distinct frequency

responses [14]. To verify anatomical locations of the cortical stimulation sites a T1-weighted

MRI of one subject was used and targeted using MNI coordinates in the Brainsight

neuronavigation system (Rogue Research Inc., Montreal, Canada). MRI scans were unavailable

for the other subjects in this study.

High-density EEG recording during TMS. TMS-evoked potentials (TMS-EP) were

recorded using an actiCHamp 64-channel amplifier (Brain Products GmbH, Germany) and TMS-

compatible actiCAP slim active electrodes (international 10-20 system), with FCz as the online

reference. BrainVision Recorder (v. 1.20.0801) was used to digitize the EEG at a sampling

frequency of 5000 Hz. Electrode impedances were kept <20 kΩ. To minimize contamination of

auditory potentials evoked by the click associated with the TMS coil discharge [17], subjects

underwent a TMS-click auditory perception (TMS-CAP) test prior to beginning the experiment.

During the test, subjects wore inserted wired silicone-tipped earplug headphones with a Noise

Reduction Rating (NRR) of 26 dB, while a masking noise with the same spectral profile of the

TMS coil click was continuously played. Recordings of the TMS coil click emitted by the

Magstim coil were used to create the masking noise and scrambled into a continuous sound file

with the same spectral properties, thus, capturing the specific time-varying frequency

components of the TMS click. For the TMS-CAP test, subjects listened to the masking noise

while a brief TMS burst was administered on top of the FCz (average reference) electrode.

Subjects were instructed to notify the experimenter if they could hear the TMS coil click. If a

subject reported hearing the click, the volume of the masking noise was raised to a level still

comfortable for the subject and/or the stimulator intensity output percentage was lowered until

the click was as imperceptible as possible without lowering the stimulator output to an

ineffective intensity (<40 V/m) [14]. Once the TMS-CAP was complete, subjects were required

to continue wearing the earplug headphones for the duration of the experiment while listening to

the masking noise at their individualized fixed volume. All TMS-induced artifacts were attended

to during offline analysis.

TMS protocol. During this experiment, subjects in both groups received a single-pulse

TMS protocol. This consisted of a series of TMS pulses that were administered one-at-a-time

over a part of the brain on the right hemisphere of the scalp, approximately over electrodes P4,

C4, and F4 (Fig 1). At each of the three electrode sites, a total of 100 pulses were administered

repetitively at each site, separated by a short period of time based on randomized experimental

group assignment (passive or active), resulting in a total of 300 pulses overall. Based on their

individual TMS-CAP results, subjects in the passive experimental group received single pulse

stimulation at a fixed intensity of between 30–50% of maximum stimulator output (MSO; range

44.7–74.5 V/m), while subjects in the active experimental group received single pulse

stimulation at a fixed intensity of between 35–50% of MSO (range 52.1–74.5 V/m,); both groups

received an average of 42.5% MSO ($\pm$ 7.23 for passive and $\pm$ 5.84 for active, respectively).

According to an independent samples t-test, MSO values were not different between the two

groups; $t(22) = 0$, $p = 1$.

We additionally calculated the modeled electrical field of our TMS stimulator across the

different MSO intensities employed. To that end, we conducted electrical field modeling using

SimNIBS software (v. 2.1.20) [18], on a standardized MNI template with a modeled Magstim

TMS coil matching our own. Similar to MSO values, we observed no differences between

groups (Passive: average 63.3±10.77 V/m; Active: average 63.3±8.71 V/m). Additionally, we

note that the lowest intensity in our tested sample (44.7 V/m) did not exceed the minimal

intensity previously reported as minimal for evoking dominant frequencies [14].

181
182   **Fig 1. Stimulation sites.** a) represents electrode stimulation sites, indicated by red X mark; top X is F4, middle X is
183   C4, and bottom X is P4. Images b) and c) represent stimulation sites on a rendered brain as determined by Brainsight
184   localization of the TMS coil on a sample subject.
185
General experimental procedures for both experimental groups. During the experiment,

subjects sat in an ergonomic chair, relaxed, and with eyes open looking at a fixation cross on a

screen. Once the selected electrode site was targeted, we stimulated it at an intensity that was set

by the subject's TMS-CAP results.

Passive group experimental procedures. Subjects in the passive experimental group did

not perform a sensorimotor task; instead, they were instructed to rest with their eyes open during

the experiment. Subjects viewed an LCD monitor with a 120 Hz refresh rate (Cambridge

Research Systems, United Kingdom) approximately 70-cm away with a black background, and

were instructed to keep their eyes fixed on a 5x5 cm black fixation cross that appeared in the

center of the screen. Each of the three electrode sites were stimulated in a counterbalanced block

design and received 100 stimuli per block at a randomized inter-stimulation-interval (ISI)

between 4–6 seconds.

Active group experimental procedures and task. Subjects in the active experimental

group performed a sensorimotor luminance detection task that relied on gradual signal detection,

and was programmed in PsychoPy (v. 1.85.6) [19]. Subjects viewed a screen approximately 70-

cm away with a grey background and a 5x5 cm fixation cross with a white outline in the center

of the screen that gradually changed from a solid black center to solid white, achieving full

luminance (Fig 2). Once the subject perceived the fixation cross to reach full luminance, they

pressed a key with their right hand, upon which a single-pulse stimulation was delivered before

the next trial automatically began. A white outline of a fixation cross was presented to each

subject at the start of each trial, during which the interior gradually increased in luminance at a

rate of 0.0025 value/frame (in HSV units). Each of the three electrode sites were stimulated in a

counterbalanced block design and received 100 stimuli per block.

209
210   **Fig 2. Schematic of the sensorimotor luminance detection task for the active experimental group.** Subjects
211   viewed a fixation point that gradually became illuminated in the center, and were required to press a button when
212   they judged the luminance to be fully saturated, thus initiating the next trial.
213
Analysis. Offline data analysis was conducted using the EEGLAB MATLAB Toolbox

[20] in MATLAB R2017b (The MathWorks Inc., Natick, MA). Continuous data were

downsampled to 500 Hz and subsequently analyzed via the *clean_rawdata* plugin (v. 0.34) [21]

to clean continuous data following the Artifact Subspace Reconstruction (ASR) method [22] to

remove bad EEG channels. To prevent result biases from potentially removing excessive

datapoints, *clean_rawdata* provided us with controlled, objective rejection criteria to eliminate

noisy channels for such artifacts as eye blinks and face/neck muscle activity. Following this, all

data were re-referenced to the grand average of all electrodes and then epoched for all three

stimulation sites from -1000 to +1000 ms around the TMS pulse; the data for each epoch was

baseline-corrected to the mean of the entire epoch span.

TMS-artifact removal. Given the emergence of concurrent TMS and EEG as an important

tool for assessing cortical properties, TESA—an open-source extension for EEGLAB—was

created with the purpose of improving and standardizing analysis across the field of TMS-EEG

research [23]. We applied the TESA toolbox to all three stimulation site epochs to remove

artifacts; all steps adhered to the TESA pipeline [23]. This process involved 1) removing all data

around the TMS pulse from -10 to +10 ms, 2) interpolating removed data within the TMS pulse,

3) removing noisy trials via EEGLAB's built-in joint probability detection, 4) running a first

round of independent component analysis (ICA) using the FastICA algorithm, 5) removing

artifact components via visual inspection, 6) applying a first-order Butterworth filter with a

bandpass of 1–100 Hz, as well as a notch filter to remove 60 Hz electrical line interference, 7)

running a second round of FastICA with subsequent artifact component rejection. Following the

above steps, data were again filtered between 1 and 50 Hz and segregated into separate, site-

specific epochs.

Time/frequency analysis. To analyze time-frequency domain responses we calculated the

event-related spectral perturbation (ERSP) values based on Morlet wavelets, via the EEGLAB

newtimef function, by convolving a mother wavelet at 100 linearly-spaced frequencies spanning

5 to 50 Hz, with 3.5 cycle wavelets and a 0.5 scaling factor. Baseline correction was applied to

the average power across trials by subtracting the mean baseline power. Analysis of

time/frequency data thus proceeded at both the "local" and "global" levels, following the

convention of previous experiments [14]. Accordingly, global effects were determined by

averaging, for each subject, the time/frequency spectrogram across all electrodes to form a single

representation of the ERSP across the scalp. In contrast, local effects were determined by

examining the ERSP underneath each stimulation electrode across all three sites of stimulation.

To minimize the effect of possible artifacts occurring at the time of stimulation, natural

frequencies were calculated by averaging the ERSP values in a time window between 20 and 200

ms (see below).

The above parameters were chosen to replicate the results of prior work; yet, we also

included an additional analysis in which ERSPs were baseline-corrected at the single-trial level,

rather than the average, via single-trial normalization, which has been shown to decrease

sensitivity to noise artifacts [24]. All of the calculations outlined below were also calculated on

this secondary data set.

Global field power. In addition to the analysis of ERSP data, we also calculated global

field power (GFP), defined as the reference-independent response strength, and calculated as the

standard deviation across all electrodes at each timepoint [25]. GFP data were analyzed across all

three sites of stimulation, separately for passive and active groups, in order to determine if there

were any differences in evoked activity following TMS at any site.

Natural frequencies. Our analysis of natural frequencies proceeded according to the

description from previous reports [14–15]. To determine the natural frequency for each subject at

each stimulation site, the global ERSP (gERSP) response was analyzed by calculating the sum of

power values for each frequency within the 20–200 ms time window, and then determining

which frequency had the highest value. In this way, the max frequency would not be driven

alone by a single frequency with a very high peak, but could instead be provided by a frequency

with a moderate yet sustained response that was larger than at other frequency bands. Natural

frequencies were calculated for each stimulation site for each subject in both groups.

268   *Statistical analysis*. All statistical analysis of behavioral data and natural frequencies

were carried out in SPSS (v. 19, IBM Corporation). For the analysis of global and local effects,

we employed cluster-level corrections for significance ($p < 0.05$) [26] and implemented via

Fieldtrip using the *statcondfieldtrip* command in EEGLAB. For both local and global effects, we

determined regions of significant deviation from baseline for each of the sites, for each of the

two groups. In addition, we compared the gERSP between groups, by averaging across all three

sites within each group and comparing the overall responses.

Results

Global response to TMS

277   Our initial analyses set out to attempt to reproduce the methods used by previous studies

[14–15], for reporting dominant frequencies in specific cortical regions. In these studies, an

increase in power is observed following TMS that is maximal at a particular frequency band,

dependent on the site of stimulation. These changes reflect the spectral properties of the TMS-

evoked oscillations, which consists of a number of repeated positive and negative deflections

[27]. When examining the gERSP response, averaged across all electrodes, we observed a

combination of increases and decreases in power following TMS. Notably, only the *decreases* in

power survived our cluster-corrected significance threshold, in contrast to the original findings.

This finding was observed across both passive and active groups. However, we note that our

study used a different design and methods to these previous reports. Most notably, these

previous studies stimulated cortical regions at a much higher intensity than ours; in the present

study, we sought to reduce the impact of artifact peripheral components in the TMS-EP [28]. In

doing so, our stimulation intensities were far lower than that used previously. Nevertheless, the

difference in evoked frequency response between conditions, our modeling findings, and

behavioral differences in the active group between motor cortex stimulation and other sites

suggest that our stimulation intensities were sufficient for inducing activity in cortical columns.

Crucial differences were observed between passive and active groups, as well as between

the different sites of stimulation. For the passive group, we observed decreases in power that

were synchronous with the TMS pulse in the gamma frequency band (40–50 Hz) across all three

sites. Across stimulation sites, the gamma desynchronization became longer lasting from

posterior to frontal regions, and was further accompanied at the frontal site by a significant

decrease in the high beta range (20–30 Hz) approximately 100–300 ms after the TMS pulse. In

contrast, the active group exhibited a larger desynchronization response across all three sites,

extending from the beta to gamma range (Fig 3).

Fig 3. Global plots for all subjects illustrating the three cortical sites targeted by TMS. Butterfly plots (top panels) of all electrode time courses with the black trace line highlighting the electrode directly underlying the stimulator. ERSP plots (bottom panels) display saturated color areas representing significant frequency (Hz) activation compared to baseline. (a) P4 for passive group, (b) C4 for passive group, (c) F4 for passive group. (d) P4 for active group, (e) C4 for active group, (f) F4 for active group. Significance was determined via cluster-based permutation testing.

In addition to the spectral response, we also calculated and measured the GFP. Here, as well, we observed no differences between site in the evoked response, nor was there any difference between passive and active groups (Fig 4).

312
313 **Fig 4. Global field power (GFP) across stimulation sites and groups.** Shaded regions display standard error. No
314 differences between stimulation site or group were detected.
315

Natural frequencies

The major finding of the previous work was that the so-called "natural frequency,"

characterized as the frequency band with the largest sustained response to TMS, increases in a

rostro-caudal gradient. Calculating the natural frequency using the same method outlined by the

previous authors yielded a range of values across all three sites. Though the individual maximum

frequencies showed a large oscillatory range, there were no outliers. Yet, no linear effects were

observed in these values across all three sites, for either the passive or active groups. However,

along with the gERSP responses, we observed that the active task group exhibited significantly

higher natural frequencies evoked by TMS than the passive group. A two-way ANOVA revealed

that group type had a significant effect on mean activation in Hz ($F(1, 22) = 4.557$, $p = 0.044$,

$\eta p^2 = 0.172$), with higher frequencies reported during the active experimental group ($M = 22.43$,

$SD = 12.24$) compared to the passive experimental group ($M = 15.66$, $SD = 12.6$).

328
329   **Fig 5. Mean natural frequencies, based on stimulation site and group assignment.** Individual data points
330   represent the maximal evoked frequency for each subject in the resting (blue) and active (red) state subject groups.
331   Consistently higher evoked frequencies were observed across all stimulation sites. Error bars represent standard
332   error.
333

Local response to TMS

Passive group

During P4 stimulation, observations for the P4 site showed significantly evoked

activations in the gamma-band that were recorded from stimulation onset to 100 ms post-

stimulation, in addition to beta-band activations from 100 ms to 200 ms. No significantly evoked

activations were observed for the C4 site during C4 stimulation. During F4 stimulation,

observations for the F4 site showed significantly evoked activations in the gamma-band that

were recorded from stimulation onset to 300 ms post-stimulation.

Active group

Though similar to the observations made in the P4 for the passive group, the significantly evoked activations in the active group, during active task engagement, showed a longer post-stimulation effect. At the P4 site, significantly evoked activations in the gamma-band that were recorded from stimulation onset to 250 ms post-stimulation, with an overlap of beta-band activations from 50 ms post-stimulation until 300 ms. No significantly evoked activations were observed for the C4 site during C4 stimulation, nor for the F4 site during F4 stimulation (Fig 6; both groups).

350
351 **Fig 6. Local responses to stimulation. ERSP plots of local responses (under each electrode) to TMS**
352 **stimulation at each site under each condition.** Panels (a–c) represent Passive group and (d-f) represent Active
353 group. A red star at the lower left corner of a panel indicates the ERSP plot under the electrode that was stimulated.
354 Saturated color areas represent significant frequency (Hz) activation compared to baseline.
355

Active group response times. Response times (RTs) were calculated in seconds (s) for the

active group, which performed the illumination detection task; P4 (2.20±1.24), C4 (2.44±1.27),

F4 (2.26±1.26) (Fig 7). A nonparametric Friedman test showed a significant difference in RTs

between stimulation sites; $\chi^2(2, 12) = 8.667$, $p = 0.013$. Post hoc analysis with Wilcoxon signed-

rank tests were conducted with a Bonferroni correction applied, resulting in a significance level

set at $p < 0.05$. Median (IQR) RTs based on stimulation sites were 2.13 s (1.19 to 2.74) for P4,

2.05 s (1.52 to 2.92) for C4, and 2.06 s (1.17 to 2.76) for F4, respectively. There were no

significant differences between RTs for P4 and F4 ($Z = -0.784$, $p = 0.433$) or between RTs for C4

and F4 ($Z = -1.412$, $p = 0.158$). However, there was a significant difference between RTs for C4

compared to P4, with slower RTs overall for C4 ($Z = -1.961$, $p = 0.05$).

366
367   **Fig 7. Average active group task response times (RTs) in seconds, based on electrode stimulation site.** Blue
368   circles represent for mean RTs for each active task subject, per stimulation site. Subjects were moderately slower in
369   responding following single-pulse stimulation over C4. Error bars represent standard error.
370

Discussion

In the current study, we used concurrent TMS-EEG to investigate cortical reactivity

differences between passive state and active sensorimotor task performance. Our results revealed

a complexity of patterns in global and local changes that occur in response to stimulation, in

addition to conspicuously broader and distinct patterns between the passive and active states.

These findings suggest cortical regions exhibit complex frequency-specific profiles. Specifically,

our findings suggest that oscillatory mechanisms are characterized by more complex, state-

dependent patterns than have previously been appreciated. Additionally, our findings suggest

that patterns of ongoing spontaneous activity are modified by task performance, and differ based

on individual activation patterns.

Endeavoring to explain the complexity of varying oscillatory bands, further

investigation into whether timescales of different frequency bands correlate with a sensory-to-

higher processing hierarchy was conducted [29]. However, no strong biases toward specific

timescales across cortical regions was observed, nor exclusivity of lower areas biased toward

faster frequencies or higher areas biased toward slower frequencies. Thus, it was contended that

dominant higher-frequency bands observed in the frontal cortex during concurrent TMS-EEG

studies may be a result of that brain region's involvement in higher level cognition. This

explanation has potential to elucidate our findings of higher frequency evoked power across the

brain during sensorimotor task engagement. Furthermore, frequency bands may serve as

channels of communication across brain regions, though dependent on the activation in multiple

bands within each region [30].

Perhaps the most notable aspect of our findings is the difference in evoked frequency

between the experimental groups. While performing the sensorimotor task, evoked responses

became more widespread in both frequency and time; additionally, when comparing the natural

frequency between experimental groups, the sensorimotor task was observed to evoke a

consistently higher frequency than in the resting state group. This difference suggests that

cognitive engagement incorporates higher frequency oscillations, consistent with several other

known findings of brain function [31–35]. A state-dependent TMS effect might account for these

findings. State-dependency is defined as response changes according to the state of the cortex

when the stimulus, such as a TMS pulse, is applied [36]. Moreover, the state of activation has

been shown to influence the response [4–12]. The effect of small TMS pulses might be

facilitated if the cortex is already active; thus, it would be reasonable to presume that single pulse

stimulation may well enhance cortical activation while a subject is actively engaged in a task

[37]. Previous studies have shown evidence of the effect of TMS pulses varying as a function of

the state of the brain. For example, when comparing neuronal activation during resting/baseline

states to active task engagement, researchers found TMS over the motor cortex enhanced

activation during motor execution and motor imagery [38–40], while others found greater ease of

inducing phosphenes with TMS over the occipital region during visual mental imagery [41]. The

latter finding suggests distinct operational modes for the brain between resting state and task-

based networks. Consistent with this view, previous investigations comparing resting-state and

task-based network activity in functional magnetic resonance imaging (fMRI) have revealed

network reorganization between these states [42,43]; in particular, the frequency profile of fMRI

inter-connectivity shifts between resting and task-based activity, with lower frequencies

dominating the former and more broadband representation during tasks [44]. Although these

fluctuations operate on an order of magnitude below those measured by EEG in the present study

(0.01-0.1Hz), they reveal a similar pattern to our findings, suggesting a correspondence [45].

Notably, for the current study, the higher evoked frequencies did not depend on the

stimulation site, suggesting a global change in brain functioning, independent of the local

changes. Finally, our findings confirm that TMS can be a useful tool for evoking latent

oscillations in the brain [14].

Limitations

In the current study, there is a possible limitation that should be noted. We recognize that

the stimulation levels in our study are lower, on average, than used in previous reports. This was

done to avoid auditory-evoked artifacts in the EEG response. Yet, we note that TMS intensity

was above 40 V/m, previously reported as minimal for evoking dominant frequencies [14].

Furthermore, of importance is that low-intensity stimulation did not induce any effect in awake

rhesus monkeys during task performance [3]; therefore, it might suggest our intensity was high

enough, as we did observe TMS-evoked differences based on experimental group.

Conclusions

We investigated TMS-evoked cortical reactivity differences between subjects who were

either at rest (passive group) or engaged in a sensorimotor task (active group), while recording

resultant EEG responses. The observed complex patterns of global and local changes, in

response to stimulation, suggests cortical regions exhibit complex frequency-specific profiles.

Additionally, the differences in evoked responses between the two experimental groups suggests

that oscillatory mechanisms are characterized by complex, state-dependent patterns, with an

overall higher mode of frequency during active engagement. All subject data are available via

the online open access repository FigShare (https://figshare.com/account/home#/projects/57563).

Acknowledgements

We would like to acknowledge Melody Barnard, our former undergraduate research assistant, for her help during data collection.

References

1. Johnson JS, Kundu B, Casali G, Postle BR. Task-dependent changes in cortical excitability and effective connectivity: A combined TMS-EEG study. J Neurophysiol. 2012;107:2383-2392.

2. Thut G, Veniero D, Romei V, Miniussi C, Schyns P, Gross J. Rhythmic TMS causes local entrainment of natural oscillatory signatures. Curr Biol. 2011;1:1176-1185.

3. Romero MC, Davare M, Armendariz M, Janssen P. Neural effects of transcranial magnetic stimulation at the single-cell level. Nat Comm. 2019;10(1), 2642.

4. Thut G, Northoff G, Ives JR, Kamitani Y, Pfennig A, Kampmann F, et al. Effects of single-pulse transcranial magnetic stimulation (TMS) on functional brain activity: A combined event-related TMS and evoked potential study. Clin Neurophysiol. 2003;114:2071-2080.

5. Silvanto J, Muggleton NG, Cowey A, Walsh V. Neural adaptation reveals state-dependent effects of transcranial magnetic stimulation. Euro J Neurosci. 2007;25:1874-1881.

6. Romei V, Rihs T, Brodbeck V, Thut G. Resting electroencephalogram alpha-power over posterior sites indexes baseline visual cortex excitability. NeuroReport. 2008;19:203-208.

7. Silvanto J, Pascual-Leone A. State-dependency of transcranial magnetic stimulation.

Brain Topogr. 2008;21:1-10.

8. Massimini M, Ferrarelli F, Murphy MJ, Huber R, Riedner BA, Casarotto S, et al. Cortical

reactivity and effective connectivity during REM sleep in humans. Cog Neurosci.

2010;1:176-183.

9. Miniussi C, Thut G. Combining TMS and EEG offers new prospects in cognitive

neuroscience. Brain Topogr. 2010;22:249-256.

10. Romei V, Thut G, Silvanto J. Information-based approaches of noninvasive transcranial

brain stimulation. Trends in Neurosci. 2016;39:782-795.

11. Petrichella S, Johnson N, He B. The influence of corticospinal activity on TMS-evoked

activity and connectivity in healthy subjects: A TMS-EEG study. PLoS ONE; 2017;12:1-

18.

12. Massimini M, Ferrarelli F, Murphy MJ, Huber R, Esser SK, Singh H, et al. Breakdown of

cortical effective connectivity during sleep. Science. 2005;309:2228-2232.

13. Niedermeyer E. The normal EEG of the waking adult. In: Niedermeyer E, Lopes da Silva

F, editors. Electroencephalography: Basic principles, clinical applications and related

fields. 4th ed. Baltimore, MD: Williams & Wilkins; 1999.

14. Rosanova M, Casali A, Bellina V, Resta F, Mariotti M, Massimini M. Natural

frequencies of human corticothalamic circuits. J Neurosci. 2009;29:7679-7685.

15. Ferrarelli F, Sarasso S, Guller Y, Riedner BA, Peterson MJ, Bellesi M, et al. Reduced

natural oscillatory frequency of frontal thalamocortical circuits in schizophrenia. Arch

Gen Psychiatry. 2012;69:766-774.

16. Rossi S, Hallett M, Rossini PM, Pascual-Leone A, The Safety of TMS Consensus Group.

Safety, ethical considerations, and application guidelines for the use of transcranial

magnetic stimulation in clinical practice and research. Clin Neuropsych. 2009;120:2008-

2039.

17. ter Braack EM, de Vos CC, van Putten MJAM. Masking the auditory evoked potential in

TMS-EEG: A comparison of various methods. Brain Topogr. 2015;28:520-528.

18. Thielscher A, Antunes A, Saturnino GB. Field modeling for transcranial magnetic

stimulation: A useful tool to understand the physiological effects of TMS? IEEE EMBC

2015: Proceedings of the 37th Annual International Conference of the IEEE Engineering

in Medicine and Biology Society; 2015 Aug 25-29; Milano, Italy. p. 222-225.

19. Peirce, JW. Generating stimuli for neuroscience using PsychoPy. Front Neuroinfo.

2009;2:1-8.

20. Delorme A, Makeig S. EEGLAB: An open source toolbox for analysis of single-trial

EEG dynamics including independent component analysis. J Neurosci Methods.

2004;134:9-21.

21. Kothe C. Clean_rawdata() (Version 0.34) [EEGLAB extension]. 2014. Available from:

https://sccn.ucsd.edu/wiki/EEGLAB_Extensions

22. Mullen T, Kothe C, Chi YM, Ojeda A, Kerth T, Makeig S, et al. Real-time modeling and

3D visualization of source dynamics and connectivity using wearable EEG. Conf Proc

IEEE Eng Med Biol Soc. 2013;2013:2184-2187.

23. Rogasch NC, Sullivan C, Thomson RH, Rose NS, Bailey NW, Fitzgerald PB, et al.

Analysing concurrent transcranial magnetic stimulation and electrocephalographic data:

A review and introduction to the open-source TESA software. Neuroimage.

2017;147:934-951.

24. Grandchamp R, Delorme A. Single-trial normalization for event-related spectral

decomposition reduces sensitivity to noisy trails. Front Neurosci. 2011;2:1-14.

25. Murray MM, Brunet D, Michel CM. Topographic ERP analysis: A step-by-step tutorial

review. Brain Topogr. 2008;20:249-264.

26. Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. J

Neurosci Methods. 2007;164:177-190.

27. Lioumis P, Kičić D, Savolainen P, Mäkelä JP, Kähkönen S. Reproducibility of TMS–

evoked EEG responses. Hum Brain Map. 2009;30:1387-1396.

28. Conde V, Tomasevic L, Akopian I, Stanek K, Saturnino GB, Thielscher A, et al. The

non-transcranial TMS-evoked potential is an inherent source of ambiguity in TMS-EEG

studies. NeuroImage. 2019;185:300-312.

29. Mellem MS, Wohltjen S, Gotts SJ, Ghuman AS, Martin A. Intrinsic frequency biases and

profiles across human cortex. J Neurophysiol. 2017;118:2853-2864.

30. Hillebrand A, Tewarie P, van Dellen E, Yu M, Carbo EWS, Douw L, et al. Direction of

information flow in large-scale resting-state networks is frequency-dependent. Proc Natl

Acad Sci USA. 2016;113:3867-3872.

31. Crone NE, Miglioretti DL, Gordon B, Lesser RP. Functional mapping of human

sensorimotor cortex with electrocorticographic spectral analysis II. Event-related

synchronization in the gamma band. Brain. 1998;121:2301-2315.

32. Canolty RT, Edwards E, Dalal SS, Soltani M, Nagarajan SS, Kirsch HE, et al. High

gamma power is phase-locked to theta oscillations in human neocortex. Science.

2006;313:1626-1628.

33. Voytek B, Canolty RT, Shestyuk A, Crone NE, Parvizi J, Knight RT. Shifts in gamma

phase-amplitude coupling frequency from theta to alpha over posterior cortex during

visual tasks. Front Human Neurosci. 2010;4:1-9.

34. Crone NE, Korzeniewska A, Franaszczuk PJ. Cortical gamma responses: Searching high

and low. Intl J Psychophysio. 2011;79:9-15.

35. Groppe DM, Bickel S, Keller CJ, Jain SK, Hwang ST, Harden C, et al. Dominant

frequencies of resting human brain activity as measured by the electrocorticogram.

NeuroImage. 2013;79:223-233.

36. Siebner HR, Hartwigsen G, Kassuba T, Rothwell JC. How does transcranial magnetic

stimulation modify neuronal activity in the brain? Implications for studies of cognition.

Cortex. 2009;45:1035-1042.

37. Matthews PBC. The effect of firing on the excitability of a model motoneurone and its

implications for cortical stimulation. J Physiol. 1999;518:867-882.

38. Kasai T, Kawai S, Kawanishi M, Yahagi S. Evidence for facilitation of motor evoked

potentials (MEPs) induced by motor imagery. Brain Res. 1997;744:147-150.

39. Fadiga L, Buccino G, Craighero L, Fogassi L, Gallese V, Pavesi G. Corticospinal

excitability is specifically modulated by motor imagery: A magnetic stimulation study.

Neuropsychologia. 1999;37:147-158.

40. Hashimoto R, Rothwell JC. Dynamic changes in corticospinal excitability during motor

imagery. Exp Brain Res. 1998;125:75-81.

41. Sparing R, Mottaghy FM, Ganis G, Thompson WL, Töpper R, Kosslyn SM, et al. Visual

cortex excitability increases during visual mental imagery–A TMS study in healthy

human subjects. Brain Res. 2002;938:92-97.

42. Spadone S, Della Penna S, Sestieri C, Betti V, Tosoni A, Perrucci MG, et al. Dynamic

reorganization of human resting-state networks during visuospatial attention. Proc Natl

Acad Sci USA. 2015;112:8112-8117.

43. Gonzalez-Castillo J, Bandettini PA. Task-based dynamic functional connectivity: Recent

findings and open questions. NeuroImage. 2018;180:526-533.

44. Ciuciu P, Abry P, He BJ. Interplay between functional connectivity and scale-free

dynamics in intrinsic fMRI networks. NeuroImage. 2014;95:248-263.

45. Bridwell DA, Wu L, Eichele T, Calhoun VD. The spatiospectral characterization of brain

networks: Fusing concurrent EEG spectra and fMRI maps. NeuroImage. 2013;69:101-

111.