Genetic risk loadings influence the susceptibility and severity of systemic lupus erythematosus

Lingyan Chen[1,2], Phil Tombleson[1], Adrianna Bielowka[1], Christopher A Odhams[1], Amy L Roberts[1], Deborah S Cunninghame Graham[1], Timothy J Vyse[1,*] and David L Morris[1,*]

1. Department of Medical & Molecular Genetics, King's College London, London, UK.

2. MRC/BHF Cardiovascular Epidemiology Unit, University of Cambridge, Cambridge, UK.

*These are corresponding authors and joint last authors.

Abstract

Systemic lupus erythematosus (SLE) is a chronic autoimmune disease with a wide range of clinical manifestations. Of these, kidney involvement is the most common cause of morbidity and mortality. Genome-wide association studies (GWAS) have identified more than 80 loci that are associated with SLE. We calculated the genetic risk score (GRS) using SNPs that are associated with SLE. We studied three GWAS sets and found that the best GRS in the prediction of SLE generated an area under the ROC curve of 0.745 (95%CI 0.735-0.754). However, it is not known whether genetic factors affect the clinical features. We further showed a significant correlation between a GRS and renal involvement in two independent European GWAS: cohort 1 (N_{Renal+} = 1,152, N_{Renal-} = 1,949) and cohort 2 (N_{Renal+} = 146, N_{Renal-} = 378) – the higher the GRS, the higher risk of renal disease ($P_{cohort1}$ = 2.44e-08; $P_{cohort2}$ = 0.00205) and the younger age of SLE onset ($P_{cohort1}$ = 1.76e-12; $P_{cohort2}$ = 0.00384). When partitioning the patients according to the age of SLE onset, we found that the GRS performed better in the prediction of renal disease in the 'late onset' group comparing to the 'early onset' group. In conclusion, age of onset incorporating a GRS may assist prediction of lupus nephritis in a clinical setting.

Key words

Genetic risk score (GRS), Systemic lupus erythematosus (SLE), lupus nephritis, age of onset, severity

Introduction

Systemic lupus erythematosus (SLE) is a chronic inflammatory autoimmune disease characterized by a wide spectrum of signs and symptoms varying among affected individuals and can involve many organs and systems, including the skin, joints, kidneys, lungs, central nervous system, and haematopoietic system [1]. A recent report underscores that SLE is among the leading causes of death in young females, particular females among ages 15-24 years, in which SLE ranked tenth in the leading causes of death in all populations and fifth for African American and Hispanic females [2]. Lupus nephritis is the most common cause of morbidity and mortality. Patients with kidney disease are likely to have more severe clinical outcomes and a shorter lifespan. 30-60% of adults and up to 70% of children with SLE have renal disease, characterized by the glomerular deposition of immune complexes and an ensuring inflammatory response [3]. Genetic ancestry influences the incidence and prevalence of SLE and kidney involvement, being more frequent in Hispanics, Africans and Asians than in European [4-7]. Currently, kidney disease in SLE is diagnosed by use of light microscopy, which drives therapeutic decision-making. However, not all patients will respond to therapy, indicating that additional information focusing on the mechanism of tissue injury is required. Moreover, early detection of kidney involvement in SLE is important because early treatment can be applied to reduce the accumulation of renal disability.

Although the exact aetiology of lupus is not fully understood, a strong genetic link has been identified through the application of family [8, 9] and twins studies [10]. SLE does not follow a Mendelian pattern of inheritance, and so it is termed a non-Mendelian disease or complex trait. Complex traits are multi-factorial with both genetic and environmental contributions. Genome-wide association studies (GWAS) have been successfully used to investigate the genetic basis of a disease and this has dramatically advanced knowledge of the genetic aetiology of SLE. Our recent review summarized a total of 84 genetic loci that are implicated as SLE risk [11]. Despite the advances in the genetics of SLE, it is not clear how to utilise genetic information for the prediction of SLE risk or severity.

A genetic risk score (GRS) summarizes risk-associated variations by aggregating information from multiple risk single nucleotide polymorphisms (SNPs). The approach to calculate the GRS is to simply count disease-associated alleles or weighting the summed alleles by log Odds Ratios. Recent studies [12, 13] have proposed methods which select SNPs from GWAS by LD (linkage disequilibrium) pruning and clumping and thresholding for GRS calculation. As the number of SNPs included in a GRS

increases, the distribution approaches normality, even when individual risk alleles are relatively

uncommon. Therefore, a GRS can be an effective means of constructing a genome-wide risk

measurement that summarises an individual's genetic predisposition to SLE. Moreover, as GRSs pool

information from multiple SNPs, each individual SNP does not strongly influence the summary

measurement. Thus, the GRS is more robust to imperfect linkage for any tag SNP and causal SNP,

and less sensitive to minor allele frequencies for individual SNPs [14-17].

In this study, we firstly tested whether a quantitative model - a GRS derived from SLE GWAS applying

a range of methods, was an effective way to distinguish SLE patients and controls in three independent

cohorts. Next, we classified SLE patients into two groups: SLE renal+ (patients with renal disease) and

SLE renal- (patients without renal disease), and performed a case-case genome-wide association study

(GWAS) in two independent SLE cohorts with available renal data for the identification of SLE renal

susceptibility loci. However, no genome-wide significant genetic risk loci were identified in the SLE

renal GWASs. We then tested whether a GRS derived from SLE GWAS was an effective way to

distinguish SLE patients with or without renal disease in two independent cohorts.

Methods

Samples source

Samples were from three previously published SLE genome-wide association studies (GWASs) in the

European population – the SLE main cohort [18], the SLEGEN cohort [19], and the Genentech cohort

[20]. The SLE main cohort [18] was the biggest SLE GWAS, which consisted of 4,036 SLE patients

and 6,959 healthy controls. A total number of 603,208 SNPs were available post quality control. The

SLEGEN cohort [19] was carried out by The International Consortium for Systemic Lupus

Erythematosus Genetics (SLEGEN) on women of European ancestry, which comprised 283,211 SNPs

genotyped for 2,542 controls and 533 SLE patients. The Genentech cohort [20] was performed by

Genentech on North American individuals of European descent, which comprised 487,208 SNPs

genotyped for 1,165 cases and 2,107 controls.

Clinical sub-phenotypes were available for the SLE main cohort and SLEGEN cohort, which were

documented according to the standard American College of Rheumatology (ACR) classification criteria.

Subgroups of patients with renal disease or without renal disease were identified according to the sub-

phenotype data using ACR classification.   Following quality control, the sample size of patients with

renal disease, lupus nephritis (LN+) were 1,152 and 146 and patients without renal disease (LN-) were

1,949 and 378 in the SLE main cohort and SLEGEN cohort, respectively. More details are presented

in **Supplementary Table 1**.

Genome-wide association study (GWAS)

SLE GWAS

SLE GWASs were performed in genotyped SNPs including principal components consistent with the

original publications in all three independent cohorts.

SLE Renal GWAS within SLE cases

The SLE Renal GWASs were performed within SLE cases, i.e., genome-wide associations of patients

with renal disease (SLE Renal+, cases) and patients without renal disease (SLE Renal-, controls) in

two independent cohorts, i.e., the SLE main cohort and the SLEGEN cohort. For Renal GWASs, we

pre-phased the genotyped data using the SHAPEIT algorithm [21] and then used IMPUTE2 [22] to

impute to the density of the 1000 Genome reference data (phase 3 integrated set, release 20130502)

[23] (data unpublished). All case-control analysis was carried out using the SNPTEST algorithm [24].

SNPs with imputation INFO scores of < 0.7 and MAF (minor allele frequency) < 0.001 were removed.

After quality control (QC), there were 21,431,070 SNPs left for further analysis. Moreover, a genome-

wide association meta-analysis of the SLE main cohort and SLEGEN cohort was performed using the

summary statistics derived from the two Renal GWASs. A standard threshold of P ≤ 5e-08 was used

to report genome-wide significance and a P ≤ 1e-05 was used to report suggestive associated signals.

Polygenic analysis

We tested for non-zero standardised effect sizes (Z scores) for SLE association in the Genentech data

for groups of SNPs stratified by their *P* values in the SLE main cohort. The Z scores in the Genentech

data were polarized with respect to the SLE main cohort in that the effect allele was set to be the same

risk allele as in the SLE main cohort. Under the null hypothesis the Z scores will have zero mean, while

under the alternative the mean will be positive. SNPs were stratified by *P* value intervals of 1-0.9, 0.9-

0.8, 0.8-0.7, 0.7-0.6, 0.6-0.5, 0.5-0.4, 0.4-0.3, 0.3-0.2, 0.2-0.1, 0.1-0.00. We would expect a positive

mean for SNPs with very small *P* values in the main SLE data as these will be enriched for true positives,

while the same is not necessarily true over other *P* values ranges unless there are more widespread

true associations with very weak effects. We also ran this analysis on renal association standardised

effect sizes (Z scores) again polarised with respect to SLE association and stratified by SLE *P* values.

In all analyses, we used an LD clumped set of SNPs with an R^2 threshold of 0.1. When comparing the

SLE main cohort to the Genentech cohort or the SLEGEN cohort, we limited the clumping to SNPs that

overlap the GWASs.

Genetic risk score derivation

A Genetic risk score (GRS) is a quantitative trait of an individual's inherited risk based on the cumulative

impact of many genetic variants, which is calculated according to the method described by Hughes et

al [25], taking the number of risk alleles (i.e., 0, 1 or 2) for a given SNP and multiplying this by its

corresponding estimated effect - β coefficient, i.e. the natural log of its odds ratio (OR). The cumulative

risk score in each subject was calculated by summing the risk scores from the target risk loci:

132
$$\text{Genetic risk score} = \sum_i^n G_i \beta_i$$

where n represents the number of SLE risk loci, G_i is the number of risk alleles at a given SNP, and β_i

is the effect size of the risk SNP i.

We used two approaches to select SNPs for GRS calculation. The first approach – a weighted GRS

was derived from all published independent SLE risk SNPs (**Supplementary Table 2**) – including 78

SLE susceptibility loci (without the X chromosome), consisting of 93 SNPs outside of the MHC region

and 2 independent tag SNPs in the MHC region for two SLE associated HLA haplotypes. The risk allele

and its effect size for each SNP is derived from its original publication, which is summarized in a recent

review [11]. Each GRS for three SLE cohorts [18, 19, 26] was generated using R version 3.4.3.

The second approach – LD clumping and thresholding – was used to build 32 GRSs. Clumping and

thresholding scores were built using a *P* value and linkage disequilibrium (LD)-driven clumping

threshold in PLINK version 1.90b (www.cog-genomics.org/plink/1.9/) [27]. In brief, the algorithm forms

clumps around SNPs with association *P* values less than a provided threshold (Index SNPs). Each

clump contains all SNPs within a specified window of the index SNP that are also in LD with the index

SNP as determined by a provided pairwise correlation threshold (r^2) in the LD reference. The algorithm

loops through all index SNPs, beginning with the smallest *P* value and only allowing each SNP to appear

in one clump. The final output should contain the most significant disease-associated SNP for each

LD-based clump across the genome. Note that when performing LD clumping, we firstly removed the

X-chromosome and the MHC extended region (24-36MB) and kept all other autosomal SNPs. Then

we included the MHC region by using two tag SNPs for two well-known HLA haplotypes in SLE, i.e.

rs2187668 for HLA-DRB1*03:01 and rs9267992 for HLA-DRB1*15:01. A GRS was built using the

genotypes for the index SNPs weighted by the estimated effect sizes (β). Specifically, when training

the GRS in the SLE main cohort and testing in the SLEGEN cohort, we performed a GWAS on the

genotyped SNPs in the SLE main cohort and generated 32 lists of clumped SNPs over a set of P values

(--clump-p1: 0.1, 0.01, 1e-03, 1e-04, 1e-05, 1e-06, 1e-07,and 5e-08), r^2 *(--clump-r2:* 0.2 and 0,5) and

clumping radius *(--clump-kb*: 250 and 1000). The 32 list of SNPs were then used to generate 32 GRSs

by summing across all variants weighted by their respective effect size for samples in the SLEGEN

cohort. We performed this cross-validation in all three cohorts, generating six training-and-testing pairs.

Receiver Operating Characteristic (ROC) curves for model evaluation

The GRS with the best discriminative capacity was determined based on the maximal Area under the

ROC curve (AUC) with SLE or RENAL as the outcome and the candidate GRS as the predictor. AUC

confidence intervals were calculated using the '*pROC*' package within R and the difference between

the ROC curves was determined with the '*roc.test'* function, which used a non-parametric approach, as

described by De Long et al [28]. To assess the degree to which the age of SLE onset contributes to

the prediction of renal involvement within SLE cases, we generated ROCs as above with the GRS and

compared to ROC curves with SLE age onset as a single predictor and the ROC with both GRS and

age onset as predictor(s).

Partitioning the genetic risk of renal disease

Since a continuous score is difficult to interpret on an individual level when a physician needs to explain

the results of the GRS to a patient, we partitioned SLE patients into quintile according to genetic dosage

(SLE GRS). We used a chi-square test to study the association of the partitioned GRS and renal risk.

The odds ratios of renal risk were then calculated compared to the reference group - the first quintile

GRS group.

To test whether the GRS correlated with renal disease independently of age-of-onset, we partitioned

SLE patients into two groups according to their age of onset, with a cut-off at age of 30 - patients with

age above 30 were defined as 'Late age onset' and others as 'Early age onset'. A two-way ANOVA

test was then performed with the function '*aov*' in R, with *aov(GRS ~ age group * renal group)*. All

statistical analyses were conducted using R version 3.4.3 software (https://www.r-project.org/).

Results

The best GRS in SLE prediction

A GRS is a simple weighted sum of SLE risk alleles, however the choice of SNPs to use for the

calculation greatly effects its performance. We used three independent cohorts for cross-validation,

generating six training-and-testing pairs. For each training-and-testing pair of cohorts, we derived 32

predictors based on a clumping and thresholding method, and one additional predictor using the set of

SNPs that were previously reported to be associated with SLE (**Supplementary Table 2**). We then

evaluated the performance of the GRS as a predictor by its AUC. We found that the best GRS in the

prediction of SLE (with highest AUC) was the one derived from the published SLE SNPs (**Figure 1** &

Supplementary Table 3), with an AUC (95% CI) of 0.729 (0.706 - 0.753), 0.692 (0.673 - 0.71), and

0.745 (0.735 - 0.754) in SLE main cohort, SLEGEN cohort, and Genentech cohort, respectively. Among

the GRSs generated from LD clumping and thresholding, the predictor with the best discriminative

capacity was the one derived from SNPs clumping at P threshold (P_{th}) of 1e-05 in the SLE main cohort

and tested in both the SLEGEN and Genentech cohorts (**Figure 1** & **Supplementary Table 3**),

suggesting there may be more true positive signals than the genome-wide significant ones involved in

the risk of SLE. In fact, the predictive performance of the GRS using all pairs of training and test data

was maximised using SNPs below the standard genome-wide threshold (**Supplementary Table 3**).

This evidence for polygenicity was also seen in an analysis of the association statistics (Z scores) in

the Genentech GWAS polarised to the risk allele in the main GWAS, partitioned by their association P

value in the main GWAS (see methods). Here, we found evidence (**Figure 2** & **Supplementary Table**

4) against a zero mean (P = 3.91e-04) for the Z scores in Genentech data for SNPs with P values

between 0.3 and 0.2 in the main GWAS.

Lupus Nephritis GWAS within SLE cases

Lupus Nephritis (LN) occurs in approximately half of all SLE patients, and its frequency ranges from

25% to 75% depending on the population studied [29]. About one third of European SLE patients

experience renal disease [30]. Until recently, one of the most common causes of death in SLE patients

was kidney failure. According to the lupus severity index (LSI) using the ACR criteria developed by

Bello et al [31], renal involvement has the highest impact and particular strongly associated with disease

severity, hence we chose LN as a proxy of SLE severity in this study.

The within case LN GWAS in the SLE main cohort, which comprised 1152 SLE patients with renal

disease (LN+) and 1949 patients without renal disease (LN-), did not identify any genome-wide

significant associated loci ($P \leqslant$ 5e-08) (**Figure 3a**). Consistently, no inflation (genomic inflation factor:

λ = 1.014) was observed in the QQ plot (**Figure 3d**). Similarly, none of the SNPs reached genome-

wide significance in the SLEGEN cohort [19] (λ = 1.023) (**Figure 3b** & **3e**). In addition, no variant

passed genome-wide significance in the meta-analysis of the SLE main cohort and SLEGEN cohort for

Renal GWAS (λ = 0.9565) (**Figure 3c** & **3f**). Summary association statistics for SNPs with $P \leqslant$ 1e-05

are provided in **Supplementary tables 5 and 6**.

We did, however, see evidence that SNPs with very strong evidence for association with SLE (P $\leqslant$ 1e-

05) were associated with LN. This was evident from an analysis of the renal association statistics (Z

scores) polarised to the risk allele for SLE. There was strong evidence (**Figure 2** & **Supplementary**

Table 4, P = 8.72e-08) against a zero mean for the Renal Z scores for SNPs with P $\leqslant$ 1e-05 for SLE

in the main cohort. This result was replicated in the SLEGEN study with P = 2.42e-03 (**Figure 2** &

Supplementary Table 4). We only found evidence of renal association with SNPs showing very strong

evidence for association with SLE. This finding could be exploited for prediction of disease progression

and we explore this below.

Genetic risk loading of SLE is significantly higher in LN+ patients

While we observed that no individual SNPs were significantly associated with renal involvement in the

SLE cases, we did show that there was a deviation from zero mean for renal Z scores taken from SNPs

with very strong evidence for association with SLE. In view of this finding, we investigated the

correlation between the SLE GRS and renal disease in all SLE cases. To accomplish this, we used the

best GRS derived from a list of published SLE associated SNPs [11] for the comparison of the SLE

genetic risk burden in patients with and without renal disease. As expected, the GRS was higher in the

SLE patients compared to healthy controls in both independent cohorts (**Figure 4**).

A significantly higher GRS was observed in the group of patients with renal disease (LN+) compared to

patients without renal disease (LN-) (**Figure 4**). In the SLE main cohort, the mean (SD) of the GRS

was 18.1 (1.64) for LN+ patients and 17.8 (1.65) for LN- patients (P = 1.60e-07); the mean for the

SLEGEN cohort was 18.2 (1.66) for LN+ patients and 17.6 (1.69) for LN- patients (P = 0.0010).

Moreover, we saw a significant increasing trend of GRS over levels of diseases: Healthy control, LN-

patients, and LN+ patients, with a trend P < 1.0e-400 in the SLE main cohort and a P = 3.81e-73 in the

SLEGEN cohort (**Figure 4).**

Genetic risk of nephritis and age of onset in SLE

We partitioned the SLE cases into five groups according to quintiles for GRS to show the risk of renal

involvement. We observed over 1.5 folds higher risk of renal disease (OR = 1.58; 95% CI: 1.25 to 1.99;

P = 0.00015) between the top and bottom quintiles of GRS in the SLE main cohort (**Figure 5a**). This is

replicated in the SLEGEN cohort (**Figure 5b**), with odds ratios of 3.16 (95% CI: 1.62 to 6.13; P =

0.00091). A significantly earlier age of SLE onset was observed in those with renal disease compared

to those without renal disease. In the main cohort (**Figure 6a**), the mean (SD) for age of disease onset

was 29yrs (12) for LN+ patients and 35yrs (13) for LN- patients (P = 2.8e-27); the means for the

SLEGEN cohort (**Figure 6b**) were 28yrs (11) and 35yrs (13) for LN+ and LN-, respectively (P = 6.05e-

09). When testing the association of GRS with age of onset in the SLE main cohort, a significant

correlation was present – the higher the GRS, the earlier age of SLE onset (P = 2.4e-07). This

correlation was also detected in the SLEGEN cohort (P = 0.021).

To test whether the GRS correlated with renal disease independently of age-of-onset, we partitioned

SLE patients into two groups according to their age of onset, i.e. 'Late age onset' and 'Early age onset'

and performed a two-way ANOVA test (See Methods). The GRS was shown to positively correlate with

both renal disease and early age-of-onset (P_{Renal} = 7.64 × 10^{-5} and $P_{age-of-onset}$ = 1.06 × 10^{-9} in the SLE

main cohort; P_{Renal} = 0.0288 and $P_{age-of-onset}$ = 0.0513 in SLEGEN cohort), while we found that there was

no statistically significant interaction between renal and early age-of-onset in either the SLE main cohort

($P_{Interaction}$ = 0.795) or the SLEGEN cohort ($P_{Interaction}$ = 0.0511) (**Supplementary Figure 1**). Notably, we

found that GRS was a better predictor of renal disease in the 'Late age onset' group (AUC = 0.621)

compared with the 'Early age onset' group (**Figure 7**).

Finally, we assessed the predictive ability of the partitioned SLE GRS (quintile GRS, see methods) over

the two age-of-onset groups. In the main SLE cohort there is a clear and significant risk effect for renal

involvement with increasing GRS in the 'Late age of onset' group, but no significant effect in the early

onset group. We observed over two fold higher risk of renal disease (OR = 2.33; 95% CI: 1.567 to

3.471; P = 3.762-05) between the upper fourth quintile and the bottom quintile in the 'Late age onset'

group in the SLE main cohort (**Figure 5a**). The results were similar in the SLEGEN cohort, with the risk

of renal disease between the top and bottom quintile of GRS being over five times (OR = 5.484; 95%

CI: 1.647 to 18.26; P = 0.006635) (**Figure 5b** & **Supplementary Table 7**) in patients of 'Late age onset'

but no significant differences in those with 'Early age onset'.

Discussion

GRS has been showed to be predictive for several diseases including cardiovascular disease

(AUC=0.81, 95%CI: 0.81-0.81) [12], inflammatory bowel disease (AUC=0.63, 95%CI: 0.62–0.64) [12]

and breast cancer (AUC=0.63, 95%CI: 0.63-0.65) [32]. However, in many of these applications the AUC

values are dependent on inclusion of age and sex for prediction and so the AUC due to genetics alone

would have been substantially lower [33]. We have shown that a SLE GRS using only SNPs has good

predictive power with AUC approaching 0.7. Our results, using three independent GWASs, shows that

a GRS using SNPs with association P values well below genome-wide levels of significance has the

best predictive performance. This is further evidence that SLE is a polygenic disease with many risk

variants as yet undiscovered, and that more powerful studies could lead to useful predictive models.

Genetic risk scores may also have utility in prediction of disease severity and we find evidence for this

to be so for SLE. Our data show that renal involvement is not related to specific genetic factors or

particular genes but simply to genetic load of risk alleles.

Until recently, the most common cause of death in SLE patients was kidney failure. Though the

frequency of death from kidney disease has decreased sharply due to better therapies (e.g. dialysis

and kidney transplantation), kidney failure is still potentially fatal in some people with SLE and causes

significant morbidity. According to the lupus severity index (LSI) using the ACR criteria developed by

Bello et al [31], renal involvement had the highest impact and particularly more strongly associated with

disease severity, hence we used renal involvement as a proxy of SLE severity in this study.

In the SLE within-case renal GWASs, we observed no genome-wide significant signals in either the

SLE main cohort or the SLEGEN cohort, or meta-analysis of these two. Both datasets had genetic

variants with less stringent P values ($P \leqslant$ 1e-05) for renal association, but none of them were replicated

in the other cohort. Considering the sample size of both cohorts are relatively small, we applied an

online genetic power calculator (http://zzz.bwh.harvard.edu/gpc/) to calculate the power of our current

sample size for the GWAS study (**Supplementary Table 8**). We assumed the effect sizes of SLE renal

risk alleles is similar to that seen in SLE GWAS, so the odds ratio (OR) of the risk allele would be

between 1.0 and 2.0. Therefore, we calculated power under a variety of parameters, including OR, risk

allele frequency (RAF) and alpha. As showed in **Supplementary Table 8**, we have a power of $\geqslant$ 0.8

to detect a genetic risk variant with an OR = 1.4 and RAF = 0.3 or an OR = 1.5 and RAF = 0.2 when

alpha = 5e-08. However, if we assume the renal associated variants are as weak as most of the SLE

associated variants (OR < 1.2), then we are under powered (< 0.8) to detect the true renal associations

at the GWAS significant threshold of P = 5e-08 in the current study.

We did however find evidence that SNPs most associated with SLE (P < 1e-05) were enriched for

associations with SLE renal and so we then tested the hypothesis that the genetic risk loading of SLE

may correlate with kidney involvement. Therefore, a genetic risk score (GRS) with the best performance

in SLE prediction was derived for the prediction of SLE renal disease. In both European cohorts, the

SLE main cohort and the SLEGEN cohort, the GRS was significantly higher in patients with renal

disease than patients without. In addition, patients with a higher GRS were more likely to have renal

involvement at a younger age, indicating the strong genetic background of SLE development. These

findings provide more evidence to support the opinion that younger-age onset lupus is generally more

severe than older-onset lupus as reported previously [34-36].

One may argue that if the severity of SLE is driven by multiple genes' contribution in a quantitative way,

the more risk alleles that are added to the model, the better the model would fit. In this study, we show

that a GRS is a useful tool for the classification of SLE renal+ and SLE renal- groups. The renal

association *P* values of the 95 SNPs (of 77 SLE risk loci) in the SLE main cohort and the SLEGEN

cohort are strongly inflated as shown in the QQ plots (**Supplementary Figure 2**), suggesting the

cumulative genetic burden from multiple SLE risk genes with modest effect.

Our analysis of Renal disease in SLE patients has shown that, while we find no SNPs significantly

associated with renal disease, the fact that SLE associated variants correlated with renal using a GRS

suggests that many SLE associated variants are also risk for renal involvement albeit with likely weaker

effects (Odds ratios). We find that the GRS and age-of-onset are correlated but the GRS is associated

with renal involvement independently of age-of-onset with no interaction observed. The GRS performs

better for predicting renal disease in patients with late age-of-onset (> 30 years). We also find that a

stratified GRS may be a more viable option for predicting renal disease, where we estimate significantly

high relative risks for those in the tails of the GRS distribution in both of our studies that had renal data.

This is the first study to investigate accumulated genetic risk and its relationship with the susceptibility

and severity of SLE. We found that the higher the GRS, the younger onset of SLE. In patients of late

onset, a higher GRS means patients are more likely to suffer from more severe disease. In brief, age

of onset incorporating a GRS may assist early prediction of lupus nephritis in a clinical setting.

Nevertheless, more clinical studies are needed to validate the usefulness of this application.

Acknowledgements The authors would like to thank Dr. Amy W.L Bulter and Mr. Akmal Droubi for

excellent clinical data administration on both SLE cohorts. This work was supported by the National

Institute for Health Research Biomedical Research Centre (NIHR BRC) at Guy's and St Thomas' NHS

Foundation and King's College London.

Funding This study is support by China Scholarship Council (CSC) and Medical Research Council

(MRC).

Competing interests None declared.

Patients consent Obtained.

Ethics approval Each participating centre has obtained approval from the local ethics committee for

including a patient's data in the SLE cohort after the patient has given written informed consent.

Provenance and peer review Not commissioned; externally peer reviewed.

References

1. Kaul, A., et al., *Systemic lupus erythematosus.* Nat Rev Dis Primers, 2016. **2**: p. 16039.
2. Yen, E.Y. and R.R. Singh, *Brief Report: Lupus-An Unrecognized Leading Cause of Death in Young Females: A Population-Based Study Using Nationwide Death Certificates, 2000-2015.* Arthritis Rheumatol, 2018. **70**(8): p. 1251-1255.
3. Davidson, A., *What is damaging the kidney in lupus nephritis?* Nat Rev Rheumatol, 2016. **12**(3): p. 143-53.
4. Feldman, C.H., et al., *Epidemiology and sociodemographics of systemic lupus erythematosus and lupus nephritis among US adults with Medicaid coverage, 2000-2004.* Arthritis Rheum, 2013. **65**(3): p. 753-63.
5. Peschken, C.A., et al., *The 1000 Canadian faces of lupus: determinants of disease outcome in a large multiethnic cohort.* J Rheumatol, 2009. **36**(6): p. 1200-8.
6. Pons-Estel, B.A., et al., *The GLADEL multinational Latin American prospective inception cohort of 1,214 patients with systemic lupus erythematosus: ethnic and disease heterogeneity among "Hispanics".* Medicine (Baltimore), 2004. **83**(1): p. 1-17.
7. Alarcon, G.S., et al., *Baseline characteristics of a multiethnic lupus cohort: PROFILE.* Lupus, 2002. **11**(2): p. 95-101.
8. Lawrence, J.S., C.L. Martins, and G.L. Drake, *A family survey of lupus erythematosus. 1. Heritability.* J Rheumatol, 1987. **14**(5): p. 913-21.
9. Kuo, C.F., et al., *Familial Aggregation of Systemic Lupus Erythematosus and Coaggregation of Autoimmune Diseases in Affected Families.* JAMA Intern Med, 2015. **175**(9): p. 1518-26.
10. Deapen, D., et al., *A revised estimate of twin concordance in systemic lupus erythematosus.* Arthritis Rheum, 1992. **35**(3): p. 311-8.
11. Chen, L., D.L. Morris, and T.J. Vyse, *Genetic advances in systemic lupus erythematosus: an update.* Curr Opin Rheumatol, 2017.
12. Khera, A.V., et al., *Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations.* Nat Genet, 2018. **50**(9): p. 1219-1224.
13. Euesden, J., C.M. Lewis, and P.F. O'Reilly, *PRSice: Polygenic Risk Score software.* Bioinformatics, 2015. **31**(9): p. 1466-8.
14. Belsky, D.W., et al., *Development and evaluation of a genetic risk score for obesity.* Biodemography Soc Biol, 2013. **59**(1): p. 85-100.
15. De Jager, P.L., et al., *Integration of genetic risk factors into a clinical algorithm for multiple sclerosis susceptibility: a weighted genetic risk score.* The Lancet Neurology, 2009. **8**(12): p. 1111-1119.
16. Karlson, E.W., et al., *Cumulative association of 22 genetic variants with seropositive rheumatoid arthritis risk.* Ann Rheum Dis, 2010. **69**(6): p. 1077-85.
17. Yarwood, A., et al., *A weighted genetic risk score using all known susceptibility variants to estimate rheumatoid arthritis risk.* Ann Rheum Dis, 2015. **74**(1): p. 170-6.
18. Bentham, J., et al., *Genetic association analyses implicate aberrant regulation of innate and adaptive immunity genes in the pathogenesis of systemic lupus erythematosus.* Nat Genet, 2015. **47**(12): p. 1457-1464.
19. International Consortium for Systemic Lupus Erythematosus, G., et al., *Genome-wide association scan in women with systemic lupus erythematosus identifies susceptibility variants in ITGAM, PXK, KIAA1542 and other loci.* Nat Genet, 2008. **40**(2): p. 204-10.
20. Hom, G., et al., *Association of systemic lupus erythematosus with C8orf13-BLK and ITGAM-ITGAX.* N Engl J Med, 2008. **358**(9): p. 900-9.
21. O'Connell, J., et al., *Haplotype estimation for biobank-scale data sets.* Nat Genet, 2016. **48**(7): p. 817-20.
22. Howie, B.N., P. Donnelly, and J. Marchini, *A flexible and accurate genotype imputation method for the next generation of genome-wide association studies.* PLoS Genet, 2009. **5**(6): p. e1000529.
23. Genomes Project, C., et al., *A global reference for human genetic variation.* Nature, 2015. **526**(7571): p. 68-74.
24. Marchini, J., et al., *A new multipoint method for genome-wide association studies by imputation of genotypes.* Nat Genet, 2007. **39**(7): p. 906-13.
25. Hughes, T., et al., *Analysis of autosomal genes reveals gene-sex interactions and higher total genetic risk in men with systemic lupus erythematosus.* Ann Rheum Dis, 2012. **71**(5): p. 694-9.

26.   Yang, W., et al., *Genome-wide association study in Asian populations identifies variants in*
ETS1 and WDFY4 associated with systemic lupus erythematosus. PLoS Genet, 2010. **6**(2): p.
e1000841.
27.   Chang, C.C., et al., *Second-generation PLINK: rising to the challenge of larger and richer*
datasets. Gigascience, 2015. **4**: p. 7.
28.   DeLong, E.R., D.M. DeLong, and D.L. Clarke-Pearson, *Comparing the areas under two or more*
correlated receiver operating characteristic curves: a nonparametric approach. Biometrics,
1988. **44**(3): p. 837-45.
29.   Cervera, R., et al., *Morbidity and mortality in systemic lupus erythematosus during a 10-year*
period - A comparison of early and late manifestations in a cohort of 1,000 patients. Medicine,
2003. **82**(5): p. 299-308.
30.   Font, J., et al., *Clusters of clinical and immunologic features in systemic lupus erythematosus:*
Analysis of 600 patients from a single center. Seminars in Arthritis and Rheumatism, 2004.
33(4): p. 217-230.
31.   Bello, G.A., et al., *Development and validation of a simple lupus severity index using ACR*
criteria for classification of SLE. Lupus Sci Med, 2016. **3**(1): p. e000136.
32.   Mavaddat, N., et al., *Polygenic Risk Scores for Prediction of Breast Cancer and Breast Cancer*
Subtypes. Am J Hum Genet, 2019. **104**(1): p. 21-34.
33.   Curtis, D., *Clinical relevance of genome-wide polygenic score may be less than claimed.* Ann
Hum Genet, 2019: p. 1-4.
34.   Lazaro, D., *Elderly-onset systemic lupus erythematosus: prevalence, clinical course and*
treatment. Drugs Aging, 2007. **24**(9): p. 701-15.
35.   Janwityanujit, S., et al., *Age-related differences on clinical and immunological manifestations*
of SLE. Asian Pac J Allergy Immunol, 1995. **13**(2): p. 145-9.
36.   Tomic-Lucic, A., et al., *Late-onset systemic lupus erythematosus: clinical features, course, and*
prognosis. Clin Rheumatol, 2013. **32**(7): p. 1053-8.
437

Figure 1. ROCs and AUCs of models in SLE prediction in European cohorts

GRSs for the prediction of SLE in the SLEGEN cohort (**a**) and Genentech cohort (**b**) were generated from SNPs of LD clumping and threshold derived from the SLE main cohort, and a list of published SLE risk SNPs (**Supplementary Table 2**). 'GRS at Pth' represented the GRS in the SLE prediction model was derived from the LD clumping at the according GWAS *P* value threshold.

Figure 2. Polygenic test of SLE and Renal disease.

Polygenic test of SLE in Genentech cohort (**a** and **b**) and polygenic test of Renal disease in the SLE main cohort (**c** and **d**) and SLEGEN cohort (**e** and **f**). The SLE main cohort is used to generate *P* value for each SNP to stratify the SNPs into groups for the *Z* score calculation of SLE association or Renal association.

Figure 3. Genome-wide scans of LN associated variants.

(Upper) Manhattan plots showing the -log10-transformed p values (y axis) against physical genomic

position (x axis) for each SNP in the SLE main cohort (**a**), the SLEGEN cohort (**b**), and the meta-

analysis of these two cohorts (**c**). The red horizontal line represents the threshold for genome-wide

significance (P $\leq$ 5e-08) and the blue horizontal line represents the threshold for suggestive

significance (P $\leq$ 1e-05).

(Lower) Quantile-quantile plots showing the observed distribution of -log10-transformed p values (y

axis) by the expected distribution (x axis) under the null hypothesis of no association (diagonal line)

for the SLE main cohort (genomic inflation factor, λ = 1.014) (**d**), the SLEGEN cohort (λ = 1.023) (**e**),

and the meta-analysis of these two cohorts (λ = 0.9565) (**f**).

Figure 4. GRS over levels of disease: Controls / SLE Renal (-) / SLE Renal (+).

The violin-and-box plots show the summary GRS for each level of the disease in the SLE main cohort

(**a**) and the SLEGEN cohort (**b**). The violins show the distribution of the GRS across each group. The

bottom line of the box inside the violin is the 1st quantile, the top line is the 3rd quantile, and the box

is divided at the median. Sample size (N) of each group is showed within brackets below the group

name. Note that GRS for SLE main cohort and SLEGEN cohort are generated by 93 non-MHC SNPs

and 2 MHC tag SNPs - a total of 95 SNPs.

Figure 5. Relationship of quintiles of the GRS and risk of renal disease within SLE patients.

Plots show the odds ratios of Renal disease for the SLE main cohort (**a**) and the SLEGEN cohort (**b**),

comparing each of the upper four GRS quintiles with the lowest quintile; dotted lines represent the

95% confidence intervals; horizontal black dotted lines represent OR = 1.

Figure 6. Age of SLE onset in patients of Renal(-) / Renal(+).

The violin-and-box plots show the age of SLE onset for each level of the disease in the SLE main

cohort (**a**) and the SLEGEN cohort (**b**). The violins show the distribution of the Age of SLE onset

across each group. The bottom line of the box inside the violin is the 1st quantile, the top line is the

3rd quantile, and the box is divided at the median. Sample size (N) of each group is showed within

brackets below the group name.

ROCs: Train in SLE.main.cohort, Predict in SLEGEN.cohort

AUC of Renal Prediction by GRS
■ All Ageonset SLE, AUC = 0.576 (0.518 – 0.634)
■ Early Ageonset SLE, AUC = 0.486 (0.404 – 0.568)
■ Late Ageonset SLE, AUC = 0.621 (0.527 – 0.714)

Figure 7. ROC Curves for models predicting a diagnosis of Renal disease in SLE patients using GRS, split
by age-of-onset.
The models were trained in the SLE main cohort and tested in the SLEGEN cohort. The plots
showed the ROC curves in the prediction of renal disease in SLE patients with GRS as a predictor,
The ROC curve in black was trained and tested with all SLE samples, the purple curve was trained
and tested in the 'Early age onset' patients, and the red curve was trained and tested in the 'Late age
onset' group. AUC, area under the ROC curve is showed with 95% CI in brackets.
488

Supplementary Figure 1. Relationship of GRS and age onset in Renal disease.

The age of SLE onset <= 30 years was defined as "Early onset" and > 30 years was defined as "Late

onset". For each age onset and renal group, the GRS was plotted with mean and 95% CI for the SLE

main cohort.

Supplementary Figure 2 . Quantile-quantile plots of Renal association results

QQ plots showed the observed distribution of -log10-transformed p values (y axis) by the expected

distribution (x axis) under the null hypothesis of no association (diagonal line) for the SLE main cohort

(**a**) and the SLEGEN cohort (**b**). The *P* values for the QQ plots were derived from Renal association

test of the 95 SNPs (of 77 SLE risk loci) which used for the GRS calculation.

499