CCDC186 recruits EARP and controls dense-core vesicle cargo sorting by exit

Jérôme Cattin-Ortolá[1†‡], Irini Topalidou[1†], Ho-Tak Lau[2], Blake Hummer[3], Cedric S. Asensio[3], Shao-En Ong[2] and Michael Ailion[1*]

[1]Department of Biochemistry, University of Washington, Seattle, United States

[2]Department of Pharmacology, University of Washington, Seattle, United States

[3]Department of Biological Sciences, University of Denver, Denver, United States

[†]These authors contributed equally to this work.

[‡]Present address: MRC Laboratory of Molecular Biology, Cambridge, United Kingdom

*Correspondence: mailion@uw.edu, Twitter: @michaelailion

Running Title: CCDC186 is a DCV cargo-sorting factor

Key words: coiled-coil, dense-core vesicle, regulated secretion, insulin.

ABSTRACT

The regulated release of peptide hormones depends on their packaging into dense-core vesicles (DCVs). Two models have been proposed for DCV cargo sorting. The "sorting by entry" model proposes that DCV cargos selectively enter nascent DCVs at the trans-Golgi network (TGN). The "sorting by exit" model proposes that sorting occurs by the post-TGN removal of non-DCV cargos and retention of mature DCV cargos. Here we show that the coiled-coil protein CCDC186 controls sorting by exit. *Ccdc186* KO insulinoma cells secrete less insulin, fail to retain insulin and carboxypeptidase E in mature DCVs at the cell periphery, and fail to remove carboxypeptidase D from immature DCVs. A mutation affecting the endosome-associated recycling protein (EARP) complex causes similar defects in DCV cargo sorting, and overexpression of CCDC186 recruits EARP to the site of DCV maturation. CCDC186 and EARP may act together to control the post-Golgi retention of cargos in mature DCVs.

INTRODUCTION

Neurons and endocrine cells contain specialized secretory vesicles called dense-core vesicles (DCVs) that undergo regulated secretion in response to external stimuli. DCVs contain neuropeptides, growth factors, biogenic amines, and peptide hormones that control numerous biological processes, ranging from growth and metabolism to mental state. DCVs are generated at the trans-Golgi network (TGN) as immature DCVs (iDCVs) and undergo a maturation process that includes vesicle acidification, proteolytic processing of cargos, and the acquisition of proper compartmental identity and release competency (Borgonovo et al., 2006; Gondré-Lewis et al., 2012; Kim et al., 2006; Tooze et al., 2001). Peptide precursors are processed in acidic iDCVs by several distinct processing enzymes. For example, the insulin precursor proinsulin is processed into mature insulin via two proprotein convertase enzymes (PC1/3 and PC2) and carboxypeptidase E (CPE) (Steiner et al., 2011).

Two models of DCV cargo sorting have been proposed. The 'sorting by entry' model proposes that cargos are sorted in the trans-Golgi and preferentially enter nascent DCVs as they bud off; the 'sorting by exit' model proposes that cargos are sorted in a post-Golgi step in which non-DCV cargos are removed from iDCVs while mature DCV cargos are retained (Arvan and Castle, 1998; Tooze, 1998). Both mechanisms likely contribute to DCV cargo sorting. Cargos removed from iDCVs include the transmembrane proteins carboxypeptidase D (CPD) and the mannose 6-phosphate receptors (CD-MPR and CI-MPR) (Klumperman et al., 1998; Varlamov et al., 1999).

Genetic studies in the nematode *C. elegans* have identified several molecules that function in neuronal DCV biogenesis, including the small G protein RAB-2, the RAB-2 effector CCCP-1 (the *C. elegans* homolog of CCDC186), the endosome-associated recycling protein (EARP) complex (VPS-50, VPS-51, VPS-52, and VPS-53), and the EARP interactor EIPR-1 (Ailion et al., 2014; Edwards et al., 2009; Paquin et al., 2016; Sumakovic et al., 2009; Topalidou et al., 2016). Loss of any of these proteins causes a similar reduction in the levels of cargos in

mature DCVs in axons, but it is unclear how RAB-2 and CCCP-1 are connected to EARP. In both *C. elegans* and in the rat insulinoma 832/13 cell line, RAB-2 and CCCP-1 colocalize near the TGN where the early steps of DCV biogenesis take place (Ailion et al., 2014; Cattin-Ortolá et al., 2017). We showed recently that EARP is localized to two distinct compartments in 832/13 cells, a recycling endosome compartment and a compartment near the TGN where it colocalizes with CCDC186 (Topalidou et al., 2018). Additionally, *Eipr1* knockout 832/13 cells have reduced insulin secretion and an altered distribution of insulin in the cell, with relatively more insulin near the TGN and less at the cell periphery (Topalidou et al., 2018). Thus, EARP and EIPR1 may be important for sorting cargos to DCVs near their site of generation at the TGN.

Here we investigate the role of CCDC186 in DCV biogenesis, maturation, and cargo sorting in 832/13 cells. Our results suggest that CCDC186 acts together with EARP to control several aspects of DCV cargo sorting, including the post-Golgi removal of some non-DCV cargos and retention of mature DCV cargos.

RESULTS

CCCP-1 and EARP act in the same genetic pathway

C. elegans mutants in *rab-2, cccp-1, eipr-1,* and in all four subunits of the EARP complex (*vps-50, vps-51, vps-52, vps-53*) have a slow locomotion rate and reduced axonal levels of DCV cargos (Ailion et al., 2014; Edwards et al., 2009; Paquin et al., 2016; Sumakovic et al., 2009; Topalidou et al., 2016). CCCP-1, EIPR-1, and EARP all act in the RAB-2 genetic pathway (Ailion et al., 2014; Topalidou et al., 2016). Though EIPR-1 and EARP act together in a common pathway (Topalidou et al., 2016), it is unknown whether CCCP-1 and EIPR-1/EARP act in the same pathway or if they act in parallel pathways that converge with RAB-2. Thus, we built double mutants between *cccp-1* and either *eipr-1* or *vps-50* and assayed their locomotion rate (Fig S1A) and the axonal levels of the DCV luminal cargo NLP-21 tagged to Venus (Fig S1B,C). In both cases, the double mutants have phenotypes similar to the single mutants (Fig S1A-C), indicating that CCCP-1 and EIPR-1/EARP act in the same genetic pathway to control locomotion behavior and cargo sorting to DCVs.

CCDC186 is required for normal insulin secretion

The mammalian CCCP-1 homolog is known as CCDC186 or C10orf118. Using an antibody to human CCDC186, we found that CCDC186 is expressed in multiple tissues in mice, including brain, heart, liver, kidney, and spleen (Fig S2A). Additionally, CCDC186 is expressed in endocrine cell lines such as PC12 cells and rat insulinoma 832/13 cells (Cattin-Ortolá et al. 2017 and Fig S2A).

To investigate the role of CCDC186 in DCV biogenesis in insulin-secreting cells, we generated *Ccdc186* knock out (KO) 832/13 cells using the CRISPR/Cas9 technology. The absence of CCDC186 in the *Ccdc186* KO line was confirmed by Western blotting, RT-PCR, and immunofluorescence (Fig S2B-D). We measured the insulin secretion of wild type (WT) and *Ccdc186* KO cells under resting (5 mM KCl, no glucose) and stimulating (55 mM KCl, 25 mM

glucose) conditions. Though insulin secretion under resting conditions was not noticeably affected by the absence of CCDC186, stimulated insulin secretion was significantly reduced (~ 49% of WT; Fig 1A, left). We recently reported a similar reduction in insulin secretion from *Eipr1* KO cells (Topalidou et al., 2018). To verify that the effect was specifically due to loss of CCDC186, we introduced a wild type *Ccdc186* cDNA into the *Ccdc186* KO cells by lentiviral transduction (Fig S2B,D). Expression of wild type *Ccdc186* in *Ccdc186* KO cells (*Ccdc186*(+)) rescued the insulin secretion defect of the *Ccdc186* KO line, indicating that the insulin secretion defect is indeed due to loss of CCDC186 (Fig 1A, left).

The defective insulin secretion could be due to reduced insulin content. However, we found that the total insulin level was not significantly altered in *Ccdc186* KO cells (Fig 1A, middle). After normalizing resting and stimulated insulin secretion to total insulin content, we found that stimulated insulin secretion was still significantly reduced in the *Ccdc186* KO line (~ 41% of WT; Fig 1A, right).

The decrease in insulin secretion in the *Ccdc186* KO could be due to a defect in the processing of proinsulin to insulin. Thus, we measured the levels of secreted and cellular proinsulin. We found that proinsulin secretion is increased slightly in *Ccdc186* KO cells but the difference was not statistically significant (Fig 1B, left). The level of cellular proinsulin was also slightly increased (Fig 1B, middle), but again the difference was not statistically significant and the proinsulin secretion did not change significantly after normalizing to total proinsulin content (Fig 1B, right). Additionally, the ratio of total cellular proinsulin to insulin was unchanged in the *Ccdc186* KO cells (Fig 1C). Thus, loss of CCDC186 has no significant effect on the processing of proinsulin to insulin.

Because CCDC186 is expressed in tissues not known to have DCVs (Fig S2A), we asked whether CCDC186 is also required for constitutive secretion. To assay constitutive secretion, we measured the secretion of GFP fused to a signal sequence (ssGFP) under resting conditions (Hummer et al., 2017). We found no difference between WT and *Ccdc186* KO cells

(Fig 1D), suggesting that CCDC186 is required for normal levels of regulated but not constitutive secretion.

CCDC186 is required for the normal distribution of insulin in cells

We have shown that EIPR1 is needed for the proper cellular distribution of insulin in 832/13 cells (Topalidou et al., 2018). To investigate whether CCDC186 is similarly required, we examined the subcellular localization of insulin and proinsulin in *Ccdc186* KO cells. In wild type cells, insulin is found as puncta spread throughout the cytoplasm and the cell periphery (Fig 2A). By contrast, in *Ccdc186* KO cells insulin is localized primarily to a perinuclear region that partially colocalizes with the trans-Golgi (TGN) marker TGN38 (Fig 2A,B). The insulin localization phenotype was fully rescued in *Ccdc186* KO cells that stably express wild type *Ccdc186* (Fig 2A,B). Using a proinsulin-specific antibody, we found that proinsulin is localized near the TGN and not at the cell periphery in both wild type and *Ccdc186* KO cells (Fig 2E,F). Thus, loss of CCDC186 does not affect the localization of proinsulin, the total cellular levels of insulin or proinsulin, or the ratio of proinsulin/insulin, indicating that CCDC186 acts after the processing of proinsulin. By contrast, loss of CCDC186 has a strong effect on the cellular distribution of mature insulin.

Normally, proinsulin is processed into insulin only in post-Golgi compartments because the processing enzymes require an acidic pH for their activity (Davidson et al., 1988; Orci et al., 1986, 1987). Because there is increased colocalization of insulin with TGN38 in *Ccdc186* KO cells, we wondered whether there might be increased acidification of the Golgi in these mutant cells which would allow processing of proinsulin into insulin in this compartment. To measure the pH of the Golgi, we measured the fluorescence of the pH-sensitive GFP variant pHluorin directed to the Golgi by fusing it to the trans-Golgi-localized transmembrane domain of beta-galactoside alpha 2,6-sialyltransferase (Hummer et al., 2017; Wong et al., 1992). In *Ccdc186* KO cells, there was little change in the pH of the Golgi and if anything a slight increase (Fig S3).

This is inconsistent with increased acidification of the Golgi and the processing of proinsulin to insulin in the Golgi in these mutant cells. Thus, mature insulin is unlikely to be found in the TGN in the *Ccdc186* KO cells. The insulin signal that partially colocalizes with the TGN marker in our immunofluorescence experiments may represent mature insulin near the TGN in a post-TGN acidified compartment.

We used two methods to assess whether insulin is retained near the TGN in *Ccdc186* KO cells. First, we measured the ratio of TGN38-colocalized insulin to the level of insulin in the rest of the cell. This ratio increases in *Ccdc186* KO cells (Fig 2C), consistent with retention of insulin near the Golgi, a decrease in insulin in the cell periphery, or both. To measure just the level of insulin near the Golgi, we quantified the colocalization between insulin and TGN38 specifically in the TGN38-positive perinuclear area. The level of TGN38-colocalized insulin was slightly increased, though the effect was not statistically significant due to high variability in this measurement (Fig 2D). Thus, the reduced secretion of insulin is probably due mainly to the reduced levels of mature insulin in DCVs located at the cell periphery.

CCDC186 is required for the normal levels and distribution of mature DCV cargos

The processing of proinsulin to insulin is mediated by several processing enzymes, including the proprotein convertases PC1/3 (PCSK1) and carboxypeptidase E (CPE) (Naggert et al., 1995; Smeekens et al., 1992). PC1/3 and CPE are themselves processed from their immature pro forms to their enzymatically active mature forms, which are retained in mature DCVs (Fricker and Devi, 1993; Smeekens and Steiner, 1990; Song and Fricker, 1995; Vindrola and Lindberg, 1992; Zhou and Lindberg, 1993). We recently showed that EIPR1 is needed for the normal cellular levels of mature PC1/3 in 832/13 cells (Topalidou et al., 2018). Similarly, we found that loss of CCDC186 resulted in a reduced level of mature PC1/3 and this phenotype was rescued in *Ccdc186*(+) cells (Fig 3A). Also, the level of the unprocessed pro form of CPE (pro-CPE; see Materials and Methods) was not altered in *Ccdc186* KO cells (Fig 3B), but the

level of the mature processed form of CPE (recognized by the antibodies CPE-N and CPE-C) was reduced (Fig 3C,D). These results suggest that CCDC186 acts after the processing of pro-CPE.

Similar to insulin, mature CPE is found as puncta spread throughout the cytoplasm in wild type cells (Fig 3E and Fig S4). These CPE puncta almost completely colocalize with insulin, indicating that CPE and insulin are in the same vesicles (Fig S4). By contrast, in *Ccdc186* KO cells, mature CPE is localized primarily to a perinuclear region that partially colocalizes with TGN38 and this phenotype was fully rescued in *Ccdc186(+)* cells (Fig 3E,F). The redistribution of CPE in *Ccdc186* KO cells appears to be due to decreased levels of mature CPE in vesicles in the cell periphery and increased retention of CPE in or near the TGN (Fig 3G,H). In contrast to mature CPE and similar to proinsulin, pro-CPE is localized near the TGN in both WT and *Ccdc186* KO cells (Fig 3I,J). These results suggest that CCDC186 acts at a post-Golgi level and that it is needed for the proper levels or cellular distribution of several mature DCV cargos, including insulin and the insulin processing enzymes PC1/3 and CPE.

To investigate the kinetics of DCV cargo release from the Golgi, we used a pulse-chase method to monitor the exit of the exogenous DCV cargo ANF::GFP from the TGN (Kögel et al., 2013; Kögel Tanja et al., 2010). At steady state, ANF::GFP is distributed throughout the cytoplasm in WT cells, but is localized relatively more near the TGN in *Ccdc186* KO cells, similar to endogenous insulin (Fig S5A). ANF::GFP exit from the TGN was blocked by incubation at 20°C (pulse) and the synthesis of new protein was blocked by cycloheximide (Fig S5B). After the low-temperature block, cells were brought back to 37ºC (chase) and incubated for different times before fixation (Fig S5B). Cells were then counted blindly based on whether ANF::GFP was (1) localized primarily at the TGN-region ("Golgi-like" in Fig S5C, top), (2) found both at the TGN-region and at the cell periphery ("Intermediate" in Fig S5C, middle), or (3) excluded from the Golgi ("Periphery" in Fig S5C, bottom). At every time point, the localization of ANF::GFP appeared to be slightly more Golgi-like in *Ccdc186* KO cells than in WT cells, but in

both cell types, ANF::GFP moved out from the Golgi towards the cell periphery at a roughly similar rate (Fig S5D). This indicates that DCV cargo is not stuck near the Golgi in *Ccdc186* KO cells, but does exit the TGN and move out to the periphery.

CCDC186 localizes to a ring-like compartment around proinsulin

We previously reported that in *C. elegans* neurons, CCCP-1 localizes near the TGN and that in 832/13 cells, CCDC186 is localized at or close to the TGN and iDCVs (Ailion et al., 2014; Cattin-Ortolá et al., 2017). These data were based largely on overexpressed GFP-tagged constructs. To more specifically define the localization of endogenous CCDC186 in 832/13 cells, we performed costaining of CCDC186 with a number of compartmental markers. Endogenous CCDC186 is localized to an area immediately adjacent to but not tightly overlapping with the TGN marker TGN38 (Fig 4A, S6). CCDC186 is also close to but not overlapping with the TGN subdomain marked by golgin-97 (Fig S7G). By contrast, CCDC186 colocalized most closely with the SNARE protein syntaxin 6, and the DCV cargos chromogranin A (CgA) and proinsulin (Fig 4A, S6). Syntaxin 6 and proinsulin are localized to the TGN and iDCVs, but are excluded from mature DCVs (Klumperman et al., 1998; Wendler et al., 2001). CgA is localized to a perinuclear area and also to puncta throughout the cytoplasm and the cell periphery (Fig S6), suggesting that CgA is present at the TGN, iDCVs, and mature DCVs. CCDC186 does not colocalize tightly with markers of the ERGIC (ERGIC53), cis-Golgi (GM130), early endosome (EEA1, GFP::RAB5A, and Alexa568-transferrin), late endosome (GFP::RAB7A), recycling endosome (GFP::RAB4A and Alexa568-transferrin) and lysosome (LAMP1) (Fig 4A, S6, S7A-D). We conclude that CCDC186 is localized at or close to the TGN and iDCVs.

In *C. elegans* neurons, Drosophila S2 cells, and mammalian COS cells, CCDC186 and its orthologs colocalized with RAB2 (Ailion et al., 2014; Gillingham et al., 2014). Consistent with this, we found that endogenous CCDC186 in 832/13 cells colocalizes partially with both rat RAB2 paralogs (GFP::RAB2A and GFP::RAB2B, Fig S7E,F).

To determine the localization of CCDC186 at higher resolution, we used structured illumination microscopy (SIM, Fig 4B) and stimulated emission depletion microscopy (STED, Fig S8). With both techniques, we found that endogenous CCDC186 is localized to ring-like structures of variable size (~ 200-500 nm diameter) rather than the puncta seen by conventional confocal microscopy. Interestingly, the rings of CCDC186 often surrounded proinsulin, suggesting that CCDC186 localizes at or close to iDCV membranes, or to a subdomain of the TGN carrying DCV cargos.

CCDC186 recruits the EARP complex

Though our genetic experiments indicate that CCDC186 and EARP act in the same pathway to control cargo sorting to DCVs (Fig S1), we could not detect a direct physical interaction between EARP and CCDC186 using yeast two-hybrid assays or coimmunoprecipitation and mass spectrometry experiments (our unpublished data). Moreover, EARP has previously been shown to localize to recycling endosomes (Schindler et al., 2015), but our data here indicate that CCDC186 does not localize to recycling endosomes (Fig S7A,B,H). However, we recently showed that in addition to its localization at recycling endosomes, the EARP subunit VPS50 also colocalizes with CCDC186 near the TGN (Topalidou et al., 2018), suggesting a possible connection of CCDC186 to EARP. Endogenous VPS50 is localized to puncta that are spread in a region that extends beyond the perinuclear region (Fig 5A and Topalidou et al., 2018). However, in cells that strongly overexpress CCDC186, we found that VPS50 almost fully colocalizes with CCDC186 in the perinuclear region (Fig 5A). CCDC186 overexpression does not increase the total level of VPS50 in the cell (Fig 5C), and the absence of CCDC186 also does not obviously affect the VPS50 protein level (Fig 5C). Thus, overexpressed CCDC186 appears to recruit VPS50 and relocate it. In *Eipr1* KO cells, CCDC186 overexpression does not lead to recruitment of VPS50 (Fig 5B). Because EIPR1 is required for the localization of EARP to membranes (Topalidou et al., 2018) and because we have been

unable to detect any direct physical interaction between CCDC186 and either VPS50 or EIPR1 (data not shown), our data are most consistent with the model that CCDC186 overexpression recruits EARP-positive membranes.

To test whether CCDC186 recruits EARP-positive recycling endosomes, we performed colabeling experiments between overexpressed CCDC186 and recycling endosomal markers. We found that overexpression of CCDC186 does not obviously affect the localization of GFP::RAB4, GFP::RAB11 or Alexa568-labeled transferrin (Fig S7A,B and S7H). Also, CCDC186 overexpression has no obvious effect on localization of markers of other tested cell compartments such as iDCVs, the TGN, and early- and late-endosomes (Fig S7C,D and Cattin-Ortolá et al., 2017). Thus, overexpression of CCDC186 appears to specifically recruit EARP-positive membranes, but not entire endosomal compartments.

We mapped down the domain of CCDC186 required to relocate EARP. CCDC186 localizes to TGN-proximal membranes via its C-terminal CC3 domain, but its N-terminal CC1+2 domain localizes to the cytoplasm (Cattin-Ortolá et al., 2017). We found that overexpression of GFP-tagged CC3 had no effect on the localization of endogenous VPS50, but similar to full-length CCDC186, overexpression of the CC2+3 domains relocated endogenous VPS50 (Fig S9B). Thus, the CC3 domain of CCDC186 is needed for its localization to membranes but not for the actual recruitment of EARP, the CC2 domain is critical for the recruitment of EARP, and the extreme N-terminal domain CC1 is not required for CCDC186 localization or EARP recruitment.

Consistent with the localization of EARP and EIPR1 at recycling endosomes, knockdown of EIPR1 or EARP causes a delay in the recycling of cargos from endosomes back to the plasma membrane (Gershlick et al., 2016; Schindler et al., 2015; Topalidou et al., 2018). To test whether CCDC186 is similarly involved in endocytic recycling, we examined the recycling of Alexa488–labeled transferrin in WT and *Ccdc186* KO 832/13 cells. We found that *Ccdc186* KO cells have no defect in transferrin recycling (Fig S10). Thus, unlike EIPR1 and EARP, CCDC186

is not involved in endocytic recycling. This suggests that CCDC186 and EARP may function together in dense-core vesicle cargo sorting near the TGN, but that EARP also has a CCDC186-independent function in endocytic recycling.

The CCDC186 C-terminal domain is in close proximity to carboxypeptidase D

To discover new interactors of CCDC186/CCCP-1, we performed mass spectrometry of immunoprecipitated CCDC186::GFP from 832/13 cells and mass spectrometry of proteins from a *C. elegans* cell lysate that bound to purified recombinant GST-CCCP-1 attached to a column. Neither of these approaches revealed any promising candidates. As an alternative strategy, we used the proximity biotinylation BioID approach to identify new direct interactors or proteins in close proximity to CCDC186, and to also investigate the protein composition of CCDC186-positive membranes. To increase the specificity of labeling of the membranes bound by CCDC186, we used a mitochondrial relocation strategy because it should give a background composed of mitochondrial proteins that we could easily discard (Roux et al., 2012; Shin et al., 2017).

CCDC186 was ectopically expressed and localized to mitochondria through attachment to the mitochondrially-localized transmembrane domain of monoamine oxidase (MAO) (Wong and Munro, 2014) (Fig 6A and S11A). We generated constructs with full-length CCDC186, the CCDC186 C-terminal domain CC3 which binds RAB2 and membranes (Cattin-Ortolá et al., 2017), and the CCDC186 C-terminal truncation CC1+2, each fused at their C-terminus to an HA tag and to MAO (Fig 6B and S11B). Experiments using the full-length CCDC186 protein fused to MAO resulted in low expression of the construct and high toxicity in 832/13 cells, so we did not pursue further. We showed by immunofluorescence that both CC3::HA::MAO and CC1+2::HA::MAO colocalized well with the mitochondrial marker MitoTracker and that neither construct colocalized with the TGN marker TGN38 (Fig S11C). We also observed that CC3::HA::MAO colocalized with constitutively active forms of both RAB2A and RAB2B, but

CC1+2::HA::MAO did not (Fig S12). Thus, the mitochondrial relocation experiment recapitulates the known interaction of CCDC186 with RAB2.

We fused the promiscuous biotin ligase BirA* to the C terminus of CC1+2 and CC3 (CC1+2::BirA*::HA::MAO and CC3:: BirA*::HA::MAO) (Fig 6A,B). BirA*::HA::MAO was used as a negative control. We generated PC12 cell lines that stably expressed these three constructs instead of 832/13 cells for technical reasons (see Material and Methods) and performed mass spectrometry on the biotinylated material pulled down by streptavidin beads. Analysis of the mass spectrometry data revealed that peptides for CCDC186 were found in the CC3 and CC1+2 samples but not the negative control, demonstrating the efficiency of the labeling (Fig 6C and Tables 1,2 and 7,8). Comparison of CC1+2::HA::MAO with the negative control did not lead to any highly promising candidates, but comparison of CC3::HA::MAO with the negative control gave carboxypeptidase D (CPD) as a possible specific hit (Fig 6C, Fig S13, Tables 1,2 and 7,8). Immunoblotting of streptavidin-purified biotinylated proteins from cells expressing the different BirA* constructs using an anti-CPD antibody validated the mass spectrometry result (Fig 6D). Using immunofluorescence in 832/13 cells, we verified that expression of CC3::HA::MAO resulted in partial relocation of CPD to the mitochondria, but CC1+2::HA::MAO did not (Fig 6E). Thus, the C-terminal CC3 domain of CCDC186 interacts with CPD or is in close proximity to CPD in cells. Given that CPD is a transmembrane protein (Kuroki et al., 1995; Xin et al., 1997), CCDC186 might interact with a membrane carrying CPD.

CCDC186 is required for the removal of carboxypeptidase D from dense-core vesicles

CPD belongs to the family of metallocarboxypeptidases that remove arginine or lysine residues from the C-terminus of peptides (Fricker, 2013). CPD localizes mainly to the TGN and has been reported to traffic between the plasma membrane, endosomes, and the TGN (Varlamov and Fricker, 1998). Immunofluorescence in 832/13 cells showed that CCDC186 and CPD partially colocalized in a perinuclear area (Fig 7A). Importantly, although CPD colocalized

with the TGN marker TGN38 in WT cells (Fig 7B), CPD was largely mislocalized to puncta distributed throughout the cytoplasm in *Ccdc186* KO cells (Fig 7B); this phenotype was rescued by expression of *Ccdc186*(+) in *Ccdc186* KO cells (Fig 7B). Because EIPR1 and CCDC186 function in the same pathway in DCV biogenesis, we tested whether CPD localization was also affected by the absence of EIPR1. We found that CPD was also localized to puncta distributed throughout the cytoplasm in *Eipr1* KO cells, but to a lesser extent than in the *Ccdc186* KO cells (Fig 7B). Thus, CCDC186 and EIPR1 restrict CPD localization to the TGN.

Studies in AtT-20 cells have shown that CPD localizes to the TGN and to iDCVs, but is mostly absent from mature DCVs, indicating that CPD is removed from iDCVs during maturation (Varlamov et al., 1999). We reasoned that the punctate cytoplasmic pattern of CPD in *Ccdc186* KO cells and *Eipr1* KO cells could be the result of defective CPD removal from iDCVs. Indeed, most CPD-positive cytoplasmic puncta colocalized with insulin in *Ccdc186* KO cells (78 +/- 10 % of CPD cytoplasmic puncta are insulin positive) and *Eipr1* KO cells, suggesting that these CPD-positive puncta represent mature DCVs (Fig 7C). To further confirm that CPD is mislocalized to mature DCVs in the *Ccdc186* and *Eipr1* mutants, we performed cellular fractionation experiments by equilibrium sedimentation through a sucrose gradient to separate the Golgi and DCV compartments (Fig 8). The absence of CCDC186 or EIPR1 did not have a significant effect on the sedimentation profile of PC1/3, indicating that the size, shape, and density of the DCVs are not significantly altered in either knock-out cell line. However, we found that some of the CPD shifted to a heavier fraction that comigrated with the DCV marker PC1/3 in both *Ccdc186* and *Eipr1* KO cells (Fig 8), consistent with the model that CPD is retained in mature DCVs in these mutants. Together, these immunofluorescence and biochemical data indicate that CCDC186 and EIPR1 are required for the removal of CPD from iDCVs.

Similar to CPD, other proteins such as the cation-dependent mannose 6-phosphate receptor (CD-MPR) are removed from mature DCVs (Klumperman et al., 1998). Interestingly, although the CD-MPR is a transmembrane protein with similar cytosolic sorting motifs as CPD

(Bonifacino and Traub, 2003; Eng et al., 1999), its removal from iDCVs is not dependent on CCDC186 or EIPR1 (Fig S14). Thus, different mechanisms may be needed for the removal of different cargos from immature DCVs.

DISCUSSION

In this study we investigated the role of CCDC186 in DCV function in the rat pancreatic beta-cell line 832/13. Using CRISPR knock-outs and rescuing experiments, we found that CCDC186 acts in the post-Golgi maturation of DCVs to control (1) the retention of mature cargos such as insulin and CPE, (2) the removal of cargos such as CPD, and (3) the recruitment of the EARP complex. We propose that CCDC186 acts as a regulator of DCV maturation by ensuring that mature DCVs carry the proper type and amount of cargo.

CCDC186 controls retention of cargos in mature DCVs

C. elegans mutants in *rab-2, cccp-1* (the *C. elegans* homolog of *Ccdc186*), *eipr-1,* and the EARP complex subunits have reduced levels of DCV cargo (Ailion et al., 2014; Edwards et al., 2009; Paquin et al., 2016; Sumakovic et al., 2009; Topalidou et al., 2016). This conclusion was based mainly on assays using exogenous DCV cargos overexpressed in neurons. In this study, we used the rat insulin-secreting insulinoma cell line 832/13 to investigate the secretion, localization, and processing of endogenous DCV cargos. We found that cells lacking CCDC186 remain responsive to stimulated secretion, but secrete reduced levels of insulin and have reduced levels of insulin and other luminal cargos in mature DCVs at the cell periphery. Our data are most consistent with a model in which CCDC186 acts in a post-Golgi step to control retention of DCV cargos (Fig 9), rather than acting in the TGN to control sorting of cargos by entry or having a direct effect on peptide processing. First, *Ccdc186* KO cells show no defect in the levels or localization of immature DCV cargos such as proinsulin and pro-CPE, but do have reduced levels of mature cargos such as CPE. Second, the exogenous DCV cargo ANF-GFP was not stuck in the TGN in *Ccdc186* KO cells, but moved to the periphery following release of a low-temperature block. Third, there is not increased acidity of the TGN in *Ccdc186* KO cells. Fourth, there is no apparent defect in processing of proinsulin to insulin in *Ccdc186* KO cells. These data suggest that early steps in the sorting and processing of pro-proteins occur normally

in the absence of CCDC186, but that mature DCV cargos are lost in a post-Golgi step. *Eipr1* KO cells have similar defects (Topalidou et al., 2018 and this study), suggesting that CCDC186 and EIPR1/EARP act in the same pathway to regulate DCV maturation.

CCDC186 controls the removal of CPD from immature DCVs

In a proximity biotinylation screen, we identified CPD as being in close proximity to the C-terminal CC3 domain of CCDC186. CPD is a transmembrane protein with a short cytosolic tail (~ 60 amino acids) and a longer N-terminal intraluminal domain (Kuroki et al., 1995; Xin et al., 1997). CPD is an ubiquitously expressed member of the metallocarboxypeptidase family, but its role in the processing of peptides destined to the regulated secretory pathway remains unclear (Fricker, 2013).

The basis of the interaction between CCDC186 and CPD-positive membranes is unclear. In coimmunoprecipitation experiments from 832/13 detergent lysates, we did not detect an interaction between GFP-tagged CCDC186 and either endogenous CPD or overexpressed mCherry-tagged CPD cytosolic tail (data not shown), indicating that CCDC186 and CPD do not interact directly or their interaction is too weak to detect by this method. CCDC186 might bind to CPD-positive membranes via interaction with another protein or by directly binding to membranes. The CC3 domain of CCDC186 is capable of direct association with synthetic membrane liposomes and is both necessary and sufficient for CCDC186 localization to TGN/iDCV membranes (Cattin-Ortolá et al., 2017).

Studies in AtT-20 cells have shown that CPD localizes to the TGN and to iDCVs but is mostly absent from mature DCVs (Varlamov et al., 1999). We show here that in the absence of CCDC186 or EIPR1, CPD remains in mature, insulin-positive DCVs. Thus, we propose that CCDC186 and EIPR1 ensure the removal of CPD from iDCVs. Similar to CPD, other proteins such as the cation-independent and the cation-dependent mannose 6-phosphate receptors (CI-MPR and CD-MPR) are removed from iDCVs by a pathway that depends on AP-1 and clathrin

(Dittie et al., 1999; Klumperman et al., 1998). The fate of these proteins once removed is unknown, but it has been proposed that they follow the endosomal route (Arvan and Halban, 2004; Feng and Arvan, 2003). Strikingly, although the CD-MPR is also a transmembrane protein with similar cytosolic sorting motifs as CPD (Bonifacino and Traub, 2003; Eng et al., 1999), its removal from iDCVs was not affected by the absence of CCDC186 or EIPR1, suggesting that different cargos may be removed from immature DCVs by different mechanisms. Our data suggest that CCDC186 and EIPR1 ensure the removal of CPD from DCVs (Fig 9).

CCDC186 recruits EARP

Our results show that CCDC186, the EARP complex, and the EARP interactor EIPR1 function in the same genetic pathway in *C. elegans* neurons to promote locomotion and cargo sorting to DCVs. Furthermore, we found that *Ccdc186* KO 832/13 cells have similar DCV cargo sorting defects to *Eipr1* KO 832/13 cells. Overexpressed and tagged EIPR1 and VPS50 were reported to localize to Rab4-positive recycling endosomes and to regulate endosomal recycling of transferrin (Gershlick et al., 2016; Gillingham et al., 2014). We found that endogenous VPS50 localizes to two pools in 832/13 cells, a CCDC186-positive pool and a recycling endosome pool from which CCDC186 is excluded (Topalidou et al., 2018). CCDC186 partially colocalizes with EARP near the TGN but does not localize to recycling endosomes and does not function in endocytic recycling of transferrin. Also, by super-resolution microscopy, we found that CCDC186 localizes to ring-like structures that surround proinsulin, suggesting that CCDC186 localizes around iDCV membranes or proinsulin-positive TGN subdomains, consistent with a role for CCDC186 in DCV biogenesis. These results suggest that CCDC186 and EIPR1/EARP act in the same pathway for DCV biogenesis, but that EIPR1/EARP also acts independently in endocytic recycling.

Interestingly, we found that CCDC186 controls the localization of EARP since overexpression of CCDC186 recruits EARP from its recycling endosome pool in an EIPR-1-

dependent manner. Because we did not detect direct protein-protein interactions between CCDC186 and EARP, we suggest that CCDC186 recruits EARP-positive membranes. This model is supported by the observation that EIPR1 is important for EARP localization to membranes (Topalidou et al., 2018). CCDC186 has features in common with the golgin family of membrane tethers, including a flexible and extended structure (Cattin-Ortolá et al., 2017), consistent with the possibility that it may be anchored by its C-terminus in membranes near the TGN/iDCVs and recruit EARP-positive membranes via its more N-terminal domains. Consistent with this model, the C-terminal CC3 domain of CCDC186 is important for its localization to membranes near the TGN (Cattin-Ortolá et al., 2017), but the middle CC2 domain is responsible for recruiting EARP.

What is the role of CCDC186 in DCV biogenesis?

Our data suggest that CCDC186 is important for both the retention of mature DCV cargos and the removal of some non-DCV cargos from immature DCVs. These are both post-Golgi processes, but it is unclear whether they are independent or controlled coordinately by a single action of CCDC186. Here we propose that CCDC186 serves a role in ensuring that the proper amount and type of cargo remains in mature DCVs.

Our data suggest that CCDC186 acts after the processing of peptide cargos. This distinguishes CCDC186 from other proteins important for DCV biogenesis such as PICK1, ICA69, and HID1. Like CCDC186, these proteins are also required for normal levels of mature DCV cargos (Cao et al., 2013; Du et al., 2016; Holst et al., 2013; Hummer et al., 2017), but in contrast to CCDC186, loss of PICK1, ICA69, or HID1 leads to a defect in the processing of proinsulin to insulin (Cao et al., 2013; Du et al., 2016). PICK1, ICA69, and HID1 have been proposed to act at an early step of DCV biogenesis during budding of immature DCVs from the TGN; our data suggest that CCDC186 and EIPR1 (Topalidou et al., 2018) act at a post-Golgi step during DCV maturation.

The connection between CCDC186 and the EARP complex suggests there may be trafficking between EARP-positive endosomes and CCDC186-positive TGN/iDCV membranes. Such a trafficking step may be involved in the removal of CPD from iDCVs. Trafficking from endosomes to iDCVs may also be involved in the retrieval of DCV membrane proteins following exocytosis. It has been shown that after DCV exocytosis, transmembrane proteins such as phogrin and PAM are recycled and traffic through endosomes before reaching newly-generated DCVs (Bäck et al., 2010; Vo et al., 2004). CCDC186 and EARP could mediate the recycling of DCV proteins from endosomes to iDCVs; disrupting this recycling pathway could also indirectly impair the proper trafficking of sorting receptors required for retention of luminal DCV cargos. Similar to loss of CCDC186 or EIPR1, loss of other proteins such as BAIAP3 and Vti1a/Vti1b that are involved in retrograde trafficking from endosomes to Golgi also leads to reduced levels of DCV cargos (Emperador-Melero et al., 2018; Zhang et al., 2017). However, BAIAP3 and Vti1a/b appear to affect endosome to Golgi trafficking more generally because their loss also affects non-DCV cargos and perturbs Golgi morphology (Emperador-Melero et al., 2018; Zhang et al., 2017), whereas loss of CCDC186 or EIPR1 does not affect non-DCV cargos and has no apparent effect on Golgi morphology as assayed by TGN38 staining. CCDC186 and EARP may be specifically involved in endosomal trafficking of DCV-related factors or cargos.

The idea that CCDC186 and EARP affect DCV cargo levels by regulating trafficking to or from endosomes is supported by experiments in *C. elegans* that found that disrupting trafficking to the endolysosomal system could restore levels of DCV cargo in *rab-2* mutants, suggesting that cargo is lost via an endolysosomal route (Edwards et al., 2009; Sumakovic et al., 2009). Given that CCDC186/CCCP-1 and EARP act in the same genetic pathway as RAB-2 (Ailion et al., 2014; Topalidou et al., 2016), it is likely that they also act to retain cargos in mature DCVs by preventing the loss of cargos to the endolysosomal system. Interestingly, a recent study showed that cholesterol-deficient DCVs are lost through lysosomal degradation, leading to a reduction in insulin secretion (Hussain et al., 2018). This observation raises the intriguing

possibility that CCDC186 and EARP may be involved not only in protein trafficking, but also in cholesterol transfer to DCVs. Supporting this possibility, the EARP complex associates with Rab11-positive recycling endosomes (Schindler et al., 2015), and Rab11 and its effector RELCH act to promote cholesterol transfer from recycling endosomes to the TGN (Sobajima et al., 2018). Perhaps CCDC186 and EARP are also involved in this process.

MATERIALS & METHODS

C. elegans strains, molecular biology, and plasmids

C. elegans strains were cultured using standard methods (Brenner, 1974). A complete list of strains used is provided in the Strain List (Table 3). A complete list of constructs is provided in the plasmid list (Table 4). Vector backbones and PCR-amplified fragments containing 20-30 bp overlapping ends were combined by Gibson cloning (Gibson et al., 2009).

Locomotion assays

To measure *C. elegans* locomotion, first-day adults were picked to 2 to 3-day-old lawns of OP50 bacteria, stimulated by touching their tail using a worm pick, and body bends were then counted for one minute. A body bend was defined as the movement of the worm from maximum to minimum amplitude of the sine wave.

C. elegans imaging and image analysis

Young adult worms were placed in 8 µl of 50 mM sodium azide on 2% agarose pads and anesthetized for 10 min. Images of animals with dorsal side up were obtained using a Nikon Eclipse 80i wide-field compound microscope. The same region of the dorsal cord around the vulva was imaged in all worms. All strains in an experiment were imaged on the same day and the microscope settings were kept constant. Maximum-intensity projections were quantified using ImageJ. Dorsal cord fluorescence is given by the total fluorescence in a region of interest surrounding the cord minus the background fluorescence of a region of the same size adjacent to the dorsal cord.

Cell culture

The 832/13 cell line is an INS-1-derived clone that was isolated by Dr. Christopher

Newgard (Duke University School of Medicine) and obtained by Dr. Duk-Su Koh (via Dr. Ian

Sweet, both at University of Washington), who provided it to us. The *Eipr1* KO and EIPR1(+)

rescued 832/13 cell lines were generated as described (Topalidou et al., 2018). The 832/13 and

832/13-derived cell lines were grown in RPMI 1640-GlutaMAX (GIBCO) medium supplemented

with 10% FBS (RMBIO), 1 mM sodium pyruvate (GIBCO), 10 mM HEPES (GIBCO), 1X

Pen/Strep (GIBCO), and 0.0005% beta-mercaptoethanol at 5% CO_2 and 37°C. The PC12 cell

line was obtained from Dr. Duk-Su Koh (University of Washington) and was grown in DMEM-

GlutaMAX (GIBCO) medium supplemented with 10% horse serum (RMBIO), 5% FBS (RMBIO),

and 1X Pen/Strep (GIBCO) at 5% CO_2 and 37°C. All cell lines were shown to be mycoplasma-

free by DAPI staining and PCR test (ABM, G238).

Protein extraction from mouse tissues

Tissues were dissected from an adult female mouse. Bigger organs such as the liver

and brain were placed in 3 ml of lysis buffer, and smaller organs such as the heart and intestine

were placed in 1.5 ml of lysis buffer (Tissue-PE LB^{TM}, G-Biosciences #786-181). These organs

were then frozen in liquid nitrogen and kept at -80°C for 1 hour. Tissues were dissolved with the

use of a homogenizer until they became liquid and then were further homogenized with

mechanical breakage using a 22-gauge needle, applying 12-15 strokes. The tissues were then

spun down at 20,000g in a microcentrifuge for 15 minutes, and the supernatant was collected

and spun again for 5 more minutes. The supernatant was finally collected and stored at -80°C.

Immunoblotting

Protein extracts were loaded on 8, 10, or 12% SDS-PAGE gels and transferred onto

PVDF membranes. For the blots shown in Figures 5C, S2A and S2B, membranes were blocked

in 3% milk in TBST (50 mM Tris pH 7.4, 150 mM NaCl, 0.1% Tween 20) for 1 hour at room

temperature (RT) and stained with the relevant primary and secondary antibodies in 3% milk in TBST for 1 hour at RT or overnight at 4ºC, followed by three 5-minute washes in TBST. For the blots shown in Figures 3A-D, 6D and 8A, membranes were blocked with Odyssey® Blocking Buffer (PBS) (#927-4000), incubated overnight at 4ºC with primary antibody diluted in Odyssey® Blocking Buffer (PBS) supplemented with 0.2% Tween 20, washed with PBST (PBS + 0.1% Tween 20), incubated for 1 hour at RT in secondary antibody diluted in Odyssey® Blocking Buffer (PBS) supplemented with 0.2% Tween 20 and 0.01% SDS, and washed with PBST. A LI-COR processor was used to image the membranes. The antibodies used are listed in Tables 5 and 6.

Generation of *Ccdc186* KO by CRISPR

We performed Cas9-mediated genome editing in 832/13 cells as described (Ran et al., 2013). We designed guide RNAs using the online CRISPR design tool (Ran et al., 2013) and selected a guide RNA that recognizes a sequence in the first exon of rat *Ccdc186*: 5'-GCGTGAGTCGTCCTTCAACTCGG -3'.

The guide RNA was cloned into the pSpCas9(BB)-2A-GFP vector as described (Ran et al., 2013). The efficiency of the cloned guide RNA was tested using the SURVEYOR nuclease assay according to the manufacturer's instructions (Surveyor Mutation Detection kit, Transgenomic). We designed a homology-directed repair (HDR) template using the pPUR vector (Clontech) as a backbone and cloned approximately 1.5 kb *Ccdc186* homology arms upstream and downstream of the puromycin selection cassette. The HDR template was assembled using Gibson cloning.

To cotransfect the CRISPR plasmid and HDR template into 832/13 cells, 832/13 cells were grown in two 10-cm petri dishes to near confluency. Cells were cotransfected with 7 µg CRISPR plasmid and 7 µg non-linearized HDR template using Lipofectamine 3000 according to the instructions (ThermoFisher). 48 hours after transfection, we removed the culture

medium and replaced with new medium containing 1 µg/ml puromycin. The puromycin

selection was kept until individual clones could be picked, grown in individual dishes, and tested

for *Ccdc186* KO. These clones were screened for *Ccdc186* KO by immunofluorescence and

subsequently by RT-PCR and Western blot.

RT-PCR

WT and *Ccdc186* KO cells were grown in 10-cm tissue-culture dishes, washed twice

with ice-cold PBS, and harvested in 1 ml of TRIzol (Invitrogen). Total RNA was isolated

following the manufacturer's protocol and cDNA was synthesized from 1 µg total RNA using the

QuantiTekt Reverse Transcription kit (Qiagen). The primers used for detecting *Ccdc186* cDNA

were:

F: 5'-ATGAAGATCAGGAGCAGATTTGAAG-3'

R: 5'-GGCCTTCTTCGTCCTCTGCTC-3'

Lentiviral production, infection of cells, and selection of stable lines

The following method was used to create a *Ccdc186* KO cell line rescued by expression

of the *Ccdc186* cDNA. Platinum-E (PlatE) retroviral packaging cells (a gift from Suzanne

Hoppins) were grown for a couple of generations in DMEM-GlutaMAX (GIBCO) medium

supplemented with 10% FBS (RMBIO), 1X Pen/Strep (GIBCO), 1 µg/ml puromycin, and 10

µg/ml blastocidin at 5% CO_2 and 37°C. On day one, approximately 3.6 x 10^5 PlatE cells per well

were plated in a six-well dish in DMEM-GlutaMAX medium supplemented with 10% FBS and 1X

Pen/Strep. On day two, a mix of 152 µl Opti-MEM (ThermoFisher), 3 µg CCDC186_pBabe-

hygro DNA, and 9 µl Fugene HD transfection reagent (Promega) was incubated for 10 minutes

at room temperature and transfected into each well. On day three, we removed the medium and

replaced with new PlatE medium. On day four, approximately 1.5 x 10^5 *Ccdc186* KO 832/13

cells per well were plated in a six-well dish in RPMI 1640-GlutaMAX, supplemented with 10%

FBS, 1 mM sodium pyruvate, 10 mM HEPES, 1X Pen/Strep, and 0.0005% beta-mercaptoethanol. 3 µl of 8 mg/ml hexadimethrine bromide (Sigma) was added to each well. The supernatant of the PlatE cells (48-hour viral supernatant) was collected with a sterile syringe, passed through a 0.45 micron filter, and added to the *Ccdc186* KO cells. The *Ccdc186* KO cells were incubated for 5-8 hours at 5% CO_2 and 37°C, then the medium was changed and replaced with new medium and the cells were incubated overnight at 5% CO_2 and 37°C. On day five, the supernatant was removed from the *Ccdc186* KO cells and replaced with the supernatant from the PlatE cells (72-hour viral supernatant) after passing through a 0.45 micron filter. 3 µl of 8 mg/ml hexadimethrine bromide was added to each well and the cells were incubated for 5-8 hours. The medium was replaced with new 832/13 cell medium. On day six, the *Ccdc186* KO cells were collected, transferred into a 10-cm petri dish, and 200 µg/ml hygromycin was added. The cells were grown under hygromycin selection until individual clones could be picked and tested for CCDC186 expression.

Immunofluorescence

Cells were seeded onto cover slips (Thomas Scientific #121N79) placed in 24-well cell culture plates. For GFP or epitope-tagged constructs, cells were transfected at least 24 hours after seeding with 500-800 ng of plasmid using Lipofectamine 2000 (ThermoFisher), according to the manufacturer's instructions, for 24 to 48 hours. When indicated, cells were incubated for 25 minutes with 50 µg/ml Alexa488- or Alexa568-transferrin (Invitrogen #T12242 or # T23365) diluted in serum-free RPMI medium + 25 mM HEPES + 1% BSA before fixation. When indicated, cells were incubated with MitoTracker deep red FM (Thermofisher #M22426) at a final concentration of 300 nM for 15 minutes in complete medium at 37°C.

For antibody staining, cells were rinsed twice with PBS and fixed with 4% paraformaldehyde (made in PBS) for 20 minutes at room temperature, then rinsed twice with PBS and permeabilized with 0.1% Triton X-100 in PBS for 20 minutes or 0.5% Triton X-100 in

PBS for 5 minutes at room temperature, and finally rinsed twice with PBS and placed in 5% milk in PBS for 1 hour at room temperature. Cells were stained with primary antibodies in 0.5% milk in PBS at room temperature for 1 hour, washed three times with PBS, and incubated in secondary antibody for 1 hour at room temperature. The cells were washed with PBS three times, mounted onto glass slides using Vectashield (Vector laboratories H1000) or Prolong Diamond (Life Technologies P36965), sealed with transparent nail polish and examined by fluorescence microscopy. Antibodies and their working dilutions are listed in Tables 5 and 6.

Most images were obtained using an Olympus Fluoview FV1200 confocal microscope with a 60X UPlanSApo oil objective (numerical aperture = 1.35). The acquisition software was Olympus Fluoview v4.2. When indicated, a Nikon Eclipse 80i wide-field compound microscope was used. For the quantification of the insulin, proinsulin, CPE, and pro-CPE distributions in *Ccdc186* KO cells (Fig 2B, 2F, 3F, and 3J), Pearson's correlation coefficients were determined from confocal images using Fiji and the coloc-2 plugin by taking maximum-intensity projections of whole cells and drawing a line around each individual cell. For the quantification shown in Figs 2C and 3G, z-stack images were obtained using a Nikon Eclipse 80i wide-field compound microscope. Maximum-intensity projections were quantified using Fiji. The final fluorescent intensity is the total fluorescence in a region of interest around the Golgi divided by the fluorescence of a region of the same size in the cytoplasm of the cell (Golgi/Cytoplasm). For the quantification shown in Figs 2D and 3H, z-stack images were obtained using an Olympus Fluoview FV1200 confocal microscope. Maximum-intensity projections were obtained by only keeping the slices that had TGN38 signal so insulin or CPE signal coming from the top and the bottom of the cell would not impact the quantification. A line was drawn around the TGN38 marker and Pearson's correlation coefficients were obtained using FIJI and the coloc-2 plugin. For the quantification of colocalization with CCDC186 (Fig 4A), a square was drawn around the perinuclear area where CCDC186 is localized, and Pearson's correlation coefficients were determined using Fiji and the JACOP plugin.

For super-resolution imaging, cells were prepared as described above. For SIM imaging, coverslips were mounted with DAPI-free Vectashield and imaged with a 3D SIM OMX SR microscope (GE Healthcare). For STED microscopy, secondary antibodies were used at higher concentration (see Table 6). Coverslips were mounted in DAPI-free fresh Prolong Diamond, sealed with twinsil® 22, and imaged with a Leica SP8 STED microscope.

Insulin and proinsulin secretion assays

Cells were grown in 24-well plates to near confluency. Cells were washed twice with PBS and incubated for 1 hour in 200 μl per well resting buffer (5 mM KCl, 120 mM NaCl, 24 mM NaHCO$_3$, 1 mM MgCl$_2$, 15 mM HEPES pH 7.4). The medium was collected, cleared by centrifugation, and stored at -80°C. The cells were incubated for 1 hour in 200 μl per well stimulating buffer (55 mM KCl, 25 mM glucose, 70 mM NaCl, 24 mM NaHCO$_3$, 1 mM MgCl$_2$, 2 mM CaCl$_2$, 15 mM HEPES pH 7.4). After stimulation, the medium was cleared by centrifugation and stored at -80°C. The cells were washed once with PBS, harvested in PBS, and extracted in 100 μl per well acid-ethanol solution (absolute ethanol:H$_2$O:HCl, 150:47:3). The pH of the acid-ethanol solution was neutralized by addition of 20 μl of 1 M Tris base per 100 μl of acid ethanol and the samples were stored at -80°C.

Samples were assayed for insulin or proinsulin content using ELISA according to the instructions of the manufacturers (Rat/Mouse insulin ELISA, Millipore, #EZRMI-13K; Rat/Mouse proinsulin ELISA, Mercodia, #10-1232-01). Secreted insulin and proinsulin levels were normalized against total cellular protein concentration and were presented as fraction of the wild type under stimulating conditions (Fig 1A,B left panels). These values were then normalized against total cellular insulin or proinsulin levels (Fig 1A,B middle panels) to give the secretion data normalized against total insulin or proinsulin content (Fig 1A,B right panels).

Constitutive secretion assay

WT and *Ccdc186* KO cells were seeded on 12-well plates and grown to subconfluency. Cells were transfected with a plasmid expressing GFP fused to a signal peptide at its N-terminus (ssGFP) (Hummer et al., 2017). 36-48 hours later, cells were incubated for 1 hour at 37ºC in the cell culture incubator with resting medium (114 mM NaCl, 4.7 mM KCl, 1.2 mM KH_2PO_4, 1.16 mM $MgSO_4$, 20 mM HEPES, 2.5 mM $CaCl_2$, 25.5 mM $NaHCO_3$, 3 mM glucose). The 12-well plate was transferred to ice and the secretion medium was collected and centrifuged at 20,000g for 10 minutes at 4ºC to remove the cell debris. The supernatant was collected in a separate tube. The pelleted cells were lysed on ice with 50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1% Triton X-100, 1 mM EDTA, protease inhibitor cocktail without EDTA (Pierce). The lysate was clarified by centrifugation at 20,000g at 4ºC for 10 minutes and the supernatant was collected to a new tube. The fluorescence of the medium (secreted GFP) and the lysate (cellular GFP) were measured using a plate reader (Molecular Devices Spectramax Gemini XPS, excitation = 485 nm, emission = 525 nm, cutoff = 515 nm). Background fluorescence was determined using the medium and lysate from cells that were not transfected. The data are presented as the ratio of fluorescence in the medium to the fluorescence in the cell lysate.

pH measurement of the late-Golgi compartment

WT and *Ccdc186* KO cells stably expressing the 17-residue transmembrane domain of beta-galactoside alpha 2,6-sialyltransferase tagged to pHluorin (St6Gal1::pHluorin) were generated as follows. HEK293T cells were maintained in DMEM with 10% fetal bovine serum under 5% CO_2 at 37°C. Lentivirus was produced by transfecting HEK293T cells with FUGW, psPAX2, and pVSVG using Fugene HD according to the manufacturer's instructions. Two days after transfection, the medium was collected and filtered (0.45 µm). Medium containing the lentivirus was then applied to 832/13 cells in suspension at varying ratios. Transduction efficiency was screened using an epifluorescence microscope. The protocol to measure the pH of the late-Golgi compartment (Hummer et al., 2017) was adapted to a plate-reader format. An

equal number of WT and *Ccdc186* KO 832/13 cells stably expressing St6Gal1::pHluorin were plated on clear-bottom black-wall 96-well plates and grown to confluence. Cells were washed once with Tyrode buffer (119 mM NaCl, 2.5 mM KCl, 2 mM $CaCl_2$, 2 mM $MgCl_2$, 25 mM HEPES, 30 mM glucose, pH 7.4) and incubated for 10 minutes either in Tyrode buffer or in KCl-enriched buffer at different pHs (125 mM KCl, 20 mM NaCl, 0.5 mM $CaCl_2$, 0.5 mM $MgCl_2$, and 25 mM MES (at pH 5.5 or 6) or 25 mM HEPES (at pH 6.5, 7, 7.5, 8, or 8.5). The KCl-enriched buffer was supplemented with 5 µM nigericin (Sigma Aldrich) and 5 nM monensin (Sigma Aldrich). The fluorescence of each well was measured using a Varioskan Lux plate reader (Thermo Scientific) (Excitation=485 nm, Emission=520 nm). For each buffer, cell type, and independent biological replicate, the reading was repeated three times. Calibration curves were generated for each cell type and each independent repetition. The absolute pH values were extrapolated from the calibration curve using the fluorescence of the cells incubated with Tyrode buffer.

ANF-GFP pulse-chase experiment

To monitor the exit of DCV cargo from the TGN, we used a protocol similar to the one described (Kögel et al., 2013; Kögel et al., 2010). WT and *Ccdc186* KO 832/13 cells were seeded on glass coverslips in 24-well plates and after at least 24 hours, 100 ng of ANF::GFP plasmid was transfected in each well with Lipofectamine 2000 for 12-16 hours at 37ºC in complete medium. Subsequently, cells were incubated at 20ºC in PBS for 2 hours in a conventional incubator (pulse) to block protein exit from the TGN. 30 minutes before the end of the low temperature block, 10 µg/ml of cycloheximide was added to the medium to block the synthesis of new ANF::GFP. The PBS was then exchanged for growth medium and the cells were shifted to 37°C for the indicated times, then fixed with 4% PFA, and stained with an anti-GFP antibody as described (see Immunofluorescence section). Cells were scored in three categories: those that had most of the ANF::GFP concentrated at the TGN ("Golgi-like"), those that had ANF::GFP both at the TGN-region and at the cell periphery ("Intermediate") and those

where the ANF::GFP was excluded from the TGN ("Periphery"). 50 to 100 cells per time point and per genotype were imaged and counted using a Nikon Eclipse 80i wide-field compound microscope. The experimenter was blind to the genotypes of the cell lines used and to the time point. The experiment was repeated three times with similar results.

Transferrin recycling assay

WT, *Eipr1* KO*,* and *Ccdc186* KO 832/13 cells were seeded onto a 24-well plate and grown in complete medium until they reached confluence. Cells were placed in warm uptake medium for a 25-minute pulse (serum-free RPMI + 1% BSA + 25 mM HEPES + 50 µg/ml of Alexa488- or Alexa568-transferrin (Invitrogen #T12242 or # T23365). The medium was exchanged with complete medium and cells were chased for the indicated times, washed, and transferred onto ice. Cells were washed 2x with room temperature PBS, 2x with ice-cold PBS, and detached on ice with 10 mM EDTA in PBS for 30 minutes to 1 hour with manual agitation and gentle pipetting. Detached single cells were transferred to a microcentrifuge tube and fixed at 4°C with 4% PFA (final concentration ~ 3.5%) for 10 minutes on a Nutator to avoid clumping. Fixed cells were washed with PBS and analyzed by FACS using an LSRII instrument (BD Biosciences). Data were analyzed using FlowJo software. The experiment was repeated four times independently: three times with Alexa488-transferrin and once with Alexa568-transferrin.

Generation of PC12 cell lines for the mitochondria relocation assays

For the mitochondria relocation assay, PC12 cells were used instead of 832/13 cells for drug selection compatibility reasons. Moreover, toxicity of the mitochondria relocation constructs appeared to be less in PC12 cells. A 6-well plate was transfected with 4 µg of the appropriate plasmid using Lipofectamine 2000 (ThermoFisher) according to the manufacturer's instructions. After 48 hours, the cell culture medium was replaced with Pen/Strep-free selection medium (DMEM-GlutaMAX (GIBCO) supplemented with 5% FBS (RMBIO), 10% horse serum (RMBIO),

and 700 µg/ml G418). The selection medium was changed every 2-3 days until large single colonies could be seen. Colonies were transferred to a 96-well plate and expanded in selection medium and then to 24-well and 6-cm plates before being frozen in 10% DMSO. Individual clones were screened by immunostaining using an anti-HA antibody. Highly expressing clones were used for the BioID experiment.

BioID experiment and sample preparation for mass spectrometry

The protocol for isolating proteins biotinylated by BirA* was adapted from (Shin et al., 2017). PC12 stable lines expressing the BirA* constructs (BirA*::HA::MAO, CC1+2::BirA*::HA::MAO and CC3::BirA*::HA::MAO) were grown to confluence in three 175-cm^2 flasks in complete PC12 DMEM medium (DMEM-GlutaMAX (GIBCO) supplemented with 5% FBS (RMBIO) and 10% horse serum (RMBIO)). 24 hours before harvest, cells were incubated with 50 µM biotin (10 mg/ml stock in DMSO). Cells were transferred on ice, washed 2x with ice-cold PBS, and harvested by gentle pipetting. Cells were pelleted by centrifugation (10 minutes, 1850g) and resuspended in lysis buffer (50 mM Tris pH 7.4, 0.1M NaCl, 1 mM EDTA, 1% Triton X-100, 1 mM PMSF, protease inhibitor cocktail (Pierce)), vortexed briefly, incubated on ice for 20 min, and vortexed again. Lysates were clarified by centrifugation at 20,000g for 10 minutes at 4ºC and supernatants were mixed with an equal volume of 50 mM Tris pH 7.4. The total protein content was about 30 mg as measured by a BCA assay. The supernatant was incubated with 250 µl Dynabeads MyOne Streptavidin C1 beads (Invitrogen # 65002) that had been pre-washed twice in the same buffer. The beads were incubated at 4ºC overnight on a Nutator, washed twice for 8 minutes in 2% SDS–PAGE and protease inhibitor cocktail (Pierce), then three times for 8 min in 1% Triton X-100, 0.1% deoxycholate, 500 mM NaCl, 1 mM EDTA, 50 mM HEPES, and protease inhibitor cocktail at pH 7.5, and three times for 8 minutes in 50 mM Tris pH 7.4, 50 mM NaCl, and protease inhibitor cocktail. Finally, the beads were incubated for 5 minutes at 98ºC with 50 µl 1X lithium dodecyl sulfate (LDS) sample buffer (Invitrogen NuPAGE®

#NP0007) containing 10% beta-mercaptoethanol and 3 mM biotin. The supernatant was collected and the beads were discarded.

The samples were then prepared for mass spectrometry as follows. The eluted proteins in 1X LDS were reduced with 1 mM Tris(2-carboxyethyl)phosphine (TCEP) at 37ºC for 20 minutes on a shaker and alkylated with 2 mM chloroacetamide for 20 minutes at 37ºC. The reaction was quenched by incubation with 1 mM TCEP for 20 minutes at 37ºC and 60% of the eluate was run on an SDS-PAGE gel (Mini-Protean TGX, 4-15%, Bio-Rad #4561084). The gel was stained with colloidal Coomassie (Bio-Rad #1610803) overnight at room temperature and destained with Milli-Q water. Gel slices (4-5 slices per lane) were excised and destained by two 10-minute washes with a 50/50 solution of high-grade ethanol/100 mM triethylammonium bicarbonate (TEAB). The gel was dehydrated with two washes of high-grade ethanol. Gel slices were rehydrated with a solution of 12.5 ng/μl trypsin (Promega Trypsin Gold, Mass Spectrometry Grade, #PRV5280) diluted in 100 mM TEAB. Once rehydrated, 2-3 volumes of 100 mM TEAB were added and the sample was incubated overnight at 37ºC in a shaking thermomixer at 1,400 rpm. The reaction was quenched by addition of 1/5 10% trifluoroacetic acid (TFA) and digested peptides were purified by StageTips (Rappsilber et al., 2007). StageTips were washed with 50 μl methanol, 50 μl reaction buffer B (0.1% TFA, 80% acetonitrile (ACN)), and equilibrated with 50 μl of buffer A (5% ACN, 0.1% TFA in water) by centrifugation at 2,000g at room temperature. The samples were loaded onto StageTips that were then washed once with 50 μl buffer A and kept at 4ºC until mass spectrometry analysis.

Liquid chromatography mass spectrometry (LC/MS) analysis.

Peptides were eluted from StageTips using elution buffer (50% acetonitrile, 0.1% TFA) and then loaded on a self-pulled 360 μm outer diameter (OD) x 100 μm inner diameter (ID) 20 cm column with a 7-μm tip packed with 3 μm Reprosil C18 resin (Dr. Maisch, Germany). Peptides were analyzed by nanoLC-MS in a 120 minutes, 5% to 35% acetonitrile gradient in

0.1% acetic acid at 300 nl/min (Thermo Dionex RSLCnano) on an Orbitrap Elite. Orbitrap

Fourier Transform Mass Spectrometry (FTMS) spectra (R = 30,000 at 400 m/z; m/z 350–1600;

3e6 target; max 500 ms ion injection time) and Top15 data dependent collision-induced

dissociation (CID) MS/MS spectra (1e4 target; max 100 ms injection time) were collected with

dynamic exclusion for 20 sec and an exclusion list size of 500. The normalized collision energy

applied for CID was 35% for 10 msec.

Data analysis

Mass spectra were searched against the Uniprot rat reference proteome downloaded on

Feburary 12, 2017 using MaxQuant v1.5.7.4. Detailed MaxQuant settings used were "LFQ

mode" with "Fast LFQ"; and "Match between runs" mode was enabled. Other settings were kept

as default.

Data analysis was carried out in R. Briefly, label-free quantification (LFQ, (Cox et al.,

2014) intensities calculated by MaxQuant were $\log_2$ and Z-score transformed; only proteins

present in at least two out of three replicates were selected for further analysis; missing values

were replaced by random numbers drawn from a distribution with mean of -3.51 and a standard

deviation of 0.3 to represent protein abundance below the detection limit; a two-sided Student's

t-test was used to determine the *p*-values and the results are presented in the form of a volcano

plot (Fig S13). Proteins with ratios ($\log_2$ CCDC186 CC3 – $\log_2$ BirA*) higher than 1.8 are

labeled. The data for all experiments are shown in Tables 7 and 8.

Equilibrium sedimentation

Wild type, *Ccdc186* KO, and *Eipr1* KO 832/13 cells were grown in 15-cm tissue culture

plates. Cells were washed twice with ice-cold PBS, transferred to 15 ml conical tubes, and

centrifuged at 300g for 10 minutes at 4°C. Cells were washed once with 5 ml of SH buffer (10

mM HEPES pH 7.2, 0.3 M sucrose, 1 mM PMSF and protease inhibitor cocktail (SIGMAFAST))

and resuspended in SH buffer, passed through a 21-gauge needle, and homogenized using a ball bearing device (Isobiotec, 18 µm clearance). The lysate was collected and centrifuged at 1000g for 8 minutes. A 0.6-1.9 M continuous sucrose gradient was prepared in 10 mM HEPES pH 7.2 in ultracentrifuge tubes (Beckman Coulter) that were coated with Sigmacote (Sigma #SL2). The post-nuclear supernatant was loaded onto the gradient and centrifuged at 30,000 rpm in an SW41 rotor for 14-16 hours at 4ºC. Fractions (750 µl each) were collected from top to bottom. The sucrose concentration for each fraction was determined by measuring the Brix using a Misco palm abbe digital refractometer. Fractions were analyzed by immunoblotting using antibodies to PC1/3 and CPD (see above and Tables 5 and 6). Band intensity was quantified using FIJI and plotted against the sucrose concentration of the fraction.

Statistics

Data were tested for normality by a Shapiro-Wilk test. When data did not pass the normality test, we tested for statistical significance between groups by the Kruskal-Wallis test followed by Dunn's test when making comparisons among groups of three or more, and the Mann-Whitney test for comparisons between groups of two. When data passed the normality test, we tested for statistical significance by a one-way ANOVA test with Bonferroni correction when making comparisons among groups of three or more, and an unpaired t-test for comparisons between groups of two.

ACKNOWLEDGEMENTS

We thank Alex Merz for mentoring and sharing equipment; Christopher Newgard, Ian Sweet, and Duk-Su Koh for the 832/13 cell line; Suzanne Hoppins for the pBabe vector, platE cells, protocol for lentiviral production and infection, help with cell culture, and sharing equipment; Richard Palmiter for the pPUR vector; Sean Munro and John Shin for the mitochondrial relocation and BirA* plasmids, advice and protocols for BioID; Lloyd Fricker for

the anti-CPD and anti-CPE antibodies; Joshua Vaughan and Aaron Halpern for secondary antibodies used in STED microscopy; David Castaneda-Castellanos and Patrina Pellett for assistance with STED and SIM microscopy, respectively; Ngoc-Han Thi Nguyen and Donna Prunkard from the UW Pathology Flow Cytometry Core Facility; Ken Miller and Joshua Kaplan for worm strains with integrated transgenes; and Kelly Duong for the mouse tissues and protocol for protein extraction. This work used an EASY-nLC1200 UHPLC and Thermo Scientific Orbitrap Fusion Lumos Tribrid mass spectrometer purchased with funding from a National Institutes of Health SIG grant S10OD021502 (S-E.O.). Some worm strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). This work was supported by American Diabetes Association grant #1-17-JDF-064 and by NIH grant R01 GM124035 to CSA, and by a University of Washington Diabetes Research Center Pilot and Feasibility Award (NIH grant P30 DK017047) and by NIH grant R01 GM121481 to MA.

AUTHOR CONTRIBUTIONS

JC, IT, and MA conceived the project. JC and IT performed most of the experiments and analyzed the data. HL and SO performed the mass spectrometry experiments and wrote the associated methods sections. JC and BH performed the Golgi pH measurement and equilibrium sedimentation experiments under the supervision of CSA. JC, IT, and MA wrote the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1. *Ccdc186* **KO cells secrete less insulin under stimulating conditions.**

A. (Left) Insulin secretion from WT, *Ccdc186* KO, and *Ccdc186*(+) rescued 832/13 cells under resting conditions (no glucose, 5 mM KCl, white bars) and stimulating conditions (25 mM glucose, 55 mM KCl, black bars). All values were normalized to the WT value in stimulating conditions (WT high Glu). n = 8; *p<0.05, **p<0.01, ns p>0.05, error bars = SEM.

(Middle) Total insulin content in WT, *Ccdc186* KO, and *Ccdc186*(+) rescued 832/13 cells. All values were normalized to WT. n = 8; ns p>0.05, error bars = SEM.

(Right) Insulin secretion normalized to insulin content in WT, *Ccdc186* KO, and *Ccdc186*(+) rescued 832/13 cells. n = 8; *p<0.05, ns p>0.05, error bars = SEM.

We performed three biological replicates. For each replicate, the same cells were used to determine the amount of insulin secreted under resting conditions, stimulating conditions, and the amount of total cellular insulin.

B. (Left) Proinsulin secretion from WT, *Ccdc186* KO and *Ccdc186*(+) rescued 832/13 cells under resting and stimulating conditions. All values were normalized to the WT value in stimulating conditions (WT high Glu). n = 6; ns p>0.05, error bars = SEM.

(Middle) Total proinsulin content in WT, *Ccdc186* KO, and *Ccdc186*(+) rescued 832/13 cells. All values were normalized to WT. n = 6; *p<0.05, ns p>0.05, error bars = SEM.

(Right) Proinsulin secretion normalized to proinsulin content in WT, *Ccdc186* KO, and *Ccdc186*(+) rescued 832/13 cells. n = 6; ns p>0.05, error bars = SEM.

We performed three biological replicates. For each replicate, the same cells were used to determine the amount of proinsulin secreted under resting conditions, stimulating conditions, and the amount of total cellular proinsulin.

C. CCDC186 is not required for proinsulin processing. Ratio of the cellular proinsulin content to the total cellular insulin content in WT, *Ccdc186* KO, and *Ccdc186*(+) rescued 832/13 cells. We

performed two biological replicates. For each replicate, the amounts of total cellular proinsulin and of total cellular insulin were determined from the same cells. n = 6, error bars = SEM.

D. *Ccdc186* KO cells have normal constitutive secretion. Secretion of ssGFP (GFP fused at its N-terminus to the signal peptide of rat ANF (Hummer et al., 2017)). The data shown are from one representative experiment of two independent experiments with similar results. n=3, error bars = SEM.

Figure 2. Insulin is abnormally distributed in *Ccdc186* KO cells.

A. Insulin is localized less in the periphery and more at or near the TGN in *Ccdc186* KO cells. Representative confocal images of WT, *Ccdc186* KO*,* and *Ccdc186*(+) rescued 832/13 cells costained for insulin and TGN38. Maximum-intensity projections. Scale bars: 5 µm. We performed three biological replicates.

B. Quantification of the colocalization between insulin and the TGN marker TGN38. Maximum-intensity projection images were obtained and Pearson's correlation coefficients were determined by drawing a line around each cell. The data shown are from one representative experiment of three biological replicates with similar results. n=12 for WT and *Ccdc186* KO; n=11 for *Ccdc186*(+), **$p < 0.01$, ***$p < 0.001$, ns $p > 0.05$, error bars = SEM.

C. *Ccdc186* KO cells have an increased proportion of insulin localized near the TGN. Fluorescence of a region of interest around the TGN divided by the fluorescence of a region of the same size in the cytoplasm. The data shown are from one experiment. n=25 for WT and *Ccdc186* KO, ***$p < 0.001$, error bars = SEM.

D. Quantification of insulin localized at or near the TGN. Maximum-intensity projection images of z-stacks encompassing the TGN38 signal were obtained and Pearson's correlation coefficients were determined by drawing a line around the TGN38 positive signal. The data shown are from one representative experiment of three biological replicates with similar results. n=20 for WT and *Ccdc186* KO, n=14 for *Ccdc186*(+); ns $p > 0.05$; error bars = SEM.

E. The localization of proinsulin is not affected by the absence of CCDC186. Representative images of WT and *Ccdc186* KO 832/13 cells costained for endogenous proinsulin and TGN38. Single slices. Scale bars: 5 μm. We performed three biological replicates.

F. Quantification of the colocalization between proinsulin and the TGN marker TGN38. The data shown are from one representative experiment of three biological replicates with similar results. n=18 for WT and n=16 for *Ccdc186* KO, ns p>0.05, error bars = SEM.

Figure 3. The levels of mature PC1/3 and CPE are decreased and mature CPE is abnormally distributed in *Ccdc186* KO cells.

A. *Ccdc186* KO cells have reduced levels of the processed form of the proprotein convertase PC1/3. Detergent lysates from WT, *Ccdc186* KO*,* and *Ccdc186*(+) rescued 832/13 cells were blotted with an antibody to PC1/3. β-tubulin was used as a loading control. The experiment was repeated four times with similar results.

B. The levels of the unprocessed form of CPE (pro-CPE) are not affected by the loss of CCDC186. Detergent lysates from WT, *Ccdc186* KO*,* and *Ccdc186*(+) rescued 832/13 cells were blotted with an antibody to pro-CPE. β-tubulin was used as a loading control. The experiment was repeated three times with similar results.

C. *Ccdc186* KO cells have reduced levels of the processed form of CPE. Detergent lysates from WT, *Ccdc186* KO*,* and *Ccdc186*(+) rescued 832/13 cells were blotted with an antibody to the N-terminus of CPE (CPE-N). β-tubulin was used as a loading control. The experiment was repeated three times.

D. *Ccdc186* KO cells have reduced levels of the processed form of CPE. Detergent lysates from WT, *Ccdc186* KO*,* and *Ccdc186*(+) rescued 832/13 cells were blotted with an antibody to the C-terminus of CPE (CPE-C). β-tubulin was used as a loading control. The experiment was repeated three times.

E. The mature processed form of CPE is localized at or near the TGN in *Ccdc186* KO cells. Representative confocal images of WT, *Ccdc186* KO*,* and *Ccdc186*(+) rescued 832/13 cells costained with the CPE C-terminal antibody (CPE-C) and TGN38. Maximum-intensity projections. Scale bars: 5 µm. The experiment was repeated three times with similar results.

F. Quantification of the colocalization between the mature form of CPE and the TGN marker TGN38. Maximum-intensity projection images were obtained and Pearson's correlation coefficients were determined by drawing a line around each cell. The data shown are from one experiment. n=44 for WT, n=37 for *Ccdc186* KO, n=31 for *Ccdc186*(+); ***$p<0.001$, ns $p>0.05$; error bars = SEM.

G. *Ccdc186* KO cells have an increased proportion of CPE localized near the TGN. Fluorescence of a region of interest around the TGN divided by the fluorescence of a region of the same size in the cytoplasm. The data shown are from one experiment. n=27 for WT and n=23 for *Ccdc186* KO, ***$p<0.001$, error bars = SEM.

H. *Ccdc186* KO cells have an increased amount of CPE localized at or near the TGN. Maximum-intensity projection images of z-stacks encompassing the TGN38 signal were obtained and Pearson's correlation coefficients were determined by drawing a line around the TGN38 positive signal. The data shown are from one experiment. n=28 for WT, n=27 for *Ccdc186* KO, n=24 for *Ccdc186*(+); *$p<0.05$, ***$p<0.001$; error bars = SEM.

I. The unprocessed form of CPE (pro-CPE) is localized at or near the TGN in both WT and *Ccdc186* KO cells. Representative confocal images of WT and *Ccdc186* KO cells costained for pro-CPE antibody and TGN38. Single slices. Scale bars: 5 µm. The experiment was repeated twice.

J. Quantification of the colocalization between pro-CPE and the TGN marker TGN38. The data shown are from one experiment. n=20 for both WT and *Ccdc186* KO, ns $p>0.05$, error bars = SEM.

Figure 4. Endogenous CCDC186 localizes to ring-like structures around the TGN/iDCV marker proinsulin.

A. Pearson's correlation coefficient was measured using confocal images (Fig S6) to quantify the colocalization between endogenous CCDC186 and markers of the different cell compartments (ERGIC53 for the ERGIC; GM130 for the cis-Golgi; EEA1 for early-endosomes; LAMP1 for lysosomes; TGN38 for the TGN; and Syn6, CgA and proinsulin for the TGN and immature DCVs). n=12 cells for TGN38 and n=9 cells for every other marker, ***p<0.001, error bars = SEM.

B. Representative structured illumination microscopy (SIM) images of 832/13 cells costained for endogenous CCDC186 and proinsulin. Maximum-intensity projections. Upper panels, scale bars: 2 μm. Lower panels, scale bars: 200 nm.

Figure 5. Overexpression of CCDC186 recruits EARP.

A. CCDC186 overexpression recruits the EARP-specific subunit VPS50. Representative confocal images of 832/13 cells transiently transfected with CCDC186::GFP and costained for GFP and endogenous VPS50. The top cell expresses a high level of CCDC186::GFP, but the bottom cell does not express a detectable level of CCDC186::GFP and presumably was not transfected. The dashed white line marks the periphery of each cell. Single slices. Scale bars: 5 μm. The experiment was repeated three times with similar results.

B. EIPR1 is required for the recruitment of VPS50 by CCDC186. Representative confocal images of WT (top) or *Eipr1* KO (bottom) 832/13 cells transiently transfected with CCDC186::GFP and costained for GFP and endogenous VPS50. Single slices. Scale bars: 5 μm. The experiment was repeated twice with similar results.

C. Overexpression or loss of CCDC186 has no effect on VPS50 level. Protein extracts from WT 832/13, *Ccdc186* KO, *Eipr1* KO and 832/13 cells transiently transfected with CCDC186::GFP

were blotted for endogenous VPS50. β-tubulin was used as a loading control. The experiment was repeated twice.

Figure 6. The CCDC186 C-terminal domain CC3 is in close proximity to carboxypeptidase D (CPD).

A. Schematic of the BioID experiment. The promiscuous biotin ligase BirA* biotinylates proteins that are in close proximity (approximately < 10 nm).

B. Domain structure of the CCDC186 fragments. The fragments were fused to the mitochondria targeting domain MAO (in black) and to the promiscuous biotin ligase BirA* (in green). The domains marked with different shades of red are predicted to be coiled-coil domains. CC=coiled-coil.

C. Mass-spectrometric (MS) analysis of biotinylated proteins from PC12 cells stably expressing CC3::BirA*::HA::MAO or the negative control BirA*::HA::MAO. The plot compares the number of unique peptides between CC3::BirA*::HA::MAO and the negative control. The data shown are from one out of three independent experiments (See Tables 1 and 7)

D. CPD is specifically recruited by CC3::BirA*::HA::MAO. Biotinylated proteins from PC12 cell extracts stably expressing BirA*::HA::MAO, CC3::BirA*::HA::MAO or CC1+2::BirA*::HA::MAO were purified with streptavidin-coated beads and immunoblotted for CPD. IN: input. CPD was found enriched in all three independent pulldowns by MS, but it could only be detected by Western blotting in the sample that showed the highest enrichment in MS (Tables 1, 7, and 8).

E. CPD is partially relocated to the mitochondria by CC3::HA::MAO, but not detectably relocated by CC1+2::HA::MAO. Representative confocal images of 832/13 cells transiently transfected with CC3::HA::MAO or CC1+2::HA::MAO, incubated with MitoTracker and costained for HA and CPD. Single slices. Scale bars: 5 μm.

Figure 7. CPD is mislocalized to mature DCVs in *Ccdc186* KO and *Eipr1* KO 832/13 cells.

A. CPD partially colocalizes with CCDC186::GFP. Representative confocal images of 832/13 cells transiently transfected with CCDC186::GFP and costained for endogenous CPD and GFP. Single slices. Scale bars: 5 μm.

B. CCDC186 and EIPR1 are required to restrict CPD localization to the TGN. Representative confocal images of WT, *Ccdc186* KO, *Ccdc186*(+) rescued, *Eipr1* KO and *Eipr1*(+) rescued 832/13 cells costained for endogenous CPD and the TGN marker TGN38. The white arrows indicate puncta in *Eipr1* KO cells. Single slices. Scale bars: 5 μm. The experiment was repeated three times with similar results.

C. In the absence of CCDC186 or EIPR1, CPD is mislocalized to insulin-positive vesicles. Representative confocal images of WT, *Ccdc186* KO, and *Eipr1* KO 832/13 cells costained for endogenous CPD and insulin. The intensity of CPD signal in WT cells was increased to show the occasional presence of low intensity CPD-positive puncta that do not overlap with insulin. The white arrows indicate CPD-positive puncta that colocalize with insulin in *Eipr1* KO cells. Single slices. Scale bars: 5 μm. The experiment was repeated three times.

Figure 8. CPD cosediments with mature DCV cargos in *Ccdc186* KO and *Eipr1* KO cells.

A. Post-nuclear supernatants from WT, *Ccdc186* KO, and *Eipr1* KO cells were separated by equilibrium sedimentation through 0.6–1.9 M sucrose. Fractions were collected from the top of the gradient and blotted with antibodies against PC1/3 (mature DCV marker) and CPD. The intensity and contrast of each blot was adjusted to show similar band intensities between cell types. The data shown are from one representative experiment of three independent experiments with similar results.

B. Band intensity for each fraction was quantified using FIJI, presented as a percentage of total band intensity, and plotted against the sucrose concentration of the fraction. The data shown are from one representative experiment of three independent experiments with similar results.

Figure 9. Model for CCDC186 and EIPR1 function in DCV maturation.

In *Ccdc186* KO cells, insulin and other luminal DCV cargos are reduced in mature DCVs in the cell periphery and mildly retained near the TGN. Proinsulin and other unprocessed cargos are not affected. Thus, CCDC186 is required for the retention of mature DCV cargos. Additionally, CPD is retained in mature DCVs in *Ccdc186* KO cells, suggesting that CCDC186 is also required for the removal of unwanted cargo from immature DCVs. *Eipr1* KO cells have similar phenotypes to *Ccdc186* KO cells, suggesting that CCDC186 and EIPR1/EARP act together to control the post-Golgi retention and removal of DCV cargos.

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. *cccp-1* functions in the same pathway as *eipr-1* and *vps-50* to control locomotion and dense-core vesicle cargo sorting in *C. elegans*.

A. *cccp-1* acts in the same pathway as *eipr-1* and *vps-50* to control locomotion. The *cccp-1*(*ox334*) mutation does not enhance the slow locomotion phenotype of either *eipr-1*(*tm4790*) or *vps-50*(*ok2627*) mutants. Error bars = SEM; n = 10.

B. *cccp-1* acts in the same genetic pathway as *eipr-1* and *vps-50* to control DCV cargo trafficking. Representative images of NLP-21::Venus (*nuIs183* transgene) fluorescence in motor neuron axons of the dorsal nerve cord. Maximum-intensity projections. Scale bar: 10 µm.

C. Quantification of NLP-21::Venus fluorescence levels in the dorsal nerve cord. Double mutants of *cccp-1*(*ox334*) with *eipr-1*(*tm4790*) or *vps-50*(*ok2627*) are not significantly different from the single mutants (***, $p < 0.001$; ns, not significant, $p > 0.05$). Error bars = SEM; n = 10.

Figure S2. Generation of a *Ccdc186* KO 832/13 beta-cell line.

A. CCDC186 is expressed in multiple mouse tissues. Protein extracts from different mouse tissues were blotted for CCDC186 using a commercial antibody. β-tubulin was used as a loading control. In two independent blots run from the same protein samples, we saw the same small differences in the size of the CCDC186 band in different tissues.

B. *Ccdc186* KO cells do not express CCDC186. Protein extracts from WT, *Ccdc186* KO, and *Ccdc186* KO 832/13 cells stably expressing wild type CCDC186 (*Ccdc186*(+)) were blotted with anti-CCDC186 antibody. The faint band of slightly larger size in the *Ccdc186* KO sample was not seen in other repetitions of this experiment and is presumed to be a nonspecific band. β-tubulin was used as a loading control.

C. Total cDNA from WT 832/13 and *Ccdc186* KO cells was PCR amplified using primers that detect the *Ccdc186* cDNA.

D. WT, *Ccdc186* KO, and *Ccdc186*(+) rescued cells were stained with anti-CCDC186 antibody. Single slices. Wide-field fluorescence compound microscopy. Scale bars: 10 µm.

Figure S3. The acidity of the Golgi does not significantly change in *Ccdc186* KO cells.

A. Example calibration curve for Golgi pH measurements based on measuring the fluorescence of TGN-targeted pHluorin in solutions of defined pH. For each biological replicate of WT and *Ccdc186* KO 832/13 cells tested for pH, an individual calibration curve such as the one shown was obtained from WT and *Ccdc186* KO cells grown in the same 96-well plate as the test samples and exposed to buffers of decreasing pH (8.5–5.5) in the presence of nigericin and monensin (see Materials and Methods). Each data point shows the mean of the fluorescent measurements (in arbitrary units (AU)) from cells in three different wells. Error bars=SEM.

B. The late-Golgi compartment is not more acidic in *Ccdc186* KO 832/13 cells. The fluorescence of TGN-targeted pHluorin was measured in WT and *Ccdc186* KO cells. The absolute pH value of each sample was extrapolated from a paired calibration curve (as in A). The data show mean ± SEM; n=8 for WT and *Ccdc186* KO.

Figure S4. CPE and insulin almost fully colocalize.

Representative confocal images of 832/13 cells costained for endogenous insulin and endogenous CPE using the CPE-C antibody. Maximum-intensity projections. Scale bars: 5 µm.

Figure S5. Pulse-chase experiment monitoring the exit of DCV cargo from the TGN.

A. The exogenous DCV cargo ANF::GFP is mostly localized near the TGN in *Ccdc186* KO cells at steady state. Representative images of WT and *Ccdc186* KO 832/13 cells transfected with

ANF::GFP and costained for GFP and TGN38. Maximum-intensity projections. Scale bars: 10 µm.

B. Experimental set up: cells were transiently transfected with ANF::GFP. Incubation at 20ºC for 2 hours resulted in blocking ANF::GFP from exiting the TGN (pulse). 30 minutes before the end of the low temperature incubation, cycloheximide was added to inhibit synthesis of new ANF::GFP. Following the low temperature block, cells were incubated at 37ºC (chase) before fixation and imaging.

C. Representative images of the three cell categories used for qualitative assessment of TGN exit: those with most of the fluorescence concentrated at the TGN (Golgi-like), those where the TGN was still apparent but a large portion of the fluorescence was at the cell periphery (Intermediate), and those where the TGN was no longer apparent (Periphery). Single slices. Scale bar: 5 µm.

D. ~70 cells for each time point and genotype were imaged blindly. The data are plotted as percentage of each phenotype at each time point. The experiment was repeated three times with similar results. Error bars = SEM.

Figure S6. Endogenous CCDC186 localizes to a perinuclear area near the TGN and immature DCVs.

Representative confocal images of 832/13 cells costained for endogenous CCDC186 and markers for different cell compartments: the ERGIC marker ERGIC53, the cis-Golgi marker GM130, the early-endosome marker EEA1, the lysosomal marker LAMP1, the TGN marker TGN38, the TGN/iDCV markers syntaxin 6 (Syn6) and proinsulin, and the DCV markers chromogranin A (CgA) and insulin. Single slices. Scale bars: 5 µm.

Figure S7. CCDC186 colocalizes with RAB2A and RAB2B but does not localize to early, late, or recycling endosomes.

A. Representative confocal images of 832/13 cells incubated with Alexa-568 labelled transferrin (Tf) (recycling and early endosomes) or transiently transfected with GFP-tagged RAB4A (recycling endosomes) and costained for Tf or GFP and endogenous CCDC186. The intensity and contrast of the images showing GFP-tagged Rab proteins were adjusted to remove the high background in the cells overexpressing Rab proteins. Single slices. Scale bars: 5 µm.

B. Representative confocal images of 832/13 cells transiently transfected with Myc-tagged CCDC186, incubated with Alexa568-labeled transferrin (Tf, recycling and early endosomes, top panel) or cotransfected with GFP-tagged RAB4A (recycling endosomes, bottom panel). The cells were costained for Myc and Tf or Myc and GFP. The intensity and contrast of the images showing GFP-tagged Rab proteins were adjusted to remove the high background in the cells overexpressing Rab proteins. Single slices. Scale bars: 5 µm.

C. Representative confocal images of 832/13 cells transiently transfected with GFP-tagged RAB5A (early endosomes) or RAB7A (late endosomes) and costained for GFP and endogenous CCDC186. The intensity and contrast of the images showing GFP-tagged Rab proteins were adjusted to remove the high background in the cells overexpressing Rab proteins. Single slices. Scale bars: 5 µm.

D. Representative confocal images of 832/13 cells transiently cotransfected with Myc-tagged CCDC186 and GFP-tagged RAB5A (early endosomes) or RAB7A (late endosomes). Cells were costained for Myc and GFP. The intensity and contrast of the images showing GFP-tagged Rab proteins were adjusted to remove the high background in the cells overexpressing Rab proteins. Single slices. Scale bars: 5 µm.

E, F. Representative confocal images of 832/13 cells transiently transfected with GFP-tagged RAB2A (E) or RAB2B (F) and costained for GFP and endogenous CCDC186. Single slices. Scale bars: 5 µm.

G. CCDC186 does not colocalize with golgin-97. Cells were costained for endogenous CCDC186 and endogenous golgin-97. Single slices. Scale bars: 2 µm.

Figure S8. Endogenous CCDC186 localizes to ring-like structures around the TGN/iDCV marker proinsulin.

Representative stimulated emission depletion (STED) microscopy images of 832/13 cells costained for endogenous CCDC186 and endogenous proinsulin. Maximum-intensity projections. Scale bars: 2 µm.

Figure S9. CCDC186 recruits EARP-positive membranes through its middle CC2 domain.

A. Domain structure of the CCDC186 fragments. The fragments were fused to GFP at their C-terminus. The domains marked with different shades of red are predicted to be coiled-coil (CC) domains.

B. Representative confocal images of 832/13 cells transiently transfected with GFP-tagged fragments of CCDC186 and costained for endogenous VPS50 and GFP. Single slices. Scale bars: 5 µm. The experiment was repeated twice.

Figure S10. CCDC186 is not required for endocytic recycling of transferrin.

FACS analysis of Alexa488–transferrin (Tf) released from 832/13 WT, *Ccdc186* KO, and *Eipr1* KO cells after 25 minutes uptake. The plot represents the median of Alexa 488–Tf intensity of the population (~20,000 cells) as a function of time (minutes) normalized to the median intensity at t = 0. The difference between WT and *Eipr1* KO at the 25 minute time point was marginally statistically significant (p=0.0571, Mann–Whitney test). Error bars = SEM from four independent experiments.

Figure S11. Mitochondria relocation of CCDC186 fragments.

A. Schematic of the CCDC186 mitochondria relocation experiment.

B. Domain structure of CCDC186 fragments fused at their C-terminus to the transmembrane domain of monoamine oxidase (MAO, black). The domains marked with different shades of red are predicted coiled-coil domains (CC).

C. CCDC186 fragments fused to MAO are relocated to the mitochondria. Representative confocal images of 832/13 cells transiently transfected with CC3::HA::MAO (upper panel) or CC1+2::HA::MAO (lower panel), incubated with MitoTracker and costained for HA and TGN38. The intensity and contrast of the images showing MitoTracker staining were adjusted to remove the high background. Of note, we observed that overexpression of CC3::HA::MAO caused clustering of mitochondria in a perinuclear region adjacent to the TGN marker TGN38, but CC1+2::HA::MAO had no apparent effect on mitochondrial distribution or structure. Single slices. Scale bars: 5 µm. The experiment was repeated twice.

Figure S12. The CCDC186 fragment CC3 fused to MAO relocates the GTP-bound forms of RAB2A and RAB2B to mitochondria.

A. Representative confocal images of 832/13 cells transiently cotransfected with CC3::HA::MAO and GFP::RAB2A(Q65L), and costained for HA and GFP. Single slices. Scale bars: 5 µm. The experiment was repeated twice.

B. Representative confocal images of 832/13 cells transiently cotransfected with CC1+2::HA::MAO and GFP::RAB2A(Q65L), and costained for HA and GFP. Single slices. Scale bars: 5 µm. The experiment was repeated twice.

C. Representative confocal images of 832/13 cells transiently cotransfected with CC3::HA::MAO and GFP::RAB2B(Q65L), and costained for HA and GFP. Single slices. Scale bars: 5 µm. The experiment was repeated twice.

D. Representative confocal images of 832/13 cells transiently cotransfected with CC1+2::HA::MAO and GFP::RAB2B(Q65L), and costained for HA and GFP. Single slices. Scale bars: 5 µm. The experiment was repeated twice.

Figure S13. Volcano plot representation of the CC3 BioID experiment.

Volcano plot showing the Student's t-test p-value for each protein as a measure of that protein's enrichment in CC3::BirA*::HA::MAO streptavidin pulldowns compared to the negative control BirA*::HA::MAO. Lysates from cells overexpressing CC3::BirA*::HA::MAO and BirA*::HA::MAO were subjected to affinity enrichment using streptavidin. Proteins with a Log2 ratio (CC3::BirA*::HA::MAO / BirA*::HA::MAO) larger than 1.8 are labeled. See Table 8 for the raw data.

Figure S14. CCDC186 and EIPR1 are not required for CD-MPR trafficking.

Representative confocal images of WT, *Ccdc186* KO*,* and *Eipr1* KO 832/13 cells transiently transfected with CD-MPR::GFP and costained for insulin and GFP. Single slices. Scale bars: 5 µm. The experiment was repeated twice.

TABLES

Table 1. Unique peptide count organized in the descending order of the difference in number of unique peptides from CC3::BirA*::HA::MAO and the negative control BirA*::HA::MAO. The table shows the hits where the difference was 3 or greater. These data are from the BioID experiment that showed the largest amount of enrichment for CPD.

Table 2. Unique peptide count organized in the descending order of the difference in number of peptides from CC1+2::BirA*::HA::MAO and the negative control BirA*::HA::MAO. The table shows the hits where the difference was 3 or greater.

Table 3. *C. elegans* strain list

N2 Bristol wild-type

EG334 *cccp-1(ox334)* III

EG6007 *vps-50(ok2627)* III

EG6939 *eipr-1(tm4790)* I

XZ1924 *vps-50(ok2627)* III *cccp-1(ox334)* III

XZ2108 *eipr-1(tm4790)* I ; *cccp-1(ox334)* III

KG1395 *nuIs183[Punc-129::NLP-21-Venus, Pmyo-2::GFP]* III

EG5258 *nuIs183[Punc-129::NLP-21-Venus, Pmyo-2::GFP]* III *cccp-1(ox334)* III

XZ1055 *eipr-1(tm4790)* I ; *nuIs183[Punc-129::NLP-21-Venus, Pmyo-2::GFP]* III

XZ1607 *nuIs183[Punc-129::NLP-21-Venus, Pmyo-2::GFP]* III *vps-50(ok2627)* III

XZ1918 *nuIs183[Punc-129::NLP-21-Venus, Pmyo-2::GFP]* III *vps-50(ok2627)* III *cccp-1(ox334)*

III

XZ2109 *eipr-1(tm4790)* I ; *nuIs183[Punc-129::NLP-21-Venus, Pmyo-2::GFP]* III *cccp-1(ox334)*

III

Table 4. Plasmid list

Plasmid name	Vector backbone	Description
pEGFP-N1		GFP expression vector, a gift from Suzanne Hoppins
pET50	pEGFP-N1	CCDC186::GFP full length, rat CCDC186 cDNA
pJC207	pEGFP-N1	GFP::RAB5A, RAB5A cDNA was PCR amplified from 832/13 cDNA library
pJC205	pEGFP-N1	GFP::RAB4A, RAB4A cDNA PCR amplified from 832/13 cDNA library
pJC208	pEGFP-N1	GFP::RAB7A, RAB7A cDNA was PCR amplified from 832/13 cDNA library
ANF::GFP	pEGFP-N1	ANF::GFP (rat atrial natriuretic factor fused to GFP, (Hummer et al., 2017)
ssGFP	pCAGGS	signal sequence of rat ANF fused to GFP in the chicken actin-based vector pCAGGS (Hummer et al., 2017)
pET55	pEGFP-N1	GFP::RAB2A, RAB2A cDNA was PCR amplified from 832/13 cDNA library
pET54	pEGFP-N1	GFP::RAB2B, RAB2B cDNA was PCR amplified from 832/13 cDNA library
pJC239	pCMV-Myc	Myc::CCDC186
pJC271	pCDNA3.1	CC3::CCDC186(750-922)::HA::MAO
pJC272	pCDNA3.1	CC1+2::CCDC186(1-749)::HA::MAO
pJC280	pCDNA3.1	CC3::CCDC186(759-922)::BirA*::HA::MAO
pJC281	pCDNA3.1	CC1+2::CCDC186(1-749)::BirA*::HA::MAO
pJJS112	pCDNA3.1	BirA*::HA::MAO, a gift from Sean Munro
pJC279	pEGFP-N1	CD-MPR::GFP, CD-MPR cDNA was PCR amplified from 832/13 cDNA library
pPUR		A gift from Richard Palmiter
pBABE-hygro		A gift from Suzanne Hoppins
pSpCas9(BB)-2A-GFP		PX458 (Addgene)
pET221	pBABE-hygro	CCDC186 in pBABE-hygro
pET64	pSpCas9(BB)-2A-GFP	CCDC186 guide RNA in pSpCas9(BB)
pET70	pPUR	CCDC186 HR (homologous recombination cassette) in pPUR backbone
FUGW		3rd generation lentiviral plasmid with hUbC-driven EGFP; can be used for cDNA expression (Addgene #14883)
psPAX2		2nd generation lentiviral packaging plasmid. Can be used with 2nd or 3rd generation lentiviral vectors and envelope expressing plasmid (Addgene #12259)
pVSVG		Envelope protein for producing lentiviral and MuLV retroviral particles. Use in conjunction with a packaging vector such as pCMV-dR8.2 dvpr (lentiviral) or pUMVC (MuLV retroviral). (Addgene #8454)

Antibody target	Manufacturer and catalog number	Notes
St6Gal1::pHluorin	FUGW	17-residue transmembrane domain of beta-galactoside alpha 2,6-sialyltransferase fused to pHluorin.

Table 5. Primary antibody list

Antibody target	Manufacturer and catalog number	Notes
CCDC186	Novus #NBP1-90440	Rabbit polyclonal, 1:150 for IF and 1:1,000 for WB
TGN38	Sigma #T9826	Rabbit polyclonal, 1:350 for IF
EEA1	BD Biosciences #610456	Mouse monoclonal, 1:100 for IF
GM130	BD Biosciences #610822	Mouse monoclonal, 1:100 for IF, but works at 1:300
ERGIC53	Santa Cruz #sc-398893	Mouse monoclonal, 1:50 for IF
LAMP-1	DHSB	Mouse monoclonal, 1:50 for IF
GFP	Santa Cruz #sc-9996	Mouse monoclonal, 1:200 to 1:400 for IF, 1:1,000 for WB
GFP	Santa Cruz #sc-8334	Rabbit polyclonal, 1:200 to 1:400 for IF, 1:1,000 for WB
Insulin	Sigma #I2018	Mouse monoclonal, 1:400 for IF
Proinsulin	Abcam #ab8301	Mouse monoclonal, 1:200 for IF
Syntaxin 6	Abcam #ab12370	Mouse monoclonal, 1:100 for IF
TGN38	Novus #NB300-575S	Mouse monoclonal, 1:500 for IF
CgA	Santa Cruz #sc-393941	Mouse monoclonal, 1:200 for IF
Myc	Santa Cruz sc-40	Mouse monoclonal, 1:200 for IF, 1:1,000 for WB
Rab2	Santa Cruz	Rabbit polyclonal, 1:200-1:500 for WB
GFP	Roche #11814460001	Mouse monoclonal, 1:1,000 for WB
Beta-tubulin	DHSB E7	Mouse monoclonal, 1:1,000 for WB
Beta-tubulin	ThermoFisher , BT7R, #MA5-16308	Mouse monoclonal, 1:1,000 for WB
HA	Roche #11867423001	Rat monoclonal, 1:150 for IF and 1:1,000 for WB
CPD*	A gift from Lloyd Fricker AE160	Rabbit polyclonal, 1:200 for IF and 1:1,000 for WB
VPS50	Sigma HPA026679	Rabbit polyclonal, 1:100 for IF and 1:1,000 for WB
PCSK1 (PC1/3)	Sigma SAB1100415	Rabbit polyclonal, 1:1,000 for WB
CPE-N terminus**	A gift from Lloyd Fricker	Rabbit polyclonal, 1:1,000 for WB, 1:100 for IF
Pro-CPE form***	A gift from Lloyd Fricker	Rabbit polyclonal, 1:1,000 for WB, 1:100 for IF
CPE-C terminus****	A gift from Lloyd Fricker	Rabbit polyclonal, 1:1,000 for WB, 1:100 for IF

* Antiserum was generated to the C-terminal 57 amino acids of mammalian CPD (cytosolic tail,

(Song and Fricker, 1996).

** Antiserum was raised to the N-terminal 15 amino acids of bovine CPE (Fricker et al., 1990).

*** Antiserum to pro-CPE was generated to the peptide QEPGAPAAGMRRC coupled to maleimide-activated keyhole limpet hemocyanin (KLH). This peptide corresponds to the 12 residues of mouse/rat/human pro-CPE with an added Cys on the C-term (for coupling to the carrier protein, KLH).

**** Antiserum was raised against the 9-residue peptide KMMSETLNF corresponding to the C terminus of mouse CPE (Fricker et al., 1996).

Table 6. Secondary antibody list

Antibody target	Manufacturer and catalog number	Notes
Anti-mouse Alexa488	Jackson immunoresearch #115-545-146	1:1000 for IF
Anti-mouse Rhodamine Rex-X	Jackson immunoresearch #715-295-150	1:1000 for IF
Anti-rabbit Alexa488	Jackson immunoresearch #711-545-152	1:1000 for IF, (1:100 for STED microscopy)
Anti-rabbit Rhodamine TRITC	Jackson immunoresearch #111-025-144	1:1000 for IF
Anti-rat Dylight550	ThermoFisher #SA5-10027	Highly cross-adsorbed, 1:200 to 1:500 for IF
Anti-rabbit Atto647	A gift from Joshua Vaughan's lab	1:200 for IF
Anti-mouse Dylight488	ThermoFisher #SA5-10166	Highly cross-adsorbed, used at 1:200 to 1:500
Anti-mouse Alexa555	A gift from Joshua Vaughan's lab	Used at 1:10 for STED microscopy
Goat anti-rabbit Alexa680	Jackson immunoresearch #111-625-144	1:20,000 for western
Goat anti-mouse Alexa680	Jackson immunoresearch #115-625-166	1:20,000 for western
Donkey anti-mouse Alexa790	Jackson immunoresearch #715-655-150	1:20,000 for western

Alexa790 Streptavidin	Jackson immunoresearch #016-650-084	1:10,000 for western

Table 7

The table contains all the data from the experiments shown in Tables 1 and 2

Table 8

The table contains Z-score normalized log2 LFQ intensities from MaxQuant output, Student's t-test p-values and ratios (CC3::BirA*::HA::MAO vs BirA*::HA::MAO). For calculation of p-values and ratios, see Materials and Methods

REFERENCES

Ailion, M., Hannemann, M., Dalton, S., Pappas, A., Watanabe, S., Hegermann, J., Liu, Q., Han, H.-F., Gu, M., Goulding, M.Q., et al. (2014). Two Rab2 Interactors Regulate Dense-Core Vesicle Maturation. Neuron *82*, 167–180.

Arvan, P., and Castle, D. (1998). Sorting and storage during secretory granule biogenesis: looking backward and looking forward. Biochem. J. *332*, 593–610.

Arvan, P., and Halban, P.A. (2004). Sorting Ourselves Out: Seeking Consensus on Trafficking in the Beta-Cell. Traffic *5*, 53–61.

Bäck, N., Rajagopal, C., Mains, R.E., and Eipper, B.A. (2010). Secretory Granule Membrane Protein Recycles through Multivesicular Bodies. Traffic *11*, 972–986.

Bonifacino, J.S., and Traub, L.M. (2003). Signals for Sorting of Transmembrane Proteins to Endosomes and Lysosomes. Annu. Rev. Biochem. *72*, 395–447.

Borgonovo, B., Ouwendijk, J., and Solimena, M. (2006). Biogenesis of secretory granules. Curr. Opin. Cell Biol. *18*, 365–370.

Brenner, S. (1974). The Genetics of Caenorhabditis Elegans. Genetics *77*, 71–94.

Cao, M., Mao, Z., Kam, C., Xiao, N., Cao, X., Shen, C., Cheng, K.K.Y., Xu, A., Lee, K.-M., Jiang, L., et al. (2013). PICK1 and ICA69 Control Insulin Granule Trafficking and Their Deficiencies Lead to Impaired Glucose Tolerance. PLoS Biol *11*, e1001541.

Cattin-Ortolá, J., Topalidou, I., Dosey, A., Merz, A.J., and Ailion, M. (2017). The dense-core vesicle maturation protein CCCP-1 binds RAB-2 and membranes through its C-terminal domain. Traffic *18*, 720–732.

Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M. (2014). Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. Mol. Cell. Proteomics *13*, 2513–2526.

Davidson, H.W., Rhodes, C.J., and Hutton, J.C. (1988). Intraorganellar calcium and pH control proinsulin cleavage in the pancreatic β cell via two distinct site-specific endopeptidases. Nature *333*, 93–96.

Dittie, A.S., Klumperman, J., and Tooze, S.A. (1999). Differential distribution of mannose-6-phosphate receptors and furin in immature secretory granules. J. Cell Sci. *112*, 3955–3966.

Du, W., Zhou, M., Zhao, W., Cheng, D., Wang, L., Lu, J., Song, E., Feng, W., Xue, Y., Xu, P., et al. (2016). HID-1 is required for homotypic fusion of immature secretory granules during maturation. eLife *5*, e18134.

Edwards, S.L., Charlie, N.K., Richmond, J.E., Hegermann, J., Eimer, S., and Miller, K.G. (2009). Impaired dense core vesicle maturation in Caenorhabditis elegans mutants lacking Rab2. J. Cell Biol. *186*, 881–895.

Emperador-Melero, J., Huson, V., Weering, J. van, Bollmann, C., Mollard, G.F. von, Toonen, R.F., and Verhage, M. (2018). Vti1a/b regulate synaptic vesicle and dense core vesicle secretion via protein sorting at the Golgi. Nat. Commun. *9*, 3421.

Eng, F.J., Varlamov, O., and Fricker, L.D. (1999). Sequences within the Cytoplasmic Domain of Gp180/Carboxypeptidase D Mediate Localization to the Trans-Golgi Network. Mol. Biol. Cell *10*, 35–46.

Feng, L., and Arvan, P. (2003). The Trafficking of α1-Antitrypsin, a Post-Golgi Secretory Pathway Marker, in INS-1 Pancreatic Beta Cells. J. Biol. Chem. *278*, 31486–31494.

Fricker, L.D. (2013). Chapter 235 - Carboxypeptidases E and D. In Handbook of Biologically Active Peptides (Second Edition), A.J. Kastin, ed. (Boston: Academic Press), pp. 1715–1720.

Fricker, L.D., and Devi, L. (1993). Posttranslational processing of carboxypeptidase E, a neuropeptide-processing enzyme, in AtT-20 cells and bovine pituitary secretory granules. J. Neurochem. *61*, 1404–1415.

Fricker, L.D., Das, B., and Angeletti, R.H. (1990). Identification of the pH-dependent membrane anchor of carboxypeptidase E (EC 3.4.17.10). J. Biol. Chem. *265*, 2476–2482.

Fricker, L.D., Berman, Y.L., Leiter, E.H., and Devi, L.A. (1996). Carboxypeptidase E Activity Is Deficient in Mice with the fat Mutation Effect on Peptide Processing. J. Biol. Chem. *271*, 30619–30624.

Gershlick, D.C., Schindler, C., Chen, Y., and Bonifacino, J.S. (2016). TSSC1 is novel component of the endosomal retrieval machinery. Mol. Biol. Cell *27*, 2867–2878.

Gibson, D.G., Young, L., Chuang, R.-Y., Venter, J.C., Hutchison, C.A., and Smith, H.O. (2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat. Methods *6*, 343–345.

Gillingham, A.K., Sinka, R., Torres, I.L., Lilley, K.S., and Munro, S. (2014). Toward a Comprehensive Map of the Effectors of Rab GTPases. Dev. Cell *31*, 358–373.

Gondré-Lewis, M.C., Park, J.J., and Loh, Y.P. (2012). Cellular mechanisms for the biogenesis and transport of synaptic and dense-core vesicles. Int. Rev. Cell Mol. Biol. *299*, 27–115.

Holst, B., Madsen, K.L., Jansen, A.M., Jin, C., Rickhag, M., Lund, V.K., Jensen, M., Bhatia, V., Sørensen, G., Madsen, A.N., et al. (2013). PICK1 Deficiency Impairs Secretory Vesicle Biogenesis and Leads to Growth Retardation and Decreased Glucose Tolerance. PLoS Biol *11*, e1001542.

Hummer, B.H., Leeuw, N.F. de, Burns, C., Chen, L., Joens, M.S., Hosford, B., Fitzpatrick, J.A.J., and Asensio, C.S. (2017). HID-1 controls formation of large dense core vesicles by influencing cargo sorting and trans-Golgi network acidification. Mol. Biol. Cell *28*, 3870–3880.

Hussain, S.S., Harris, M.T., Kreutzberger, A.J.B., Inouye, C.M., Doyle, C.A., Castle, A.M., Arvan, P., and Castle, J.D. (2018). Control of insulin granule formation and function by the ABC transporters ABCG1 and ABCA1 and by oxysterol binding protein OSBP. Mol. Biol. Cell *29*, 1238–1257.

Kim, T., Gondré-Lewis, M.C., Arnaoutova, I., and Loh, Y.P. (2006). Dense-Core Secretory Granule Biogenesis. Physiology *21*, 124–133.

Klumperman, J., Kuliawat, R., Griffith, J.M., Geuze, H.J., and Arvan, P. (1998). Mannose 6–Phosphate Receptors Are Sorted from Immature Secretory Granules via Adaptor Protein AP-1, Clathrin, and Syntaxin 6–positive Vesicles. J. Cell Biol. *141*, 359–371.

Kögel, T., Rudolf, R., Hodneland, E., Copier, J., Regazzi, R., Tooze, S.A., and Gerdes, H.-H. (2013). Rab3D Is Critical for Secretory Granule Maturation in PC12 Cells. PLoS ONE *8*, e57321.

Kögel, T., Rudolf, R., Hodneland, E., Hellwig, A., Kuznetsov, S., Seiler, F., Söllner T.H., Barroso, J., and Gerdes, H.-H. (2010). Distinct Roles of Myosin Va in Membrane Remodeling and Exocytosis of Secretory Granules. Traffic *11*, 637–650.

Kuroki, K., Eng, F., Ishikawa, T., Turck, C., Harada, F., and Ganem, D. (1995). gp180, a host cell glycoprotein that binds duck hepatitis B virus particles, is encoded by a member of the carboxypeptidase gene family. J. Biol. Chem. *270*, 15022–15028.

Naggert, J.K., Fricker, L.D., Varlamov, O., Nishina, P.M., Rouille, Y., Steiner, D.F., Carroll, R.J., Paigen, B.J., and Leiter, E.H. (1995). Hyperproinsulinaemia in obese fat/fat mice associated with a carboxypeptidase E mutation which reduces enzyme activity. Nat. Genet. *10*, 135–142.

Orci, L., Ravazzola, M., Amherdt, M., Madsen, O., Perrelet, A., Vassalli, J.D., and Anderson, R.G. (1986). Conversion of proinsulin to insulin occurs coordinately with acidification of maturing secretory vesicles. J. Cell Biol. *103*, 2273–2281.

Orci, L., Ravazzola, M., Storch, M.J., Anderson, R.G., Vassalli, J.D., and Perrelet, A. (1987). Proteolytic maturation of insulin is a post-Golgi event which occurs in acidifying clathrin-coated secretory vesicles. Cell *49*, 865–868.

Paquin, N., Murata, Y., Froehlich, A., Omura, D.T., Ailion, M., Pender, C.L., Constantine-Paton, M., and Horvitz, H.R. (2016). The Conserved VPS-50 Protein Functions in Dense-Core Vesicle Maturation and Acidification and Controls Animal Behavior. Curr. Biol. *26*, 862–871.

Ran, F.A., Hsu, P.D., Wright, J., Agarwala, V., Scott, D.A., and Zhang, F. (2013). Genome engineering using the CRISPR-Cas9 system. Nat. Protoc. *8*, 2281–2308.

Rappsilber, J., Mann, M., and Ishihama, Y. (2007). Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. Nat. Protoc. *2*, 1896–1906.

Roux, K.J., Kim, D.I., Raida, M., and Burke, B. (2012). A promiscuous biotin ligase fusion protein identifies proximal and interacting proteins in mammalian cells. J. Cell Biol. *196*, 801–810.

Schindler, C., Chen, Y., Pu, J., Guo, X., and Bonifacino, J.S. (2015). EARP is a multisubunit tethering complex involved in endocytic recycling. Nat. Cell Biol. *17*, 639–650.

Shin, J.J.H., Gillingham, A.K., Begum, F., Chadwick, J., and Munro, S. (2017). TBC1D23 is a bridging factor for endosomal vesicle capture by golgins at the *trans*-Golgi. Nat. Cell Biol. *19*, 1424–1432.

Smeekens, S.P., and Steiner, D.F. (1990). Identification of a human insulinoma cDNA encoding a novel mammalian protein structurally related to the yeast dibasic processing protease Kex2. J. Biol. Chem. *265*, 2997–3000.

Smeekens, S.P., Montag, A.G., Thomas, G., Albiges-Rizo, C., Carroll, R., Benig, M., Phillips, L.A., Martin, S., Ohagi, S., and Gardner, P. (1992). Proinsulin processing by the subtilisin-related proprotein convertases furin, PC2, and PC3. Proc. Natl. Acad. Sci. *89*, 8822–8826.

Sobajima, T., Yoshimura, S., Maeda, T., Miyata, H., Miyoshi, E., and Harada, A. (2018). The Rab11-binding protein RELCH/KIAA1468 controls intracellular cholesterol distribution. J. Cell Biol. 217, 1777-1796.

Song, L., and Fricker, L. (1995). Processing of procarboxypeptidase E into carboxypeptidase E occurs in secretory vesicles. J. Neurochem. *65*, 444–453.

Song, L., and Fricker, L.D. (1996). Tissue Distribution and Characterization of Soluble and Membrane-bound Forms of Metallocarboxypeptidase D. J. Biol. Chem. *271*, 28884–28889.

Steiner, D. F., Chan, S. J. and Rubenstein, A. H. (2011). Biosynthesis of Insulin. In Comprehensive Physiology, R. Terjung (Ed.). doi:10.1002/cphy.cp070203

Sumakovic, M., Hegermann, J., Luo, L., Husson, S.J., Schwarze, K., Olendrowitz, C., Schoofs, L., Richmond, J., and Eimer, S. (2009). UNC-108/RAB-2 and its effector RIC-19 are involved in dense core vesicle maturation in Caenorhabditis elegans. J. Cell Biol. *186*, 897–914.

Tooze, S.A. (1998). Biogenesis of secretory granules in the trans-Golgi network of neuroendocrine and endocrine cells. Biochim. Biophys. Acta - Mol. Cell Res. *1404*, 231–244.

Tooze, S.A., Martens, G.J.M., and Huttner, W.B. (2001). Secretory granule biogenesis: rafting to the SNARE. Trends Cell Biol. *11*, 116–122.

Topalidou, I., Cattin-Ortolá, J., Pappas, A.L., Cooper, K., Merrihew, G.E., MacCoss, M.J., and Ailion, M. (2016). The EARP Complex and Its Interactor EIPR-1 Are Required for Cargo Sorting to Dense-Core Vesicles. PLOS Genet. *12*, e1006074.

Topalidou, I., Cattin-Ortolá, J., and Ailion, M. (2018). EIPR1 controls dense-core vesicle cargo sorting and EARP complex localization in insulinoma cells. BioRxiv 374488. https://doi.org/10.1101/374488

Varlamov, O., and Fricker, L.D. (1998). Intracellular trafficking of metallocarboxypeptidase D in AtT-20 cells: localization to the trans-Golgi network and recycling from the cell surface. J. Cell Sci. *111*, 877–885.

Varlamov, O., Eng, F.J., Novikova, E.G., and Fricker, L.D. (1999). Localization of Metallocarboxypeptidase D in AtT-20 Cells. Potential Role in Prohormone Processing. J. Biol. Chem. *274*, 14759–14767.

Vindrola, O., and Lindberg, I. (1992). Biosynthesis of the prohormone convertase mPC1 in AtT-20 cells. Mol. Endocrinol. *6*, 1088–1094.

Vo, Y.P., Hutton, J.C., and Angleson, J.K. (2004). Recycling of the dense-core vesicle membrane protein phogrin in Min6 β-cells. Biochem. Biophys. Res. Commun. *324*, 1004–1010.

Wendler, F., Page, L., Urbé, S., and Tooze, S.A. (2001). Homotypic Fusion of Immature Secretory Granules During Maturation Requires Syntaxin 6. Mol. Biol. Cell *12*, 1699–1709.

Wong, M., and Munro, S. (2014). The specificity of vesicle traffic to the Golgi is encoded in the golgin coiled-coil proteins. Science *346*, 1256898.

Wong, S.H., Low, S.H., and Hong, W. (1992). The 17-residue transmembrane domain of beta-galactoside alpha 2,6-sialyltransferase is sufficient for Golgi retention. J. Cell Biol. *117*, 245–258.

Xin, X., Varlamov, O., Day, R., Dong, W., Bridgett, M.M., Leiter, E.H., and Fricker, L.D. (1997). Cloning and Sequence Analysis of cDNA Encoding Rat Carboxypeptidase D. DNA Cell Biol. *16*, 897–909.

Zhang, X., Jiang, S., Mitok, K.A., Li, L., Attie, A.D., and Martin, T.F.J. (2017). BAIAP3, a C2 domain–containing Munc13 protein, controls the fate of dense-core vesicles in neuroendocrine cells. J Cell Biol *216*, 2151–2166.

Zhou, Y., and Lindberg, I. (1993). Purification and characterization of the prohormone convertase PC1(PC3). J. Biol. Chem. *268*, 5615–5623.

Figure 1

Figure 2

Figure 3

A

PC1/3 — 65 kDa
WT: 1 Ccdc186 KO: 0.3 CCDC186(+): 0.82
β-tub — 50 kDa

B

Pro-CPE — 55 kDa
WT: 1 Ccdc186 KO: 0.92 CCDC186(+): 0.91
β-tub — 50 kDa

C

CPE-N — 55 kDa
WT: 1 Ccdc186 KO: 0.33 CCDC186(+): 0.83
β-tub — 50 kDa

D

CPE-C — 55 kDa
WT: 1 Ccdc186 KO: 0.34 CCDC186(+): 1.23
β-tub — 50 kDa

E

CPE-C | TGN38 | Merge
WT

CPE-C | TGN38 | Merge
Ccdc186 KO

CPE-C | TGN38 | Merge
CCDC186(+)

F

Total CPE-C – TGN38

Pearson's correlation coefficient CPE-C – TGN38

ns
*** ***

WT   Ccdc186 KO   CCDC186(+)

G

Golgi CPE-C/ Cytoplasmic CPE-C

Normalized intensity

***

WT   Ccdc186 KO

H

Golgi CPE-C – TGN38

Pearson's correlation coefficient

*** *

WT   Ccdc186 KO   CCDC186(+)

I

Pro-CPE | TGN38 | Merge
WT

Pro-CPE | TGN38 | Merge
Ccdc186 KO

J

Pearson's correlation coefficient Pro-CPE – TGN38

ns

WT   Ccdc186 KO

Figure 4

A

B

Figure 5

Figure 6

Figure 7

Figure 8

A

B

Figure 9

WT

TGN

Plasma membrane

CPD

Proinsulin

Insulin

Immature DCV compartment

Mature DCV

Ccdc186 KO

Eipr1 KO