1 **Title:** Split-Belt walking induces changes in active, but not passive, perception of step
2 length
3

4 **Authors:** Dr. Carly Sombric[1], Marcela Gonzalez-Rubio[1], Dr. Gelsy Torres-Oviedo[1]
5 [1]Department of Bioengineering, University of Pittsburgh, Pittsburgh, PA
6
Corresponding Author:
Dr. Gelsy Torres-Oviedo
Department of Bioengineering
4420 Bayard Street
Suite 110, Pitt
Pittsburgh, PA 15213
Univerity of Pittsburgh
gelsyto@pitt.edu

Abstract

The estimation of limbs' position is critical for motor control. While motor adaptation

changes the estimation of limb position in volitional arm movements, this has not been

observed in locomotion. We hypothesized that split-belt walking with the legs moving at

different speeds changes the estimation of the legs' position when taking a step. Thus, we

assessed young subjects' perception of step length (i.e., inter-feet distance at foot

landing) when they moved their legs (active perception) or these were moved by the

experimenter (passive perception). Step length's active, but not passive perception was

altered by split-belt walking; indicating that adapted efferent inputs changed the

perceived limbs' position without changing information from sensory signals. These

perceptual shifts were sensitive to how they were tested: they were observed in the

trailing, but not the leading leg, and they were more salient when tested with short than

long step lengths. Our results suggest that sensory changes following motor adaptation

might arise from mismatched limb position estimates from different sensory sources (i.e.,

proprioception and vision), which is less prominent in walking. We also speculate that

split-belt walking could improve the deficient perception of step length post-stroke,

which contributes to their gait asymmetry impairing patients' mobility.

Introduction

Successful motor control requires accurate estimation of limb position, which is achieved

through a combination of afferent and efferent signals. Afferent information from sensory

signals is used to estimate limb position. In addition, efferent information also

contributes to the estimation of limb position through prediction of sensory consequences

prior to movement execution. The integration of actual and predicted sensory

information, from afferent and efferent signals (respectively), is beneficial because it

results in more accurate limb's estimation in space (e.g., [1–4]). It has been shown that

subjects' estimation of limb position is altered following motor adaptation to a persistent

stimulus (e.g., a constant force [5]). These adjustments to the estimation of limb position

can be due to either adaptation of afferent or efferent information. Shifts in sensory

information after motor adaptation have been observed with passive perception tasks

during which subjects use afferent information to localize their limb after it has been

passively moved (e.g., [5–12]). Moreover, predicted sensory consequences (from efferent

information) rely on internal models that are recalibrated during motor adaptation. Thus,

the integration of adapted sensory predictions with actual sensory information leads to

changes in limb estimation that are assessed through active perceptual tasks, which

require subjects to localize their limb after active movements [5,10,11,13–15]. In summary, the

critical ability to estimate the position of our limbs in space using afferent and efferent

information is altered by motor adaptation.

Interestingly, the effects of motor adaptation on the estimation of limb position in

walking remains elusive. Namely, a recent study probing the perception of one leg

position with respect to the body did not find passive or active perceptual after-effects

despite robust motor after-effects [16]. This might be due to how perceptual after-effects

were tested given evidence that passive and active perceptual changes post-adaptation are

sensitive to the condition in which they are evaluated [17–19]. For example, passive

perceptual after-effects depend on the limb that is tested [13], raising the possibility that

clearer perceptual changes could be observed when probing both limbs contributing to

each step length. Similarly, active perceptual after-effects might be regulated by test

condition such as walking speed; given that this factor alters the magnitude of motor

after-effects [20], and hence the recalibration of internal models underlying active

perceptual shifts. It is important to understand the influence of sensorimotor adaptation

on perception of limb position because stroke survivors have difficulty perceiving that

their step lengths are asymmetric [21]. Thus, it is of clinical interest to determine the extent

to which the estimation of limb position could be shifted following sensorimotor

adaptation in locomotion.

In summary, we investigated the effects of sensorimotor adaptation on the estimation

of limb position in walking. We hypothesized that sensorimotor adaptation alters the

estimation of step length, which requires concurrent localization of each leg's position.

We further hypothesized that walking condition, such as walking speed and/or step size,

would regulate the shifts in active perception of limb positions since walking speed alters

motor after-effects [20] and walking speed also alters step size (e.g., [22,23]). To test these

hypotheses, we separately investigated shifts in passive and active perception of step

length following sensorimotor adaptation in walking. Passive perception was assessed by

externally moving subjects' legs, whereas active perception was evaluated by instructing

subjects to take steps of different lengths. Importantly, the active perception task was

performed under distinct walking speeds and step sizes to test our hypothesis that active

perceptual effects depend on the condition in which they are tested. We assessed both

passive and active perception since it is possible that sensorimotor adaptation only

induces changes in the integration of afferent and efferent information, but not in afferent

information alone.

Results

Passive Perception of Step Length was Not Adapted Following Split-belt Walking

We aimed to determine the extent to which locomotor adaptation induced a change in the

estimation of limb position purely based on afferent information. To this end, adaptation

effects in passive step length perception (i.e., passive perceptual after-effects) and

movements (i.e., motor after-effects) were assessed in a group of unimpaired young

subjects (n=8, 2 females, 24.8±4.8 yrs.) following a split-belt adaptation paradigm known

to induce sensorimotor recalibration (e.g., [24]). We found motor, but not perceptual after-

effects. After-effects were defined as differences in step length (Figure 1a-b) or step

length asymmetry (Figure 1c) before and after the Adaptation epoch. These after-effects

were significantly different between the split and tied sessions in the motor domain

(Figure 1b: paired t-tests: slow leg's step length: $p<0.001$; fast leg's step length: $p<0.001$;

Figure 1c: step length asymmetry: $p<0.001$). In contrast, we did not observe perceptual

after-effects based on sensory information alone. Subjects' passive perception of their

step length was assessed with a task in which participants indicated their estimated step

length after the legs were moved by the experimenter to different step length sizes

(Figure 1a: short, comfortable, long). We did not observe significant changes in subjects'

perception of the slow leg's step length across testing sessions (Figure 1a: one-way

repeated measures ANOVA testing the effect of session; p=0.98). While we found a

significant session effect on the perception of fast leg's step length (Figure 1a: p=0.035),

this constituted a very small perceptual shift (0.8 ± 1.5 cm) compared to the significant

motor after-effects in the fast leg's step length (11.7 ± 5.9 cm) and the slow leg's step

length (-20.1 ± 6.7 cm). Thus, we observed a significant, but marginal adaptation effect

on the perception of the fast leg's step length. In sum, split-belt walking produced robust

motor after-effects, but did not change subjects' estimation of step lengths based on

sensory (afferent) information alone.

Figure 1 Split-belt walking did not induce changes in passive perception of
step lengths | (a-b) In all plots, step length is defined as the distance between the
ankles at heel strike (i.e., when the foot hits the ground). Change in step length was
computed as the difference in step length before and after the Adaptation epoch,

which during the control session was an extended period of tied walking (light grey)
and during the split session was an extended period of split walking (dark grey).
These changes were expressed as a percentage of each subject's mean step length
during baseline walking at 1 m/s to account for different step sizes across
individuals. Negative and positive values indicate that step lengths were either
perceived shorter or longer, respectively, after the Adaptation epoch compared to
before. **(a) Passive Perception After-Effects**: Bars' height indicates group mean
changes in perception after the Adaptation epoch relative to baseline behavior. Dots
indicate values for each individual and error bars indicate standard errors. Passive
perception was quantified by probing subjects' perception of three step length sizes
(short, comfortable, and long) imposed by passively moving subjects' legs. Each
participant's weight was equally distributed between the legs when their step
length perception was probed. **(b) Motor After-Effects of individual legs:** Bars'
height indicates group mean changes in step length after either the tied session or
the split session. Standard errors and individual values (dots) are also displayed. **(c)**
Motor After-Effects of step length asymmetry: Step length asymmetry is defined
as the differences between step lengths of two consecutive steps normalized by the
sum of these step lengths. Positive values indicate that steps taken with the
dominant leg are longer compared to those taken with the non-dominant leg; and
vice versa for negative values. Bar height indicates group mean changes in step
length symmetry after the Adaptation epoch. Standard errors and individual values
are also displayed.

Active Perception of Step Length was Modulated in Both Legs After Split-Belt Walking

Unlike passive perception, subjects' active perception of step length based on efferent

information changed after split-belt walking compared to after tied walking. We

evaluated subjects' (n=27, 16 females, 25.1±5.4 yrs.) active perception of step length by

prompting step lengths of two distinct magnitudes (i.e., long or short) on a VR headset

(see Methods section). We specifically assessed whether participants overshot or

undershot these target step lengths after an Adaptation epoch, which consisted of an

extended period of tied walking during the tied session and split-belt walking during the

split session. Active perception after-effects were quantified as differences in executed

step lengths before and after the Adaptation epoch (i.e., %Change in Step Length).

Perceptual after-effects were subsequently compared between the two testing sessions to

determine if split-belt adaptation induced changes in the active perception of step length.

Importantly, perceptual and motor after-effects were tested under distinct walking speeds

(i.e., mid and slow) to determine if walking condition had a similar effect on active

perceptual after-effects, as it does on motor after-effects [20]. Thus, we had a total of three

groups labeled by the step length size that was prompted first in the perceptual task and

the speed at which it was tested (Long-MidSpeed, Short-MidSpeed, Short-SlowSpeed).

Figure 2a shows that following split-belt walking, all groups undershot (negative values)

or overshot (positive values) target step lengths with the leg that walked slow (slow leg)

or fast (fast leg), respectively, during the Adaptation epoch. This indicated that subjects

perceived their step lengths to be either longer or shorter than actual when stepping with

the slow or fast leg, respectively. Consistently, we found a significant effect of session on

the active perception of the slow ($p<0.001$) and fast step lengths ($p<0.001$) with one-way

repeated measures ANOVAs. These shifts in perception of individual leg's step length

led to significant differences in step length asymmetries during the active perceptual task

across sessions (Figure 2c: session effect: $p<0.001$).

Active Perception of Step Length was Modulated by Step Size, but Not Walking Speed

Subsequent analysis on the split session indicated that step length size, but not walking

speed, had an effect on active perception after-effects. Specifically, a two-way repeated

measures ANOVA indicated a significant interaction between leg and step length size

(leg#step length size: p =0.004) for the groups starting the perceptual task with different

step lengths sizes, but the same walking speed (i.e., Short-MidSpeed vs. Long-

MidSpeed). This interaction was driven by greater after-effects in the perception of the

fast leg's step length when taking short (Figure 2a, red bar) compared to long (Figure 2a,

blue bar) step lengths (two sample t-test: p=0.024). This might be due to greater

similarities between training and testing in the short vs. long conditions. Namely, the

executed step lengths during the training condition (i.e., split-belt walking, Late

Adaptation in Figure 2a insert, white background) were qualitatively more similar to the

target step lengths in the Short-MidSpeeds group (Figure 2a insert, red background) than

in the Long-MidSpeeds group (Figure 2a insert, blue background). Conversely, we did

not find an interaction between leg and walking speed (leg#speed: p =0.16) for the groups

that walked at different speeds, but same step lengths in the active perceptual task (i.e.,

Short-MidSpeed vs. Short-SlowSpeed). This contrasted the effect of walking speed on

motor after-effects (Figure 2b). Specifically, the slow step length motor after-effects were

larger when tested at slow than mid walking speeds (Figure 2b: two sample t-test between

pooled mid speed groups vs. a slow speed group; p=0.008), whereas they were the same

for fast step lengths (p=0.83). Thus, the effect of speed on step length asymmetry (Figure

2b: p=0.001) was due to the slow step length after-effects. In summary, split-belt walking

induces motor and active perceptual after-effects that are regulated by the way they are

tested: motor after-effects are altered by walking speed, whereas perceptual after-effects

are altered by step length size.

Active Perception of Step Length Decayed Without Fully Washing Out Motor After-

Effects

We observed that the active perceptual after-effects decayed as they were tested, but their

washout did not fully eliminate motor after-effects. Namely, we found that perceptual

after-effects decayed exponentially as people walked in the active perceptual task (Figure

3a: $y = c + ae^{(-x/\tau)}$). This exponential decay was clearly observed in the active

perception of the slow step length in all groups (Figure 3a, top panel: τ [95% CI]; Short-

MidSpeed: 24.8 [31.0, 20.8], Long-MidSpeed: 17.7 [14.1, 23.6], Short-SlowSpeed: 27.1

[19.4, 45.0]), but only in the fast step length for groups tested with the short target first

(Figure 3a, bottom panel: τ [95% CI]; Short-MidSpeed: 25.7 [19.4, 37.8], Long-

MidSpeed: 2.4 [-0.86, 3.0], Short-SlowSpeed: 53.2 [37.2, 93.1]). These findings further

support that subjects did not have perceptual shifts of the fast leg's step length when

probed with long steps. Interestingly, the active perceptual task did not fully washout

motor after-effects. This was tested by comparing perceptual after-effects at the end of

the active perceptual task (Figure 3b: Late Perception) and the remaining motor after-

effects tested immediately afterwards (Figure 3b: Early Post-Adaptation). Of note, the

belts moved at the same tied speed in both instances. The only experimental difference is

that in Late Perception subjects aimed at their comfortable step length (recorded during

baseline walking) while wearing the VR headset, whereas in Early Post-Adaptation they

walked unconstrained (i.e., without any specific instruction) and without VR headset.

Thus, washout of motor after-effects by the active perceptual task is expected, as

indicated by the qualitatively smaller motor after-effects during Early Post-Adaptation

**Figure 2 Split-belt walking induces changes in active perception | (a) Step
Length Active Perception After-Effects**: Bars' height indicates group mean
changes in step length before and after the Adaptation epoch, which consisted of
tied (empty bars) or split walking (filled bars). These changes were expressed as a
percentage of each subject's mean step length during the baseline active perceptual
task tested at the specific step length size and walking speed for each group. For
example, we used the mean step lengths during the baseline perceptual task aiming
at short step lengths and walking at slow speed for the Short-SlowSpeed group.
Negative or positive values indicate that subjects either undershot or overshot the
target step lengths, respectively because they perceived step lengths to be longer or
shorter than the actual value. The insert illustrates the %Change in Step Length that
was executed by the slow (red dot) and fast (blue dot) step lengths at steady state
during split-belt walking (Late Adaptation; white background). We also show the
target step lengths (red and blue dots) for probing active perception during baseline
in the Short-MidSpeed group (red background) and the Long-MidSpeed group (blue

background). Standard errors are displayed in all plots. **(b) Step Length Motor**
After-Effects: Bars' height indicates group mean changes in step length relative to
baseline walking during the Catch trial tested at either 1.0m/s (Short-MidSpeed and
Long-MidSpeed) or 0.5 m/s (Short-SlowSpeed). The mid speed groups were pooled
for this analysis since these two groups experienced exactly the same protocol prior
to the catch trial. **(c) Step Length Asymmetry Perceptual and Motor After-**
Effects: Bars' height indicates group mean changes in step length asymmetry
relative to baseline walking during the Perceptual trials and during the Catch trials.
The left panel illustrates the mean step length asymmetry that results from the
change in individual step lengths during the active perceptual task for each group.
The right panel illustrates the mean step length asymmetry that results from
changes during the Catch trial tested at the Mid (1.0 m/s) or Slow (0.5 m/s) walking
speed. Standard errors are displayed in all plots.
253

compared to those during the Catch trial (colored dotted lines in Figure 3b). Two-way

repeated measures ANOVAs (factors: epoch and speed) were used for each step length to

show that motor after-effects are present after perceptual after-effects are washed out and

that the motor after-effects are influenced by speed. Specifically, the slow and fast leg

had significant resurgence of the motor after-effects following perceptual testing (Figure

3b: slow step length: $p_{epoch}<0.001$; fast step length: $p_{epoch}<0.001$) and the magnitudes of

after-effects were regulated by walking speed (Figure 3b: slow step length: $p_{speed}=0.04$,

$p_{speed\#epoch}=0.018$; fast step length: $p_{speed}=0.041$, $p_{speed\#epoch}=0.43$). In summary, active

perception after-effects decayed as subjects walked, but this did not fully washout motor

after-effects.

Figure 3 Active perception after-effects, but not motor after-effects, are fully
washed out by the end of the perceptual task | (a) Decay of active perception
after-effects. We show the %Change in Step Length (i.e., after-effects) for the slow
(Top panel) and fast step length (Bottom Panel) over the course of the active
perceptual task. Note that all groups exhibit an exponential decay of active
perception after-effects in both legs except for the fast leg in the Long-MidSpeed
group whose τ was not significantly different from zero (lack of significance
indicated with a dotted line). This further indicates that subjects did not have
significant fast step length perceptual shifts when probed with long step lengths.
Perceptual after-effects assessed at target step lengths that were different from the
initial probe were not included in the analysis (white spaces). **(b) Remaining**
motor after-effects following the active perceptual task. Colored lines indicate
the motor after-effects recorded during the Catch trial (prior to the active
perceptual task, Figure 2b). Bars indicate group means for the slow and fast
%Change Step Length at the end of the perceptual task (Late Perception) and at the
beginning of the Post-Adaptation period (Early Post-Adaptation). Black lines
indicate Fisher's LSD post Hoc results. Subjects walked at the exact same tied-belt
speeds during the Late Perception and Early Post-Adaptation periods. The only
difference in these time points was that subjects were prompted to take their
comfortable step length during Late Perception, but during Early Post-Adaptation
subjects walked unconstrained (i.e., without any specific instruction of where to

step). Standard errors are displayed in all plots. Note that the mid speed groups are
pooled for this analysis since they experienced the same Adaptation protocol and
target step lengths during Late Perception.
289

Active Perception After-Effects are Mediated by Changes in Trailing Leg Perception

Step lengths during active perceptual trials were further decomposed into perceived

leading (α) and trailing (X) leg positions (Figure 4a) to determine which leg's position

perception underlies changes in step length perception following split-belt adaptation.

Our results show that only the trailing leg's position (X) exhibited active perception after-

effects (Figure 4b). These were quantified with symmetry measures of each leg's X, such

that values different from zero indicated active perception after-effects. We separately

characterized α and X perceptual after-effects for short and long target step lengths with

single exponential fits ($y = c + ae^{(-x/\tau)}$). A separate fit was performed for individuals

walking slow vs. those walking at mid speeds. Fits of the data indicate that only the

trailing leg position experiences large changes in perception that decayed as subjects

experienced the active perceptual task. This is indicated by the decay of perceptual after-

effects probed at either the short step length target (Figure 4b: τ [95% CI]; Short Target:

MidSpeed= 12.5 [10.8, 14.9], SlowSpeed=36.9 [32.6, 42.7]) or long step length target

(MidSpeed=37.5 [33.4, 42.8], SlowSpeed=41.0 [33.5, 52.9]). In contrast, the leading

leg's position exhibited perceptual after-effects that were very short lived, regardless of

whether they were probed with the short step length target (Figure 4c: τ [95% CI]; Short

Target: MidSpeed= 8.4 [6.4, 12.4], SlowSpeed=1.2 [-0.01, 2.4]) or the long step length

target (MidSpeed=7.2 [4.7, 15.2], SlowSpeed=7.3 [3.8, 90.0]). Taken together, these

results indicate that active perception shifts in step lengths are due to changes in the

perception of the trailing leg's position, and not the leading leg's position.

313
Figure 4 Split-belt Walking Induces Active Perception Changes in the Trailing Leg
Position | (a) Schematic of the decomposition of step length into the leading (α) and
trailing (X) leg relative to the hips. Both α and X are normalized by the speed specific
baseline walking behavior, such that positive values mean the distance from the body is
greater following Adaptation and vice versa. Symmetry measures were computed for
both X **(b)** and α **(c)** and were defined as the differences between these measures on two
consecutive steps (slow leg - fast leg). Thus, positive values of %Change X Symmetry
here indicates that the slow leg (X_{slow}) is trailing farther behind than fast leg (X_{fast})
relative to baseline behavior. Similarly, positive values of %Change α Symmetry would

indicate that the slow leg (α_{slow}) is farther in front of the body than the fast leg (α_{fast})
relative to baseline behavior. Note that the X position perception is altered following
split-belt walking, but that the α position perception is not altered, as indicated by
negligible %Changes in α Symmetry following split-belt walking.
327

Discussion

Summary

We investigated the effects of locomotor adaptation on passive and active step

length perception. We observed active, but not passive, perceptual after-effects of step

length. Thus, locomotor adaptation did not induce sensory shifts, but altered the

estimation of limb position based on integration of afferent and updated efferent

information. In addition, we found that active perception after-effects were only observed

on the trailing limb's position, indicating that the perceived position of the trailing, but

not the leading leg, with respect to the body is recalibrated after locomotor adaptation.

Moreover, perceptual and motor after-effects were regulated by how they were tested.

Specifically, perceptual after-effects were altered by step length size, whereas motor

after-effects were altered by walking speed. Lastly, the observed perceptual after-effects

decayed as subjects walked in the active perceptual task, but motor after-effects

remained, raising the possibility that after-effects in these two domains are originated by

partially distinct processes.

Motor recalibration in locomotion induces changes in active, but not passive perception

of limb position.

Our results show that the perception of limb position is altered following motor

adaptation in locomotion, which supports prior findings that the perception of limb

position is susceptible to motor adaptation (e.g., [5,6,15,7–14]). However, we only observed

perceptual after-effects when subjects actively generated motor commands and not when

their legs were passively moved, which is consistent with prior work indicating that

afferent signals encoding position remain intact following split-belt walking [16]. These

findings contrast multiple reports of shifts in the estimation of hand position following

motor adaptation when the arms are moved by the experimenter (e.g., [5–12]). We believe

that this discrepancy between reaching and locomotion suggests that sensory changes

post-adaptation arise from mismatched position estimates from different sensory sources

(i.e., proprioception and vision), which is less prominent in walking. We also found that

active perceptual effects were not observed in the leading leg's position, as previously

reported [16], but they were clearly observed in the trailing leg's position, which had not

been assessed before. We posit two potential explanations for the altered estimation of

the trailing leg's position. First, subjects might maintain the expectation that one leg will

move faster than the other post-adaptation. As a result, they will stand longer on the leg

that walked slow compared to the other as they take steps in the active perceptual task,

leading to differences in trailing leg position that are consistent to the ones we see (i.e.,

$X_{slow} > X_{fast}$). Alternatively, the shifts in position might be due to changes in how subjects

perceived the environment. It is known that the perception of symmetric walking speeds

is shifted following split-belt walking [16,25]. More specifically, individuals during post-

adaptation perceive the fast leg to move slower and the slow leg to move faster than their

actual speed. Subjects might act according to this perception by standing longer on their

trailing leg when taking steps with the fast leg and vice versa for steps with the slow leg,

resulting in the observed larger X_{slow} than X_{fast} values. In sum, subjects' estimation of

limb position is altered following sensorimotor adaptation. This shift is not due to

changes to afferent information, but might be due to updated motor commands or altered

perception of the environment following adaptation.

Motor and perceptual after-effects are susceptible to how they are tested.

It has been shown that split-belt motor after-effects are sensitive to walking speed [20] and

perceptual after-effects are sensitive to movement distance [18]. Thus, our observations that

motor after-effects are regulated by walking speed and perceptual after-effects are

regulated by step length are consistent with previous literature. However, it should be

noted that walking speed and step length are not independent (e.g., [22,23]) raising the

possibility that step length is the factor that regulates both motor and perceptual after-

effects. Should that be the case, we speculate that similarity in step lengths sizes during

training and testing is a critical feature to maximize after-effects, given the well-known

effect of similarity on the generalization of learned movements across conditions [26–28]. In

sum, our results highlight the importance of considering how both motor and perceptual

after-effects are tested.

Why did motor after-effects remain after perceptual ones were fully washed out?

It is interesting that walking in the active perceptual task did not fully eliminate motor

after-effects. We believe this could be due to three possible reasons. First, the perceptual

and motor effects were tested under different peripheral vision because subjects were

wearing a virtual reality headset in the active perceptual task. This distinct situation

constitutes a contextual difference known to reduce the washout of the pattern specific to

the split-belt treadmill by walking in a symmetric environment [29]. Second, the

constrained nature of walking in the perceptual task might also influence the remaining

motor after-effects. Namely, perceptual after-effects were tested by instructing subjects

"where" to step, whereas motor after-effects were tested in unconstrained walking.

Constrained and unconstrained movements are thought to be differently controlled (e.g.

[30]). Thus, the fact that individuals were aiming to a target in the perceptual task might

have altered the extent to which implicit actions were washed out. Lastly, the limited

washout of motor after-effects could also indicate that motor and perceptual after-effects

are originated by partially distinct processes. This is supported by previous research

showing that motor and perceptual effects exhibit different patterns of saving [25] and

retention [11], distinct adaptation rates [6,15] and uncorrelated amounts of adaptation [8,9,12,13].

They also exhibit different patterns of generalization to other limbs [17] or novel targets

[18,19]. Thus, while the limited washout of motor after-effects by the perceptual task might

be due to testing differences, it could also indicate that the processes underlying motor

and perceptual after-effects might be partially independent.

Clinical implications

The split-belt walking may be a viable tool to address perceptual deficits associated with

stroke. Notably, stroke survivors' inability to perceive that their gait is asymmetric is

associated with limited recovery of symmetric gait [21]. Our results indicate that active

perception of step lengths can change following split-belt walking, suggesting that a

similar effect might be observed post-stroke. It has been shown that split-belt training can

correct motor deficits [31,32] and it would be interesting to determine if it also corrects

perceptual deficits as well. Of note, recent work indicates that perceptual changes in the

estimation of speed are reduced with repeated exposure to the split condition [25],

suggesting that similar reduction would be observed in the shifts of limb position. Thus,

there might be a limited long term effect of split-belt walking on perceptual changes.

Therefore, future work is needed to understand if split-belt walking has similar effects in

the lesioned motor system and whether training protocols could lead to long lasting

improvements in motor, as well as, perceptual deficits post-stroke.

Methods

The influence of split-belt walking on passive (Experiment 1: n=8, 2 females, 24.8±4.8

yrs.) and active (Experiment 2: n=27, 16 females, 25.1±5.4 yrs.) step length perception

was characterized in separate groups of subjects. This study was approved by the

University of Pittsburgh Institutional Review Board, was carried out in accordance with

the approved guidelines, and all subjects provided written informed consent.

Experiment 1: Passive Perception

We tested subjects' perception of their limb position by moving their legs in order to

determine if afferent information was altered by split-belt walking. This was done before

and after a tied and a split session, which occurred on different days. Both the tied and

split sessions included perceptual and locomotor epochs (Figure 5a).

Locomotor Task

Subjects experienced a tied and split session that differed only in the locomotor

Adaptation and Re-Adaptation epochs (Figure 5a). The first session was used to

determine if an extended period of tied belt walking (i.e., walking with both belts moving

1.0 m/s) during the Adaptation epoch changed subjects' passive perception of step length.

During the second session, subjects walked with the legs moving at different speeds

during the Adaptation epoch to determine if locomotor adaptation altered subjects'

passive perception of step length. The split-belt perturbation was gradually introduced

during the Adaptation epoch for consistency with prior work studying passive perception

(e.g., [6,8,9,12,33]). More specifically, belts started at 1.0 m/s and linearly diverged for 600

strides until a 3:1 split-belt ratio (i.e., 1.5 m/s: 0.5 m/s) was achieved and maintained for

200 strides resulting in a total of 800 strides for the Adaptation epoch. Subjects' passive

perception of step length was assessed immediately before and after the Adaptation

epoch. We re-exposed individuals to either the tied condition (i.e., both legs at 1 m/s) or

split condition (i.e., 1.5 m/s: 0.5 m/s) during a Re-Adaptation epoch to subsequently

evaluate motor after-effects in each subject. We chose to evaluate motor after-effects

following the perceptual task to avoid changes in perception due to multiple transitions

between tied and split environments, which are known to alter perceptual effects [25]. The

Re-Adaptation epoch lasted 300 strides to ensure individuals reached a similar steady

state as in the Adaptation epoch prior to assessing motor after-effects. Then, we evaluated

motor after-effects during a Post-Adaptation epoch (450 strides) when the two legs

moved at the same speed (1 m/s). Subjects walked in this tied environment for 450 strides

to fully wash out any split-belt after-effects before testing subjects' retention of the

mapping between the imposed step lengths and the analog scale used in the perceptual

task (see section below).

Passive Perceptual Tasks

During the passive perception task, subjects wore a virtual reality headset (Oculus,

Facebook Technologies, Irvine, CA, United States) while standing with their weight

evenly distributed between the two feet and with each foot placed on one belt of the split-

belt treadmill. The sequence of a passive perceptual trial is shown in Figure 5b. First, legs

were passively moved to a specific step length value (ankle to ankle distance) (Figure 5b,

first panel). Then, belts were stopped and subjects were asked to distribute their weight

equally between their feet by maintaining their center of pressure (green dot) within a

target region (blue rectangle) (Figure 5b, second panel). Subjects maintained this

balanced position for 5 seconds during which they had to indicate their perceived step

length using a clicker that moved a cursor (red dot) along an analog scale displayed in the

virtual reality headset (Figure 5b, third panel). Then, a new perceptual trial was initiated

by moving the legs to a different step length value (Figure 5b, fifth panel). Subjects

received feedback of the actual step length in the analog scale (dark green dot) only

during perceptual training (see full description below) (Figure 5b, fourth panel). Subjects

did not lift their feet off the belts when the belts were moving or still. We alternated the

leg that was placed forward after every perceptual probe/trial.

Perceptual Training

Subjects trained extensively on the passive step length task to learn the mapping between

the imposed step lengths and the analog scale. Training occurred at the beginning of

every session. Three out of the 8 participants had an additional session with only training

trials prior to the tied session to determine if improved accuracy at the task would allow

us to identify small perceptual shifts. The analog display observed by all subjects is

shown in Figure 5b The middle of the analog scale corresponded to a medium size step

length computed as 75% of each subject's averaged step length during baseline walking

at 1.0 m/s. Subjects additionally trained on targets 8 cm shorter (short step length) and

longer (long step length) than the medium size step length. Six out of the eight

participants on this task also trained with step lengths that were 4 cm shorter and longer

than the medium size step length. During training, subjects experienced the same

perceptual probe sequence as during testing: 1) legs were passively moved to one of the

target step length values (e.g., medium, short, and long step length sizes), 2) subjects

indicated their perceived step length while maintaining an equal distribution of their

weight between their feet, and 3) subjects received feedback on their actual step length

along the same analog scale. Subjects experienced 10 or 20 iterations of each of the target

step lengths in a random order. Our results did not change when either more step length

sizes were assessed (e.g., 3, rather than 5, target step lengths) or more instances of each

step length size were tested (e.g., 21, rather than 10).

Perceptual Trials

We evaluated subjects' capacity to classify step length sizes with the analog scale

immediately following training (i.e., Baseline Perceptual epoch) and right after the

Adaptation epoch during each session (i.e., Post-Adaptation epoch). Each epoch

consisted of several iterations (10 or 21) of the target step sizes presented in a random

order. Subjects performance was computed as the mean value that they selected in the

analog scale across all perceptual trials at each target size. In all trials, subjects were told

that they would experience step lengths all along the analog scale, including values that

were shorter or longer than those they experienced during training. In reality, subjects

were only exposed to the step length values that they trained on. Feedback was not

provided in any trial. The performance of the perceptual trials for each leg were used to

determine if subjects could differentiate the distinct step length sizes with the analog

scale (i.e., Baseline Perceptual epoch performance), and if passive perception of step

length was altered by the Adaptation epoch (Baseline Perceptual vs. Post-Adaptation

epoch performance). First, the Baseline Perceptual performance for each leg was

compared across all target sizes and sessions to determine if subjects could differentiate

the distinct step length sizes with the analog scale for each session. Figure 5c indicates

that subjects learned to discern the three step length values they experienced when their

legs were passively moved (Figure 5c: two-way repeated measures ANOVA testing the

effect of target and session on perceived step length; Slow Leg: $p_{target}<0.001$, $p_{session}=0.59$,

$p_{target\#session}=0.56$; Fast Leg: $p_{target}<0.001$, $p_{session}=0.32$, $p_{target\#session}=0.71$). Next, it was

determined if passive perception of step length size was altered by split belt walking by

comparing the Baseline Perceptual performance and Post-Adaptation performance (see

results). Lastly, we probed subjects' passive perception at the very end of the split

session in a Post-Washout epoch, during which subjects' perception of the short and long

step length were probed 7 or 10 times each. This was done to ensure that changes in the

reported step length with the analog scale were due to perceptual changes, rather than

forgetting of the mapping between the sensed step length and the analog scale. Thus,

subjects' perception of the short and long step lengths were compared between the

Baseline Perceptual and Post-Washout epochs. We found no differences between epochs

or target step lengths (Figure 5c: two-way repeated measures ANOVA testing the effect

of epoch and target on perceived step lengths: Slow Leg: $p_{epoch}=0.80$, $p_{target}<0.001$,

$p_{epoch\#target}=0.13$; Fast Leg: $p_{epoch}=0.88$, $p_{target}<0.001$, $p_{epoch\#target}=0.83$). This indicated that

individuals could classify well the distinct step length values throughout the duration of

the experiment.

Figure 5 Locomotor and Passive Perception Protocol | (a) Locomotor epochs (colored blocks) and Passive Perceptual epochs (grey scale blocks) were included in both the control and testing session. Note that subjects were standing during the passive perceptual trials and walking during the locomotor trials. **(b) Passive Step Length Tasks:** Subjects' legs were passively moved to a target step length. Subjects maintain an equal distribution of their weight between their feet using visual feedback displayed in the virtual reality headset. Subjects indicated their perceived step length size using an analog scale that was also displayed. Only during training, participants received feedback on their perceived vs. actual step length along the analog scale. This was done such that individuals could learn the mapping between what they felt and what was displayed in the analog scale. **(c) Baseline and Post-Washout Perceptual Trials:** Subjects' performance during the Baseline Perceptual Trials for both sessions and the Post-Washout Trials for the split session are shown for each leg. Note that subjects could discern three distinct step lengths from the beginning (Baseline) until the end (Post-Washout) of the experiment. Thus, any performance changes between the Baseline vs.

569 Post-Adaptation epochs were attributed to shifts in subjects' passive perception of step
570 length, rather than subjects' inability to maintain the mapping between the imposed step
571 lengths and the analog scale.
572

Experiment 2: Active Perception

Subjects' estimation of step length upon active motion of their legs was evaluated to

determine if this estimation was altered by split-belt walking. This was done before and

after a tied and a split session, which occurred on different days. Both the tied and split

sessions included perceptual and locomotor epochs (Figure 6a). Importantly, perceptual

and motor after-effects were tested under distinct walking speeds (i.e., mid and slow)

since this factor has been shown to modulate the generalization of sensorimotor

recalibration across walking conditions [20]. Moreover, speed naturally regulates the

magnitude of step lengths [22,23]. Therefore, we evaluated the possibility that active

perception of step length would be distinct for different step length sizes (i.e., long and

short). Of note, both step length sizes were tested in all groups, but we altered the order at

which the perception of step lengths were probed (i.e., long first vs. short first) given our

expectation that active perceptual after-effects would decay as subjects performed the

active perceptual task. In sum, motor and active perceptual after-effects were tested at

either a slow (0.5 m/s) or mid (1 m/s) walking speeds, and active perceptual after-effects

were first measured at either short or long step lengths. These experimental conditions

resulted in three groups designated first by testing speed and second by the first step

length size at which active perception was probed (Long-MidSpeed, Short-MidSpeed,

Short-SlowSpeed).

Locomotor Task

Similar to Experiment 1, subjects experienced a tied and split session that differed

only in the locomotor Adaptation and Re-Adaptation epochs (Figure 6a). The first session

ensured that tied walking (i.e., walking with both belts moving 1.0 m/s) during the

Adaptation and Re-Adaptation epochs did not induce active changes in step length

perception. During the second session, subjects walked for 600 strides at a 3:1 split-belt

perturbation (1.5 m/s and 0.5 m/s) during the Adaptation epoch. Unlike Experiment 1, we

introduce the split-belt perturbation abruptly because it has been shown that active

perception adapts more slowly than motor recalibration [15]. Thus, we wanted to maximize

the exposure of the full split condition before assessing perceptual after-effects. Motor

after-effects were characterized in a short tied Catch trial at the group-specific walking

speed. During the split session's Re-Adaptation epoch, subjects again experienced 3:1

split-belt walking for 300 strides in order to return subjects to steady state split-belt

walking before testing active perception of step length in the Post-Adaptation epoch.

Following the active perceptual assessment during the Post-Adaptation epoch, subjects

walked for 150 strides at the group-specific walking speed in order to evaluate any

remaining motor after-effects. Once these motor after-effects were fully washed out, we

performed one last active perceptual task without feedback to assess subjects' retention of

the map between the target and executed step lengths.

General Approach for all Active Perception Trials

Subjects wore a virtual reality headset (Oculus, Facebook Technologies, Irvine, CA,

United States) to eliminate the possibility of using visual information to bias subjects'

estimate of step length based on their body motion with respect to the treadmill. Subjects

received limited feedback on their step length on a numbered grid that was displayed in

the virtual reality headset (Figure 6). Each grid line corresponded to 2 cm in real space.

The cursors on the screen indicated the step length when taking a step with the right

(orange cursor) or left leg (green cursor). The cursor position for each leg was only

updated at heel-strike. The target step length was indicated with the blue highlighted line

on the grid. The comfortable step length (grid #10) corresponded to the average step

length during the speed-specific baseline walking. Subjects were also prompted to take a

short (grid #7, 6 cm shorter than comfortable step length) and long (grid #13, 6 cm longer

than comfortable step length) target step lengths. Importantly, the short and long step

lengths used in the perceptual task were different from those regularly taken during

baseline walking or during the catch trail. This was done to test active perceptual effects

with step lengths that were distinct from those that subjects would normally take when

assessing motor after-effects. Subjects performed the active perceptual task as they

walked while lightly touching (<2N, enforced with verbal feedback) an instrumented

handrail located in front of the treadmill. This was done to ensure that individuals would

maintain their position on the treadmill while walking with the virtual reality headset [34].

Recall that the Short-MidSpeed and Long-MidSpeed groups performed the step length

tasks at a mid-walking speed (1.0 m/s), whereas the Short-SlowSpeed group performed

the step length tasks at a slow walking speed (0.5 m/s).

Training for active perceptual testing

A step length Training epoch was first conducted in order to train subjects to take three

distinct step lengths. During training, subjects received feedback on 100% of their step

length error, where step length error is defined as the distance between the target step

length and the executed step length. Subjects experienced targets in 4 sets where each set

included each of the three targets in a random order. Each presentation of the target

included 50 strides with visual feedback followed by 10 strides with no visual feedback.

This was done to habituate the subjects to performing the perceptual task without

feedback and favor the learning and retention process (e.g., [35]). In total, subjects trained

for 720 strides.

Map Test

Map Tests were used to ensure that subjects had successfully learned and maintained a

mapping between the target step lengths prompted by each number on the grid. During

this map test, subjects performed two sets of 25 strides for each target (total of 150

strides) without any feedback. This was done right before the Adaptation epoch and at the

end of each session following the extinction of any locomotor after-effects during the

Post-Adaptation epoch. The perceptual training protocol was effective as indicated by the

fact that subjects took distinct step lengths for each target step length value (Figure 6c:

two-way repeated measures ANOVA testing the effect of target step length and session

on step length for each leg; Slow Leg: $p_{Target}<0.001$, $p_{Session}=0.20$, $p_{Target\#Session}=1.00$; Fast

Leg: $p_{Target}<0.001$, $p_{Session}=0.35$, $p_{Target\#Session}=0.80$) that were retained for the duration of

each session except for the fast leg during the tied session (two-way repeated measures

ANOVA with effect of target step length and epoch; Tied Session, slow step length:

$p_{Target}=0.82$, $p_{Epoch}=0.79$, $p_{Epoch\#Target}=0.68$; Tied Session, fast step length: $p_{Target}=0.92$,

$p_{Epoch}=0.009$, $p_{Epoch\#Target}=0.37$; Split Session, slow step length: $p_{Target}=0.89$, $p_{Epoch}=0.069$,

$p_{Epoch\#Target}=0.80$; Split Session, fast step length: $p_{Target}=0.76$, $p_{Epoch}=0.13$, $p_{Epoch\#Target}$

$=0.72$). While the mapping between prompted and taken step lengths was not well

maintained for the fast leg during the tied session, qualitatively, subjects still maintained

distinct step lengths. Therefore, any changes observed between the Baseline and Post-

Adaptation Perceptual trials are due to changes in perception, rather than subjects

forgetting of the step length value that they were required to take when prompted by each

number on the grid.

681
Figure 6 Locomotor and Active Perception Protocol | (a) Locomotor (colored blocks)
and Active Perceptual (grey scale blocks) epochs were included in both the control and
testing session. Note that subjects were walking during the active perceptual trials. **(b)**
Active Step Length Task: Subjects walked on the treadmill while wearing a virtual
reality headset. Note that subject's walking speed depended on the group. Subjects saw
a grid where each grid corresponded to 2 cm in real space. The blue highlighted grid line
indicated a target step length that subjects should take. Grid 7, 10, and 13 corresponded to

short, medium, and long step lengths, respectively. During training trials, subjects
received accurate endpoint feedback on each step length. During Map Tests, subjects did
not receive any feedback on their performance. During perceptual trials, subjects
received reduced error feedback (i.e., subjects only saw 35% of the error that they made).
(c) Map Test: Subjects performance during the Map Test indicated how well subjects
learned and retained the spatial mapping between step lengths prompted with
biofeedback and taken step lengths. Note that subjects learned distinct step lengths that
were well maintained from the beginning until the end of the experiment. Thus, any
changes in the active perception of step length are due to perceptual changes rather than
subject's inability to learn or maintain a mapping between prompted and taken step
lengths.
700
Active Perception Trials

To determine if active perception was altered, subjects needed to be able to have large

step length errors without receiving feedback that would illicit strategic movement

corrections that could override perceptual shifts. However, pilot data indicated that

subjects needed some feedback to stay on task for the duration of the perceptual trials.

Therefore, 35% of subjects step length error was projected during the active perceptual

trials directly before and after the Adaptation epoch without the subjects' knowledge in

order to observe active perceptual shifts and their natural decay (e.g., if subjects had a 10

cm step length error, the feedback only indicated that there was a 3.5 cm step length

error). Recall that the Short-MidSpeed and Short-SlowSpeed group started with short

targets whereas the Long-MidSpeed group started with long targets. Subjects alternated

between the short and long target step lengths in a predefined manner in order to

characterize the decay of any temporally persistent perceptual shifts (455 strides total: 10

strides first target, 15 strides of the second target, one set of the short and long target for

15 strides, one set of the short and long target for 25 strides, three sets of the short and

long target for 50 strides, 50 strides of the comfortable step length). Note that because all

groups took the same number of strides, that it took the group walking slow (0.5 m/s)

twice as long to complete the perceptual trials as the groups walking at the mid speed (1

m/s).

Data collection

Kinematic data were collected to characterize subjects' locomotor behaviors while

walking on the treadmill during locomotor and perceptual trials. A motion analysis

system (Vicon Motion Systems, Oxford, UK) was used to collect kinematic data at 100

Hz. A quintic spline interpolation was used to fill gaps in the raw kinematic data

(Woltring; Vicon Nexus Software, Oxford, UK). Subjects' movements were tracked via

passive reflective markers placed bilaterally over the hip (greater trochanter) and ankle

(lateral malleoulous) and asymmetrically on the thigh and shank to distinguish the legs.

The duration of treadmill trials was defined by real time kinetic detection of heel strikes.

Heel strikes were identified with raw vertical kinetic data collected from the instrumented

treadmill (Bertec, Columbus, OH, United States).

Perceptual Task

Custom written Vizard (WoldViz, Santa Barbara, CA, United States) and MATLAB (The

MathWorks, Inc., Natick, MA, United States) code was used to control the treadmill in

order to impose desired step lengths during the passive perception tasks and to provide

appropriate feedback during all perceptual tasks.

Statistical Analysis

The statistical analyses were performed as reported in the text. A significance level of α = 0.05 was used for all analysis. Stata was used to perform all statistical analysis (StataCorp LP, College Station, TX).

Competing Interests

The authors declare that there are no competing interests.

Author Contributions

G.T.-O. and C.S. were involved with the conception and design of the work. C.S. and M.G.-R. collected and analyzed the data. C.S. and G.T.-O. interpreted the results. C.S. drafted the manuscript, which was carefully revised by all authors. The final version of the manuscript has been approved by all the authors who agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All authors qualify for authorship and all those who qualify for authorship are listed.

Funding

This work is funded by NSF1535036 and C.S is funded by a NSF fellowship (GRFP)

Bibliography

1.   Vaziri, S., Diedrichsen, J. & Shadmehr, R. Why Does the Brain Predict Sensory

Consequences of Oculomotor Commands? Optimal Integration of the Predicted

and the Actual Sensory Feedback. *J. Neurosci.* **26,** 4188–4197 (2006).

2.   Coslett, H. B., Buxbaum, L. J. & Schwoebel, J. Accurate reaching after active but

not passive movements of the hand: Evidence for forward modeling. *Behav.*

Neurol. **19,** 117–125 (2008).

3.   Laufer, Y., Hocherman, S. & Dickstein, R. Accuracy of reproducing hand position

when using active compared with passive movement. *Physiother. Res. Int.* **6,** 65–

75 (2001).

4.   Paillard, J. & Brouchon, M. Active and passive movements in the calibration of

position sense. *Neuropsychol. Spat. oriented Behav.* **11,** 37–55 (1968).

5.   Haith, A., Jackson, C. P., Miall, R. C. & Vijayakumar, S. Unifying the Sensory

and Motor Components of Sensorimotor Adaptation. *Adv. Neural Inf. Process.*

Syst. 593–600 (2009).

6.   Mattar, A. A. G., Darainy, M. & Ostry, D. J. Motor learning and its sensory

effects : time course of perceptual change and its presence with gradual

introduction of load. *J. Neurophysiol.* **109,** 782–791 (2013).

7.   Van Beers, R. J., Sittig, A. C. & Denier van der Gon, J. J. Integration of

proprioceptive and visual position-information: An experimentally supported

model. *J. Neurophysiol.* **81,** 1355–1364 (1999).

8.   Cressman, E. K. & Henriques, D. Y. P. Sensory Recalibration of Hand Position

Following Visuomotor Adaptation. *J. Neurophysiol.* **102,** 3505–3518 (2009).

9.   Cressman, E. K. & Henriques, D. Y. P. Reach Adaptation and Proprioceptive

Recalibration Following Exposure to Misaligned Sensory Input. *J. Neurophysiol.*

103, 1888–1895 (2010).

10. t' Hart, B. M., Henriques, D. Y. P., Marius, B. & Henriques, D. Y. P. Separating

predicted and perceived sensory consequences of motor learning. *PLoS One* **11,** 1–

15 (2016).

11. Ostry, D. J., Darainy, M., Mattar, A. A. G., Wong, J. & Gribble, P. L.

Somatosensory Plasticity and Motor Learning. *J. Neurosci.* **30,** 5384–5393 (2010).

12. Salomonczyk, D., Cressman, E. K. & Henriques, D. Y. P. Neuropsychologia

Proprioceptive recalibration following prolonged training and increasing

distortions in visuomotor adaptation. *Neuropsychologia* **49,** 3053–3062 (2011).

13. Salomonczyk, D., Henriques, D. Y. P. & Cressman, E. K. Proprioceptive

recalibration in the right and left hands following abrupt visuomotor adaptation.

Exp. Brain Res. **217,** 187–196 (2012).

14. Salomonczyk, D., Cressman, E. K. & Henriques, D. Y. P. The role of the cross -

sensory error signal in visuomotor adaptation. *Exp. Brain Res.* **228,** 313–325

(2013).

15. Zbib, B., Henriques, D. Y. P. & Cressman, E. K. Proprioceptive recalibration

arises slowly compared to reach adaptation. *Exp. Brain Res.* **234,** 2201–2213

(2016).

16. Vazquez, A., Statton, M. A., Busgang, S. A. & Bastian, A. J. Split-belt walking

adaptation recalibrates sensorimotor estimates of leg speed but not position or

force. *J. Neurophysiol.* **114,** 3255–3267 (2015).

17.   Mostafa, A. A., Salomonczyk, D., Cressman, E. K. & Henriques, D. Y. P.

Intermanual transfer and proprioceptive recalibration following training with

translated visual feedback of the hand. *Exp. Brain Res.* **232,** 1639–1651 (2014).

18.   Mostafa, A. A., Kamran, R., Bahari, G., Erin, K. & Henriques, D. Y. P.

Generalization of reach adaptation and proprioceptive recalibration at different

distances in the workspace. *Exp. brain Res.* **233,** 817–827 (2015).

19.   Cressman, X. E. K. & Henriques, D. Y. P. Generalization patterns for reach

adaptation and proprioceptive recalibration differ after visuomotor learning. *J.*

Neurophysiol. **114,** 354–365 (2015).

20.   Vasudevan, E. V. L. & Bastian, A. J. Split-belt treadmill adaptation shows

different functional networks for fast and slow human walking. *J. Neurophysiol.*

103, 183–91 (2010).

21.   Wutzke, C. J., Faldowski, R. A. & Lewek, M. D. Individuals Poststroke Do Not

Perceive Their Spatiotemporal Gait Asymmetries as Abnormal. *Phys. Ther.* **95,**

1244–1253 (2015).

22.   Orendurff, M. S., Bernatz, G. C., Schoen, J. A. & Klute, G. K. Kinetic mechanisms

to alter walking speed. *Gait Posture* **27,** 603–610 (2008).

23.   Dewolf, A. H., Ivanenko, Y., Zelik, K. E., Lacquaniti, F. & Willems, P. A.

Kinematic patterns while walking on a slope at different speeds. *J. Appl. Physiol.*

125, 642–653 (2018).

24.   Torres-Oviedo, G. & Bastian, A. J. Natural error patterns enable transfer of motor

learning to novel contexts. *J. Neurophysiol.* **107,** 346–56 (2012).

25.   Leech, K. A., Roemmich, R. T., Day, K. A. & Bastian, A. J. Movement and

perception recalibrate differently across multiple days of locomotor learning. *J.*

Neurophysiol. **114,** 608–623 (2018).

26.   Bouton, M. E., Nelson, J. B. & Rosas, J. M. Stimulus generalization, context

836   change, and forgetting. *Psychol. Bull.* **125,** 171–186 (1999).

837   27.   Spear, N. E. *The processing of memories: forgetting and retention.* (Hillsdale,

838   1978).

839   28.   Tulvig, E. & Thomson, D. M. Encoding specificity and retrieval processes in

840   episodic memory. *Psychol. Rev.* **80,** 352–373 (1973).

841   29.   Torres-Oviedo, G. & Bastian, A. J. Seeing Is Believing : Effects of Visual

842   Contextual Cues on Learning and Transfer of Locomotor Adaptation. *J. Neurosci.*

843   **30,** 17015–17022 (2010).

844   30.   Desmurget, M., Jordan, M., Prablanc, C. & Jeannerod, M. Constrained and

845   Unconstrained Movements Involve Different Control Strategies. *J. Neurophysiol.*

846   **77,** 1644–1650 (1997).

847   31.   Reisman, D. S., McLean, H., Keller, J., Danks, K. A. & Bastian, A. J. Repeated

848   split-belt treadmill training improves poststroke step length asymmetry.

849   *Neurorehabil. Neural Repair* **27,** 460–468 (2013).

850   32.   Lewek, M. D., Braun, C. H. & Wutzke, C. The role of movement errors in

851   modifying spatiotemporal gait asymmetry post stroke: a randomized controlled

852   trial. *Clin. Rehabil.* **32,** 161–172 (2018).

853   33.   Cressman, E. K., Salomonczyk, D. & Henriques, D. Y. P. Visuomotor adaptation

854   and proprioceptive recalibration in older adults. *Exp. Brain Res.* **205,** 533–544

855   (2010).

856   34.   Forero, J. & Misiaszek, J. E. The contribution of light touch sensory cues to

857   corrective reactions during treadmill locomotion. *Exp. Brain Res.* **226,** 575–584

858   (2013).

859   35.   Winstein, C. J. & Schmidt, R. A. Reduced frequency of knowledge of results

860   enhances motor skill learning. *J. Exp. Psychol. Learn. Mem. Cogn.* **16,** 677 (1990).