1 **Evaluation of DNA extraction methods on individual helminth egg and larval stages**
2 **for whole genome sequencing**
3

Stephen R. Doyle[1*], Geetha Sankaranarayan[1], Fiona Allan[2], Duncan Berger[1], Pablo D.
Jimenez Castro[3], James Bryant Collins[3], Thomas Crellen[1,10], María A. Duque-Correa[1],
Peter Ellis[1], Tegegn G. Jaleta[3], Roz Laing[4], Kirsty Maitland[4], Catherine McCarthy[1],
Tchonfienet Moundai[6], Ben Softley[1], Elizabeth Thiele[7], Philippe Tchindebet Ouakou[6],
John Vianney Tushabe[1,11], Joanne P. Webster[8], Adam J. Weiss[9], James Lok[4], Eileen
Devaney[5], Ray M. Kaplan[3], James A. Cotton[1], Matthew Berriman[1], Nancy Holroyd[1*]
10

11 * Correspondence:
12 Stephen R. Doyle: stephen.doyle@sanger.ac.uk
13 Nancy Holroyd: neh@sanger.ac.uk
14

[1.] Wellcome Sanger Institute, Hinxton, Cambridgeshire, United Kingdom
[2.] Department of Life Sciences, Natural History Museum, London, United Kingdom
[3.] Department of Infectious Diseases, College of Veterinary Medicine, University of
Georgia, Athens, Georgia, USA
[4.] Department of Pathobiology, School of Veterinary Medicine, University of
Pennsylvania, Philadelphia, Pennsylvania, USA
[5.] Institute of Biodiversity Animal Health and Comparative Medicine, College of
Medical, Veterinary and Life Sciences, University of Glasgow, Garscube Campus,
Glasgow, United Kingdom
[6.] Department of Biology, Vassar College, Poughkeepsie, New York, USA
[7.] Ministry of Public Health, N'Djamena, Chad
[8.] Centre for Emerging, Endemic and Exotic Diseases, Department of Pathology and
Population Sciences, Royal Veterinary College, University of London, Herts, United
Kingdom
[9.] Guinea Worm Eradication Program, The Carter Center, Atlanta, Georgia, USA
[10.] Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom
[11.] Medical Research Council/Uganda Virus Research Institute and London School of
Hygiene & Tropical Medicine Uganda Research Unit, P.O. Box 49, Entebbe, Uganda

Keywords: Helminth, genomics, whole genome sequencing, DNA extraction, low
input, diagnostics

Abstract

Whole genome sequencing is being rapidly applied to the study of helminth
genomes, including *de novo* genome assembly, population genetics, and diagnostic

applications. Although late-stage juvenile and adult parasites typically produce

sufficient DNA for molecular analyses, these parasitic stages are almost always

inaccessible in the live host; immature life stages found in the environment for which

samples can be collected non-invasively offer a potential alternative, however, these

samples are typically yield very low quantities of DNA, can be environmentally

resistant, and are susceptible to contamination, often from bacterial or host DNA.

Here, we have tested five low-input DNA extraction protocols together with a low-

input sequencing library protocol to assess the feasibility of whole genome

sequencing of individual immature helminth samples. These approaches do not use

whole genome amplification, a common but costly approach to increase the yield of

low input samples. We first tested individual parasites from two species spotted

onto FTA cards - egg and L1 stages of *Haemonchus contortus* and miracidia of

Schistosoma mansoni - before further testing on an additional six species -

Ancylostoma caninum, Ascaridia dissimilis, Dirofilaria immitis, Dracunculus

medinensis, Strongyloides stercoralis, and Trichuris muris - with an optimal protocol.

Whole genome sequencing followed by analyses to determine the proportion of on-

and off-target mapping revealed successful sample preparations for six of the eight

species tested with variation between species, and within species but between life

stages, described. These results demonstrate the feasibility of whole genome

sequencing of individual parasites, and highlight a new avenue towards generating

sensitive, specific, and information-rich data for the diagnosis and surveillance of

helminths.

Introduction

Accurate methods for diagnosis and surveillance of helminth infections are of increasing interest in both human and animal health settings. Such approaches are typically proposed to monitor the presence and ultimately decline of populations targeted by large-scale control measures, such as mass drug administration (MDA) for the prevention and/or treatment of human helminth infections, or prophylactic treatment of domesticated animals. An ideal diagnostic will be sensitive, to detect the parasite if in fact present, and specific, to identify the targeted parasite species in the presence of non-target material such as other parasite species or the host. Ideally, samples taken for diagnostic purposes can be used to gather additional information beyond the presence or absence of a specific parasite, so the same material could be used for example, to predict how well the infection will respond to drug treatment, or how the parasite is related to other endemic or imported parasites. As most parasitic stages of helminths of humans and animals are naturally inaccessible *in vivo* (not accounting for potential availability of some mature stages of helminths following chemo-expulsion, for example, *Ascaris lumbricoides* and *Trichuris trichiura*), a diagnostic should also be informative on non-invasive stages of the parasite, such as eggs deposited in faeces, or intermediate stages of the parasite's life cycle that exist in the external environment.

A key challenge of working with environmental intermediate stages is that they are often immature, for example, eggs or early stage larvae, and extremely small (for example, *Haemonchus contortus* eggs are approximately 75 × 44 µm and *Schistosoma mansoni* miracidia approximately 140 × 55 µm), limiting the amount of accessible material (e.g., DNA) available to perform the diagnostic assay. They are often environmentally resistant, and the same features that naturally protect the DNA from damage prior to reinfection make it difficult to extract DNA. In many cases they are isolated from host faeces and so are susceptible to bacterial contamination, or from host tissues and so become contaminated with host DNA. Furthermore, samples may need to be transported efficiently to a laboratory setting without a significant loss of this already limited material. A number of approaches have been tested to preserve macromolecules from individual parasites for transport and

storage, including ethanol, RNAlater and Whatman® FTA® cards, from which robust

PCR and microsatellite data could be profiled (Boué et al., 2017; Campbell et al.,

2017; Gower et al., 2007; Marek et al., 2014; Webster, 2009; Webster et al., 2012;

Xiao et al., 2013). Although under ideal conditions the detection of a single DNA

molecule is possible, the limited material available per parasite has to date largely

restricted assaying to a small number of loci, limiting the amount of information

obtained from any individual parasite.

Genomic approaches offer an information-rich technology for diagnostic and

surveillance applications. Increasing throughput and decreasing costs of whole

genome sequencing has resulted in the recent and steadily growing application of

genomics in helminth parasitology; for example, for diagnostic applications, high

throughput amplicon sequencing for helminth species identification and community

composition (Avramenko et al., 2015) and the presence of drug resistance alleles

(Avramenko et al., 2019) have been described. Although low DNA concentrations are

typically prohibitive for genome-wide approaches on individual parasites, a number

of studies have successfully used whole genome amplification on DNA extracted

from single larval stages to perform reduced representation (Shortt et al., 2017) and

exome (Le Clec'h et al., 2018; Platt et al., 2019) sequencing on miracidia of

Schistosoma spp., and whole genome sequencing of *Haemonchus contortus* L3 stage

larvae (Doyle et al., 2018) and microfilaria of *Wuchereria bancrofti* (Small et al.,

2018). Whole genome amplification protocols do, however, add considerable

expense per sample, and can introduce technical artefacts such as uneven and/or

preferential amplification (potentially of contaminant sequences), chimeric

sequences, and allele dropout (Sabina and Leamon, 2015; Tsai et al., 2014), that may

lead to a reduction in genetic diversity, and in turn, relevance to the original

unamplified material. The field of genomics is, however, rapidly advancing towards

very low minimum sample input requirements, and single cell approaches for DNA

and RNA sequencing are now available. Such approaches have begun to be used on

parasitic species such as *Plasmodium* spp. (Howick et al., 2019; Ngara et al., 2018;

Reid et al., 2018; Trevino et al., 2017), but are yet to be adopted by helminth

parasitologists. Although these low-input, high-throughput approaches are not

designed – and perhaps not currently suitable – for diagnostic applications, the

developments in molecular biology techniques for low input sequencing can benefit

the use of genomics for helminth applications. Here, we test a number of low-input

DNA extraction approaches for individual helminth samples stored on Whatman®

FTA® cards, followed by low-input library preparation without whole genome

amplification, and whole-genome sequencing. A total of five DNA extraction

approaches were initially tested, after which the most promising approach was

applied to multiple life stages from eight helminth species. The results presented

here demonstrate the advancement of low-input whole genome sequencing, and are

discussed in the context of their utility for helminth diagnostics and surveillance.

Methods

Sample collection

Samples representing accessible, immature life stages of a total of eight helminth

species were tested, the collection of which is described below.

Ancylostoma caninum: Fresh feces from a research purpose-bred laboratory beagle

(University of Georgia AUP # A2017 10-016-Y1-A0) infected with the Barrow isolate

(drug-susceptible isolate from Barrow County Georgia, USA) were collected and

made into a slurry with water, filtered through 425 μm and 180 μm sieves, and

centrifuged at 2500 rpm for 5 min after which the supernatant was discarded. Kaolin

(Sigma-Aldrich, St. Louis, MO) was then added and resuspended in sodium nitrate

(SPG 1.25–1.3) (Feca-Med®, Vedco, Inc. St Joseph, MO, USA). The tube was then

centrifuged at 2500 rpm for 5 min, after which the supernatant was passed through

a 30 μm sieve and rinsed with distilled water, and reduced to a volume of 10-15 mL.

The volume was adjusted to 1 egg per 5 μL using distilled water. The eggs were

stored at room temperature for 2 h before placing them onto the Whatman® FTA®

cards. Eggs were also placed onto Nematode growth medium (NGM) plates (Sulston

and Hodgkin, 1988) and incubated at 26°C to obtain the first-stage (L1) larvae. After

48 h, larvae were rinsed off the plate with distilled water and centrifuged at 1000

rpm for 5 mins. Larvae were counted and the concentration adjusted to 1 larva per 5

μL. The larvae were stored at room temperature for 2 h before placing them onto

the Whatman® FTA® cards. To obtain third-stage (L3) larvae, eggs were isolated from

fresh feces from a research purpose-bred laboratory beagle (University of Georgia

AUP # A2017 10-016-Y1-A0) infected with the Worthy isolate (Worthy 3.1F3Pyr;

multiple-drug resistant isolate originally isolated from a greyhound dog, Florida,

USA). Eggs were placed onto NGM plates (Sulston and Hodgkin, 1988) and incubated

at 26°C. After seven days, larvae were rinsed off the plate with distilled water and

centrifuged at 1000 rpm for 5 mins. Larvae were counted and the concentration

adjusted to 1 larva per 5 μL. The larvae were stored at room temperature for 2 h

before placing them onto the Whatman® FTA® cards.

Ascaridia dissimilis: Eggs of *A. dissimilis* (Isolate Wi: North Carolina, USA) were

isolated from excreta of experimentally infected turkeys. Water was added to the

excreta and made into a slurry, which was filtered using a 425 µm and 180 µm sieve

to remove large debris. The remaining particulates were placed into 50 mL

centrifuge tubes and centrifuged at 433 g for 7 mins. Supernatant was removed and

the pellet was resuspended in a saturated sucrose solution with a specific gravity of

1.15. The suspension was centrifuged as before and eggs were isolated from the top

layer. Eggs were rinsed over a 20 µm sieve with water to remove residual sucrose

and then concentrated to 1 egg per 5 µL using deionized water. Multiple 5 µL

aliquots of the egg solution were dispensed using a micropipette onto the

Whatman® FTA® card.

Dirofilaria immitis: Blood was taken from a dog infected with the macrocyclic lactone

(ML)-resistant Yazoo strain (Yazoo: originally isolated from a dog in Yazoo City,

Mississippi, USA; See Maclean *et al.* (2017) for complete history). To obtain

microfilariae, blood was collected in heparin tubes and centrifuged for 30 min at

2500 rpm after which the supernatant was discarded. The pellet was suspended in

3.8% sodium citrate (Sigma-Aldrich, St. Louis, MO) and 15% saponin (Sigma-Aldrich,

St. Louis, MO) was added in a 1:7 dilution. The tube was then vortexed and

centrifuged for 30 min at 2500 rpm after which the supernatant was discarded and

the pellet resuspended in 3.8% sodium citrate to the original blood volume,

vortexed, and then centrifuged for 4 min at 2500 rpm. The pellet was then

resuspended and mixed in a 1:9 solution of 10⊠ phosphate buffered saline (PBS;

Thermo Fisher Scientific, Waltham, MA) and distilled water. The tube was then

centrifuged for 4 min at 2500 rpm and the pellet resuspended in PBS. The

microfilariae were then counted and adjusted accordingly to have one microfilaria

per 5 µL and stored at room temperature for 2 h before placing them onto the

Whatman® FTA® cards.

Dracunculus medinensis: Individual L1 samples were obtained as progeny of an adult

female worm manually extracted from an infected dog in Tarangara village, Chad

(9.068611 N, 18.708611 E) in 2016. This extraction forms part of the standard

containment and treatment procedure for Guinea worm infections, as agreed upon

and sanctioned by the World Health Organisation and country ministries of health.

The adult worm was submerged in ethanol in a microcentrifuge tube for storage; L1

stage progeny that were found settled on the bottom of the tube were collected for

analysis.

Haemonchus contortus: Eggs representing the F5 generation of a genetic cross

(described in Doyle *et al.* (2018)) were collected from fresh faeces from

experimentally infected sheep housed at the Moredun Research Institute, UK. All

experimental procedures were examined and approved by the Moredun Research

Institute Experiments and Ethics Committee and were conducted under approved

UK Home Office licenses (PPL 60/03899) in accordance with the Animals (Scientific

Procedures) Act of 1986. Briefly, faeces were mixed with tap water and passed

through a 210 μm sieve, then centrifuged at 2500 rpm for 5 mins in polyallomer

tubes. The supernatant was discarded, before adding kaolin to the faecal pellet,

vortexing, and resuspending in a saturated salt solution. After centrifugation at 1000

rpm for 10 mins, the polyallomer tube was clamped to isolate eggs, which were

collected on a 38 μm sieve and rinsed thoroughly with tap water. Eggs were

incubated on NGM plates at 20°C for 48 h to hatch to L1 stage larvae. In addition to

freshly collected material, eggs collected in the same manner then stored at -20°C,

from a previous generation of the cross, were also tested. Eggs and L1 larvae were

resuspended in PBS and spotted onto Whatman FTA cards in 3 μL per egg/L1.

Schistosoma mansoni: Three collections of *S. mansoni* samples were used in this

work. The first were field samples collected from humans on Lake Victoria fishing

villages in Uganda as part of the LaVIISWA trial (Sanya et al., 2018). Ethical approval

for this trail was given by the Uganda Virus Research Institute (reference number

GC127), Uganda National Council for Science and Technology (reference number HS

1183) and London School of Hygiene & Tropical Medicine (reference number 6187).

Parasite eggs were collected from participants' stool samples using a Pitchford-Visser

funnel, washed with mineral water until clean, and transferred into a petri dish with

water to be hatched in direct sunlight. After hatching, the miracidia were picked in 2

μL water using a pipette and placed on a Whatman® FTA® card for storage. The
second were field samples collected as part of a repeated cross-sectional study of
MDA exposure in school children in Uganda. Patient enrolment, including written
consent, and sample collection have been described previously (Crellen et al., 2016).
Ethical approvals for this study were granted by the Uganda National Council of
Science and Technology (MoU sections 1.4, 1.5, 1.6) and the Imperial College
Research Ethics Committee (EC NO: 03.36. R&D No: 03/SB/033E). Host stool was
sampled 1-3 days prior to treatment with praziquantel (40 mg/kg) and albendazole
(400 mg). A Pitchford-Visser funnel was used to wash and filter stool to retain
parasite eggs. The filtrate was kept overnight in water and hatched the following
morning in sunlight. Individual miracidia were isolated with a 20 μL pipette and
transferred into petri-dishes of nuclease-free water twice before spotting onto
Whatman® FTA® cards. The third source of miracidia were derived from the livers of
experimentally infected mice kept to maintain the *S. mansoni* life cycle at Wellcome
Sanger Institute. Mouse infection protocols were approved by the Welfare and
Ethical Review Body (AWERB) of the Wellcome Sanger Institute. The AWERB is
constituted as required by the UK Animals (Scientific Procedures) Act 1986
Amendment Regulations 2012. BalbC mice (6-8 week old) were infected with 250
cercariae, after which livers were collected on day 40 post infection. Eggs were
isolated from the liver tissues using collagenase digestion followed by percoll
gradient, and were washed well with sterile PBS, before being hatched in sterile
conditioned water. The hatched individual miracidia were spotted onto Whatman®
FTA® cards.
262
Strongyloides stercoralis: The *S. stercoralis* UPD strain and the isofemale isolate
PVO1 were maintained in purpose-bred, prednisone-treated mix breed dogs
according to protocol 804883 approved by the University of Pennsylvania
Institutional Animal Care and Use Committee (IACUC), USA. IACUC-approved
research protocols and all routine husbandry care of the animals was conducted in
strict accordance with the Guide for the Care and Use of Laboratory Animals of the
National Institutes of Health, USA. Faeces were collected, moisturized and mixed
with equal volume of charcoal and cultured at 22°C in 10 cm plates (Lok, 2007). Post

parasitic stage one and two larvae (L1/L2), free-living male and female adults, and

infective third-stage (L3) were isolated by the Baermann technique after 24 h, 48 h,

and 6 days in these charcoal cocultures, respectively (Jaleta et al., 2017; Lok,

2007). For this study, free living females, L1 and L3 larvae were collected from the

Baermann funnel sediments and washed three times using PBS.

Trichuris muris: Infection and maintenance of *T. muris* was conducted as described

(Wakelin, 1967). The care and use of mice were in accordance with the UK Home

Office regulations (UK Animals Scientific Procedures Act 1986) under the Project

license P77E8A062 and were approved by the institutional Animal Welfare and

Ethical Review Body. Female SCID mice (6–10 wk old) were orally infected under

anaesthesia with isoflurane with a high dose (n = 400) of embryonated eggs from *T.*

muris E-isolate. Mice were monitored daily for general condition and weight loss. At

day 35 post infection, mice were killed by exsanguination under terminal anesthesia,

after which adult worms were harvested from cecums. Adult worms were cultured in

RPMI 1640 supplemented with 10% fetal calf serum (v/v), 2 mM L-glutamine,

penicillin (100 U/mL), and streptomycin (100 mg/mL; all Invitrogen), for 4 h or

overnight, and eggs were collected. The eggs were allowed to embryonate for at

least 6 weeks in distilled water, and infectivity was established by worm burden in

SCID mice. *T. muris* eggs were hatched to produce sterile L1 larvae using 32% sodium

hypochlorite in sterile water for 2 h at 37°C with 5% CO_2. Eggs were washed with

RPMI 1640 supplemented with 10% fetal calf serum (v/v), 2 mM L-glutamine,

penicillin (100 U/mL), and streptomycin (100 mg/mL; all Invitrogen), and incubated

at 37°C with 5% CO_2 for 4 to 5 days until they hatched.

For each species, unless otherwise stated, pools of individuals were washed in sterile

PBS, before being transferred to a petri dish. Individuals were identified under the

microscope, after which 5 μL PBS containing an individual parasite was transferred

onto a Whatman® FTA® card and dried for a minimum of 20 mins at room

temperature prior to storage or shipping to the Wellcome Sanger Institute, UK. The

Whatman® FTA® cards with samples spotted were stored in a clean plastic bag in the

dark at room temperature prior to analysis.

DNA extraction

The sample spots on FTA cards were punched out manually into 96-well plates using either a Harris Punch or autonomously using robotics. The DNA extraction was carried out for each method as described below:

1. *Nexttec* (NXT): Extraction using the Nexttec 1-step DNA Isolation Kit for Tissues & Cells (cat: 10N.904; Waendel Technology Limited, UK) was performed according to manufacturer's guidelines. 75 µl of proteinase lysis buffer was used for digestion.

2. *Bloodspot* (BSP): The bloodspot extraction was derived from the QIAamp DNA Investigator Kit (cat: 51104/51106; Qiagen), following the "DNA Purification from Dried Blood Spots (QIAamp DNA Mini Kit)" protocol according to manufacturer's guidelines.

3. *CGP* (CGP): Extraction using the CGP protocol (Moore et al., 2018) involved adding 30 µL of lysis buffer (1.25 µg/mL of Protease reagent (Qiagen; cat# 19155) in Tris HCl pH 8.0, 0.5% Tween 20, 0.5% NP40) to individual Whatman® FTA® punches in a 96-well PCR plate. The samples were incubated at 50°C for an hour followed by protease inactivation by heating the samples to 75°C for 30 min.

4. *ForensicGem* (FGM): Extraction using the ForensicGEM Universal DNA extraction kit (cat: FUN0100; ZyGem) was performed according to the manufacturer's guidelines.

5. *PicoPure* (PIP): Extraction using the ARCTURUS® PicoPure® DNA Extraction Kit (cat: KIT0103; ThermoFisher) was performed according to the manufacturer's guidelines, using 75 µL of the extraction buffer.

When the extracted DNA from all the above methods was present in a volume greater than 25 µL, samples were cleaned with Agencourt AMPure XP beads (Beckman-Coulter) and eluted in 25 µL nuclease-free H_2O. The entire DNA samples were used downstream to make sequencing libraries. A summary of the number of species, life stages, and conditions tested, is presented in **Table 1**.

338   **Table 1.** Summary of the number of species, life stages, and conditions tested

Species	Life stage	Extraction kit	Sample number
Ancylostoma caninum	Eggs	CGP	8
Ancylostoma caninum	L1	CGP	8
Ancylostoma caninum	L3	CGP	8
Ascaridia dissimilis	Eggs	CGP	8
Dirofilaria immitis	microfilaria	CGP	8
Dracunculus medinensis	L1	Nexttec	129
Haemonchus contortus	Egg	Bloodspot	10
Haemonchus contortus	Egg	CGP	10
Haemonchus contortus	Egg	ForensicGem	4
Haemonchus contortus	Egg	Nexttec	10
Haemonchus contortus	Egg	Picopure	10
Haemonchus contortus	Frozen Egg	Bloodspot	6
Haemonchus contortus	Frozen Egg	CGP	6
Haemonchus contortus	Frozen Egg	Nexttec	6
Haemonchus contortus	Frozen Egg	Picopure	6
Haemonchus contortus	L1	Bloodspot	10
Haemonchus contortus	L1	CGP	10
Haemonchus contortus	L1	ForensicGem	4
Haemonchus contortus	L1	Nexttec	10
Haemonchus contortus	L1	Picopure	10
Schistosoma mansoni	miracidia	Bloodspot	59
Schistosoma mansoni	miracidia	CGP	36
Schistosoma mansoni	miracidia	ForensicGEM	10
Schistosoma mansoni	miracidia	Nexttec	51

Schistosoma mansoni	miracidia	PicoPure	12
Strongyloides stercoralis	FL	CGP	8
Strongyloides stercoralis	iL3	CGP	8
Strongyloides stercoralis	L1	CGP	8
Trichuris muris	bleached eggs	CGP	8
Trichuris muris	eggs	CGP	8
Trichuris muris	L1	CGP	8

Library preparation and Sequencing

DNA sequencing libraries for all samples were prepared using a protocol designed for library preparation of laser capture microdissected biopsy (LCMB) samples using the Ultra II FS enzyme (New England Biolabs) for DNA fragmentation as previously described (Lee-Six et al., 2019). A total of 12 cycles of PCR were used (unless otherwise stated in **Table S2**) to amplify libraries and to add a unique 8-base index sequence for sample multiplexing.

Multiplexed libraries were sequenced using the Illumina MiSeq system with V2 chemistry 150 bp paired end (PE) reads. The *D. medinensis* samples were sequenced as part of a different study using the HiSeq 2500 with V4 chemistry 125 bp PE reads. In general, we performed sufficient low coverage sequencing on each sample to enable us to identify: (i) the proportion of on-target mapped reads, (ii) the proportion of duplicate reads, i.e. library artefacts, and (iii) the proportion of off-target contaminant reads.

Metadata for each sample, including sample IDs, sequencing lane IDs, ENA sample accession numbers, and data generated are described in **Table S2**. Raw sequence data will be made available under ENA study ID ERP114942.

Analysis

Reference genomes from each of the test species were obtained from WormBase

Parasite (Howe et al., 2017) Release 12. Raw sequence data for each species were

mapped to their respective reference genome using BWA-MEM (Li, 2013), after

which duplicate reads were marked using Picard (v2.5.0;

https://github.com/broadinstitute/picard). Samtools flagstats and bamtools stats

were used to characterise the outcome of the mapping, the results of which were

collated using MultiQC (Ewels et al., 2016). Data were manipulated and visualised in

the R (v3.5.0) environment using the following packages: ggplot2

(https://ggplot2.tidyverse.org/), patchwork

(https://github.com/thomasp85/patchwork), and dplyr

(https://dplyr.tidyverse.org/).

The code to reproduce the analysis and figures for this manuscript is described in

https://github.com/stephenrdoyle/helminth_extraction_wgs_test.

Results & Discussion

Results & Discussion

The aim of this work was to determine the feasibility and efficiency of using a low

input DNA extraction and library preparation approach for whole-genome

sequencing of individual helminth parasites. We have targeted immature life stages

that are found in the environment for which it is possible to collect samples non-

invasively. We first tested our approaches on the nematode *H. contortus* and the

trematode *S. mansoni*. Five approaches were tested using single egg (fresh and

frozen) and L3 of *H. contortus*, and miracidia of *S. mansoni*. We determined the

success of the library preparation approach by comparing the proportion of reads

mapped (**Figure 1 & 2**; top plots), representing on-target mapping as a measure of

specificity, as well as the proportion of duplicate reads (**Figure 1 & 2**; bottom plots),

which typically represent library preparation artefacts due to over-amplification of

DNA during PCR. Finally, we determined the proportion of reads associated with off-

target contamination by comparison of raw reads to a kraken contamination

database (**Table S2**; kraken_unassigned).

There were distinct differences in overall mapping between *H. contortus* and *S.*

mansoni; while some of this variation may reflect differences in extracting DNA from

different species, the *S. mansoni* samples generally had a greater proportion of

contaminating DNA present relative to the *H. contortus* samples (**Table S2**;

kraken_unassigned). We achieved some on-target mapping under all optimisation

conditions tested, however, significant variation was observed between approaches

(**Figure 1**). BSP and NXT generally performed poorly, with either low (median =

19.42; median absolute deviation (MAD) = 16.50) or significant variance (median =

66.20; MAD = 26.18) in mapping frequency between samples observed, respectively.

PIP performed consistently well with high mapping rates across all stages in *H.*

contortus (median = 90.30; MAD = 3.96), however, it was poor in *S. mansoni* (median

= 11.00; MAD = 12.73). The duplication rate of all conditions were within an

acceptable low range (median = 0.46, MAD = 0.4), with only 2% of samples having

greater than 5% duplicate reads. CGP and FGM performed most consistently

between stages and species; FGM had higher variance and duplication rates relative

to CGP across all samples tested, and therefore, CGP was chosen to explore further.

Figure 1. Comparison of library preparation approaches on *Haemonchus contortus* eggs (EGG), frozen eggs (EGGf), and L1 (LS1) stages, and *Schistosoma mansoni* miracidia (MIR). The top plot presents the percentage of reads that mapped to each reference genome, and the bottom plot presents the proportion of duplicate reads identified. Boxplots are coloured by the library preparation approach used. The number of samples in each comparison is presented in Table 1.

We expanded our analysis to a total of eight helminth species for which samples were available, including a total of six distinct life stages (**Figure 2**). All tested samples were extracted with the CGP protocol, except for *D. medinensis*, which were

extracted using NXT and included for comparison. High variability in mapping was

observed between species, with 50% or greater mapping frequency achieved in at

least one life stage of 6 of the 8 species tested. Clear differences were observed

between multiple life stages tested within a species, likely reflecting differences in

extraction efficiency, for example: for *A. canium,* reads from eggs (median = 54.98)

mapped much more effectively than L1 (median = 3.61) or L3 stages (median = 7.57),

and in *S. stercoralis,* full-length females (median = 67.84) and L1 (median = 47.19)

performed better than L3 (median = 9.01). Interestingly, L1 larvae (median = 41.85)

and bleached eggs (median = 51.35) performed much better than untreated eggs

(median = 2.72) of *T. muris*; bleaching is experimentally used for promoting hatching

of *T. muris*, by dissolving the egg shell layers and in turn improving access to DNA

within. However, bleached eggs were embryonated and developmentally more

advanced than untreated unembryonated eggs, and therefore would have more

DNA available for library preparation. Similar to untreated *T. muris* eggs, *A. dissimilis*

eggs performed poorly (median = 0.16); for both species, few if any nematode

sequences were recovered and the majority of sequencing reads were

contaminating bacterial-derived contaminants, perhaps indicative of the challenge of

accessing material with the environmentally-resistant egg. Analysis of *A. dissimilis*

was further limited by the lack of a reference genome for this species; we mapped

against the available genome of *Ascaris lumbricoides*, the nearest species for which a

reference was available, and therefore at best, would have expected suboptimal

mapping due to sequence divergence. The duplication rates remain low for all

species and conditions tested, apart from *D. medinensis*, which was much higher

(median = 36.90). This was unexpected, given NXT did not produce high duplication

rates under the initial conditions tested with *H. contortus* or *S. mansoni*, nor were

there excessive PCR cycles used in this instance.

Figure 2. Comparison of library preparation from multiple life stages of 8 helminth species. All libraries were prepared with the CGP approach, except for *D. medinensis*, which was prepared with NXT and presented alongside for comparison. As in Figure 1, the top plot presents the percentage of reads that mapped to each reference genome, and the bottom plot presents the proportion of duplicate reads identified. Boxplots are coloured by the life stage assayed. Egg = untreated egg; EGGb = bleached egg; EGGf = frozen egg; FL = full-length female; LS1 = larval stage 1; LS3 = larval stage 3; MFL = microfilaria; MIR = miracidia. The number of samples in each comparison is presented in Table 1.

In summary, we present successful DNA extraction followed by whole-genome sequencing of individual parasites from 6 of 8 species examined. These results significantly extend the possibility of genomic analyses for life stages for which, at

best, were limited to low-resolution, low-throughput PCR based assays without the

addition of whole genome amplification. Whatman® FTA® cards provide a

convenient substrate for sample collection and storage, and do not limit the

application of direct DNA extraction and whole genome sequencing of parasite

samples, even for field samples as demonstrated for *S. mansoni* miracidia that were

collected and processed in Uganda before they were transported to the UK. Further

optimisation is required to improve the DNA recovery from eggs, for example, from

A. dissimilis and *T. muris*, to provide greater applicability of our approaches to

species that generate particularly environmentally-resistant stages, such as the soil

transmitted helminths. The application of whole genome sequencing to diagnose

and monitor helminth infections at scale is largely limited by the costs of library

preparation and sequencing, and therefore, will be restricted to niche applications of

the technology. However, targeting the mitochondrial genome, for example, by

whole genome sequencing may be a viable and cost-effective alternative, potentially

providing greater diagnostic information than low throughput PCR-based diagnostics

(**Table 2**). Continued development of genomic technologies and the associated

reduction in sequencing and library preparation costs will make screening large

samples by genome sequencing more routine as in viral (Dudas et al., 2017) and

bacterial (Domman et al., 2018) population studies. In doing so, the ability to derive

information-rich data for diagnostic and surveillance purposes using genomics

(Cotton et al., 2018) will be particularly informative as efforts to control human

infective helminths using MDA move from control to elimination.

Table 2. Breakdown of sequencing strategies per species based on whole genome sequencing at 30X coverage and whole genome sequencing to achieve 100X whole mitochondrial genome coverage

Species / stage	Genome assembly size [1]	Number of samples that can be multiplexed [2]	mtDNA/ nuclear ratio [3]	Sampled multiplexed targeting mtDNA at 100X coverage [2]	Nuclear genome coverage per 100X mtDNA genome
S. mansoni miracidia	409	173	142.77	7418	0.70
H. contortus egg	283	250	70.76	5313	1.41
H. contortus L1	283	250	44.55	3345	2.24
D. medinensis L1	103	688	70.81	14608	1.41
D. immitis microfilaria	88	805	14.27	3446	7.01
S. stercoralis FL	42	1687	11.06	5597	9.04

1. Genome size: Obtained from Wormbase Parasite v12
2. Multiplex: total number of samples per NovaSeq 6000 sequencing run with S4 2x150 bp PE chemistry generating 2.5 Tb of data and accounting for 85% mapping rate. Theoretically possible, but may be limited by barcoding available.
3. mtDNA/nuclear ratio: based on coverage of mitochondrial and nuclear derived sequencing reads, normalised to the nuclear genome coverage

Acknowledgments

Work performed at the Wellcome Sanger Institute is supported by Wellcome Trust
(grant 206194) and by the Biotechnology and Biological Sciences Research Council
(BB/M003949/1). We thank Alison Elliot for access to *S. mansoni* samples collected
from Uganda, the collection of which was supported by Wellcome Trust (grant
095778/Z/11/Z). *S. mansoni* samples were also obtained from the Schistosomiasis
Collection at the Natural History Muesem (NHM) [SCAN], which is funded with
support from the Wellcome Trust (grant no. 104958/Z/14/Z). We thank the Carter
Center for supporting molecular work on *D. medinensis,* and the Guinea worm
eradication program for making samples available.

Author contributions

Conceptualization: SRD, JAC, NH

Methodology: SRD, GS, NH

Software: SRD

Formal analysis: SRD

Investigation: SD, GS

Resources: FA, TJ, JL, JBC, PJC, JPW, TC, ET, TC, MAD-C, PE, RL, KM, CM, TM, BS, PTO,

JVT, AW, ED, RK, DB, MB

Data Curation: SRD, NH

Writing – original draft preparation: SRD

Writing – review and editing: All authors

Visualisation: SRD

Supervision: SRD, JAC, NH

Project administration: NH

Conflict of Interests

The authors declare no conflict of interests.

Supplementary Data

Table S1 - Summary of total samples analysed by species, life stage, and DNA

extraction protocol (excel workbook)

Table S2 - Complete metadata per sample (excel workbook)

References

Avramenko, R. W., Redman, E. M., Lewis, R., Yazwinski, T. A., Wasmuth, J. D., and Gilleard, J.
S. (2015). Exploring the Gastrointestinal "Nemabiome": Deep Amplicon Sequencing to
Quantify the Species Composition of Parasitic Nematode Communities. *PLoS One* 10,
e0143559. doi:10.1371/journal.pone.0143559.

Avramenko, R. W., Redman, E. M., Melville, L., Bartley, Y., Wit, J., Queiroz, C., et al. (2019).
Deep amplicon sequencing as a powerful new tool to screen for sequence
polymorphisms associated with anthelmintic resistance in parasitic nematode
populations. *International Journal for Parasitology* 49, 13–26.
doi:10.1016/j.ijpara.2018.10.005.

Boué, F., El Berbri, I., Hormaz, V., Boucher, J.-M., El Mamy, A. B., Traore, A., et al. (2017). Use
of FTA® card methodology for sampling and molecular characterization of Echinococcus
granulosus sensu lato in Africa. *Exp. Parasitol.* 173, 29–33.

Campbell, S. J., Stothard, J. R., O'Halloran, F., Sankey, D., Durant, T., Ombede, D. E., et al.
(2017). Urogenital schistosomiasis and soil-transmitted helminthiasis (STH) in
Cameroon: An epidemiological update at Barombi Mbo and Barombi Kotto crater lakes
assessing prospects for intensified control interventions. *Infect Dis Poverty* 6, 49.

Cotton, J. A., Berriman, M., Dalén, L., and Barnes, I. (2018). Eradication genomics-lessons for
parasite control. *Science* 361, 130–131.

Crellen, T., Walker, M., Lamberton, P. H. L., Kabatereine, N. B., Tukahebwa, E. M., Cotton, J.
A., et al. (2016). Reduced Efficacy of Praziquantel Against Schistosoma mansoni Is
Associated With Multiple Rounds of Mass Drug Administration. *Clin. Infect. Dis.* 63,
1151–1159.

Domman, D., Chowdhury, F., Khan, A. I., Dorman, M. J., Mutreja, A., Uddin, M. I., et al.
(2018). Defining endemic cholera at three levels of spatiotemporal resolution within
Bangladesh. *Nat. Genet.* 50, 951–955.

Doyle, S. R., Laing, R., Bartley, D. J., Britton, C., Chaudhry, U., Gilleard, J. S., et al. (2018). A
Genome Resequencing-Based Genetic Map Reveals the Recombination Landscape of an
Outbred Parasitic Nematode in the Presence of Polyploidy and Polyandry. *Genome Biol.*
Evol. 10, 396–409.

Dudas, G., Carvalho, L. M., Bedford, T., Tatem, A. J., Baele, G., Faria, N. R., et al. (2017). Virus
genomes reveal factors that spread and sustained the Ebola epidemic. *Nature* 544,
309–315.

Ewels, P., Magnusson, M., Lundin, S., and Käller, M. (2016). MultiQC: summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics* 32, 3047–3048.

Gower, C. M., Shrivastava, J., Lamberton, P. H. L., Rollinson, D., Webster, B. L., Emery, A., et
al. (2007). Development and application of an ethically and epidemiologically
advantageous assay for the multi-locus microsatellite analysis of Schistosoma mansoni.
Parasitology 134, 523–536.

Howe, K. L., Bolt, B. J., Shafie, M., Kersey, P., and Berriman, M. (2017). WormBase ParaSite -
a comprehensive resource for helminth genomics. *Mol. Biochem. Parasitol.* 215, 2–10.

Howick, V. M., Russell, A., Andrews, T., Heaton, H., Reid, A. J., Natarajan, K. N., et al. (2019).
The Malaria Cell Atlas: a comprehensive reference of single parasite transcriptomes
across the complete Plasmodium life cycle. *bioRxiv*, 527556. doi:10.1101/527556.

Jaleta, T. G., Zhou, S., Bemm, F. M., Schär, F., Khieu, V., Muth, S., et al. (2017). Different but
overlapping populations of Strongyloides stercoralis in dogs and humans-Dogs as a
possible source for zoonotic strongyloidiasis. *PLoS Negl. Trop. Dis.* 11, e0005752.

Le Clec'h, W., Chevalier, F. D., McDew-White, M., Allan, F., Webster, B. L., Gouvras, A. N., et
al. (2018). Whole genome amplification and exome sequencing of archived schistosome
miracidia. *Parasitology* 145, 1739–1747.

Lee-Six, H., Ellis, P., Osborne, R. J., Sanders, M. A., Moore, L., Georgakopoulos, N., et al.
(2019). The landscape of somatic mutation in normal colorectal epithelial cells. *bioRxiv*.
doi:10.1101/416800.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. *arXiv [q-bio.GN]*. Available at: http://arxiv.org/abs/1303.3997.

Lok, J. B. (2007). Strongyloides stercoralis: a model for translational research on parasitic
nematode biology. *WormBook*, 1–18.

Maclean, M. J., Savadelis, M. D., Coates, R., Dzimianski, M. T., Jones, C., Benbow, C., et al.
(2017). Does evaluation of in vitro microfilarial motility reflect the resistance status of
Dirofilaria immitis isolates to macrocyclic lactones? *Parasit. Vectors* 10, 480.

Marek, M., Zouhar, M., Douda, O., Maňasová, M., and Ryšánek, P. (2014). Exploitation of
FTA cartridges for the sampling, long-term storage, and DNA-based analyses of plant-
parasitic nematodes. *Phytopathology* 104, 306–312.

Moore, L., Leongamornlert, D., Coorens, T. H. H., Sanders, M. A., Ellis, P., Dawson, K., et al.
(2018). The mutational landscape of normal human endometrial epithelium. *bioRxiv*.
doi:10.1101/505685.

Ngara, M., Palmkvist, M., Sagasser, S., Hjelmqvist, D., Björklund, Å. K., Wahlgren, M., et al.
(2018). Exploring parasite heterogeneity using single-cell RNA-seq reveals a gene
signature among sexual stage Plasmodium falciparum parasites. *Exp. Cell Res.* 371, 130–
138.

Platt, R. N., McDew-White, M., Le Clec'h, W., Chevalier, F. D., Allan, F., Emery, A. M., et al.
(2019). Ancient hybridization and introgression of an invadolysin gene in schistosome
parasites. *bioRxiv*. doi:10.1101/539353.

Reid, A. J., Talman, A. M., Bennett, H. M., Gomes, A. R., Sanders, M. J., Illingworth, C. J. R., et
al. (2018). Single-cell RNA-seq reveals hidden transcriptional variation in malaria
parasites. *Elife* 7. doi:10.7554/eLife.33105.

Sabina, J., and Leamon, J. H. (2015). Bias in Whole Genome Amplification: Causes and
Considerations. *Methods Mol. Biol.* 1347, 15–41.

Sanya, R. E., Nkurunungi, G., Hoek Spaans, R., Nampijja, M., O'Hara, G., Kizindo, R., et al.
(2018). The impact of intensive versus standard anthelminthic treatment on allergy-
related outcomes, helminth infection intensity and helminth-related morbidity in Lake
Victoria fishing communities, Uganda: results from the LaVIISWA cluster randomised

trial. *Clin. Infect. Dis.* doi:10.1093/cid/ciy761.

Shortt, J. A., Card, D. C., Schield, D. R., Liu, Y., Zhong, B., Castoe, T. A., et al. (2017). Whole
Genome Amplification and Reduced-Representation Genome Sequencing of
Schistosoma japonicum Miracidia. *PLoS Negl. Trop. Dis.* 11, e0005292.

Small, S. T., Labbé, F., Coulibaly, Y. I., Nutman, T. B., King, C. L., Serre, D., et al. (2018).
Human Migration and the Spread of the Nematode Parasite Wuchereria bancrofti.
bioRxiv. doi:10.1101/421248.

Sulston, J., and Hodgkin, J. (1988). "Methods," in *The Nematode Caenorhabditis elegans*, ed.
W. B. Wood (Cold Spring Harbor, NY, Cold Spring Harbor Laboratory Press).

Trevino, S. G., Nkhoma, S. C., Nair, S., Daniel, B. J., Moncada, K., Khoswe, S., et al. (2017).
High-Resolution Single-Cell Sequencing of Malaria Parasites. *Genome Biol. Evol.* 9,
3373–3383.

Tsai, I. J., Hunt, M., Holroyd, N., Huckvale, T., Berriman, M., and Kikuchi, T. (2014).
Summarizing specific profiles in Illumina sequencing from whole-genome amplified
DNA. *DNA Res.* 21, 243–254.

Wakelin, D. (1967). Acquired immunity to Trichuris muris in the albino laboratory mouse.
Parasitology 57, 515–524.

Webster, B. L. (2009). Isolation and preservation of schistosome eggs and larvae in
RNAlater® facilitates genetic profiling of individuals. *Parasit. Vectors* 2, 50.

Webster, B. L., Emery, A. M., Webster, J. P., Gouvras, A., Garba, A., Diaw, O., et al. (2012).
Genetic diversity within Schistosoma haematobium: DNA barcoding reveals two distinct
groups. *PLoS Negl. Trop. Dis.* 6, e1882.

Xiao, N., Remais, J. V., Brindley, P. J., Qiu, D.-C., Carlton, E. J., Li, R.-Z., et al. (2013).
Approaches to genotyping individual miracidia of Schistosoma japonicum. *Parasitol.*
Res. 112, 3991–3999.

655