Single-cell DNA and RNA sequencing reveals the dynamics of intra-tumor heterogeneity in a colorectal cancer model

AUTHORS

Hanako Ono[1], Yasuhito Arai[2], Eisaku Furukawa[1], Daichi Narushima[1], Tetsuya Matsuura[3], Hiromi Nakamura[2], Daisuke Shiokawa[4], Momoko Nagai[1], Toshio Imai[3], Koshi Mimori[5], Koji Okamoto[6], Yoshitaka Hippo[3,7], Tatsuhiro Shibata[2,8], and Mamoru Kato[1,*]

AFFILIATIONS

[1] Department of Bioinformatics, National Cancer Center Research Institute, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

[2] Division of Cancer Genomics, National Cancer Center Research Institute, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

[3] Department of Animal Experimentation, National Cancer Center Research Institute, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

[4] Division of Cancer Differentiation, National Cancer Center Research Institute, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

[5] Department of Surgery, Kyushu University Beppu Hospital, 101 Hasamamachiidaigaoka, Yufu, Oita 879-5593, Japan

[6] Division of Cancer Differentiation, National Cancer Center Research Institute, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

[7] Division of Biochemistry and Molecular Carcinogenesis, Chiba Cancer Center Research Institute, 666-2 Nitona-cho, Chiba Chuo-ku, Chiba 260-8717, Japan

[8] Laboratory of Molecular Medicine, Human Genome Center, The Institute of Medical Science, The University of Tokyo, 4-6-1 Shiroganedai, Minato-ku, Tokyo 108-8639, Japan

28 * Corresponding author

29 Tel: +81-3-3542-2511

30 E-mail: mamkato@ncc.go.jp

Abstract

Intra-tumor heterogeneity (ITH) encompasses cellular differences in tumors and is related to clinical

outcomes, such as drug resistance. However, little is known about the dynamics of ITH, owing to the lack

of time-series analysis at the single-cell level. We performed single-cell exome and transcriptome

sequencing of 200 cells and investigated how ITH is generated from one single cell in a mouse colorectal

cancer model. The ITH of the transcriptome increased after transplantation from cultured organoids, while

that of the exome decreased. Mutations generated in the culture did not greatly change at the

transplantation at the bulk-cell level. The RNA ITH increase was due to the emergence of new

transcriptional subpopulations. In contrast to the initial cells expressing mesenchymal-marker genes, new

subpopulations repressed these genes at transplantation. Analyses of colorectal cancer data from The

Cancer Genome Atlas revealed a high proportion of metastatic cases in human subjects with expression

patterns similar to the new cell subpopulations in mouse. These results suggest that the birth of

transcriptional subpopulations may be a key for adaptation to drastic micro-environmental changes when

cancer cells have sufficient genetic alterations at later tumor stages. This study revealed an evolutionary

dynamics of single-cell RNA and DNA changes in tumor progression, giving insights into the

mesenchymal-epithelial transformation of tumor cells at metastasis in colorectal cancer.

Keywords

Single-cell sequencing, cancer genomics, colorectal cancer, tumorigenesis, intra-tumor heterogeneity

Background

It is well established that cancer is pathologically composed of different types of cells [1]; however, intra-tumor heterogeneity (ITH) has only been recently addressed at the genomic level [2]. ITH is clinically important. For example, elevated copy-number heterogeneity is related to an increased risk of recurrence or death in non-small-cell lung cancer [3]. High levels of ITH ultimately provide the seeds for the emergence of anti-cancer drug resistance [4]. High levels of genetically-characterized heterogeneity in Barrett's esophagus are associated with incidence of esophageal adenocarcinoma [5].

ITH essentially stands for the cellular differences in tumor tissue arising from genetic changes, called clonal evolution, or non-genetic changes, such as cancer stem cells and simple transcriptional responses to the environment. In clonal evolution, as in Darwinian evolution, cancer cells with advantageous genetic mutations evolve into different types of cancer cells [6]. In contrast, cancer stem cells, like normal stem cells, produce a variety of differentiated daughter cells that constitute phenotypically distinct cancer cells without genetic differences through epigenetic and the resultant transcriptional mechanisms [7, 8].

A flood of studies have addressed ITH through the variant allele frequencies (VAFs) of tumor cells in bulk, which are calculated from sequence reads with variants identified through next-generation sequencing (reviewed in [2, 9]). In this bulk-cell sequencing approach, the presence of minor clones is often reflected on lower VAFs than the VAF of the major clone [10]. However, this bulk-cell DNA sequencing approach is limited in revealing genetic ITH because it only infers the presence of clones, not directly observing individual cells. In addition, the bulk-cell approach is generally not suitable to resolve transcriptomic ITH, where transcript mixtures from different cells are sequenced.

Single-cell sequencing is a powerful technology for investigating ITH by identifying genomic alterations and distinct transcriptomic states in single tumor cells [11-19]. For example, in clinical samples of glioblastoma, single-cell RNA sequencing showed that individual tumor cells vary in terms of their degree of stemness-related gene expression from extremely stem-like to differentiated states [13]. Additionally, the existence of cancer stem cells that continuously differentiate into astrocyte- and oligodendrocyte-like cells has been demonstrated in oligodendrogliomas by single-cell RNA sequencing

[14]. Single-cell DNA sequencing has also been applied to breast cancer samples to evaluate ITH

originating in genomic DNA, leading to the suggestion of stepwise/sweepstake or gradual evolution of

cancer cells from single nucleotide variation (SNV) data [11, 12, 20]. However, these types of ITH and

their respective evolutionary mechanisms are based on snapshot data at one time-point. Furthermore,

either RNA or DNA was solely examined. It is necessary to address both RNA and DNA over time for the

full elucidation of tumor evolutionary dynamics associated with ITH.

Mouse models are more useful than human clinical samples for examining changes in genomic

and transcriptomic states over time. In a breast tumor xenograft mouse model, single-cell DNA

sequencing of serially passaged samples identified tumor cell subpopulations and suggested that tumor

cells in the same initial state followed the same evolutionary trajectory [21]. In the present study, we

employed a modified version of the mouse colorectal cancer model that we previously established [22]

and sequenced both single-cell DNA and RNA. We thus investigated how ITH based on the exome and

transcriptome changes over time at the single-cell level.

Results

Colorectal cancer mouse model

The colorectal cancer mouse model was established by knocking down *APC* expression in normal

epithelial cells taken from mouse intestinal crypts using short hairpin RNA (shAPC; **Fig. 1A**) [22]. In the

previous system, we used bulk cells from a tissue for culture; however, in this study, we cultured

organoids from *one single cell* so that heterogeneity observed in these cultures could not be confused with

heterogeneity originating from the knock-down efficiency or intestinal crypts [23].

We grew organoids for a period of five months so that a single cell having only artificial *APC*

intervention could naturally obtain mutations to transform into tumor cells. Cultured cells were

subcutaneously transplanted into a nude mouse. One month after transplantation, the mouse was

sacrificed, and the tumor was harvested; half of the tumor tissue was re-cultured in our three-dimensional

(3D) culture system for one month for the removal of stromal cells. Using half-samples preserved the

same genetic lineage over time. The process was repeated once more. Cells were sampled immediately

before the first transplantation at time point T1 and at two time points T2 and T3 following the first and

second transplantations, respectively (**Fig. 1A**). We sequenced single-cell RNA and DNA separately taken

from the different single cells of multiple organoids, which descended from one single cell. We also

obtained a DNA sample before T1, at T0.5, 1.5 months after culture initiation (**Fig. 1A**).

Hematoxylin-eosin (HE) staining revealed that subcutaneously transplanted organoids formed

tumors consisting of both glandular and non-glandular structures (HE *a* and *b* in **Fig. 1A**). Glandular

components in HE *a* were mainly lined with single-layered epithelia, while those in HE *b* were

characterized by increased multi-layered regions, loss of cellular polarity, and nuclear enlargement. Non-

glandular components had a stromal/medullary structure consisting of spindle-shaped or round to

polygonal cells, were characteristically gelatinous/fibrous, and had an abundance of fibrous stroma.

The *APC* expression was decreased in the *APC* knockdown samples (**Fig. 1B**). Out of the 31

significantly mutated genes (excluding *TTN*) defined by The Cancer Genome Atlas (TCGA) colorectal

cancer study [24], we found two mutations in *KRAS* and *TP53* by bulk-cell DNA sequencing in our model

(**Fig. 1C**) , though the *KRAS* mutation was located outside of, but close (9 bps) to, an exon and the position

was evolutionary conserved as much as exons (Additional file 1: **Figure S1**). The *KRAS* mutation occupied

only a small fraction (2.5%) of the population at T0.5 but increased to 46.4% at T3. Additionally, we found

nonsynonymous mutations in six, *CLTC*, *LRP1B*, *ALK*, *GRIN2A*, *MSH2*, and *SALL4* out of the cancer-

related genes in COSMIC Gene Census [25] (**Fig. 1C**). It seems that a major clone with the *TP53* mutation

were extinct between T0.5 and T1 and instead minor clones with the *ALK* and *KRAS* mutations became

major clones.

Single-cell transcriptome analysis

We checked various indices of single-cell transcriptome data to filter 42, 42, and 51 cells out of the 50 T1,

43 T2, and 52 T3 cells, respectively (Additional file 1: **Figure S2**). The median (± inter-quartile range)

number of mapped reads, mapping rate, and number of expressed genes across selected cells were $6.2 \times$

10^6 ($\pm 2.0 \times 10^6$), 61.9% (± 5.39%), and 3814 (± 889.5), respectively. There was a strong correlation

between gene expression levels in the bulk sequencing data and average expression levels across single

cells (Additional file 1: **Figure S2**; $R^2 = 0.9$). The expression levels of housekeeping genes (GAPDH and

mtATP6) [26] were well reproduced across T1, T2, and T3 (Additional file 1: **Figure S2**).

A principal component analysis (PCA) plot of cells based on expression levels revealed increased

diversity from T1 to T2 (**Fig. 2A**). This was quantitatively confirmed by the diversity index (distance from

the centroid in the PCA space) (**Fig. 2B**). In the plot, T2 and T3 cells partly overlapped but were separate

from T1 cells. We identified genes whose expression levels varied greatly across cells at each time point;

that is, these genes had high corrected coefficient of variation (c*CV*) values (Additional file 1: **Figure S3**),

and were thus referred to as highly variable genes. There were eight, 14, and 16 highly variable genes at

T1, T2, and T3, respectively, reflecting an increase in variability from T1 to T2.

A cluster analysis of highly variable genes identified three gene groups (Additional file 1: **Figure**

S4); expression levels were correlated within two of the groups, but not within the third group. Gene set

enrichment analysis showed that one of the correlated groups was associated with negative regulation of

keratinocyte differentiation (referred to as Anti-Epithelial genes) ($P = 3.80 \times 10^{-3}$), whereas the other was

associated with positive regulation of cGMP and guanylate cyclase (GC) activity (referred to as cGMP/GC

genes) ($P = 1.30 \times 10^{-3}$), which are known to be associated with negative regulation of β-catenin signaling

and matrix metalloproteinase activity in colorectal cancer [27, 28].

A heatmap generated from the cluster analysis revealed that T1 cells were relatively homogenous

and formed one group that highly expressed Anti-Epithelial genes but showed negligible expression of

cGMP/GC genes (**Fig. 2C**). This group was therefore termed Anti-Epithelial. In addition to an Anti-

Epithelial cell group, two new groups appeared at T2: one showing the opposite pattern, repression of

Anti-Epithelial and activation of cGMP/GC gene expression, referred to as the cGMP/GC cell group; the

other showed repression of both Anti-Epithelial and cGMP/GC genes, referred to as the Dormant group

for the marker analysis described below. Notably, as shown in the heat map, bulk-cell sequencing analysis

alone could not have identified these cell groups, where their distinct expression patterns were offset by

bulk-cell expression levels (labeled as T1, T2, and T3 bulk in **Fig. 2C**).

Using a PCA plot based on highly variable gene expression, we confirmed that T1 cells were

relatively homogeneous and T2 cells showed similar grouping to T3 cells (**Fig. 2D**). Looking at the PCA

closer, we found that cells of the Anti-Epithelial group seemed close together across all time points, but

seemed to form two groups—i.e., T1 main (referred to as T1 main) and T1/T2/T3 mixture (T1+T2+T3).

The cGMP/GC and Dormant groups seemed close together and contained T2 and T3 only (T2+T3 only).

Marker gene expression

We examined the expression of several types of marker genes. We first looked at proliferation/cell cycle

markers (Additional file 1: **Figure S5**) and performed PCA to summarize the multiple expression levels

(**Fig. 3A**). Remarkably, most cells in the Anti-Epithelial group at T1 expressed high levels of proliferation-

and cell cycle-related genes according to the PCA loading plot. In contrast, nearly all cells in the Dormant

group at T3 showed a downregulation of the marker genes (so the cell group was termed Dormant). At T2,

about half of the cells showed a downregulation of the proliferation/cell-cycle genes.

We next examined epithelial and mesenchymal markers (Additional file 1: **Figure S5**). A PCA

plot of the markers showed that expression of mesenchymal cell-related genes decreased with time (T2

and T3), with cells forming two groups (**Fig. 3B**): one (upper left) overlapping with some T1 Anti-

Epithelial cells with decreased mesenchymal N-cadherin (*CDH2*) and fibronectin (*FN1*) levels; the other

(middle right) group was composed only of T3 cells with decreased mesenchymal vimentin (*VIM*) and

increased epithelial E-cadherin (*CDH1*) levels. These results suggest a similarity between the processes

occurring from T1 to T2 and mesenchymal-epithelial transition (MET).

Stem cell and differentiation markers showed that over time, cells generally expressed more

differentiation than stem cell markers (**Fig. 3C**; Additional file 1: **Figure S5**), though a remarkable

variation across individual cells was also observed. Fractions of cells with the expressions of stem cell

markers decreased with time. Among the markers for crypt base stem cells, *SOX9* appeared to be the most

influential; *LGR5*, *OLFM4*, and *MSI1* were not substantially expressed. It seems that with time, cells

differentiated into those expressing a marker for absorption cells (*KRT20*) and those for secretion cells

(*MUC2*) in the digestive tract.

There was no remarkable change in the expression of drug efflux genes [29, 30] at any time point

(**Fig. 2C**), although *ABCB1* expression was slightly lower in the T3 Dormant group (Additional file 1:

Figure S5) and *ABCE1* was downregulated at T2 and T3. There was variable expression of glycolysis-

related gene *PDK1* [30] across all cells, irrespective of groups (**Fig. 2C**; Additional file 1: **Figure S5**).

Single-cell exome analysis

Based on several indices from single-cell exome sequencing (Additional file 1: **Figure S6**), we selected

21, 23, and 23 cells out of the 23 T1, 24 T2, and 24 T3 cells for analysis. On average (expressed as the

median [± inter quartile range] across selected cells), the number of mapped reads was 1.2×10^8 (± $2.2 \times$

10^7), mapping rate was 76.6% (± 4.9%), coverage with > 0 depth regions was 76.9% (± 34.2%), average

depth was 43 (± 34.5), Gini coefficient was 0.85 (± 0.15), allelic drop-out (ADO) rate was 47.0 (± 36.1),

and number of called SNVs was 462 (± 313.5). The false positive rate in single-cell sequencing was

estimated to be $0.1–1.1 \times 10^{-7}$ per chromosomal site, based on normal intestinal tract tissue samples from

two mice and four single cells obtained from one of these samples (**Additional file 1: Supplementary**

Results). We compared the fractions of single cells with SNVs to the variant allele frequencies (VAFs) of

the bulk-cell sequencing; in theory, the single-cell fractions should be equal to half of the VAFs. We

confirmed a good concordance between these variables, although the cell fractions were slightly lower

than those expected from bulk VAFs (Additional file 1: **Figure S6**).

We first examined the bulk-cell sequence data. The T0.5 tissue had much fewer SNVs than the

later stages (**Fig. 4A**), which suggests that DNA heterogeneity only weakly appeared soon (1.5 months)

after culture initiation. The numbers of SNVs increased markedly from T0.5 to T1, a five-month period

(**Fig. 4A**). Although these numbers decreased slightly at T2 before recovering at T3, they were all mostly

saturated at T1, T2, and T3. Thus, new SNVs were largely generated from T0.5 to T1, and most of these

SNVs remained in the genome after T1 at the bulk-cell level (**Fig. 4B**).

We then used single-cell sequencing data to draw a multi-dimensional scaling (MDS) plot based

on single-cell SNVs at polymorphic SNV sites (defined as SNVs with 10–35% bulk VAFs) (**Fig. 4C**). T1

cells showed the greatest genetic diversity, whereas T2 and T3 cells showed less diversity. This decrease in

diversity was confirmed by a statistical significance of the diversity index (average distance from the

centroid), where the bias due to ADO rates was taken into account by a bootstrapping test (**Fig. 4D**).

Interestingly, this diversity tendency was the complete opposite of the transcriptomic pattern (**Fig. 2A**, **B**).

Although transitional, cells can be classified into three groups composed of T1 cells only (T1 main); T1,

T2, and T3 cells (T1+T2+T3); and T2 and T3 cells (T2+T3 only) (**Fig. 4C**), which could correspond to the

three groups in the single-cell transcriptome analysis (**Fig. 2D**)

Association with human cancer

We used 244 TCGA samples of colorectal cancer [24] to identify human samples with gene

expression patterns similar to the groups of mouse single cells by MDS (**Fig. 5A**). In the MDS plot, the

human samples were separated from the mouse cell groups, but we found that 94 (38.5%), 42 (17.2%),

and 13 (5.3%) human samples were respectively close to the Anti-Epithelial, cGMP/GC, and Dormant

mouse cell groups. TCGA Anti-Epithelial samples showed enhanced *REG* and repressed cGMP/GC gene

expression; TCGA cGMP/GC samples showed the opposite pattern; and TCGA Dormant samples had

both repressed *REG* and GC-related gene expression (**Fig. 5B**). TCGA cGMP/GC and TCGA Dormant

samples tended to be more closely associated with metastasis than those with patterns similar to the Anti-

Epithelial group (two-sided Fisher's exact test $P = 0.04$; **Fig. 5C**). We determined that our mouse tumor

cells were molecularly similar to cells of human colon adenocarcinoma classified as high microsatellite

instability type (Additional file 1: **Figure S7**; Additional file 1; **Figure S8**; Additional file 1:

Supplementary Results).

Discussion

Our mouse model was close to human colon adenocarcinoma of the MSI-high hyper-mutation type. In this

model, once cancer cells accumulate a sufficient number of genetic alterations (SNVs/indels), they may be

able to adapt to drastic environmental changes, such as the shift from a 3D culture to a live mouse, by only

altering their transcriptional profiles without further genetic changes. Such transcriptional adaptation may

cause the generation of new subpopulations, leading to increased transcriptional heterogeneity.

Meanwhile, genetic heterogeneity decreased, possibly as a result of microscale natural selection that

occurred during the environmental transition. Though expected, it is nonetheless surprising to see that this

diversity was indeed generated from *one single cell*.

T1 cells had the saturated number of genetic mutations, expressing active cell-cycle,

mesenchymal, and stem cell markers. Thus, the cells are considered as those at a late tumor stage when

they move out from the niches or microenvironment of intestinal crypts [31]. Moreover, the emergence of

the Dormant and cGMP/GC groups at T2 and T3 was associated with metastasis in the analysis using

TCGA human samples. Therefore, our observation that cells lose their mesenchymal-like phenotype and

acquire epithelial-like characteristics after subcutaneous transplantation may be analogized to MET during

metastasis, though this implication should be tested by further investigations of clinical samples including

single-cell sequencing of TCGA samples.

Classically, cells that generate a tumor by subcutaneous transplantation are called tumor-initiating

cells or cancer stem cells (CSCs) [30]. In this classical model, it is expected that differentiated cells die

while CSCs can survive at the start of subcutaneous transplantation and 3D culture; then, CSCs re-

generate differentiated cells. We initially expected that fractions of cells expressing stem cell markers

increased over the serial transplantation, because we simply thought that it is CSCs, not differentiated

cells, that can survive at transplantation. However, our observation was the opposite. CSCs that efficiently

generate differentiated cells may be more adaptive for the merit of obtaining mutual benefits between

different types of cells. Alternatively, contrary to the classical expectations, CSCs may not necessarily

express high levels of stem cell markers: the Dormant and cGMP/GC cells with low expression levels of

stem cell markers may be also CSCs or tumor-initiating cells that survived at the transplantation and 3D

culturing.

Recently, more fine-scale single-cell sequencing technology, such as 10X/Drop-Seq, has emerged

for RNA-seq, enabling researchers to capture tens of thousands of cells. Although the number of cells we

addressed was relatively small compared to that technology, we believe that we successfully captured a

major part of the heterogeneity constructed by cell clones, constituting as small as ~2% (an inverse

number of 42, 42, and 51 cells at T1, T2, and T3) of the tumor cell population. Nevertheless, 10X/Drop-

Seq will be needed to investigate rarer cells.

We demonstrated that time-series ITH analysis by single-cell DNA and RNA sequencing for a

mouse model is able to deepen our understanding of the evolutional processes of cancer cells and raise

issues on CSCs from the genomic and transcriptomic viewpoints. The birth of transcriptional

subpopulations of cells may be a key for adaptation to drastic micro-environmental changes when cancer

cells have sufficient genetic alterations at later tumor stages. It will be crucial to examine how such

genetic changes accumulate in the earlier stages of tumorigenesis and how transcriptional subpopulations

develop to increase malignancy in the further later stages of tumor progression.

Materials and Methods

Ethics approval and consent to practice

Animal studies were carried out according to the Guideline for Animal Experiments established by the

Committee for Ethics in Animal Experimentation of the National Cancer Center (T10-033-M05), which

meets the ethical standards required by law and guidelines for animal experimentation in Japan. All

sacrificed mice were anesthetized by inhalation of isoflurane. And cervical dislocation was used as a

euthanasia method.

Organoid culture of small intestinal cells and lentiviral transduction

C57BL/6J mice and BALB/cAnu/nu immune-deficient nude mice were purchased from CLEA Japan

(Tokyo, Japan). The small intestine was harvested from wild-type male C57BL/6J mice at 3–5 weeks of

age. Crypts were purified and dissociated into single cells, which were then put into serum-free Matrigel-

based organoid culture as previously described [22, 32]. Transduced organoids were maintained in culture

medium lacking R-spondin 1. Single cell-derived shAPC-transduced organoids were obtained by limiting

dilution of dissociated organoids in a 96-well plate. Organoids composed of 5×10^5 cells were mixed with

200 µl of Matrigel and injected into one side of the dorsal skin of nude mice, while uninjected cells were

maintained in 3D cultures for later use.

Analysis of subcutaneous tumors in nude mice

Palpable tumors from the injection sites were harvested for histological examination or cell culture. Half

of the subcutaneous tumors were formalin-fixed, paraffin-embedded, and sectioned at a thickness of 5 µm,

followed by HE staining to assess histological features. The other half of the tumors were minced and

digested to recover cells as described previously (22), then seeded onto solidified Matrigel to resume

organoid culture.

Single-cell transcriptome and exome sequencing

Cultured mouse organoids derived from a single cell were harvested and treated with Accumax
(Innovative Cell Technologies, AM105) to generate a single-cell suspension. To capture cells and extract
RNA or DNA from a single cell, the cell suspensions (4.4×10^5 cells/ml) were loaded on a C1 Single Cell
Auto Prep System (Fluidigm, C1) with medium-sized (10–17 μm) microfluidic chips for 96 cells.
Approximately 1300 cells were applied to each chip. Captured cells were imaged on a BZ-9000 digital
microscope (Keyence, BZ-9000) and a visual inspection was performed to determine whether a single cell
was captured in each well of the chip. Capture efficiency for a single cell was determined as 71–82%.

For single-cell whole transcriptome (RNA) sequencing, captured cells were lysed on the chip
using a C1 Single-Cell Auto Prep Reagent Kit for mRNA Seq (Fluidigm, 100-6201). Full-length cDNA
fragments were transcribed and amplified from poly-A RNA in each single cell using the SMARTer Ultra
Low RNA kit (Takara Bio, 634832). Tagmentation of cDNA was performed and sequencing libraries were
prepared using the Nextera XT DNA sample preparation kit (Illumina, FC-131-1096) according to the
manufacturer's protocol. Up to 52 independent single-cell RNA-seq libraries were prepared for
sequencing.

For single-cell DNA sequencing, genomic DNA was prepared from single cells using the C1
Single-cell Auto Prep Reagent Kit for DNA Seq (Fluidigm, 100-7357) and whole genome amplification
was achieved by multiple displacement amplification with Phi29 DNA polymerase and the Illustra
GenomiPhi v.2 kit (GE Healthcare, 25660032). Amplified genomic DNA (70 ng) was used to generate
exome sequence libraries using the Hyper Prep kit (Kapa Biosystems, KK8504), SureSelect Target
Enrichment kit (Agilent Technologies, 931171), and SureSelect XT Mouse All Exon v.1 probe (Agilent
Technologies, 5190-4642).

Bulk-cell transcriptome and exome sequencing

Among the cells that were not used for single-cell capture with the C1 system, suspensions of about 200
cells were subjected to whole transcriptome (RNA) sequencing for bulk-cell RNA-seq (T1, T2, and T3
samples). The sequencing libraries were prepared using the same reagents as the single cell RNA-seq. As
control bulk cells, normal intestinal crypt epithelial cells from two wild-type mice of the same strain were

grown in the 3D culture system for seven days, then harvested and lysed for total RNA preparation using

the miRNAeasy Mini kit (Qiagen, 217004). RNA-seq libraries for control bulk RNA were generated using

the SureSelect Strand-specific kit (Agilent Technologies, G9691B). Bulk DNA from over 1×10^5 cells was

obtained from the cell culture (T0.5 sample, 1.5 months after culture initiation) and the remaining cells in

single-cell capture (T1, T2, and T3 samples) using the QIAamp DNA Mini kit (Qiagen, 51304), and 500

ng of DNA were used to construct exome sequencing libraries with the same reagents as the single cell

DNA-seq.

Sequencing

All the sequencing libraries were subjected to paired-end sequencing of 101-bp fragments on the

HiSeq2500 DNA sequencer (Illumina, SY–401–2501).

Transcripts per kilobase million (TPM) calculation for single and bulk cells

The procedure for calculating TPM is summarized in Additional file 1: **Figure S9.** Specifically, sequence

reads were quality-filtered and trimmed using PrinSeq [33], and then used as input for quality-check by

FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). We used the following parameters:

--min_len 30 (removing reads ≤30 bases); --min_qual_mean 20 (average read quality ≤ 20); --

trim_tail_right 5, --trim_tail_left 5 (trim bases if the 3' and 5' end poly A/Ts are ≥ five bases); and --

trim_qual_right 20, --trim_qual_left 20 (trim 3' or 5' end for read quality ≤ 20). Paired-end reads were

mapped to the University of California Santa Cruz mouse genome sequence (mm10) using Bowtie2 [34]

built in RSEM [35]. Expression levels (in TPM) were estimated by RSEM using the command rsem-

calculate-expression with the parameters --estimate-rspd, --paired-end, --bowtie2, -p 30, and --ci-memory

10192. We removed eight T1 cell samples due to an excessive number of genes (≥ 5,200) with TPM ≥ 10

or with too few unique mapping reads ($< 2.2 \times 10^6$). We also removed two samples with unique mapping

rates that were too low (< 20%) and discarded genes with low expression levels (≤ 10 TPM) across all cell

samples, leaving 14,258 out of 32,732 genes for analysis.

Detection of highly variable genes

To detect genes with variable expression levels across cells, we defined highly variable genes according to

the *CV*, corrected in the locally weighted scatterplot smoothing (LOWESS) method using the "lowess"

function in R. To fit a single LOWESS curve across all ranges, we divided average expression level data

into three ranges: < 4, 4–8.5, and > 8.5. c*CV* values were yielded by dividing *CV* values by the value of

the upper variability band (± 1.96 times the standard deviation) of smoothed curve estimated using

"loess.sd" in the "msir" package. Because of the large bias in original *CV* values against low average

expression levels, only those with c*CV* values > 1.3 and high average expression levels ($\log_2 [\text{TPM+1}] \geq$

4) were defined as highly variable genes.

PCA of RNA data

PCA was carried out for gene expression levels ($\log_2 [\text{TPM} + 1]$) without scaling. For the loading analysis

of marker genes, we used the following genes; *MKI67* and *PCNA* for positive markers and *CDKN1A* for a

negative marker for cell proliferation in colorectal cancer [36]. *CCND2* and *CCND3* for positive markers

for cell cycle in this cancer [37]. E-cadherin (*CDH1*) for an epithelial marker; N-cadherin (*CDH2*),

vimentin (*VIM*), and fibronectin (*FN1*) for mesenchymal markers [38]. *LGR5*, *ASCL2*, *OLFM4*, *MSI1*, and

SOX9 for crypt base stem cell markers, *HOPX*, *BMI1*, and *LRIG1* for +4 (position from the crypt base)

stem cell markers, *AQP8*, *CAR1*, *CEACAM1*, *KRT20*, and *SLC26A3* for differentiation makers for

absorption cells, and *MUC2*, *SPINK1* for differentiation markers for secretion cells [39].

Hierarchal clustering, correlation plot, and heatmap analysis

For hierarchal clustering, we used the "hclust" function in the "stats" package of R software, where we

calculated the Euclidean distance of expression levels ($\log_2 [\text{TPM} +1]$) of all highly variable genes

between cells and used the Ward method for agglomeration. We generated correlation plots of highly

variable genes using the "corrplot" function in the R "corrplot" package, where we used the Ward method

for agglomeration. We divided genes into three clusters based on these hierarchical clustering results using

the "addrect = 3" option. A heatmap was generated using the "heatmap.2" function in the "ggplot2"

package. In the heatmap, cells were arranged according to their order in the dendrogram described above

and genes were arranged according to their order in the correlation plot of highly variable genes.

Gene set enrichment analysis

DAVID [40] was used to identify gene ontologies (biological processes) in which genes of an identified

group were enriched ($P < 0.01$).

SNV detection for single and bulk cells

For bulk-cell data, we used a previously described method for SNV/indel calling [41] by cisCall with cell-

line/frozen parameters [42], mapping reads to the mouse genome (mm9) by BWA [43]. We filtered out

PCR-duplicated reads as well as reads and bases with low mapping and base qualities. The remaining

variants were further filtered statistically using Fisher's exact test to compare fore- and background

samples, which came from two different individuals of the same pure C57BL/6J strain. We verified the

negligible effects of using a different individual for the background sample (**Supplementary Results**). A

series of filters was used to remove suspicious variant calls, such as those related to misalignments.

Variants for which allele frequencies were significantly greater than 1% in the binomial test were retained.

The procedure is summarized in Additional file 1: **Figure S9**.

For single-cell sequencing data, we called SNVs only at SNV sites called in bulk-cell sequencing

data. Specifically, we counted nucleotide bases with high qualities (mapQ $\geq$ 20, BaseQ $\geq$ 10) in single-cell

sequencing data as well as in background data (same as those used in bulk-cell SNV calling) with the

Samtools mpileup function [43, 44]. We then selected variants with the largest χ^2 test statistic (with Yates's

correction) among all possible variants at each position to identify those that were most likely to differ

between single-cell and background data. We required a variant count $\geq$ 2 and VAF $\geq$ 2% for such variants

in single-cell data. We then selected variants that overlapped with SNV sites called in bulk-cell data.

Detecting mutation in cancer-related genes

We investigated nonsynonymous mutations in cancer-related genes contained in Tier1 in COSMIC Cancer

Gene Census [25].

MDS of DNA data and the diversity index

We performed MDS from the cell × site matrix composed of zero and one, which respectively represent

the absence and presence of SNVs (both synonymous and non-synonymous SNVs) and NA, which

represents an undetermined call due to shallow depth. We assigned zero to non-called sites as the true

negative when those sites had depths $\geq$ 30 and assigned NA to non-called sites when the depth was $<$ 30.

We only used SNV sites where a variant was called in at least one cell and the VAFs at the same sites in

bulk data ranged from 10–35% (polymorphic) for at least one time point. We removed cells and sites (two

each) with too few or too many NAs, yielding 104 sites and 69 cells. Using this 0/1/NA matrix, we

calculated the p-distance (proportion of different sites) used in molecular evolution without using NA, and

then performed MDS.

The diversity index was calculated as the average Euclidian distance from the centroid over cells

in the MDS space, where we used up to the sixth dimension because of a sudden drop in the eigenvalues

over this dimension. To calculate the statistical significance of differences between cell groups, we used a

bootstrapping approach in which we randomly re-sampled cells' sequences from the 0/1/NA matrix of

each cell group 10,000 times and performed the same MDS as in the observed data for each replicate. We

then calculated the diversity index for each replicate to determine the 95% confidence interval and

standard deviation for each cell group.

Lorenz curve and Gini coefficients

433   A Lorenz curve was generated with read depth at each site using the "Lc" function in the "ineq" package

434   of R software. To quantify uniformity, the Gini coefficient was calculated using the "Gini" function in the

435   "ineq" package.

ADO rate

439 The ADO rate was defined as the rate of homozygous sites in single-cell samples where the bulk sample

440 was heterozygous (defined as sites where VAFs were 45–55%) at the same nucleotide site. We removed

441 outlier cells with high ADO rates at each time point (one cell each with an ADO rate > 80% at T2 and T3).

443 **Average copy number**

444 The average copy number (ACN) was calculated as follows:

445
$$\text{ACN} = 2 \times \left\{ \left(2^{\frac{\sum (\log_2 R_i \times L_i)}{\sum L_i}} \right) \times \left(\frac{\sum L_i}{GL} \right) + \left(1 - \frac{\sum L_i}{GL} \right) \right\} , \qquad (1),$$

where $\log_2 R_i$, L_i, and GL represent the log-ratio of CNA segment i, length of CNA segment i, and genome

length (50 Gb for mouse, 40 Gb for human), respectively. CNAs of mouse bulk data were detected as

previously described [41]. Briefly, segments were called for the same fore- and background BAM files as

those used in SNV with Exome CNV [45] and Varscan2 [46]. Overlapping segments called by both tools

were used as CNA segments.

Random Forest

Random Forest was used to generate the classifier for the histological type and MSI status of human

cancer. We used gene expression levels, number of SNVs in each gene, total mutation (SNV/indel)

number, and mutation density (total number of SNVs/indels divided by target region size) as explanatory

variables. Using TCGA data [24], we first filtered out unimportant explanatory variables using the two-

sided Kruskal-Wallis test with P values of 5.00×10^{-5} and 1.00×10^{-9} yielding 171 and 78 variables for

histological type and MSI status, respectively. These were used to train a Random Forest classifier with

the "randomForest" function in the "randomForest" package of R software, with the options ntree = 10000

(setting the number of trees to grow to 1000) and mtry = 5 (setting the number of variables randomly

sampled to five). Using the created classifier, the same explanatory variables for mouse data were used to

classify each feature in the mouse model.

MDS of mouse cell and TCGA samples

We first identified TCGA samples with gene expression patterns similar to the mouse single-cell groups.

For that purpose, we calculated a normalized 1-*r* distance as follows:

$$d_{h,G} = \frac{1 - r_{h,m^G}}{\text{MADN}(1 - r_{m_i^G, m^G})} \quad , \tag{2}$$

where $r_{i,j}$ is a Pearson correlation coefficient between vectors *i* and *j* of expression levels in log across

highly variable genes, *h* represents a human TCGA sample, *G* represents a mouse single-cell group, m_i^G

represents mouse single cell *i* in group *G*, m^G represents the centroid of m_i^G that was calculated by the

median, and MADN represents the median absolute deviation adjusted by a factor for asymptotically

normal consistency. We calculated this distance from a TCGA sample to every mouse group and selected a

TCGA sample for those whose minimum distance across the groups was less than 4.05 and the difference

between the first and second minimum distances was larger than 0.31. For selected TCGA and all mouse

single-cell samples, MDS was performed based on the distance of 1-*r*.

List of abbreviations

ITH: intra-tumor heterogeneity; PCA: principal component analysis; VAFs: variant allele frequencies;

MDS: multi-dimensional scaling; SNV: single nucleotide variation; TPM: transcripts per kilobase million;

LOWESS: locally weighted scatterplot smoothing; ACN: average copy number.

Declarations

Ethics approval and consent to practice

Animal studies were carried out according to the Guideline for Animal Experiments established by the Committee for Ethics in Animal Experimentation of the National Cancer Center (T10-033-M05), which meets the ethical standards required by law and guidelines for animal experimentation in Japan.

Consent for publication

Not applicable.

Availability of data and materials

Sequence data used in this study are available in the DDBJ Sequenced Read Archive under Accession Nos. DRX100507-DRX100729. [These data are held until the acceptance. URL will be added after the data release.]

Competing interests

None to be declared.

Funding

This work was supported by MEXT (25134721 and 25830141 to M.K., and 15K06916 to Y.A., Y.H., and M.K.); AMED (16ck0106013h0003 to T.S. and M.K., and 16ck0106115h0003 to K.O. and M.K.); National Cancer Center Research and Development Funds (29-A-6 to H.O. and T.S.); the Foundation for the Promotion of Cancer Research (to M.K.); and the Kato Memorial Bioscience Foundation (to M.K.).

Authors' contributions

Y.A., Y.H., T.S., and M.K. designed the study. Y.A., T.M., T.I., and Y.H. performed the experiments. H.O., E.F., D.N., H.N., M.N., and M.K. analyzed the data. H.O., Y.A., E.F., T.I., Y.H., and M.K. wrote the manuscript. D. S., K.M., K.O., and T.S. reviewed the manuscript. M.K. led the project.

Acknowledgements

We thank Joe Miyamoto, Asmaa Elzawahry, Iku Orihashi, Masako Ochiai, and Wakako Mukai for technical assistance, and Ryuichi Sugino and Daniel A. Vasco for useful suggestions.

References

1.   Almendro V, Marusyk A, Polyak K: **Cellular heterogeneity and molecular evolution in cancer**.

Annu Rev Pathol 2013, **8**:277-302.

2.   McGranahan N, Swanton C: **Clonal Heterogeneity and Tumor Evolution: Past, Present, and**

the Future. *Cell* 2017, **168**(4):613-628.

3.   Jamal-Hanjani M, Wilson GA, McGranahan N, Birkbak NJ, Watkins TBK, Veeriah S, Shafi S,

Johnson DH, Mitter R, Rosenthal R *et al*: **Tracking the Evolution of Non-Small-Cell Lung**

Cancer. *N Engl J Med* 2017, **376**(22):2109-2121.

4.   Dagogo-Jack I, Shaw AT: **Tumour heterogeneity and resistance to cancer therapies**. *Nat Rev*

Clin Oncol 2018, **15**(2):81-94.

5.   Maley CC, Galipeau PC, Finley JC, Wongsurawat VJ, Li X, Sanchez CA, Paulson TG, Blount PL,

Risques RA, Rabinovitch PS *et al*: **Genetic clonal diversity predicts progression to esophageal**

adenocarcinoma. *Nat Genet* 2006, **38**(4):468-473.

6.   Nowell PC: **The clonal evolution of tumor cell populations**. *Science* 1976, **194**(4260):23-28.

7.   Fulawka L, Donizy P, Halon A: **Cancer stem cells--the current status of an old concept:**

literature review and clinical approaches. *Biol Res* 2014, **47**:66.

8.   Kreso A, Dick JE: **Evolution of the cancer stem cell model**. *Cell Stem Cell* 2014, **14**(3):275-291.

9.   Davis A, Gao R, Navin N: **Tumor evolution: Linear, branching, neutral or punctuated?**

Biochim Biophys Acta Rev Cancer 2017, **1867**(2):151-161.

10.   Williams MJ, Werner B, Heide T, Curtis C, Barnes CP, Sottoriva A, Graham TA: **Quantification**

of subclonal selection in cancer from bulk sequencing data. *Nat Genet* 2018, **50**(6):895-903.

11.   Navin N, Kendall J, Troge J, Andrews P, Rodgers L, McIndoo J, Cook K, Stepansky A, Levy D,

Esposito D *et al*: **Tumour evolution inferred by single-cell sequencing**. *Nature* 2011,

472(7341):90-94.

12.   Wang Y, Waters J, Leung ML, Unruh A, Roh W, Shi X, Chen K, Scheet P, Vattathil S, Liang H *et*

al: **Clonal evolution in breast cancer revealed by single nucleus genome sequencing**. *Nature*

2014, **512**(7513):155-160.

13.   Patel AP, Tirosh I, Trombetta JJ, Shalek AK, Gillespie SM, Wakimoto H, Cahill DP, Nahed BV,

Curry WT, Martuza RL *et al*: **Single-cell RNA-seq highlights intratumoral heterogeneity in**

primary glioblastoma. *Science* 2014, **344**(6190):1396-1401.

14.   Tirosh I, Venteicher AS, Hebert C, Escalante LE, Patel AP, Yizhak K, Fisher JM, Rodman C,

Mount C, Filbin MG *et al*: **Single-cell RNA-seq supports a developmental hierarchy in human**

oligodendroglioma. *Nature* 2016, **539**(7628):309-313.

15.   Suzuki A, Matsushima K, Makinoshima H, Sugano S, Kohno T, Tsuchihara K, Suzuki Y: **Single-**

cell analysis of lung adenocarcinoma cell lines reveals diverse expression patterns of

individual cells invoked by a molecular target drug treatment. *Genome Biol* 2015, **16**:66.

16.   Gawad C, Koh W, Quake SR: **Dissecting the clonal origins of childhood acute lymphoblastic**

leukemia by single-cell genomics. *Proc Natl Acad Sci U S A* 2014, **111**(50):17947-17952.

17.   Kim KT, Lee HW, Lee HO, Kim SC, Seo YJ, Chung W, Eum HH, Nam DH, Kim J, Joo KM *et al*:

Single-cell mRNA sequencing identifies subclonal heterogeneity in anti-cancer drug

responses of lung adenocarcinoma cells. *Genome Biol* 2015, **16**:127.

18.   Hou Y, Song L, Zhu P, Zhang B, Tao Y, Xu X, Li F, Wu K, Liang J, Shao D *et al*: **Single-cell**

exome sequencing and monoclonal evolution of a JAK2-negative myeloproliferative

neoplasm. *Cell* 2012, **148**(5):873-885.

19.   Xu X, Hou Y, Yin X, Bao L, Tang A, Song L, Li F, Tsang S, Wu K, Wu H *et al*: **Single-cell exome**

sequencing reveals single-nucleotide mutation characteristics of a kidney tumor. *Cell* 2012,

148(5):886-895.

20.   Kato M, Vasco DA, Sugino R, Narushima D, Krasnitz A: **Sweepstake evolution revealed by**

population-genetic analysis of copy-number alterations in single genomes of breast cancer. *R*

Soc Open Sci 2017, **4**(9):171060.

21.   Eirew P, Steif A, Khattra J, Ha G, Yap D, Farahani H, Gelmon K, Chia S, Mar C, Wan A *et al*:

Dynamics of genomic clones in breast cancer patient xenografts at single-cell resolution.

Nature 2015, **518**(7539):422-426.

22.   Onuma K, Ochiai M, Orihashi K, Takahashi M, Imai T, Nakagama H, Hippo Y: **Genetic**

reconstitution of tumorigenesis in primary intestinal cells. *Proc Natl Acad Sci U S A* 2013,

110(27):11127-11132.

23.   Itzkovitz S, Blat IC, Jacks T, Clevers H, van Oudenaarden A: **Optimality in the development of**

intestinal crypts. *Cell* 2012, **148**(3):608-619.

24.   Cancer Genome Atlas N: **Comprehensive molecular characterization of human colon and**

rectal cancer. *Nature* 2012, **487**(7407):330-337.

25.   Futreal PA, Coin L, Marshall M, Down T, Hubbard T, Wooster R, Rahman N, Stratton MR: **A**

census of human cancer genes. *Nature Reviews Cancer* 2004, **4**:177.

26.   Janssens N, Janicot M, Perera T, Bakker A: **Housekeeping genes as internal standards in**

cancer research. *Mol Diagn* 2004, **8**(2):107-113.

27.   Fajardo AM, Piazza GA, Tinsley HN: **The role of cyclic nucleotide signaling pathways in**

cancer: targets for prevention and treatment. *Cancers (Basel)* 2014, **6**(1):436-458.

28.   Lubbe WJ, Zhou ZY, Fu W, Zuzga D, Schulz S, Fridman R, Muschel RJ, Waldman SA, Pitari

GM: **Tumor epithelial cell matrix metalloproteinase 9 is a target for antimetastatic therapy**

in colorectal cancer. *Clin Cancer Res* 2006, **12**(6):1876-1882.

29.   Li CJ, Zhang X, Fan GW: **Updates in colorectal cancer stem cell research**. *J Cancer Res Ther*

2014, **10 Suppl**:233-239.

30.   Qureshi-Baig K, Ullmann P, Haan S, Letellier E: **Tumor-Initiating Cells: a criTICal review of**

isolation approaches and new challenges in targeting strategies. *Mol Cancer* 2017, **16**(1):40.

31.   Batlle E, Clevers H: **Cancer stem cells revisited**. *Nat Med* 2017, **23**(10):1124-1134.

32.   Maru Y, Orihashi K, Hippo Y: **Lentivirus-Based Stable Gene Delivery into Intestinal**

Organoids. *Methods Mol Biol* 2016, **1422**:13-21.

33.   Schmieder R, Edwards R: **Quality control and preprocessing of metagenomic datasets**.

Bioinformatics 2011, **27**(6):863-864.

34.   Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2**. *Nat Methods* 2012,

9(4):357-359.

35.   Li B, Dewey CN: **RSEM: accurate transcript quantification from RNA-Seq data with or**

without a reference genome. *BMC Bioinformatics* 2011, **12**:323.

36.   Whitfield ML, George LK, Grant GD, Perou CM: **Common markers of proliferation**. *Nat Rev*

Cancer 2006, **6**(2):99-106.

37.   Tominaga O, Nita ME, Nagawa H, Fujii S, Tsuruo T, Muto T: **Expressions of cell cycle**

regulators in human colorectal cancer cell lines. *Jpn J Cancer Res* 1997, **88**(9):855-860.

38.   Weinberg R: **The biology of cancer**: Garland science; 2013.

39.   Hong SN, Dunn JC, Stelzner M, Martin MG: **Concise Review: The Potential Use of Intestinal**

Stem Cells to Treat Patients with Intestinal Failure. *Stem Cells Transl Med* 2017, **6**(2):666-676.

40.   Huang da W, Sherman BT, Lempicki RA: **Bioinformatics enrichment tools: paths toward the**

comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 2009, **37**(1):1-13.

41.   Takahashi T, Elzawahry A, Mimaki S, Furukawa E, Nakatsuka R, Nakamura H, Nishigaki T,

Serada S, Naka T, Hirota S *et al*: **Genomic and transcriptomic analysis of imatinib resistance**

in gastrointestinal stromal tumors. *Genes Chromosomes Cancer* 2017, **56**(4):303-313.

42.   Kato M, Nakamura H, Nagai M, Kubo T, Elzawahry A, Totoki Y, Tanabe Y, Furukawa E,

Miyamoto J, Sakamoto H *et al*: **A computational tool to detect DNA alterations tailored to**

formalin-fixed paraffin-embedded samples in cancer clinical sequencing. *Genome Med* 2018,

10(1):44.

43.   Li H, Durbin R: **Fast and accurate short read alignment with Burrows-Wheeler transform**.

Bioinformatics 2009, **25**(14):1754-1760.

44.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,

Genome Project Data Processing S: **The Sequence Alignment/Map format and SAMtools**.

Bioinformatics 2009, **25**(16):2078-2079.

45.   Sathirapongsasuti JF, Lee H, Horst BA, Brunner G, Cochran AJ, Binder S, Quackenbush J, Nelson

SF: **Exome sequencing-based copy-number variation and loss of heterozygosity detection:**

ExomeCNV. *Bioinformatics* 2011, **27**(19):2648-2654.

46.   Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding L,

Wilson RK: **VarScan 2: somatic mutation and copy number alteration discovery in cancer by**

exome sequencing. *Genome Res* 2012, **22**(3):568-576.

 Figure Legends

Fig. 1. The mouse model. (A) The experimental procedure and HE staining of subcutaneously transplanted

tumors. *One single cell* was 3D-cultured in a compartment in a 96-well plate, and single-cell derived

organoids were taken to separate single cells. RNA and DNA were separately extracted from the different

single cells of multiple organoids and then sequenced. The numbers of cells for RNA and DNA sequencing

in boxes are those obtained after quality control of data. (B) The *APC* gene expression from bulk- and single-

cell RNA sequencing. "APC_ctrl" indicates control samples that were cultured in our 3D culture system and

derived from normal cells without *APC* knockdown at bulk-cell sequencing. "SC average" indicates the

average across single cells at each time point. (C) Variant allele frequencies of mutations found in the

significantly mutated genes of colorectal cancer by bulk-cell DNA sequencing. See Additional file 1: **Figure**

S1 for the annotations of the mutations.

Fig. 2. Transcriptome analysis. (A) PCA plot of single cells based on expression levels (genes with TPM $\geq$

10 in at least one cell). T1, at the time of 3D culturing; T2 and T3, after the first and second

transplantations, respectively. (B) Euclidean distance from the centroid in the PCA space (using full

dimensions). *$P < 0.01$ (two-sided Wilcoxon rank sum test). (C) Heatmap of gene expression levels (in

TPM). The rows represent single cells or bulk-cell samples (in the bottom), and the columns represent

highly variable genes and several types of marker genes. The cell and gene groups were determined as

shown in Additional file 1: Figure S4. The red, blue, and green codes in the rows correspond to T1, T2,

and T3. "Diff." and "Prol./cell-cyc." represents differentiation and proliferation/cell cycle. "APC_ctrl"

indicates control samples that were cultured in our 3D culture system and derived from normal cells

without *APC* knockdown. (D) PCA plot of cells grouped based on expression levels of highly variable

genes.

Fig. 3. PCA and overlaid loading plots based on expression levels of markers (A) about the

proliferation/cell cycle. The arrow indicates the direction from negative to positive markers in the loading

plot; cells positioned in that direction in the PCA plot had higher expression levels of positive marker

genes. (B) About the epithelial and mesenchymal. The arrows along the *x* and *y* axes represent projected

loadings in the loading analysis, where cells positioned in that direction in the PCA plot had higher marker

gene expression levels. (C) About stem cell and differentiation.

Fig. 4. Exome analysis. (A) Number of SNVs called in bulk-cell sequencing. (B) Comparison of VAFs of

SNVs called in bulk-cell sequencing at successive time points. One point indicates one SNV. Numbers

represent the number of points. (C) MDS plot based on single-cell exome sequencing. "No SNV" and "All

SNV" represent sequences with no SNVs and with SNVs at all sites, respectively, which were artificially

generated as a reference. Error bars represent the standard deviation for each dimension calculated with a

bootstrapping approach that took into account ADO rates. (D) Median Euclidean distance from the

centroid over cells in the MDS space. The black and red bars represent the observed value and 95%

confidence interval calculated with the bootstrapping approach. $*P < 0.05$ (bootstrapping test).

Fig. 5. Analysis of TCGA human samples with gene expression patterns similar to mouse cell groups. (A)

MDS plot of mouse single-cell samples and such TCGA samples on the basis of a similarity of gene

expression patterns. (B) Heatmap of the samples. Genes are highly variable genes shown in Fig. 2C. (C)

The fraction of patients with metastatic tumor in TCGA samples with expression patterns similar to mouse

cell groups

A

Intestinal crypts → Normal cells

shAPC → Lentivirus

C57BL/6J mouse

- shAPC transduced cells
- Diluted to <1 single cell per well

3D culture

A single cell grows into organoids in 3D culture

In 3D culture, Stromal cells are removed

Graft

Tumor develops in nude mouse 1

Tumor develops in nude mouse 2

HE *a*

HE *b*

■ Organoid
➢ For RNA
  • 42 cells
  • Bulk cells
➢ For DNA
  • 21 cells
  • Bulk cells

■ Organoid
➢ For RNA
  • 42 cells
  • Bulk cells
➢ For DNA
  • 23 cells
  • Bulk cells

■ Organoid
➢ For RNA
  • 51 cells
  • Bulk cells
➢ For DNA
  • 23 cells
  • Bulk cells

0   1.5   5   6   7   8   9   months
     T0.5   **T1**   **T2**   **T3**

B

APC expression

log2(TPM + 1)

APC ctrl1, APC ctrl2, T1 bulk, T1 SC average, T2 bulk, T2 SC average, T3 bulk, T3 SC average

C

Variant allele frequency (%)

KRAS — TP53
CLTC — ALK
LRP1B — GRIN2A
MSH2 — SALL4

T0.5   T1   T2   T3

A

Legend:
- ○ Anti-epi, (mouse)
- □ cGMP/GC (mouse)
- △ Dormant (mouse)
- + Anti-epi, (TCGA)
- × cGMP/GC (TCGA)
- ◇ Dormant (TCGA)

Dim2 (vertical axis): 0.6, 0.4, 0.2, 0.0, -0.2, -0.4
Dim1 (horizontal axis): -0.5, 0.0, 0.5

B

Anti-epi.   cGMP/GC   Non-correlation

Anti-epi.

cGMP/GC

Dormant

Mouse Cells

Anti-epi.

cGMP/GC

Dormant

TCGA human samples

Gene labels:
24166_REG1B, 24167_REG3A, 24162_REG3G, 24162_REG3A, 1888_C6orf58, 25176_MGST1, 192_COX5B, 5714_ID2, 15659_FAM129B, 16937_B2M, 14531_MS4A8, 21016_RNF186, 20627_GUCA2B, 19299_CLCA1, 20626_GUCA2A, 24747_CD9, 17747_WFDC2, 18672_S100A6, 11650_MPC1, 19200_OSTC, 28743_COPE, 30963_GSTA1, 30963_GSTA5, 1741_SDC4, 27788_ALDOA, 18442_SPTSSB, 19128_F3

C

Proportion of patients with metastatic cancer

Patient proportion (%): 0, 10, 20, 30, 40

- TCGA Anti-Epithelial: 11/95
- TCGA cGMP/GC: 9/41
- TCGA Dormant: 4/10