An alanine permease mutation or exposure to D-cycloserine increases the susceptibility of

MRSA to β-lactam antibiotics

Laura A. Gallagher[1], Rebecca K. Shears[2], Claire Fingleton[1], Laura Alvarez[3], Elaine M.

Waters[1,2], Jenny Clarke[2], Laura Bricio-Moreno[2], Christopher Campbell[1], Akhilesh K. Yadav[3],

Fareha Razvi[4], Eoghan O'Neill[5], Alex J. O'Neill[6], Felipe Cava[3], Paul D. Fey[4], Aras Kadioglu[2] and

James P. O'Gara[1]*

[1]School of Natural Sciences, National University of Ireland, Galway, Ireland.

[2]Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and
Global Health, University of Liverpool, UK.

[3]MIMS-Molecular Infection Medicine Sweden, Molecular Biology Department, Umeå
University, Umeå, Sweden.

[4]Department of Pathology and Microbiology, University of Nebraska Medical Center,
Omaha, Nebraska, USA.

[5]Department of Clinical Microbiology, Royal College of Surgeons in Ireland, Connolly
Hospital, Dublin 15, Ireland.

[6]Antimicrobial Research Centre, School of Molecular and Cellular Biology, Faculty of
Biological Sciences, University of Leeds, Leeds, UK.

*Correspondence: jamesp.ogara@nuigalway.ie

Running title: Re-sensitization of AMR pathogens to β-lactams

26
27

Abstract. Prolonging the clinical effectiveness of β-lactams, which remain first-line antibiotics for many infections, is an important part of efforts to address antimicrobial resistance. We report here that inactivation of the predicted D-cycloserine (DCS) transporter gene *cycA* re-sensitizes MRSA to β-lactam antibiotics. The *cycA* mutation also resulted in hyper-susceptibility to DCS, an alanine analogue antibiotic that inhibits the alanine racemase and D-alanine ligase enzymes required for D-alanine incorporation into peptidoglycan. Amino acid transport studies showed that CycA functions as an alanine permease, and that impaired alanine transport in the *cycA* mutant correlated with increased susceptibility to oxacillin and DCS. The *cycA* mutation was accompanied by increased accumulation of muropeptides with tripeptide stems lacking the terminal D-ala-D-ala, and reduced peptidoglycan cross linking. Exposure of MRSA to DCS was also associated with a dose-dependent accumulation of muropeptides with tripeptide stems and reduced PG crosslinking. Because impaired alanine transport or DCS-treatment have similar effects on PG structure, synergism between β-lactams and DCS was investigated. DCS re-sensitised MRSA to β-lactam antibiotics *in vitro* and significantly enhanced MRSA eradication by oxacillin in a mouse bacteraemia model. The susceptibility of several other ESKAPE group pathogens to β-lactams was also increased by DCS. These data reveal that impaired uptake of alanine or DCS-meditated inhibition of D-alanine incorporation into peptidoglycan increases the susceptibility of MRSA to β-lactam antibiotics.

Importance. Treatment options for infections caused by ESKAPE group pathogens continue to dwindle with increasing antimicrobial resistance. Particularly important are the β-lactams, which remain first line antibiotics for many infections, and finding new ways to maintain their clinical effectiveness is a priority. Here we report that impaired alanine transport in an MRSA *cycA* mutant was associated with hyper-susceptibility to β-lactam antibiotics and D-cycloserine (DCS). DCS, which is used in the treatment of multi-drug resistant tuberculosis infections, re-sensitised MRSA to oxacillin and other β-lactams *in vitro* and significantly enhanced oxacillin activity in a mouse model of bacteraemia. DCS also sensitised other ESKAPE pathogens to β-lactams. These findings reveal important roles for alanine transport and the pathway for D-alanine incorporation into peptidoglycan in controlling the susceptibility of MRSA to β-lactam antibiotics.

Introduction

Whilst many bacteria can exhibit resistance to select antimicrobials, isolates of the human pathogen *Staphylococcus aureus* can express resistance to all licensed anti-staphylococcal drugs. This results in significant morbidity and mortality, with up to 20% of patients with systemic methicillin resistant *S. aureus* (MRSA) infections dying, despite receiving treatment with anti-staphylococcal drugs (1). Although aggressive hospital infection prevention and control initiatives appear to be having a positive impact on hospital-acquired MRSA rates in developed countries, *S. aureus* infections caused by community-associated MRSA strains and strains that are currently methicillin susceptible remain stubbornly high (2),(3).

As part of our efforts to identify improved therapeutic approaches for MRSA infections, we recently described the novel use of β-lactam antibiotics to attenuate the virulence of MRSA-induced invasive pneumonia and sepsis (4). We demonstrated that oxacillin-induced repression of the Agr quorum-sensing system and altered cell wall architecture resulted in downregulated toxin production and increased MRSA killing by phagocytic cells, respectively (4). Supporting this *in vitro* data, a randomised controlled trial involving 60 patients showed that the β-lactam antibiotic flucloxacillin in combination with vancomycin shortened the duration of MRSA bacteraemia from 3 days to 1.9 days (5, 6).

Because expression of methicillin resistance in *S. aureus* impacts fitness and virulence and is a regulated phenotype, further therapeutic interventions may also be possible. The complexity of the methicillin resistance phenotype is evident among clinical isolates of MRSA, which express either low-level, heterogeneous (HeR) or homogeneous, high-level methicillin resistance (HoR) (7-9). Exposure of HeR isolates to β-lactam antibiotics induces expression of *mecA,* which encodes the alternative penicillin binding protein 2a (PBP2a) and can select for mutations in auxiliary genes resulting in a HoR phenotype, including mutations that affect the stringent response and c-di-AMP signalling (10-14). Because auxiliary genes can influence the expression of methicillin resistance in MRSA, targeting the pathways associated with such genes may identify new ways to increase the susceptibility of MRSA to β-lactams. To pursue this, we performed a forward genetic screen to identify loci that impact the expression of resistance to β-lactam antibiotics in MRSA. Using the Nebraska Transposon Mutant Library, which comprises 1,952 sequence-defined transposon insertion mutants (15), inactivation of an amino acid permease gene was found to reduce resistance to cefoxitin, the β-lactam drug recommended by the Clinical and Laboratory Standards

Institute for measuring *mecA*-mediated methicillin resistance in MRSA isolates. Impaired alanine transport in the permease mutant also correlated with hyper-susceptibility to D-cycloserine (DCS), an alanine analogue antibiotic that interferes with two enzymatic steps in the D-alanine pathway of peptidoglycan biosynthesis. DCS exposure was shown to increase the susceptibility of MRSA and other ESKAPE pathogens to β-lactam antibiotics. Peptidoglycan analysis revealed that the *cycA* mutation or exposure to DCS results in loss of the terminal D-ala-D-ala from the stem peptide and reduced cross-linking of the cell wall. Given that β-lactam antibiotics remain among the most effective, safe and affordable antibiotics, DCS deserves further consideration as a potential agent to restore the therapeutic efficacy of β-lactam antibiotics.

Results

Mutation of *cycA* increases the susceptibility of MRSA to β-lactam antibiotics. To identify new ways of controlling expression of methicillin resistance, we sought to identify novel mutations involved in this phenotype. To achieve this, an unbiased screen of the Nebraska Transposon Mutant Library (NTML) was performed using a disk diffusion assay to identify mutants with increased susceptibility to cefoxitin. The parent strain of the library (JE2, a derivative of USA300 LAC cured of plasmids p01 and p03 (15)) was used as a positive control. Phage 80α-mediated backcross of transposon mutations into the parent strain JE2, together with genetic complementation were used to confirm the role of putative genes involved in cefoxitin resistance. Only mutants verified by both transduction and complementation were examined further. Among the mutants identified was NE810 (SAUSA300_1642) (Fig. 1A), which exhibited significantly increased susceptibility to cefoxitin as determined by a disk diffusion assay (Fig. S1A) and a >500-fold increase in susceptibility to oxacillin as determined by E-test (Fig. 1B). Consistent with this, NE810 was previously reported to be one of several NTML mutants shown to be more susceptible to amoxicillin (16), but was not investigated further. RT-qPCR analysis revealed that expression of *mecA* was not significantly affected in NE810 compared to JE2 (Fig. S1B), whilst whole genome sequence analysis of NE810 further revealed that the SCC*mec* element was fully intact and that no other mutations were present (data not shown). These findings implied that increased cefoxitin susceptibility was not attributable to a failure of this strain to express a functional *mecA* gene product or disruption/mutation of an auxiliary gene essential for expression of resistance. Transduction of the *cycA* transposon insertion into several MRSA strains from a number of clonal complexes and with different SCC*mec* types (USA300 strains JE2, FPR3757 and LAC-13C, DAR173, DAR22 and DAR169 (17)) was also accompanied by significant increases in cefoxitin and oxacillin susceptibility (Table 1).

Mutation of *cycA* increases susceptibility to ᴅ-cycloserine. SAUSA300_1642 is annotated as a ᴅ-serine/ʟ- and ᴅ-alanine/glycine transporter with homology to CycA in *Mycobacterium tuberculosis* (18, 19). Nonsynonymous point mutations in *cycA* contribute, in part, to increased ᴅ-cycloserine (DCS) resistance in *Mycobacteria* (18, 19), prompting us to investigate the susceptibility of the *S. aureus cycA* mutant to DCS. In contrast to the observations in *Mycobacteria*, our data showed that NE810 was significantly more

susceptible to DCS than the wild type JE2 and was successfully complemented by the wild-type *cycA* gene (Fig. 2A). By broth dilution, the DCS MIC decreased from 32 μg/ml in JE2 to 2 μg/ml in NE810 (Table 1). The *cycA* mutants of the MRSA strains USA300, DAR173, DAR22 and DAR169 exhibited similar and significant increases in susceptibility to DCS (Table 1). Transduction of the *cycA* allele from NE810 into the MSSA strains 8325-4 and ATCC29213 was also associated with a reduction in the DCS MIC from 32 to 4 μg/ml. DCS is a broad-spectrum antibiotic produced by several *Streptomyces* species used as a second line drug in the treatment of TB infections in humans. This drug is a cyclic analogue of alanine and serine that inhibits the alanine racemase (Alr) enzyme that converts L-alanine to D-alanine, as well as the ligase enzyme that links two D-alanine amino acids together (20). These are important enzymes in both peptidoglycan (cell wall cross-linking) and teichoic acid biosynthesis in Gram-positive bacteria. A mutant in the putative D-alanyl:D-alanine ligase (*ddl*, SAUSA300_2039) is not available in the NTML library, suggesting that this gene may be essential. However, the NTML library *alr* mutant NE1713 was associated with significantly increased susceptibility to cefoxitin (Fig. S1A) and hyper-susceptibility to DCS (Fig. S1C), which is consistent with the important role for D-alanine in *S. aureus* peptidoglycan biosynthesis and β-lactam resistance.

CycA is an alanine permease that is required for D-ala-D-ala incorporation into the peptidoglycan stem peptide. To investigate the role of CycA as a potential permease, JE2 and NE810 were grown for 8 h in chemically defined media containing 14mM glucose (CDMG) and amino acid consumption in spent media was measured. Although no growth rate or yield difference was noted between JE2 and NE810 in CDMG (Fig. 2B), alanine uptake by NE810 was significantly impaired compared to JE2 (Fig. 2C). Utilisation of other amino acids by NE810 and JE2, including serine and glycine, were similar (Figs. 2D and 2E and Fig. S2). Impaired alanine transport in the *cycA* mutant grown in CDMG correlated with increased susceptibility to oxacillin (1 μg/ml) (Fig. 2F) and DCS (1 μg/ml) (Fig. 2G). These data demonstrate for the first time that CycA in *S. aureus* functions as an alanine permease. Scanning and transmission electron microscopy revealed gross differences in whole cell morphology (Fig. 3A and B) and cell wall thickness (Fig. 3C and D) in NE810 compared to JE2. NE810 exhibited a highly irregular cell shape, and many cells appeared collapsed (Fig. 3B). TEM imaging revealed that NE810 cells had a wrinkled cell surface (Fig. 3D), a thinner cell

wall (19 ± 2.5nm) compared to JE2 (26 ± 1.8nm) and increased rupturing, although cell size (mean cell diameter) was unaffected (Table S2).

Quantitative peptidoglycan compositional analysis was performed using UPLC analysis of muramidase-digested muropeptide fragments extracted from exponential phase cultures of JE2 and NE810 grown for 220 mins in TSB media (Fig. S3). The PG profile of the *cycA* mutant revealed a significant accumulation of tripeptides compared to wild-type JE2 (Fig. 4A-C), which was associated with a significant reduction in crosslinking (Fig. 4D). In NE810, the dimer, trimer and tetramer fractions were decreased, which was accompanied by a concomitant increase in the monomer fraction (Fig. 4E). Consistent with this data, exposure of JE2 to DCS 8μg/ml was also associated with a similar accumulation in muropeptides with tripeptide stems (Fig. 4C), reduced cross-linking (Fig. 4D), increased muropeptide monomers and reduced dimers, trimers and tetramers (Fig. 4E). DCS had a strong dose-dependent effect, and at concentrations of 20 and 32 μg/ml, led to amplified effects on the accumulation of muropeptides with tripeptide stems, reduced cross-linking and accumulation of monomers (Fig. 4 A-E). These data are consistent with the previously reported impacts of sub-inhibitory (0.25× MIC) and 4× MIC DCS concentrations, which were accompanied by incorporation of an incomplete stem peptide (tripeptide) (20) and a 4-fold reduction in D-ala-D-ala levels (21), respectively. Collectively these data indicate that impaired D-ala incorporation into the PG stem peptide in the *cycA* mutant or following exposure to DCS is accompanied by reduced PG cross-linking and increased β-lactam susceptibility.

Mutation of *cycA* or exposure to D-cycloserine increases the susceptibility of MRSA to β-lactam antibiotics. Previously reported synergy between DCS and β-lactam antibiotics (20, 22) suggests that impaired alanine uptake in the *cycA* mutant may have the same impact on cell wall biosynthesis as DCS-mediated inhibition of Alr and Ddl. To further investigate this, we compared the activity of DCS and β-lactam antibiotics, alone and in combination, against JE2 and NE810. Checkerboard microdilution assay fractional inhibitory concentration indices (ΣFICs ≤0.5) revealed synergy between DCS and several licensed β-lactam antibiotics with different PBP selectivity against JE2 and USA300 FPR3757 (Table 1). Oxacillin and nafcillin were not included in checkerboard assays because measurement of their MICs involves supplementing the media with 2% NaCl, which distorts the MIC of DCS (data not shown).

Using the MRSA strains JE2, USA300, DAR173, DAR22, DAR169 and their corresponding *cycA* mutants, the kinetics of killing by DCS, oxacillin and cefoxitin, alone and in combination was measured over 24h using antibiotic concentrations corresponding to 0.125×, 0.25× and 0.5× MICs. Recovery of growth in media supplemented with oxacillin or cefoxitin alone was evident after 8 h (Fig. 5A-E), reflecting the selection and expansion of HoR mutants as described previously (4, 21, 23). Recovery of growth in cultures exposed to DCS alone was also evident (Fig. 5A-E), which may correlate with our observation that mutants resistant to DCS (on BHI agar supplemented with 128 μg/ml DCS) arise at a rate of approximately 5.5×10^{-8} per cell per generation (data not shown). Using combinations of DCS and oxacillin or cefoxitin at 0.125× MIC did not achieve a $\geq 2 \log^{10}$ reduction in the number of CFU/ml (data not shown). However, at 0.5× MIC for strains JE2, USA300, DAR73 and DAR22, DCS (16 μg/ml)/oxacillin (32 μg/ml) and DCS (16 μg/ml)/cefoxitin (32 μg/ml) combinations achieved a $\geq 5 \log^{10}$ reduction in the number of CFU/ml compared to oxacillin, cefoxitin or DCS alone (Fig. 5A-D). For strain DAR169, DCS/β-lactam combinations at 0.25× MIC was sufficient to achieve a $\geq 5 \log^{10}$ reduction in CFUs recovered compared to the individual antibiotics (Fig. 5E). DCS/β-lactam combinations at 0.5× MIC were also able to achieve $\geq 5 \log^{10}$ reduction in the number of CFU/ml against the methicillin resistant *S. epidermidis* (MRSE) strain RP62A (24) compared to either antibiotic alone (Fig. S4). Checkerboard experiments with fourteen MRSA strains and MRSE strain RP62A further revealed synergy (ΣFICs ≤0.5) between DCS and a range of β-lactam antibiotics with different penicillin binding protein (PBP) specificity, namely cefoxitin (PBP4), cefaclor (PBP3), cefotaxime (PBP2), piperacillin-tazobactin (PBP3/β-lactamase inhibitor) and imipenem (PBP1) (Table 1).

Preliminary studies using disk diffusion assays also revealed synergy between DCS and imipenem (IMP) or piperacillin-tazobactin (Pip/Taz) against several ESKAPE pathogens. Using checkerboard assays synergy (ΣFIC values ≤0.5) was measured between DCS and Pip/Taz or IMP against *Pseudomonas aeruginosa* LES431, *Acinetobacter baumannii* DSM 30007, *K. pneumoniae* 511232.1 (ESBL[+]), and two vancomycin resistant *Enterococcus* strains (Table 2). This synergy appears to be specific to β-lactams and no synergy (ΣFICs > 0.5) was measured between DCS and several antibiotics that are used topically or systemically for the decolonization or treatment of patients colonized/infected with *S. aureus* or MRSA (clindamycin, trimethoprim, mupirocin, ciprofloxacin), or several antibiotics to which *S.*

aureus isolates commonly exhibit resistance (tobramycin, kanamycin and spectinomycin) (Table S1). Furthermore the *cycA* mutation had no impact on susceptibility to any of these non-β-lactam antibiotics (apart from clindamycin resistance which is encoded by *ermB* on the transposon).

Combination therapy with DCS and oxacillin significantly reduces the bacterial burden in the kidneys and spleen of mice infected with MRSA. The virulence of the NE810 mutant and the therapeutic potential of oxacillin in combination with DCS in the treatment of MRSA infections were assessed in mice. Mice infected with NE810 displayed more piloerection than those infected with wild-type JE2 suggestive of more severe infection, and one mouse infected with NE810 required euthanization during the experiment. The average number of CFUs recovered from the kidneys of mice infected with NE810 ($1.5 \times 10^8 \pm 1.49 \times 10^8$) was higher than for mice infected with JE2 ($5.7 \times 10^7 \pm 6.3 \times 10^7$), but did not reach significance. Interestingly, NE810 colonies exhibited visibly increased haemolytic activity on sheep blood agar (Fig. S5A). We previously reported that increased haemolytic activity in MSSA strains was associated with increased expression of the cell-density controlled, global accessory gene regulator (*agr*) system (4, 21, 25). Comparison of *agr* (RNAIII) temporal expression using RT-qPCR revealed a more significant activation of the *agr* system after 6 hrs growth in NE810 compared to JE2 (Fig. S5B). This difference in *agr* activation was not related to any change in growth of NE810 compared to JE2 (Fig. S5C). Taken together these data indicate that increased β-lactam susceptibility in the NE810 *cycA* mutant is accompanied by both increased *agr* expression and haemolytic activity, and a trend towards increased virulence in this model.

Treatment with oxacillin or DCS alone significantly reduced the number of CFUs recovered from the kidneys of mice infected with JE2 (Fig. 6). Furthermore the oxacillin/DCS combination was significantly more effective than either antibiotic alone and the combination was equally effective in reducing the bacterial burden in the kidneys of animals infected with JE2 or NE810 when compared to no treatment ($p \leq 0.0001$) (Fig. 6) demonstrating the capacity of DCS to significantly potentiate the activity of β-lactam antibiotics against MRSA under *in vivo* conditions. Unexpectedly, oxacillin- or DCS-mediated eradication of NE810 infections was similar to JE2 (Fig. 6). In the spleen, only oxacillin/DCS

combination treatment was associated with a significant reduction in the number of CFUs

recovered from mice infected with JE2 or NE810 (Fig. S6).

CycA-independent alanine transport increases oxacillin and DCS resistance in the *cycA*

mutant. The failure of oxacillin or DCS treatment to enhance the eradication of NE810

infections in the mouse bacteraemia model prompted us to further characterise the growth

conditions used for the *in vitro* antibiotic susceptibility assays. Specifically we investigated

the role of glucose, which we previously reported to increase the growth requirement for

amino acids (26), and which we reasoned may be important for CycA alanine permease

activity. Growth of JE2 and NE810 was similar in CDM lacking glucose (Fig. 7A) and uptake of

alanine (Fig. 7B) and other amino acids (Fig. S7) was unchanged in NE810 compared to JE2.

Furthermore NE810 and JE2 grew equally well in CDM supplemented with oxacillin and DSC

(Fig. 7C and D). These data reveal a strong correlation between CycA-dependent alanine

transport and susceptibility to oxacillin and DCS, and may explain in part why the *cycA*

mutant does not exhibit increased β-lactam and DCS susceptibility in the mouse

bacteraemia model.

Discussion

The exploitation of antibiotic re-purposing as part of concerted efforts to address the antimicrobial resistance crisis has been hampered by a lack of mechanistic data to explain demonstrated therapeutic potential and the perception that studies attempting to identify new uses for existing drugs are not hypothesis-driven. However, basic research discoveries that advance our understanding of bacterial virulence and antibiotic resistance mechanisms can also be used to inform targeted antibiotic re-purposing, in which the mechanism of action is understood. In this study, we identified an alanine permease (CycA) required for full expression of resistance to β-lactam antibiotics and DCS. Loss of function of this alanine transporter significantly increased the susceptibility of MRSA to β-lactam antibiotics, an outcome that could be reproduced through exposure to DCS. DCS targets the early steps in cell wall biosynthesis by inhibiting the activity of Alr and Ddl.

The potential of β-lactam / DCS combinations in the treatment of recalcitrant *S. aureus* infections is particularly important given that *S. aureus* isolates resistant to all licensed anti-staphylococcal drugs have been reported, and in light of the regulatory barriers to the introduction of new antimicrobial drugs. The recent report that DCS can potentiate the activity of vancomycin against a laboratory-generated vancomycin highly-resistant *S. aureus* (VRSA) strain *in vitro* and in a silkworm infection model (27) is also important. For MRSA infections, the excellent safety profile of β-lactam antibiotics makes these drugs particularly attractive as components of combination antimicrobial therapies. When used in the treatment of tuberculosis DCS (trade name Seromycin, The Chao Centre) is typically administered orally in 250 mg tablets twice daily for up to two years. At this dosage, the DCS concentration in blood serum is generally 25-30 μg/ml, which is similar to the concentrations used in our *in vitro* and *in vivo* experiments. The known neurological side effects associated with DCS therapy mean that this antibiotic is unlikely to be considered for the treatment of MRSA infections unless alternative therapeutic approaches have been exhausted. DCS is a co-agonist of the *N*-methyl-D-aspartic acid receptor in the central nervous system can include seizures and peripheral neuropathy (28, 29), and the use of this antibiotic is limited to the treatment of recalcitrant, multi-drug resistant tuberculosis infections, for which long-term therapy is required (30). Oxacillin/DCS combination therapy was significantly more effective than DCS or oxacillin alone over a 5-day therapeutic window

suggesting that further studies on using DCS to augment β-lactams as a treatment option for

recalcitrant staphylococcal infections is merited.

The alanine permease identified in this study, CycA, shares 46% and 53% identity with CycA

in *M. tuberculosis* and *E. coli*, respectively. Mutation of *cycA* increases the susceptibility of

MRSA to β-lactam antibiotics and results in hyper-susceptibility to D-cycloserine, whereas a

cycA point mutation in *M. bovis* contributes, in part, to increased DCS resistance presumably

by interfering with DCS transport into the cell (19). In *E. coli*, *cycA* null mutations can also

result in increased resistance or have no effect in DCS susceptibility depending on the

growth media (31-35), suggesting that CycA is primarily important for DCS resistance under

conditions when its contribution to amino acid transport is also important. The opposite

effects of *cycA* mutation on DCS susceptibility in *S. aureus* and *E. coli* or *Mycobacteria*

indicates that DCS uptake is not significantly blocked by the *cycA* mutation in *S. aureus*.

Under growth conditions where CycA is required for alanine transport (in nutrient/glucose-

replete media), mutation of *cycA* or DCS-exposure have similar effects on the structure of *S.*

aureus peptidoglycan (Fig. 8). Consistent with previous studies in *S. aureus* (20) and in *M.*

tuberculosis (36), our studies showed a dose-dependent accumulation of muropeptides with

a tripeptide stem in MRSA exposed to DCS. The *cycA* mutation was also associated with the

increased accumulation of muropeptides with a tripeptide stem. These data indicate that a

reduced intracellular alanine pool or inhibition of Alr and Ddl is associated with reduced D-

ala-D-ala incorporation into the PG stem peptide. The increased accumulation of tripeptides

in turn interferes with normal PBP transpeptidase activity and offers a plausible explanation

for increased susceptibility to β-lactam antibiotics. The importance of the terminal stem

peptide D-ala-D-ala for β-lactam resistance has previously been reported. Mutation of the

murF-encoded ligase, which catalyses of the D-ala-D-ala into the stem peptide also increased

β-lactam (but not DCS) susceptibility (37, 38). Similarly growth of a HoR MRSA strain in

media supplemented with high concentrations of glycine was accompanied by replacement

of the D-ala-D-ala with D-ala-gly and decreased methicillin resistance (39).

Impaired uptake of alanine in CDMG correlated with increased susceptibility to oxacillin and

DCS, suggesting that alanine utilisation via CycA is important to make D-alanine available for

cell wall biosynthesis and consequently resistance to β-lactam antibiotics. Consistent with

this, NE810 also exhibited increased oxacillin susceptibility in BHI, TSB and MH media.

However no change in alanine transport or susceptibility to oxacillin and DCS was measured

in CDM lacking glucose, which was consistent with the failure of oxacillin and DCS to more

efficiently eradicate NE810 infections in the mouse bacteraemia model. The availability of

nutrients such as glucose and amino acids varies in different niches colonised by *S. aureus*

during infection ranging from glucose-rich in organs such as the liver (40), to glucose-

depleted in established abscesses (41), which in turn impacts the role of amino acids as

carbon sources (26, 42), and potentially the activity of CycA in alanine transport and β-

lactam susceptibility. Furthermore, alanine transport was unaffected in the *cycA* mutant

grown in CDM indicating that an alternative alanine transport mechanism(s) may be active

under these growth conditions (Fig. 8). Identification of this alternative alanine permease

may be important in the development of therapeutic strategies targeting alanine transport

to increase β-lactam susceptibility in MRSA, while elucidation of the role of glucose in the

control of alanine transport should provide new insights into β-lactam resistance.

Materials and Methods

Bacterial strains, growth conditions and antimicrobial susceptibility testing. Bacterial strains used in this study (Table S3) were grown in Luria Bertoni (LB) (Sigma), brain heart infusion (BHI) (Oxoid), Mueller Hinton (MH) (Oxoid), nutrient (Oxoid), sheep blood BHI or chemically defined media (CDM).

The minimum inhibitory concentrations (MICs) of oxacillin, cefaclor, cefataxime, nafcillin, cefoxitin, imipenem, piperacillin-tazobactam (Pip/Taz), clindamycin, tobramycin, trimethoprim, mupirocin, ciprofloxacin, spectinomycin, kanamycin and D-cycloserine were determined in accordance with the Clinical Laboratory Standards Institute (CLSI) guidelines using plate and broth dilution assays in MH or MH containing 2% NaCl for oxacillin and nafcillin. Oxacillin MICs were also measured using E-tests strips from Oxoid on MH agar containing 2% NaCl. Quality control strains ATCC29213 and ATCC25923 were used for broth dilution and disk diffusion oxacillin and cefoxitin MIC assays, respectively.

Identification of cefoxitin susceptible MRSA mutant NE810. The Nebraska transposon mutant library (NTML) comprising 1,952 strains harbouring transposon insertions in all non-essential genes was constructed in the USA300 derivative JE2, which has been cured of plasmids p01 and p03 (15). Cefoxitin susceptibility of individual mutants was determined using the disk diffusion method in accordance with the CLSI guidelines using cefoxitin (30μg) disks (Oxoid) on Mueller Hinton agar (Oxoid). Briefly, each library mutant was revived from the freezer on BHI agar supplemented with 10μg/ml erythromycin and grown overnight for 18 h at 37°C. The wild-type parent strain of the library JE2 and the library mutant containing a transposon insertion in *mecA* (NE1868) were used as positive (cefoxitin resistant) and negative (cefoxitin susceptible) controls, respectively. The cefoxitin susceptible *S. aureus* strain ATCC25923 was also used as a negative control. Several colonies were used to prepare a bacterial suspension in sterile phosphate-buffered saline (PBS), which was standardised to 0.5 McFarland and used to inoculate a MH agar plate before the application of the cefoxitin-impregnated disc. The MH agar plates were then incubated at 35°C for 18 h and the zone sizes measured. The zone diameter for JE2 was 18mm and >35mm for NE1868 (*mecA*). The cefoxitin susceptible mutant NE810 exhibited a zone diameter = 22mm. The entire library was screened in duplicate and putative cefoxitin susceptible mutants were

tested at least five times. This susceptibility of cefoxitin sensitive mutants to oxacillin was also determined by E-test, as well as broth and plate dilution assays.

PCR was used to verify the presence of the transposon insertion NE810 using the primers NE810_Fwd and NE810_Rev (Table S4). Phage 80α was used to transduce the transposon mutation from NE810 back into JE2 and USA300 to ensure that secondary mutations were not involved in increased cefoxitin susceptibility phenotype. DNA extracted from transductants selected on BHI agar supplemented with erythromycin 10μg/ml were verified by PCR using primers NE810_Fwd and NE810_Rev. Comparative whole genome sequencing of JE2 and NE810 (MicrobesNG, UK) using the USA300_FPR3757 genome as a reference revealed the absence of other mutations in NE810.

To complement NE810, the *cycA* gene was amplified from JE2 on a 1608 bp PCR product using primers NE810F1_Fwd and NE810F1_Rev (Table S4) and cloned into the *E. coli-Staphylococcus* shuttle plasmid pLI50 using the Clontech In-fusion HD cloning kit. The primers were designed using the CloneAmp In-fusion tool with 15 bp extensions that were complementary to the ends of the linearized vector. The recombinant plasmid generated was transformed into *E. coli* HST08 and verified by Sanger sequencing (Source Biosciences) before being transformed by electroporation into the restriction-deficient laboratory strain RN4220, and subsequently NE810. All plasmid-harbouring strains were cultured in medium supplemented with 100 μg/ml ampicillin (*E. coli*) or 10 μg/ml chloramphenicol (*S. aureus*) to maintain plasmid selection.

Growth measurements of JE2 strains in chemically defined media and amino acid analysis. Overnight cultures of JE2 and NE810 were grown at 37^{o}C in TSB, washed in PBS and inoculated at a starting cell density of A_{600} = 0.05 into chemically defined medium (CDM) or CDM supplemented with 14mM glucose (CDMG) (43). The bacteria were grown aerobically (250 rpm; 10:1 flask to volume ratio, 37°C) and the A_{600} was determined every two hours for 8-12 hours. One ml of the bacterial culture was collected every two hours and pelleted by centrifugation at 14,000x g for 3 minutes. The spent media was filtered using an Amicon Ultra centrifugal filter (Millipore; 3,000 molecular weight cutoff [MWCO]). Amino acid analysis was performed by the Protein Structure Core Facility, University of Nebraska Medical Center, using a Hitachi L-8800 amino acid analyzer as described previously (26).

mecA gene expression analysis. Quantitative reverse transcription PCR (RT-qPCR) was used to measure *mecA* transcription on the Roche LightCycler 480 instrument using the LightCycler 480 Sybr Green Kit (Roche) with primers mecA1_Fwd and mecA1_Rev (Table S4). The following cycling conditions were used: $95^{o}C$ for 5 minutes and followed by 45 cycles of $95^{o}C$ for 10 seconds, $58^{o}C$ for 20 seconds and $72^{o}C$ for 20 seconds. Melt curve analysis was performed at $95^{o}C$ for 5 seconds followed by $65^{o}C$ for one minute up to 97 C at a ramp rate of 0.11c/sec with five readings taken for every degree of temperature increase. The *gyrB* gene amplified with primers gyrB_Fwd and gyrB_Rev (Table S4) served as an internal standard for all reactions. For each reaction, the ratio of *mecA* and *gyrB* transcript number was calculated as follows: $2^{(Ct\ gyrB\ -\ Ct\ mecA)}$. Each RT-qPCR experiment was performed three times and presented as average data with standard errors.

Scanning Electron Microscopy (SEM). Cell shape and cell size analysis were carried out using SEM imaging. Overnight BHI cultures were diluted 1:200 in BHI broth supplemented with 1% glucose and incubated for 24h at $37^{o}C$. Subsequently, the slides were rinsed three times with distilled water, dried at $60^{o}C$ for 1h, before being rinsed with 0.1M phosphate buffer, fixed in 2.5% glutaraldehyde for 2h, rinsed again with 0.1M phosphate buffer, dehydrated in ethanol (20%, 30%, and 50% for 10 min each step; 70% ethanol with 0.5% uranyl acetate for 30 min; and then 90%, 96%, and 100% ethanol), soaked in hexamethyldisilazane (HMDS) for 30 minutes and finally dried overnight. The slides were fixed to metal stubs before being coated in gold and imaged using a Hitachi S2600N Variable Pressure Scanning Electron Microscope.

Transmission Electron Microscopy (TEM). Overnight cultures were diluted 1:200 in 10ml BHI media and grown at 37°C to A_{600} = 1. The cells were pelleted at 8,000 × *g* before being re-suspended in fixation solution (2.5% glutaraldehyde in 0.1 M cacodylate buffer [pH 7.4]) and incubated overnight at 4°C. The fixed cells were further treated with 2% osmium tetroxide, followed by 0.25% uranyl acetate for contrast enhancement. The cell pellets were then dehydrated in increasing concentrations of ethanol as described above for the SEM cell preparation, followed by pure propylene oxide, and transferred to a series of resin and propylene oxide mixtures (50:50, 75:25, pure resin) before being embedded in Epon resin. Thin sections were cut on an ultramicrotome. Images were analysed using a Hitachi H7000

instrument. At least 3 to 5 measurements of cell wall thickness were performed on each cell

and approximately 30 cells were measured for each sample.

Analysis of peptidoglycan composition in NE810 and JE2 treated with D-cycloserine. Using

overnight cultures of JE2 and NE810, independent quadruplicate 50ml broth cultures were

set up at a starting cell density of A_{600}=0.05 and grown for approximately 2 hours to

≈A_{600}=0.5, before being left untreated or dosed with DCS at a final concentration of 8, 20 or

32 µg/ml for a further 100 mins before the cells in the entire 50 ml culture were harvested

by centrifugation and resuspended in 5ml PBS (Fig. S3). The cell samples were slowly

dropped into an equal volume of boiling 10% (wt/vol) SDS, vigorously stirred for 2h in a

boiling water bath and left stirring overnight at room temperature. The insoluble fractions

were recovered by high speed centrifugation (150,000 x g, 15 min, 25°C) and washed until

the fractions were free from SDS. The insoluble fractions were purified for peptidoglycan

according to the protocol described previously (44). Purified peptidoglycan was re-

suspended in 50 mM NaPO$_4$ buffer, pH 4.9 and treated with 80 µg/mL muramidase (Cellosyl)

for 16 h at 37°C. Muramidase digestion was stopped by incubation in a boiling water bath,

and coagulated proteins were removed by centrifugation (21,000 x g, 10 min). The

supernatants were adjusted to pH 9.0 with sodium borate and reduced with sodium

borohydride for 30 min at room temperature. Finally, samples were adjusted to pH 3.5 with

orthophosphoric acid and filtered prior to UPLC analysis with a Waters Acquity UPLC H-class

system (Waters) equipped with a Waters Xevo G2-XS QTof MS (Waters MS

Technologies). Chromatographic separation was achieved on acquity UPLC® BEH C18, 1.7

µm, 150 x 2.1 mm id (Waters) column. The mobile consisted of solvent A: 0.1% formic acid

in Milli-Q water and B: 0.1% formic acid in acetonitrile. The gradient was set as follows: 0-3

min 2-5% B, 3-6 min 5-6.8% B, 6-7.5 min 6.8-9% B, 7.5-9 min 9-14% B, 9-11 min 14-20% B,

11-12 min hold at 20% B keeping the constant flow of 0.250µl/min, 12-12.1 min 20-90% B

with a flow rate ramp of 0.250µl/min to 0.300µl/min further holding the same flow rate and

% B till 13.5 min, 13.5-13.6 min 90-2% B, 13.6-16 min hold at 2% B. The flow rate was

ramped back again to 0.250µl/min at 16.1 min and column was equilibrated form next

analysis till 18min. The chromatographic separation was recorded at 204nm.

Mass spectrometry was performed on a Xevo G2-XS QTof quadrupole time-of-flight mass

spectrometer (Waters MS Technologies). The scan range was from 10 to 2000 m/z with a

scan rate of 0.25s. Instrument was operated in positive electro-spray ionization mode. The capillary and sample cone voltages were 3kV and 40 V, respectively. Gas flows were set at 100 and 500 l/hr for cone gas and desolvation gas, respectively. The source temperature was 120°C and desolvation temperature was 350°C. For MS^E low collision energy was set at 6 eV while ramping the high collision energy from 15-40 eV. Leucine-enkephalin reference was used as the lockmass at a concentration of 200 pg/mL with a continuous flow of 5μL/min and 0.25s scan time, to maintain the accuracy of the analysis. All the acquisition and analysis of data were controlled by Waters UNIFI software. Structural characterization of muropeptides was performed based on their MS data and MS/MS fragmentation pattern, which matched with the PG composition and structure reported in previous publications (39, 45-47).

Antibiotic synergy analysis using the microdilution checkerboard assay. The synergistic activity of D-cycloserine and other antibiotics was measured using the checkerboard microdilution method in 96-well plates. The final inoculum in each well was 5×10^5 CFU/ml, and the results (growth or no growth) were determined after 24 h incubation at 37°C. Un-inoculated MH broth supplemented with the antibiotics was used as a negative control. Inoculated MH broth with no antibiotics was used as a positive control. The fractional inhibitory concentration index (ΣFIC) was calculated for each drug combination. An FIC index of ≤0.5 was considered synergistic, one of >0.5 to <2 was considered indifferent, and one of >2 was considered antagonistic. All experiments were performed in triplicate.

Kill curve assays. Suspensions of bacterial cells from overnight cultures adjusted to 10^7 CFU/ml were exposed to increasing concentrations (0.125×, 0.25×, and 0.5× MIC) of oxacillin, cefoxitin and DCS alone or in combination, and the number of colony forming units (CFU)/ml enumerated at 0, 2, 4, 8 and 24 h. Data is presented at the antibiotic concentrations where synergy was measured i.e. 0.5× MIC for JE2, USA300, DAR173, DAR22, DAR113, BH1CC, and RP62A, and 0.25× MIC for DAR169. Antibiotic synergism was defined as a ≥2 log^{10} decrease in the number of CFU/ml in cell suspensions exposed to DCS/β-lactam combinations compared to the most effective individual drug after 8 h.

Mouse infection experiments. 6- to 8-week-old, age matched, outbred CD1 female mice (Charles River, UK) were used in a non-lethal model of bacteremia. JE2 and NE810 cultures were grown to A_{600}=0.5 in BHI, washed in PBS, adjusted to 1×10^8 CFU/ml. Mice were

infected intravenously (via the tail vein) with 5×10^6 CFU ($n = 10$ mice per group). The infections were left untreated (PBS control) or treated with either 75 mg oxacillin/Kg/12 hours, 30 mg DCS/Kg/12 hours or a combination of both (first antibiotic dose administered 16 hours post infection), before being sacrificed after 5 days. Bacteria present in homogenised spleens and kidneys recovered from the mice were enumerated on blood agar. Statistical significance was assessed using one-way ANOVA with Kruskal-Wallis test followed by Dunn's multiple comparisons test in the GraphPad Prism application.

Ethics Statement. Mouse experiments were approved by the United Kingdom Home Office (Home Office Project License Number 40/3602) and the University of Liverpool Animal Welfare and Ethics Committee. This study was carried out in strict accordance with the United Kingdom Animals (Scientific Procedures) Act 1986. All efforts were made to minimize suffering.

Statistical analysis. Two-tailed Student's t-Tests and one-way ANOVA with Kruskal-Wallis test followed by Dunn's multiple comparisons test (*in vivo* experiments) were used to determine statistically significant differences in assays performed during this study. A *p* value <0.05 was deemed significant.

518 **Acknowledgements:**

519 We thank Craig Winstanley and Kate Reddington for generously providing bacterial strains.

Figure 1. Mutation of *cycA* increases the susceptibility of MRSA to β-lactam antibiotics. A. Chromosomal location of *cycA* and neighbouring genes *icd* (isocitrate dehydrogenase), *gltA* (citrate synthase), *pyk* (pyruvate kinase) and *pfkA* (6-phosphofructokinase). The locations of transposon insertions in NE810, NE491, NE594 and NE1497 mutants from the Nebraska library are indicated. **B.** E-test measurement of oxacillin minimum inhibitory concentrations (MICs) in JE2 (wild type), NE810 (*cycA*::Tn), NE810 carrying pLI50 (control) and p*cycA*.

Figure 2. Mutation of *cycA* increases susceptibility to D-cycloserine in MRSA and impairs alanine uptake. A. Comparison of zones of inhibition around D-cycloserine 30μg disks on lawns of JE2, NE810 (*cycA*::Tn), NE810 pLI50 (control) and NE810 p*cycA* grown on Mueller Hinton (MH) agar. **B.** Growth of JE2 and NE810 in chemically defined media supplemented with glucose (CDMG). Cell density was measured at A_{600}. **C, D and E.** Alanine (C), Serine (D) and Glycine (E) consumption by JE2 and NE810 grown aerobically in CDMG. Residual amino acid was measured in spent media after 0, 4, 6 and 8 h growth. **F and G.** Growth of JE2 and NE810 cultures for 12 h in CDMG supplemented with 1 μg/ml oxacillin (F) or 1 μg/ml D-cycloserine (G). Cell density was measured at A_{600}.

Figure 3. Mutation of *cycA* reduces cell wall thickness in MRSA. A and B. Comparison of JE2 (A) and NE810 (B) cell morphology by scanning electron microscopy (SEM). **C and D.** Comparison of JE2 (C) and NE810 (D) cell wall thickness by transmission electron microscopy (TEM). TEM magnifications are ×15,000 (top) and ×70,000 (bottom) and cell wall thickness measurements are provided in Table S2.

Figure 4. Mutation of *cycA* or D-cycloserine (DCS) treatment impacts peptidoglycan peptide stem length and reduces cell wall crosslinking. A. Representative UV chromatograms of peptidoglycan from wild-type JE2, NE810 and JE2 treated with increasing concentrations of DCS (8, 20 and 32 μg/ml). Muropeptides with tripeptide stems are numbered 1-3. **B.** Proposed structures of the three muropeptides with tripeptide stems identified in NE810 and DCS-treated JE2 cells. NAG, N-acetylglucosamine; NAM, N-acetylmuramic acid; L-Ala, L-alanine; D-Gln, D-glutamine; D-Glu, D-glutamic; L-Lys, L-lysine. The theoretical and observed neutral masses determined by MS are indicated. **C.** Relative abundance of muropeptides with tripeptides in the stem. **D.** Relative crosslinking efficiency. **E.** Relative proportions of cell wall muropeptide fractions based on oligomerization. All errors bars represent 95% confidence interval (n = 4). Significant differences determined using Students t-test are denoted using asterisks (* p<0.05; ** p<0.01; *** p<0.001).

Figure 5. *In vitro* kill curves for D-cycloserine (DCS), oxacillin and cefoxitin with JE2 (A), USA300 FPR3757 (B), DAR173 (C), DAR22 (D), DAR169 (E) and their isogenic *cycA* mutants. Antibiotics at the concentrations indicated (μg/ml) were added to suspensions of overnight bacterial cultures adjusted to 10^7 CFU/ml in BHI (A_{600}=0.05), incubated at 37°C and the number of CFU/ml enumerated at 0, 2, 4, 8 h. The data presented are the mean of three independent experiments, and standard error of the mean is shown. Antibiotic synergism

was defined as a ≥ 2 log^{10} decrease in the number of CFU/ml in cell suspensions exposed to
DCS/β-lactam combinations compared to the most effective individual antibiotic alone.

Figure 6. Combination therapy with D-cycloserine and oxacillin significantly reduces the
bacterial burden in the kidneys of mice infected with MRSA. The number of colony-forming
units (CFU) recovered from the kidneys of mice infected by tail vein injection with 5×10^6 JE2
or NE810 (CycA) and left untreated or treated with 75mg of oxacillin (Ox)/kg, 30mg of
DCS/kg or a combination of both Ox and DCS delivered subcutaneously every 12 hours for 5
days. The first antibiotic dose was given 16 hours after infection. Significant differences
determined using one-way ANOVA with Kruskal-Wallis test followed by Dunn's multiple
comparisons test are denoted using asterisks (*$p \leq 0.05$, **$p \leq 0.01$, ****$p \leq 0.0001$). The limit
of detection (50 colonies) is indicated with a hashed red line.

Figure 7. CycA-independent alanine transport increases oxacillin and D-cycloserine
resistance in the *cycA* mutant. **A.** Growth of JE2 and NE810 in chemically defined medium
lacking glucose (CDM). Cell density was measured at A_{600}. **B.** Alanine consumption by JE2
and NE810 grown aerobically in CDM. Residual amino acid was measured in spent media
after 0, 4, 6 and 8 h growth. **C and D.** Growth (cell density at A_{600}) of JE2 and NE810 cultures
for 12 h in CDM supplemented with 1 μg/ml oxacillin (C) or 1 μg/ml DCS (D).

Figure 8. Proposed model depicting how impaired alanine transport associated with
mutation of CycA or exposure to DCS can inhibit the D-alanine pathway for peptidoglycan
biosynthesis leading to increased susceptibility to β-lactam antibiotics.

References

1. **Clatworthy AE, Pierson E, Hung DT.** 2007. Targeting virulence: a new paradigm for antimicrobial therapy. Nat Chem Biol **3:**541-548.
2. **Jones M, Jernigan JA, Evans ME, Roselle GA, Hatfield KM, Samore MH.** 2019. Vital Signs: Trends in Staphylococcus aureus Infections in Veterans Affairs Medical Centers - United States, 2005-2017. MMWR Morb Mortal Wkly Rep **68:**220-224.
3. **Kourtis AP, Hatfield K, Baggs J, Mu Y, See I, Epson E, Nadle J, Kainer MA, Dumyati G, Petit S, Ray SM, Emerging Infections Program Mag, Ham D, Capers C, Ewing H, Coffin N, McDonald LC, Jernigan J, Cardo D.** 2019. Vital Signs: Epidemiology and Recent Trends in Methicillin-Resistant and in Methicillin-Susceptible Staphylococcus aureus Bloodstream Infections - United States. MMWR Morb Mortal Wkly Rep **68:**214-219.
4. **Waters EM, Rudkin JK, Coughlan S, Clair GC, Adkins JN, Gore S, Xia G, Black NS, Downing T, O'Neill E, Kadioglu A, O'Gara JP.** 2017. Redeploying beta-lactam antibiotics as a novel antivirulence strategy for the treatment of methicillin-resistant Staphylococcus aureus infections. J Infect Dis **215:**80-87.
5. **Davis JS, Sud A, O'Sullivan MV, Robinson JO, Ferguson PE, Foo H, van Hal SJ, Ralph AP, Howden BP, Binks PM, Kirby A, Tong SY, Combination Antibiotics for MRSAsg, Australasian Society for Infectious Diseases Clinical Research N.** 2016. Combination of Vancomycin and beta-Lactam Therapy for Methicillin-Resistant Staphylococcus aureus Bacteremia: A Pilot Multicenter Randomized Controlled Trial. Clin Infect Dis **62:**173-180.
6. **Tong SY, Nelson J, Paterson DL, Fowler VG, Jr., Howden BP, Cheng AC, Chatfield M, Lipman J, Van Hal S, O'Sullivan M, Robinson JO, Yahav D, Lye D, Davis JS, group Cs, the Australasian Society for Infectious Diseases Clinical Research N.** 2016. CAMERA2 - combination antibiotic therapy for methicillin-resistant Staphylococcus aureus infection: study protocol for a randomised controlled trial. Trials **17:**170.
7. **Chambers HF.** 1997. Methicillin resistance in staphylococci: molecular and biochemical basis and clinical implications. Clin Microbiol Rev **10:**781-791.
8. **Chambers HF, Hackbarth CJ.** 1987. Effect of NaCl and nafcillin on penicillin-binding protein 2a and heterogeneous expression of methicillin resistance in Staphylococcus aureus. Antimicrob Agents Chemother **31:**1982-1988.
9. **Sabath LD, Wallace SJ.** 1971. The problems of drug-resistant pathogenic bacteria. Factors influencing methicillin resistance in staphylococci. Ann N Y Acad Sci **182:**258-266.
10. **Griffiths JM, O'Neill AJ.** 2012. Loss of function of the gdpP protein leads to joint beta-lactam/glycopeptide tolerance in Staphylococcus aureus. Antimicrob Agents Chemother **56:**579-581.
11. **Dengler V, McCallum N, Kiefer P, Christen P, Patrignani A, Vorholt JA, Berger-Bachi B, Senn MM.** 2013. Mutation in the C-di-AMP cyclase dacA affects fitness and resistance of methicillin resistant Staphylococcus aureus. PLoS One **8:**e73512.
12. **Geiger T, Goerke C, Fritz M, Schafer T, Ohlsen K, Liebeke M, Lalk M, Wolz C.** 2010. Role of the (p)ppGpp synthase RSH, a RelA/SpoT homolog, in stringent response and virulence of Staphylococcus aureus. Infect Immun **78:**1873-1883.
13. **Geiger T, Kastle B, Gratani FL, Goerke C, Wolz C.** 2014. Two small (p)ppGpp synthases in Staphylococcus aureus mediate tolerance against cell envelope stress conditions. J Bacteriol **196:**894-902.
14. **Mwangi MM, Kim C, Chung M, Tsai J, Vijayadamodar G, Benitez M, Jarvie TP, Du L, Tomasz A.** 2013. Whole-genome sequencing reveals a link between beta-lactam resistance and synthetases of the alarmone (p)ppGpp in Staphylococcus aureus. Microb Drug Resist **19:**153-159.

628   15.   **Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, Bayles KW.** 2013. A genetic
629   resource for rapid and comprehensive phenotype screening of nonessential Staphylococcus
630   aureus genes. MBio **4:**e00537-00512.
631   16.   **Mlynek KD, Callahan MT, Shimkevitch AV, Farmer JT, Endres JL, Marchand M, Bayles KW,**
632   **Horswill AR, Kaplan JB.** 2016. Effects of Low-Dose Amoxicillin on Staphylococcus aureus
633   USA300 Biofilms. Antimicrob Agents Chemother **60:**2639-2651.
634   17.   **O'Neill E, Pozzi C, Houston P, Smyth D, Humphreys H, Robinson DA, O'Gara JP.** 2007.
635   Association between methicillin susceptibility and biofilm regulation in Staphylococcus
636   aureus isolates from device-related infections. J Clin Microbiol **45:**1379-1388.
637   18.   **Desjardins CA, Cohen KA, Munsamy V, Abeel T, Maharaj K, Walker BJ, Shea TP, Almeida**
638   **DV, Manson AL, Salazar A, Padayatchi N, O'Donnell MR, Mlisana KP, Wortman J, Birren**
639   **BW, Grosset J, Earl AM, Pym AS.** 2016. Genomic and functional analyses of Mycobacterium
640   tuberculosis strains implicate ald in D-cycloserine resistance. Nat Genet **48:**544-551.
641   19.   **Chen JM, Uplekar S, Gordon SV, Cole ST.** 2012. A point mutation in cycA partially
642   contributes to the D-cycloserine resistance trait of Mycobacterium bovis BCG vaccine strains.
643   PLoS One **7:**e43467.
644   20.   **Sieradzki K, Tomasz A.** 1997. Suppression of beta-lactam antibiotic resistance in a
645   methicillin-resistant Staphylococcus aureus through synergic action of early cell wall
646   inhibitors and some other antibiotics. J Antimicrob Chemother **39 Suppl A:**47-51.
647   21.   **Pozzi C, Waters EM, Rudkin JK, Schaeffer CR, Lohan AJ, Tong P, Loftus BJ, Pier GB, Fey PD,**
648   **Massey RC, O'Gara JP.** 2012. Methicillin resistance alters the biofilm phenotype and
649   attenuates virulence in Staphylococcus aureus device-associated infections. PLoS Pathog
650   **8:**e1002626.
651   22.   **Soper TS, Manning JM.** 1976. Synergy in the antimicrobial action of penicillin and beta-
652   chloro-D-alanine in vitro. Antimicrob Agents Chemother **9:**347-349.
653   23.   **Plata KB, Riosa S, Singh CR, Rosato RR, Rosato AE.** 2013. Targeting of PBP1 by beta-lactams
654   determines recA/SOS response activation in heterogeneous MRSA clinical strains. PLoS One
655   **8:**e61083.
656   24.   **Conlon KM, Humphreys H, O'Gara JP.** 2002. icaR encodes a transcriptional repressor
657   involved in environmental regulation of ica operon expression and biofilm formation in
658   Staphylococcus epidermidis. J Bacteriol **184:**4400-4408.
659   25.   **Rudkin JK, Edwards AM, Bowden MG, Brown EL, Pozzi C, Waters EM, Chan WC, Williams P,**
660   **O'Gara JP, Massey RC.** 2012. Methicillin resistance reduces the virulence of healthcare-
661   associated methicillin-resistant Staphylococcus aureus by interfering with the agr quorum
662   sensing system. J Infect Dis **205:**798-806.
663   26.   **Halsey CR, Lei S, Wax JK, Lehman MK, Nuxoll AS, Steinke L, Sadykov M, Powers R, Fey PD.**
664   2017. Amino Acid Catabolism in Staphylococcus aureus and the Function of Carbon
665   Catabolite Repression. MBio **8**.
666   27.   **Tabuchi F, Matsumoto Y, Ishii M, Tatsuno K, Okazaki M, Sato T, Moriya K, Sekimizu K.**
667   2017. D-cycloserine increases the effectiveness of vancomycin against vancomycin-highly
668   resistant Staphylococcus aureus. J Antibiot (Tokyo) **70:**907-910.
669   28.   **Fujihira T, Kanematsu S, Umino A, Yamamoto N, Nishikawa T.** 2007. Selective increase in
670   the extracellular D-serine contents by D-cycloserine in the rat medial frontal cortex.
671   Neurochem Int **51:**233-236.
672   29.   **Batson S, de Chiara C, Majce V, Lloyd AJ, Gobec S, Rea D, Fulop V, Thoroughgood CW,**
673   **Simmons KJ, Dowson CG, Fishwick CWG, de Carvalho LPS, Roper DI.** 2017. Inhibition of D-
674   Ala:D-Ala ligase through a phosphorylated form of the antibiotic D-cycloserine. Nat Commun
675   **8:**1939.
676   30.   **Chen J, Zhang S, Cui P, Shi W, Zhang W, Zhang Y.** 2017. Identification of novel mutations
677   associated with cycloserine resistance in Mycobacterium tuberculosis. J Antimicrob
678   Chemother doi:10.1093/jac/dkx316.

31. **Baisa G, Stabo NJ, Welch RA.** 2013. Characterization of Escherichia coli D-cycloserine
transport and resistant mutants. J Bacteriol **195:**1389-1399.
32. **Curtiss R, 3rd, Charamella LJ, Berg CM, Harris PE.** 1965. Kinetic and genetic analyses of D-
cycloserine inhibition and resistance in Escherichia coli. J Bacteriol **90:**1238-1250.
33. **Russell RR.** 1972. Mapping of a D-cycloserine resistance locus in escherichia coli K-12. J
Bacteriol **111:**622-624.
34. **Wargel RJ, Hadur CA, Neuhaus FC.** 1971. Mechanism of D-cycloserine action: transport
mutants for D-alanine, D-cycloserine, and glycine. J Bacteriol **105:**1028-1035.
35. **Feher T, Cseh B, Umenhoffer K, Karcagi I, Posfai G.** 2006. Characterization of cycA mutants
of Escherichia coli. An assay for measuring in vivo mutation rates. Mutat Res **595:**184-190.
36. **Prosser GA, Rodenburg A, Khoury H, de Chiara C, Howell S, Snijders AP, de Carvalho LP.**
2016. Glutamate Racemase Is the Primary Target of beta-Chloro-d-Alanine in
Mycobacterium tuberculosis. Antimicrob Agents Chemother **60:**6091-6099.
37. **Sobral RG, Ludovice AM, Gardete S, Tabei K, De Lencastre H, Tomasz A.** 2003. Normally
functioning murF is essential for the optimal expression of methicillin resistance in
Staphylococcus aureus. Microb Drug Resist **9:**231-241.
38. **Sobral RG, Ludovice AM, de Lencastre H, Tomasz A.** 2006. Role of murF in cell wall
biosynthesis: isolation and characterization of a murF conditional mutant of Staphylococcus
aureus. J Bacteriol **188:**2543-2553.
39. **de Jonge BL, Chang YS, Gage D, Tomasz A.** 1992. Peptidoglycan composition of a highly
methicillin-resistant Staphylococcus aureus strain. The role of penicillin binding protein 2A. J
Biol Chem **267:**11248-11254.
40. **Li C, Sun F, Cho H, Yelavarthi V, Sohn C, He C, Schneewind O, Bae T.** 2010. CcpA mediates
proline auxotrophy and is required for Staphylococcus aureus pathogenesis. J Bacteriol
192:3883-3892.
41. **Thurlow LR, Joshi GS, Richardson AR.** 2018. Peroxisome Proliferator-Activated Receptor
gamma Is Essential for the Resolution of Staphylococcus aureus Skin Infections. Cell Host
Microbe **24:**261-270 e264.
42. **Lehman MK, Nuxoll AS, Yamada KJ, Kielian T, Carson SD, Fey PD.** 2019. Protease-Mediated
Growth of Staphylococcus aureus on Host Proteins Is opp3 Dependent. MBio **10**.
43. **Hussain M, Hastings JG, White PJ.** 1991. A chemically defined medium for slime production
by coagulase-negative staphylococci. J Med Microbiol **34:**143-147.
44. **Alvarez L, Hernandez SB, de Pedro MA, Cava F.** 2016. Ultra-Sensitive, High-Resolution Liquid
Chromatography Methods for the High-Throughput Quantitative Analysis of Bacterial Cell
Wall Chemistry and Structure. Methods Mol Biol **1440:**11-27.
45. **De Jonge BL, Gage D, Xu N.** 2002. The carboxyl terminus of peptidoglycan stem peptides is a
determinant for methicillin resistance in Staphylococcus aureus. Antimicrob Agents
Chemother **46:**3151-3155.
46. **Kuhner D, Stahl M, Demircioglu DD, Bertsche U.** 2014. From cells to muropeptide structures
in 24 h: peptidoglycan mapping by UPLC-MS. Sci Rep **4:**7494.
47. **Boneca IG, Xu N, Gage DA, de Jonge BL, Tomasz A.** 1997. Structural characterization of an
abnormally cross-linked muropeptide dimer that is accumulated in the peptidoglycan of
methicillin- and cefotaxime-resistant mutants of Staphylococcus aureus. J Biol Chem
272:29053-29059.
48. **Diep BA, Gill SR, Chang RF, Phan TH, Chen JH, Davidson MG, Lin F, Lin J, Carleton HA,
Mongodin EF, Sensabaugh GF, Perdreau-Remington F.** 2006. Complete genome sequence of
USA300, an epidemic clone of community-acquired meticillin-resistant Staphylococcus
aureus. Lancet **367:**731-739.
49. **Gill SR, Fouts DE, Archer GL, Mongodin EF, Deboy RT, Ravel J, Paulsen IT, Kolonay JF,
Brinkac L, Beanan M, Dodson RJ, Daugherty SC, Madupu R, Angiuoli SV, Durkin AS, Haft
DH, Vamathevan J, Khouri H, Utterback T, Lee C, Dimitrov G, Jiang L, Qin H, Weidman J,**

Tran K, Kang K, Hance IR, Nelson KE, Fraser CM. 2005. Insights on evolution of virulence and
resistance from the complete genome analysis of an early methicillin-resistant
Staphylococcus aureus strain and a biofilm-producing methicillin-resistant Staphylococcus
epidermidis strain. J Bacteriol **187:**2426-2438.
50.   **Robinson DA, Enright MC.** 2003. Evolutionary models of the emergence of methicillin-
resistant Staphylococcus aureus. Antimicrob Agents Chemother **47:**3926-3934.
51.   **Horsburgh MJ, Foster TJ, Barth PT, Coggins JR.** 1996. Chorismate synthase from
Staphylococcus aureus. Microbiology-Uk **142:**2943-2950.
52.   **Salunkhe P, Smart CH, Morgan JA, Panagea S, Walshaw MJ, Hart CA, Geffers R, Tummler B,**
Winstanley C. 2005. A cystic fibrosis epidemic strain of Pseudomonas aeruginosa displays
enhanced virulence and antimicrobial resistance. J Bacteriol **187:**4908-4920.

Table 1. Antibacterial activity (minimum inhibitory concentrations, MIC) and drug synergy (fractional inhibitory concentration indices, ΣFIC) of D-cycloserine (DCS) and several β-lactam antibiotics with different PBP specificity, namely oxacillin (OX; PBP1, 2, 3), nafcillin (NAF; PBP1), cefoxitin (FOX; PBP4), cefaclor (CEC; PBP3), cefotaxime (CTX; PBP2), piperacillin-tazobactin (TZP; PBP3/β-lactamase inhibitor) and imipenem (IMP; PBP1), alone and in β-lactam combinations, against fourteen *S. aureus* strains and *S. epidermidis* RP62A.

Antibiotic → ↓ Strain	OX*	NAF*	DCS*	FOX*	DCS/FOX** (ΣFIC)***	CEC*	DCS/CEC** (ΣFIC)***	CTX*	DCS/CTX** (ΣFIC)***	TZP*	DCS/TZP** (ΣFIC)***	IPM*	DCS/IPM** (ΣFIC)***
JE2	64	32	32	64	8/8 (0.37)	64	8/2 (0.28)	64	8/4 (0.31)	32	8/0.5 (0.26)	1	ND
NE810 (*cycA*)	0.25	0.5	2	8	ND	4	ND	8	ND	2	ND	0.125	ND
USA300	64	32	32	64	8/8 (0.37)	64	8/8 (0.37)	64	8/8 (0.37)	64	8/2 (0.28)	1	ND
USA300 *cycA*	0.25	1	2	8	ND	4	ND	8	ND	4	ND	0.125	ND
DAR173	128	128	32	256	4/32 (0.25)	128	8/4 (0.28)	512	8/4 (0.25)	128	4/4 (0.15)	64	4/2 (0.15)
DAR173 *cycA*	0.5	8	4	32	ND	16	ND	16	ND	4	ND	1	ND
DAR22	128	128	32	128	8/8 (0.31)	128	8/8 (0.31)	512	8/4 (0.25)	128	4/4 (0.15)	128	4/1 (0.13)
DAR22 *cycA*	0.5	8	0.5	16	ND	16	ND	16	ND	4	ND	0.5	ND
DAR169	32	32	32	32	8/2 (0.31)	128	4/32 (0.37)	64	4/8 (0.25)	8	4/1 (0.25)	2	ND
DAR169 *cycA*	16	4	0.5	16	ND	64	ND	16	ND	2	ND	0.5	ND
COL	512	256	64	512	8/128 (0.37)	128	16/32 (0.5)	>2048	ND	128	32/0.5 (0.5)	256	8/64 (0.37)
DAR113	128	64	32	128	8/8 (0.3)	64	8/4 (0.3)	256	8/8 (0.2)	128	8/4 (0.2)	16	8/0.5 (0.2)
BH1CC	256	512	32	256	8/32 (0.3)	128	8/8 (0.3)	1028	8/16 (0.2)	256	8/4 (0.2)	64	8/1 (0.2)
BH14B(04)	128	64	16	128	4/32 (0.5)	128	4/32 (0.5)	512	4/32 (0.31)	64	4/1 (0.26)	64	2/8 (0.25)
BH8(03)	128	128	32	128	8/32 (0.5)	256	8/64 (0.5)	128	8/16 (0.25)	128	8/8 (0.3)	64	8/0.5 (0.1)
BH6	128	128	32	128	8/16 (0.37)	256	8/64 (0.5)	512	8/32 (0.31)	128	8/4 (0.28)	32	8/0.5 (0.26)
DAR202	64	64	32	128	4/32 (0.37)	128	4/32 (0.37)	64	8/16 (0.5)	64	8/2 (0.28)	4	8/1 (0.5)
DAR45	2	0.5	32	4	ND	32	8/2 (0.31)	4	ND	2	ND	0.5	ND
DAR13	128	32	32	128	4/32 (0.37)	128	8/4 (0.28)	128	4/32 (0.37)	32	8/1 (0.28)	8	4/2 (0.37)
RP62A	128	2	32	64	6/16 (0.5)	64	6/16 (0.5)	32	8/8 (0.5)	8	8/2 (0.5)	32	8/1 (0.2)

* MIC values for each antibiotic when measured individually; μg/ml

** MIC values for each antibiotic when measured in combination, also known as the fractional inhibitory concentration (FIC); μg/ml

*** FIC indices (ΣFIC) for the antibiotic combination. ΣFIC = FIC A + FIC B, where FIC A is the MIC of DCS in combination with the β-lactam/MIC of DCS alone, and FIC B is the MIC of the β-lactam in combination with DCS/MIC of the β-lactam alone. The combination is considered synergistic when the ΣFIC is ≤0.5, indifferent when the ΣFIC is >0.5 to <2.

‡ND. Not determined if strain is susceptible (or hyper-resistant) to β-lactam antibiotic or for *cycA* mutants with reduced DCS & β-lactam MICs.

Table 2. Antibacterial activity (minimum inhibitory concentrations, MIC) and drug synergy (fractional inhibitory concentration indices, ΣFIC) of D-cycloserine (DCS) and the β-lactam antibiotics piperacillin-tazobactin (TZP) and imipenem (IMP), alone and in β-lactam/DCS combinations, against selected strains of *Pseudomonas aeruginosa*, vancomycin resistant *Enterococcus, Klebsiella pneumoniae and Acinetobacter baumanii*

Strain	DCS*	TZP*	DCS/TZP** (ΣFIC)***	IMP*	DCS/IPM** (ΣFIC)***
Pseudomonas aeruginosa LES431	256	128	64/8 (0.312)	16	32/4 (0.375)
Vancomycin resistant *Enterococcus* CIP104105	64	512	32/1 (0.502)	256	16/16 (0.5)
Vancomycin resistant *Enterococcus* DSM17050	64	256	16/64 (0.5)	128	16/1 (0.257)
Klebsiella pneumoniae 511232.1 (ESBL[+])	128	32	8/32 (0.5)	1	ND
Acinetobacter baumanii DSM30007	128	32	32/8 (0.5)	1	ND

* MIC values for each antibiotic when measured individually; μg/ml

** MIC values for each antibiotic when measured in combination, also known as the fractional inhibitory concentration (FIC); μg/ml

*** FIC indices (ΣFIC) for the antibiotic combination. ΣFIC = FIC A + FIC B, where FIC A is the MIC of DCS in combination with the β-lactam/MIC of DCS alone, and FIC B is the MIC of the β-lactam in combination with DCS/MIC of the β-lactam alone. The combination is considered synergistic when the ΣFIC is ≤0.5 and indifferent when the ΣFIC is >0.5 to <2. Green shading indicates synergy, orange shading indicates no synergy.

[‡]ND. Not determined where the bacterial strain is susceptible to the β-lactam antibiotic.

Supplementary Tables

Table S1. Antibacterial activity (minimum inhibitory concentrations, MIC) and drug synergy (fractional inhibitory concentration indices, ΣFIC) of D-cycloserine (DCS), and clindamycin (CLI), tobramycin (TOB), trimethoprim (TMP), mupirocin (MUP), ciprofloxacin (CIP), kanamycin (KAN) and spectinomycin (SPT) alone and in DCS/antibiotic combinations, against fourteen *S. aureus* strains and *S. epidermidis* RP62A.

Antibiotic → ↓ Strain	DCS*	CLI*	DCS/CLI** (ΣFIC)***	TOB	DCS/TOB (ΣFIC)	TMP	DCS/TMP (ΣFIC)	CIP	DCS/CIP (ΣFIC)	KAN	DCS/KAN (ΣFIC)	SPT	DCS/SPT (ΣFIC)	MUP	DCS/MUP (ΣFIC)
JE2	32	0.25	ND	1	ND	1	ND	32	16/16 (1)	2	ND	64	16/32 (1)	0.25	ND
USA300	32	2048	32/2048 (2)	0.5	ND	1	ND	32	16/16 (1)	2	ND	64	16/32 (1)	2048	32/2048 (2)
DAR173	32	0.5	ND	512	32/512 (2)	0.5	ND	256	16/128 (1)	512	16/256 (1)	64	16/32 (1)	0.5	ND
DAR22	32	0.5	ND	512	16/256 (1)	0.5	ND	256	16/128 (1)	256	16/128 (1)	64	16/32 (1)	0.5	ND
DAR169	32	0.5	ND	0.5	ND	0.5	4/8 (0.25)	1	ND	2	ND	64	16/32 (1)	0.5	ND
COL	64	0.5	ND	0.5	ND	0.5	ND	1	ND	0.5	ND	64	16/32 (1)	0.5	ND
DAR113	32	0.25	ND	0.5	ND	1	ND	0.5	ND	2	ND	64	16/32 (1)	0.25	ND
BH1CC	32	1024	32/1024 (2)	32	32/32 (2)	256	32/256 (2)	32	16/16 (1)	>1024	ND	>1024	ND	32	16/8 (0.75)
BH14B(04)	16	0.5	ND	0.5	ND	0.5	ND	128	16/128 (1.5)	2	ND	64	8/32 (1)	0.5	ND
BH8(03)	32	1024	32/1024 (2)	0.5	ND	0.5	ND	256	16/128 (1)	8	32/4 (2)	128	16/64 (1)	4	32/4 (2)
BH6	32	0.5	ND	0.5	ND	0.5	ND	>512	16/256 (1)	1	ND	128	16/64 (1)	0.5	ND
DAR202	32	0.5	ND	128	16/64 (1)	0.5	ND	2	ND	>1024	ND	>1024	ND	0.5	ND
DAR45	32	1024	32/1024 (2)	512	16/256 (1)	0.5	ND	128	16/64 (1)	128	16/64 (1)	512	32/1024 (2)	0.5	ND
DAR13	32	0.5	ND	0.5	ND	32	16/16 (1)	32	16/16 (1)	2048	ND	>1024	ND	0.5	ND
RP62A	32	2048	32/2048 (2)	16	16/8 (1)	256	32/256 (2)	0.25	ND	>1024	ND	>1024	ND	0.25	ND

 * MIC values for each antibiotic when measured individually; μg/ml

** MIC values for each antibiotic when measured in combination, also known as the fractional inhibitory concentration (FIC); μg/ml

*** FIC indices (ΣFIC) for the antibiotic combination. ΣFIC = FIC A + FIC B, where FIC A is the MIC of DCS in combination with the antibiotic/MIC of DCS alone, and FIC B is the MIC of the antibiotic in combination with DCS/MIC of the antibiotic alone. The combination is considered synergistic when the ΣFIC is ≤0.5, indifferent when the ΣFIC is >0.5 to <2.

‡ND. Not determined if strain is susceptible (or hyper-resistant) to the antibiotic or for *cycA* mutants with reduced DCS & antibiotic MICs.

Table S2. Cell diameter and cell wall thickness of JE2 and NE810 (*cycA* mutant).

	Mean cell diameter (µm) ± SD	Mean cell wall thickness (nm) ± SD
JE2	0.89 ± 0.075	25.92 ± 1.79
NE810	0.88 ± 0.05	18.95 ± 2.5

Table S3. Bacterial strains and plasmids used in this study

Strains/plasmids	Relevant Details
RN4220	Restriction-deficient laboratory *S. aureus*
USA300 FPR3757	Community associated MRSA isolate of the USA300 lineage (48). SCCmec type IV. CC8.
JE2	USA300 cured of p01 & p03. Parent of Nebraska Transposon Mutant Library (NTML).
13C	USA300 derivative cured of plasmid p03. Erm[s].
NE810	JE2 NTML *cycA* (SAUSA300_1642) mutation. Erm[r].
USA300 *cycA*	USA300 FPR3757 *cycA*. Constructed by transduction of *cycA*::Tn allele from NE810.
13C *cycA*	13C *cycA* mutant (*cycA*::Tn allele from NE810).
NE1868	JE2 NTML *mecA* mutation.
NE1713	JE2 NTML *alr* (SAUSA300_2027) mutation.
NE799	JE2 NTML *alr2* (SAUSA300_1292) mutation.
NE923	JE2 NTML *ynfA* (SAUSA300_2286) mutation.
NE1025	JE2 NTML *thrE* (SAUSA300_0729) mutation.
BH1CC	MRSA clinical isolate; SCC*mec* type II; CC8 (17)
COL	MRSA reference strain; SCC*mec* type I; CC8 (49)
BH14(04)	MRSA clinical isolate; SCC*mec* type IV; CC22 (17)
BH8(03)	MRSA clinical isolate; SCC*mec* type IV; CC22 (17)
BH6(03)	MRSA clinical isolate; SCC*mec* type II; CC8 (17)
DAR113	MRSA reference isolate; SCC*mec* type IV; CC22 (17, 50)
DAR13	MRSA reference isolate; SCC*mec* type IV; CC8 (17, 50)
DAR45	MRSA reference isolate; SCC*mec* type II; CC30 (17, 50)
DAR202	MRSA reference isolate; SCC*mec* type III; CC239 (17, 50)
DAR173	MRSA reference isolate; SCC*mec* type IV; CC5 (17, 50)
DAR173 *cycA*	DAR173 *cycA* mutant (*cycA*::Tn allele from NE810)
DAR22	MRSA reference isolate; SCC*mec* type III; CC5 (17, 50)
DAR22 *cycA*	DAR22 *cycA* mutant (*cycA*::Tn allele from NE810)
DAR169	MRSA Reference strain; SCCmec type I; CC8 (17, 50)
DAR169 *cycA*	DAR169 *cycA* mutant (*cycA*::Tn allele from NE810)
8325-4	NCTC 8325 derivative cured of prophages (51), methicillin susceptible, CC8.
8325-4 *cycA*	8325-4 *cycA* mutant (*cycA*::Tn allele from NE810).
ATCC 29213	Methicillin susceptible *S. aureus* strain for antibiotic susceptibility testing.
ATCC 29213 *cycA*	ATCC 29213 *cycA* mutant (*cycA*::Tn allele from NE810).
ATCC 25923	Methicillin susceptible *S. aureus* strain for antibiotic susceptibility testing.
S. epidermidis RP62A	ATCC 35984. Methicillin resistant, biofilm positive. (24)
E. coli	*E. coli* HST08
Vancomycin resistant *Enterococcus* CIP104105	Reference strain
Vancomycin resistant *Enterococcus* DSM17050	Reference strain
Acinetobacter baumanii DSM 30007	Reference strain
Klebsiella pneumoniae 511232.1	Clinical Isolate (Dublin Hospital).
Pseudomonas aeruginosa LES431	Liverpool epidemic strain (LES) of *Pseudomonas aeruginosa* (52).
Plasmids	
pLI50	*E. coli-Staphylococcus* shuttle vector. Ap[r] (*E. coli*), Cm[r]. (*Staphylococcus*)
p*cycA*	pLI50 carrying *cycA* from JE2

Table S4. Oligonucleotide primers used in this study

Target Gene	Primer Name	Primer Sequence (5'-3')
cycA	NE810_Fwd	ACAGAATAGCCACAAATAGCACC
	NE810_Rev	ACAGAATAGCCACAAATAGCACC
cycA	NE810F1_Fwd	GTCTTCAAGAATTCGGCCACAAATAGCACCATTAA
	NE810F1_Rev	CGACTCTAGAGGATCATGTCCCAAGCCCTAAAAC
mecA	mecA1_Fwd	TGCTCAATATAAAATTAAAACAAACTACGGTAAC
	mecA1_Rev	GAATAATGACGCTATGATCCCAA
gyrB	gyrB_Fwd	CCAGGTAAATTAGCCGATTGC
	gyrB_Rev	AAATCGCCTGCGTTCTAGAG
	RNAIII FWD	GAAGGAGTGATTTCAATGGCACAAG
RNAIII	RNAIII REV	GAAAGTAATTAATTATTCATCTTATTTTTTAGTGAATTTG

A

B

| JE2 | NE810 | NE810 pLI50 | NE810 pcycA |

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Kidneys

Fig. 7

Fig. 8

Figure S1. A. Susceptibility of JE2 (wild type), NE1868 (*mecA*::Tn), NE810 (*cycA*::Tn), and NE1713 (*alr*::Tn) grown on MH agar to cefoxitin (FOX, 30μg disks). **B.** Comparison of relative *mecA* gene expression by LightCycler RT-PCR in JE2 and NE810 grown to A_{600}=3 in BHI media. The data are the average of three independent experiments and standard deviations are shown. **C.** Susceptibility of JE2 (wild type) and NE1713 (*alr*::Tn) grown on MH agar to D-cycloserine (DCS, 30μg disks).

Figure S2. Amino acid consumption by JE2 and NE810 grown aerobically in chemically defined media containing 14mM of glucose (CDMG). Residual amino acids were measured in spent media after 0, 4, 6 and 8 h growth.

Figure S3. Preparation of cell suspensions for peptidoglycan extraction and structural analysis by UPLC-MS. 50 ml flask cultures were inoculated into fresh BHI media from overnight cultures at a starting cell density of A_{600}=0.05 and incubated at 37^{o}C. The number of CFU/ml was enumerated every 1-2 h for 6 h and again after 24 hours. For JE2 cultures being dosed with DCS, the antibiotic was added after approximated 2 h (A_{600}≈0.5) and the cells collected after a further 100 mins. Cells from untreated JE2 and NE810 control cultures were collected at the same time point.

Figure S4. *In vitro* kill curves for D-cycloserine (DCS), oxacillin and cefoxitin with methicillin resistant *S. epidermidis strain* RP62A. Antibiotics at the concentrations indicated (equivalent to 0.5× MIC) were added to suspensions of overnight bacterial cultures adjusted to 10^7 CFU/ml in BHI, incubated at 37°C and the number of CFU/ml enumerated at 0, 2, 4, 8 and 24 h. The data presented are the mean of three independent experiments. Antibiotic synergism was defined as a ≥2 $\log^{10}$ decrease in the number of CFU/ml in cell suspensions exposed to DCS/β-lactam combinations compared to the most effective individual antibiotic alone.

Figure S5. A. Zones of haemolysis of JE2 and NE810 grown on sheep blood agar for 24 hours. **B**. Comparison of relative *agr (RNAIII)* gene expression by LightCycler RT-PCR in JE2 and NE810 grown for 3, 4, 5 and 6 h in BHI media. The data are the average of at least three independent experiments and standard deviations are shown. **C.** Growth of JE2 and NE810 cultures for 6 h in BHI. Cell density was measured at A_{600}.

Figure S6. Combination therapy with D-cycloserine and oxacillin significantly reduces the bacterial burden in the spleen of mice infected with MRSA. The number of colony-forming units (CFU) recovered from the spleens of mice infected by tail vein injection with 5×10^6 JE2 or NE810 (CycA) and left untreated or treated with 75mg of oxacillin (Ox)/kg, 30mg of DCS/kg or a combination of both Ox and DCS delivered subcutaneously every 12 hours for 5 days. The first antibiotic dose was given 16 hours after infection. Significant differences determined using one-way ANOVA with Kruskal-Wallis test followed by Dunn's multiple comparisons test are denoted using asterisks (*$p \leq 0.05$). The limit of detection (50 colonies) is indicated with a hashed red line.

Figure S7. Amino acid consumption by JE2 and NE810 grown aerobically in chemically defined media lacking glucose(CDM). Residual amino acids were measured in spent media after 0, 4, 6 and 8 h growth.