Impaired alanine transport or exposure to D-cycloserine increases the susceptibility of

MRSA to β-lactam antibiotics

Laura A. Gallagher[1], Rebecca K. Shears[2], Claire Fingleton[1], Laura Alvarez[3], Elaine M.

Waters[1,2], Jenny Clarke[2], Laura Bricio-Moreno[2], Christopher Campbell[1], Akhilesh K. Yadav[3],

Fareha Razvi[4], Eoghan O'Neill[5], Alex J. O'Neill[6], Felipe Cava[3], Paul D. Fey[4], Aras Kadioglu[2] and

James P. O'Gara[1]*

[1]School of Natural Sciences, National University of Ireland, Galway, Ireland.

[2]Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and
Global Health, University of Liverpool, UK.

[3]MIMS-Molecular Infection Medicine Sweden, Molecular Biology Department, Umeå
University, Umeå, Sweden.

[4]Department of Pathology and Microbiology, University of Nebraska Medical Center,
Omaha, Nebraska, USA.

[5]Department of Clinical Microbiology, Royal College of Surgeons in Ireland, Connolly
Hospital, Dublin 15, Ireland.

[6]Antimicrobial Research Centre, School of Molecular and Cellular Biology, Faculty of
Biological Sciences, University of Leeds, Leeds, UK.

*Correspondence: jamesp.ogara@nuigalway.ie

Running title: Re-sensitization of MRSA to β-lactams

Abstract. Prolonging the clinical effectiveness of β-lactams, which remain first-line antibiotics for many infections, is an important part of efforts to address antimicrobial resistance. We report here that inactivation of the predicted D-cycloserine (DCS) transporter gene *cycA* re-sensitized MRSA to β-lactam antibiotics. The *cycA* mutation also resulted in hyper-susceptibility to DCS, an alanine analogue antibiotic that inhibits alanine racemase and D-alanine ligase required for D-alanine incorporation into cell wall peptidoglycan (PG). CycA functions as an alanine permease, and impaired alanine transport correlated with increased susceptibility to oxacillin and DCS. The *cycA* mutation or exposure to DCS were both associated with the accumulation of muropeptides with tripeptide stems lacking the terminal D-ala-D-ala and reduced PG crosslinking, prompting us to investigate synergism between β-lactams and DCS. DCS re-sensitised MRSA to β-lactams *in vitro* and significantly enhanced MRSA eradication by oxacillin in a mouse bacteraemia model. These findings reveal alanine transport as a new therapeutic target to enhance the susceptibility of MRSA to β-lactam antibiotics.

Abstract word count: 161

Introduction

Whilst many bacteria can exhibit resistance to select antimicrobials, isolates of the human

pathogen *Staphylococcus aureus* can express resistance to all licensed anti-staphylococcal

drugs. This results in significant morbidity and mortality, with up to 20% of patients with

systemic methicillin resistant *S. aureus* (MRSA) infections dying, despite receiving treatment

with anti-staphylococcal drugs [1]. As part of our efforts to identify improved therapeutic

approaches for MRSA infections, we recently described the novel use of β-lactam antibiotics

to attenuate the virulence of MRSA-induced invasive pneumonia and sepsis [2]. We

demonstrated that oxacillin-induced repression of the Agr quorum-sensing system and

altered cell wall architecture resulted in downregulated toxin production and increased MRSA

killing by phagocytic cells, respectively [2]. Supporting this *in vitro* data, a randomised

controlled trial involving 60 patients showed that the β-lactam antibiotic flucloxacillin in

combination with vancomycin shortened the duration of MRSA bacteraemia from 3 days to

1.9 days [3, 4].

Because expression of methicillin resistance in *S. aureus* impacts fitness and virulence and is

a regulated phenotype, further therapeutic interventions may also be possible. The

complexity of the methicillin resistance phenotype is evident among clinical isolates of MRSA,

which express either low-level, heterogeneous (HeR) or homogeneous, high-level methicillin

resistance (HoR) [5-7]. Exposure of HeR isolates to β-lactam antibiotics induces expression of

mecA, which encodes the alternative penicillin binding protein 2a (PBP2a) and can select for

mutations in accessory genes resulting in a HoR phenotype, including mutations that affect

the stringent response and c-di-AMP signalling [8-12]. Because accessory genes can influence

the expression of methicillin resistance in MRSA, targeting the pathways associated with such

genes may identify new ways to increase the susceptibility of MRSA to β-lactams. To pursue

this, we performed a forward genetic screen to identify loci that impact the expression of

resistance to β-lactam antibiotics in MRSA. Using the Nebraska Transposon Mutant Library,

which comprises 1,952 sequence-defined transposon insertion mutants [13], inactivation of

a putative amino acid permease gene, *cycA*, was found to reduce resistance to cefoxitin, the

β-lactam drug recommended by the Clinical and Laboratory Standards Institute for measuring

mecA-mediated methicillin resistance in MRSA isolates. Amino acid transport and

susceptibility to oxacillin and D-cycloserine (DCS) were compared in the wild-type and *cycA*

mutant grown in chemically defined media (CDM), CDM supplemented with glucose (CDMG)

and other complex media. The activity of DCS and β-lactams, alone and in combination,

against MRSA was measured *in vitro* and in a mouse model of bacteraemia. Peptidoglycan

analysis was performed to compare the impact of the *cycA* mutation or exposure to DCS on

cell wall structure and crosslinking. Our experiments suggest that therapeutic strategies

targeting alanine transport, which was required for resistance to β-lactams and DCS, and a re-

evaluation of DCS may be important as part of efforts to restore the efficacy of β-lactam

antibiotics against MRSA.

     Results

     Mutation of *cycA* increases the susceptibility of MRSA to β-lactam antibiotics and D-

     cycloserine. To identify new ways of controlling expression of methicillin resistance, we

     sought to identify novel mutations involved in this phenotype. An unbiased screen of the

     NTML to identify mutants with increased susceptibility to cefoxitin identified NE810

     (SAUSA300_1642) (Fig. S1A), which also exhibited a >500-fold increase in susceptibility to

     oxacillin (Fig. S1B). NE810 was previously identified among several NTML mutants reported

     to be more susceptible to amoxicillin [14], but was not investigated further. Expression of

     mecA was not affected in NE810 and genome sequence analysis revealed an intact SCC*mec*

     element and the absence of any other mutations (data not shown). NE810 was successfully

     complemented (Fig. S1B), and transduction of the SAUSA300_1642 allele into several MRSA

     strains from a number of clonal complexes and with different SCC*mec* types was also

     accompanied by increased cefoxitin and oxacillin susceptibility (Table 1).

     SAUSA300_1642 is annotated as a serine/alanine/glycine transporter with homology to CycA

     in *Mycobacterium tuberculosis* [15, 16] required for D-cycloserine (DCS) resistance in

     Mycobacteria [15, 16]. In contrast to the observations in *Mycobacteria*, our data showed that

     NE810 and several unrelated MRSA strains carrying the *cycA* mutation were significantly more

     susceptible to DCS than the wild type JE2 (Fig. S1C, Table 1). The *cycA* mutation also reduced

     the DCS MIC of the MSSA strains 8325-4 and ATCC29213 from 32 to 4 μg/ml. DCS inhibits

    alanine racemase (Alr) that converts L-alanine to D-alanine and the Ddl D-alanyl:D-alanine

    ligase [17]. A mutant in the putative *ddl* SAUSA300_2039 gene is not available in the NTML

    library, suggesting that it may be essential. However, the *alr* mutant NE1713 was significantly

    more susceptible to cefoxitin and DCS (data not shown), consistent with an important role for

    D-alanine in resistance to both antibiotics.

    CycA is an alanine permease required for D-ala-D-ala incorporation into the peptidoglycan

    stem peptide. To investigate the role of CycA as a potential permease, JE2 and NE810 were

    grown for 8 h in chemically defined media containing 14mM glucose (CDMG) and amino acid

    consumption in spent media was measured. Although no growth rate or yield difference was

    noted between JE2 and NE810 in CDMG (Fig. 1A), alanine uptake by NE810 was significantly

    impaired compared to JE2 (Fig. 2B). Utilisation of other amino acids by NE810 and JE2,

    including serine and glycine, were similar (Fig. S2). Impaired alanine transport in the *cycA*

mutant grown in CDMG correlated with increased susceptibility to oxacillin (1 μg/ml) (Fig. 1C)

and DCS (1 μg/ml) (Fig. 1D). These data demonstrate for the first time that CycA in *S. aureus*

functions as an alanine permease.

Quantitative peptidoglycan compositional analysis was performed using UPLC analysis of

muramidase-digested muropeptide fragments extracted from exponential phase cultures of

JE2 and NE810 grown for 220 mins in TSB media (Fig. S3). The PG profile of the *cycA* mutant

revealed a significant accumulation of tripeptides compared to wild-type JE2 (Fig. 2A,B),

which was associated with a significant reduction in crosslinking (Fig. 2C). In NE810, the dimer,

trimer and tetramer fractions were decreased, which was accompanied by a concomitant

increase in the monomer fraction (Fig. 2D). Consistent with this data, exposure of JE2 to DCS

8μg/ml was also associated with a similar accumulation in muropeptides with tripeptide

stems (Fig. 2B), reduced cross-linking (Fig. 2C), increased muropeptide monomers and

reduced dimers, trimers and tetramers (Fig. 2D). DCS had a strong dose-dependent effect on

the accumulation of muropeptides with tripeptide stems, reduced cross-linking and

accumulation of monomers (Fig. 2A-D). Sub-inhibitory (0.25× MIC) and 4× MIC DCS

concentrations, were previously shown to be associated with incorporation of an incomplete

stem peptide (tripeptide) [17] and reduced D-ala-D-ala levels [18], respectively. These data

show that impaired D-ala incorporation in the *cycA* mutant or following exposure to DCS is

accompanied by reduced PG cross-linking and increased β-lactam susceptibility.

Mutation of *cycA* or exposure to D-cycloserine increases the susceptibility of MRSA to β-

lactam antibiotics. Previously reported synergy between DCS and β-lactam antibiotics [17,

19] suggests that impaired alanine uptake in the *cycA* mutant may have the same impact on

cell wall biosynthesis as DCS-mediated inhibition of Alr and Ddl. To further investigate this,

we compared the activity of DCS and β-lactam antibiotics, alone and in combination, against

JE2 and NE810. Checkerboard microdilution assay fractional inhibitory concentration indices

(ΣFICs ≤0.5) revealed synergy between DCS and several licensed β-lactam antibiotics with

different PBP selectivity against JE2 and USA300 FPR3757 (Table 1). Oxacillin and nafcillin

were not included in checkerboard assays because measurement of their MICs involves

supplementing the media with 2% NaCl, which distorts the MIC of DCS (data not shown).

Using the MRSA strains JE2, USA300, DAR173, DAR22, DAR169 and their corresponding *cycA*

mutants, the kinetics of killing by DCS, oxacillin and cefoxitin, alone and in combination was

measured over 24h using antibiotic concentrations corresponding to 0.125×, 0.25× and 0.5×

MICs. Recovery of growth in media supplemented with oxacillin or cefoxitin alone was evident

after 8 h (Fig. 3), reflecting the selection and expansion of HoR mutants as described

previously [2, 18, 20]. Recovery of growth in cultures exposed to DCS alone was also evident

(Fig. 3), which may correlate with our observation that mutants resistant to DCS (on BHI agar

supplemented with 128 µg/ml DCS) arise at a rate of approximately 5.5×10^{-8} per cell per

generation (data not shown). Using combinations of DCS and oxacillin or cefoxitin at 0.125×

MIC did not achieve a $\geq 2 \log^{10}$ reduction in the number of CFU/ml (data not shown). However,

at 0.5× MIC for strains JE2, USA300, DAR173 and DAR22, DCS (16 µg/ml)/oxacillin (32 µg/ml)

and DCS (16 µg/ml)/cefoxitin (32 µg/ml) combinations achieved a $\geq 5 \log^{10}$ reduction in the

number of CFU/ml compared to oxacillin, cefoxitin or DCS alone (Fig. 3). For strain DAR169,

DCS/β-lactam combinations at 0.25× MIC was sufficient to achieve a $\geq 5 \log^{10}$ reduction in

CFUs recovered compared to the individual antibiotics (Fig. 3). DCS/β-lactam combinations at

0.5× MIC were also able to achieve $\geq 5 \log^{10}$ reduction in the number of CFU/ml against the

methicillin resistant *S. epidermidis* (MRSE) strain RP62A [21] compared to either antibiotic

alone (Fig. S4). Checkerboard experiments with fourteen MRSA strains and MRSE strain

RP62A further revealed synergy (ΣFICs ≤0.5) between DCS and a range of β-lactam antibiotics

with different penicillin binding protein (PBP) specificity, namely cefoxitin (PBP4), cefaclor

(PBP3), cefotaxime (PBP2), piperacillin-tazobactin (PBP3/β-lactamase inhibitor) and

imipenem (PBP1) (Table 1).

This synergy appears to be specific to β-lactams and no synergy (ΣFICs > 0.5) was measured

between DCS and several antibiotics that are used topically or systemically for the

decolonization or treatment of patients colonized/infected with *S. aureus* or MRSA

(clindamycin, trimethoprim, mupirocin, ciprofloxacin), or several antibiotics to which *S.*

aureus isolates commonly exhibit resistance (tobramycin, kanamycin and spectinomycin)

(Table S1). Furthermore the *cycA* mutation had no impact on susceptibility to any of these

non-β-lactam antibiotics (apart from *ermB*-encoded clindamycin resistance on the

transposon).

Combination therapy with DCS and oxacillin significantly reduces the bacterial burden in

the kidneys and spleen of mice infected with MRSA. The virulence of the NE810 mutant and

the therapeutic potential of oxacillin in combination with DCS in the treatment of MRSA

infections were assessed in mice. Treatment with oxacillin or DCS alone significantly reduced the number of CFUs recovered from the kidneys of mice infected with JE2 (Fig. 4). Furthermore the oxacillin/DCS combination was significantly more effective than either antibiotic alone and the combination was equally effective in reducing the bacterial burden in the kidneys of animals infected with JE2 or NE810 when compared to no treatment (p≤0.0001) (Fig. 4) demonstrating the capacity of DCS to significantly potentiate the activity of β-lactam antibiotics against MRSA under *in vivo* conditions. Unexpectedly, oxacillin- or DCS-mediated eradication of NE810 infections in the kidneys was similar to JE2 (Fig. 4). In the spleen, only oxacillin/DCS combination treatment was associated with a significant reduction in the number of CFUs recovered from mice infected with JE2 or NE810 (Fig. S5).

Alanine transport and resistance to oxacillin and DCS in chemically defined medium are not dependent on *cycA*. The failure of oxacillin or DCS treatment to enhance the eradication of NE810 infections in the mouse bacteraemia model prompted us to further characterise the growth conditions used for the *in vitro* antibiotic susceptibility assays. Specifically we investigated the role of glucose, which we previously reported to increase the growth requirement for amino acids [22], and which we reasoned may be important for CycA alanine permease activity. Growth of JE2 and NE810 was similar in CDM lacking glucose (Fig. 5A) and uptake of alanine (Fig. 5B) and other amino acids (Fig. S6) was unchanged in NE810 compared to JE2. Furthermore NE810 and JE2 grew equally well in CDM supplemented with oxacillin and DSC (Fig. 5C and D). These data explain in part why the *cycA* mutant does not exhibit increased β-lactam and DCS susceptibility in the mouse bacteraemia model and further reveal the strong correlation between alanine transport and susceptibility to oxacillin and DCS.

Discussion

The exploitation of antibiotic re-purposing as part of concerted efforts to address the antimicrobial resistance crisis has been hampered by a lack of mechanistic data to explain demonstrated therapeutic potential and the perception that studies attempting to identify new uses for existing drugs are not hypothesis-driven. In this study, we identified an alanine permease (CycA) required for full expression of resistance to β-lactam antibiotics and DCS. Loss of function of this alanine transporter significantly increased the susceptibility of MRSA to β-lactam antibiotics, an outcome that could be reproduced through exposure to DCS, which targets the Alr and Ddl enzymes in the early steps of cell wall biosynthesis.

The potential of β-lactam/DCS combinations for treatment of MRSA infections follows a recent report that DCS can also potentiate the activity of vancomycin against a laboratory-generated vancomycin highly-resistant *S. aureus* (VRSA) strain *in vitro* and in a silkworm infection model [23]. The excellent safety profile of β-lactam antibiotics makes these drugs particularly attractive as components of combination antimicrobial therapies. When used in the treatment of tuberculosis DCS (trade name Seromycin, The Chao Centre) is typically administered orally in 250 mg tablets twice daily for up to two years. At this dosage, the DCS concentration in blood serum is generally 25-30 μg/ml, which is similar to the concentrations used in our *in vitro* and *in vivo* experiments. The known neurological side effects associated with DCS therapy [24, 25] mean that this antibiotic is unlikely to be considered for the treatment of MRSA infections unless alternative therapeutic approaches have been exhausted. Oxacillin/DCS combination therapy was significantly more effective than DCS or oxacillin alone over a 5-day therapeutic window suggesting that further studies on using DCS to augment β-lactams as a treatment option for recalcitrant staphylococcal infections is merited.

Mutation of *cycA* increases the susceptibility of MRSA to β-lactam antibiotics and results in hyper-susceptibility to D-cycloserine, whereas a *cycA* point mutation in *M. bovis* contributes, in part, to increased DCS resistance presumably by interfering with transport into the cell [16]. In *E. coli*, *cycA* mutations can also result in increased resistance or have no effect on DCS susceptibility depending on the growth media [26-30], suggesting that CycA is primarily important for DCS resistance under conditions when its contribution to amino acid transport

is also important. Our data showing that mutation of *cycA* was not associated with increased DCS resistance strongly suggests that CycA has no role in uptake of this antibiotic in *S. aureus*. Under growth conditions where CycA is required for alanine transport (in nutrient/glucose-replete media), mutation of *cycA* or DCS-exposure have similar effects on the structure of *S. aureus* peptidoglycan (Fig. 8). Consistent with previous studies in *S. aureus* [17] and in *M. tuberculosis* [31], our studies showed a dose-dependent accumulation of muropeptides with a tripeptide stem in MRSA exposed to DCS. The *cycA* mutation was also associated with the increased accumulation of muropeptides with a tripeptide stem. These data indicate that a reduced intracellular alanine pool or inhibition of Alr and Ddl is associated with reduced D-ala-D-ala incorporation into the PG stem peptide. The increased accumulation of tripeptides in turn interferes with normal PBP transpeptidase activity and offers a plausible explanation for increased susceptibility to β-lactam antibiotics. The importance of the terminal stem peptide D-ala-D-ala for β-lactam resistance has previously been reported. Mutation of the *murF*-encoded ligase, which catalyses of the D-ala-D-ala into the stem peptide also increased β-lactam (but not DCS) susceptibility [32, 33]. Similarly growth of a HoR MRSA strain in media supplemented with high concentrations of glycine was accompanied by replacement of the D-ala-D-ala with D-ala-gly and decreased methicillin resistance [34].

Impaired uptake of alanine in CDMG correlated with increased susceptibility to oxacillin and DCS, suggesting that alanine utilisation via CycA is important to make D-alanine available for cell wall biosynthesis and consequently resistance to β-lactams. Consistent with this, NE810 also exhibited increased oxacillin susceptibility in BHI, TSB and MH media. However no change in alanine transport or susceptibility to oxacillin and DCS was measured in CDM lacking glucose, which may explain the failure of oxacillin and DCS to more efficiently eradicate NE810 infections in the mouse bacteraemia model. The availability of nutrients such as glucose and amino acids varies in different niches colonised by *S. aureus* during infection ranging from glucose-rich in organs such as the liver [35], to glucose-depleted in established abscesses [36]. In turn this impacts the role of amino acids as carbon sources [22, 37], and potentially the activity of CycA in alanine transport and β-lactam susceptibility. Furthermore, normal alanine transport in the *cycA* mutant grown in CDM indicates that an alternative alanine transport mechanism(s) may be active under these growth conditions (Fig. 6). Identification of this alternative alanine permease may be important in the development of therapeutic strategies

261   targeting alanine transport to increase β-lactam susceptibility in MRSA, while elucidation of

262   the role of glucose in the control of alanine transport should provide new insights into β-

263   lactam resistance.

Materials and Methods

Bacterial strains, growth conditions and antimicrobial susceptibility testing. Bacterial strains used in this study (Table S2) were grown in Luria Bertoni (LB) (Sigma), brain heart infusion (BHI) (Oxoid), Mueller Hinton (MH) (Oxoid), nutrient (Oxoid), sheep blood BHI, chemically defined media (CDM) [38] or CDM supplemented with 14mM glucose (CDMG) [38]. Minimum inhibitory concentrations (MICs) were determined in accordance with the Clinical Laboratory Standards Institute (CLSI) guidelines using plate and broth dilution assays in MH or MH containing 2% NaCl for oxacillin and nafcillin. Oxacillin MICs were also measured using E-tests strips from Oxoid on MH agar containing 2% NaCl. Quality control strains ATCC29213 and ATCC25923 were used for broth dilution and disk diffusion oxacillin and cefoxitin MIC assays, respectively.

Identification of cefoxitin susceptible MRSA mutant NE810. Cefoxitin (30μg) disks (Oxoid) were used to measure susceptibility of NTML mutants. The zone diameters for JE2 was 18mm NE1868 (*mecA*::Em^r) was >35mm and NE810 was 22mm. The *cycA* transposon insertion in NE810 was verified by PCR using the primers NE810_Fwd and NE810_Rev (Table S3). Phage 80α was used to transduce the NE810 *cycA* allele into JE2 and other strains. Genome sequencing was performed by MicrobesNG using the USA300_FPR3757 genome as a reference. To complement NE810, *cycA* was amplified from JE2 on a 1608 bp fragment using primers NE810F1_Fwd and NE810F1_Rev (Table S3) and cloned into pLI50 using the Clontech In-fusion kit.

Amino acid transport studies. Amino acid analysis in spent media from cultures grown in CDM or CDMG was performed as described previously [22].

Analysis of peptidoglycan composition in NE810 and JE2 treated with D-cycloserine. Independent quadruplicate 50ml cultures were grown to A_{600}=0.5, dosed with DCS at 0, 8, 20 or 32 μg/ml for 100 mins, then harvested and resuspended in 5ml PBS (Fig. S3) before peptidoglycan was extracted as described previously [39]. Mass spectrometry was performed on a Waters Xevo G2-XS QTof mass spectrometer. Structural characterization of muropeptides was determined based on their MS data and MS/MS fragmentation pattern, matched with PG composition and structure reported previously [34, 40-42].

Antibiotic synergy analysis using the microdilution checkerboard assay. Antibiotic synergism was measured using the checkerboard microdilution method in 96-well plates inoculated with 5×10^5 CFU/ml, and growth or no growth was assessed after 24 h at 37°C. The fractional inhibitory concentration index (ΣFIC) was calculated for each drug combination in triplicate experiments with an FIC index of ≤0.5 considered synergistic.

Kill curve assays. Overnight cultures adjusted to 10^7 CFU/ml were exposed to 0.125×, 0.25×, and 0.5× MIC of oxacillin, cefoxitin and DCS alone or in combination, and the number of colony forming units (CFU)/ml enumerated at 0, 2, 4, 8 and 24 h. Data is presented at the antibiotic concentrations where synergy was measured i.e. 0.5× MIC for JE2, USA300, DAR173, DAR22, DAR113, BH1CC, and RP62A, and 0.25× MIC for DAR169. Synergism was defined as a ≥2 $\log^{10}$ decrease in the number of CFU/ml in cell suspensions exposed to DCS/β-lactam combinations compared to the most effective individual drug after 8 h.

Mouse infection experiments. 6- to 8-week-old, age matched, outbred CD1 female mice (Charles River, UK) were used in a non-lethal model of bacteremia. JE2 and NE810 cultures were grown to A_{600}=0.5 in BHI, washed in PBS, adjusted to 1×10^8 CFU/ml. Mice were infected intravenously (via the tail vein) with 5×10^6 CFU ($n = 10$ mice per group). The infections were left untreated (PBS control) or treated with either 75 mg oxacillin/Kg/12 hours, 30 mg DCS/Kg/12 hours or a combination of both (first antibiotic dose administered 16 hours post infection), before being sacrificed after 5 days. Bacteria present in homogenised spleens and kidneys recovered from the mice were enumerated on blood agar.

Ethics Statement. Mouse experiments were approved by the United Kingdom Home Office (Home Office Project License Number 40/3602) and the University of Liverpool Animal Welfare and Ethics Committee. This study was carried out in strict accordance with the United Kingdom Animals (Scientific Procedures) Act 1986. All efforts were made to minimize suffering.

Statistical analysis. Two-tailed Student's t-Tests and one-way ANOVA with Kruskal-Wallis test followed by Dunn's multiple comparisons test in the GraphPad Prism application (for the mouse infection experiments) were used to determine statistically significant differences in assays performed during this study. A p value <0.05 was deemed significant.

Acknowledgements. We thank Craig Winstanley and Kate Reddington for generously

providing bacterial strains.

Footnotes

Financial support. This study was funded by grants from the Irish Health Research Board

(HRA_POR/2012/51, HRA-POR-2015-1158) and Science Foundation Ireland (16/TIDA/4137) to

(J.P.O'G), the UK Medical Research Council (MR/M020045/1) to (A.K), the National Institutes

of Health (AI083211) to (P.D.F) and Svenska Forskningsrådet Formas to (F.C).

Disclaimer. The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

Potential conflicts of interest. The authors declare no conflicts of interest relating to this

study.

Correspondence: Prof James P. O'Gara, Discipline of Microbiology, School of Natural

Sciences, National University of Ireland, Galway, Ireland. Tel: +353 91 492250; Email:

jamesp.ogara@nuigalway.ie

 Figure Legends

Figure 1. Mutation of *cycA* impairs alanine uptake. A. Growth of JE2 and NE810 in chemically

defined media supplemented with glucose (CDMG). Cell density was measured at A_{600}. **B.**

Alanine consumption by JE2 and NE810 grown aerobically in CDMG. Residual amino acid was

measured in spent media after 0, 4, 6 and 8 h growth. **C and D.** Growth of JE2 and NE810

cultures for 12 h in CDMG supplemented with 1 μg/ml oxacillin (F) or 1 μg/ml D-cycloserine

(G). Cell density was measured at A_{600}.

Figure 2. Mutation of *cycA* or D-cycloserine (DCS) treatment impacts peptidoglycan peptide

stem length and reduces cell wall crosslinking. A. Representative UV chromatograms of

peptidoglycan from wild-type JE2, NE810 and JE2 treated with increasing concentrations of

DCS (8, 20 and 32 μg/ml). Muropeptides with tripeptide stems are numbered 1-3. The

Proposed structures of the three muropeptides with tripeptide stems identified in NE810 and

DCS-treated JE2 cells. NAG, N-acetylglucosamine; NAM, N-acetylmuramic acid; L-Ala, L-

alanine; D-Gln, D-glutamine; D-Glu, D-glutamic; L-Lys, L-lysine. The theoretical and observed

neutral masses determined by MS are indicated. **B.** Relative abundance of muropeptides with

tripeptides in the stem. **C.** Relative crosslinking efficiency. **D.** Relative proportions of cell wall

muropeptide fractions based on oligomerization. All errors bars represent 95% confidence

interval (n = 4). Significant differences determined using Students t-test are denoted using

asterisks (* p<0.05; ** p<0.01; *** p<0.001).

Figure 3. *In vitro* kill curves for D-cycloserine (DCS), oxacillin and cefoxitin with JE2, USA300

FPR3757, DAR173, DAR22, DAR169 and their isogenic *cycA* mutants. Antibiotics at the

concentrations indicated (μg/ml) were added to suspensions of overnight bacterial cultures

adjusted to 10^7 CFU/ml in BHI (A_{600}=0.05), incubated at 37°C and the number of CFU/ml

enumerated at 0, 2, 4, 8 h. The data presented are the mean of three independent

experiments, and standard error of the mean is shown. Antibiotic synergism was defined as a

≥2 $\log^{10}$ decrease in the number of CFU/ml in cell suspensions exposed to DCS/β-lactam

combinations compared to the most effective individual antibiotic alone.

Figure 4. Combination therapy with D-cycloserine and oxacillin significantly reduces the

bacterial burden in the kidneys of mice infected with MRSA. The number of colony-forming

units (CFU) recovered from the kidneys of mice infected by tail vein injection with 5 × 10^6 JE2

or NE810 (CycA) and left untreated or treated with 75mg of oxacillin (Ox)/kg, 30mg of DCS/kg

or a combination of both Ox and DCS delivered subcutaneously every 12 hours for 5 days. The

first antibiotic dose was given 16 hours after infection. Significant differences determined

using one-way ANOVA with Kruskal-Wallis test followed by Dunn's multiple comparisons

test are denoted using asterisks (*p≤0.05, **p≤0.01, ****p≤0.0001). The limit of detection

(50 colonies) is indicated with a hashed red line.

Figure 5. Alanine transport and resistance to oxacillin and D-cycloserine in chemically

defined medium are *cycA*-independent. **A.** Growth of JE2 and NE810 in chemically defined

medium lacking glucose (CDM). Cell density was measured at A_{600}. **B.** Alanine consumption by

JE2 and NE810 grown aerobically in CDM. Residual amino acid was measured in spent media

after 0, 4, 6 and 8 h growth. **C and D.** Growth (cell density at A_{600}) of JE2 and NE810 cultures

for 12 h in CDM supplemented with 1 μg/ml oxacillin (C) or 1 μg/ml DCS (D).

Figure 6. Proposed model depicting how impaired alanine transport associated with mutation

of CycA or exposure to DCS can inhibit the D-alanine pathway for peptidoglycan biosynthesis

leading to increased susceptibility to β-lactam antibiotics.

References

1. Clatworthy AE, Pierson E, Hung DT. Targeting virulence: a new paradigm for antimicrobial therapy. Nat Chem Biol **2007**; 3:541-8.

2. Waters EM, Rudkin JK, Coughlan S, et al. Redeploying beta-lactam antibiotics as a novel antivirulence strategy for the treatment of methicillin-resistant Staphylococcus aureus infections. J Infect Dis **2017**; 215:80-7.

3. Davis JS, Sud A, O'Sullivan MV, et al. Combination of Vancomycin and beta-Lactam Therapy for Methicillin-Resistant Staphylococcus aureus Bacteremia: A Pilot Multicenter Randomized Controlled Trial. Clin Infect Dis **2016**; 62:173-80.

4. Tong SY, Nelson J, Paterson DL, et al. CAMERA2 - combination antibiotic therapy for methicillin-resistant Staphylococcus aureus infection: study protocol for a randomised controlled trial. Trials **2016**; 17:170.

5. Chambers HF. Methicillin resistance in staphylococci: molecular and biochemical basis and clinical implications. Clin Microbiol Rev **1997**; 10:781-91.

6. Chambers HF, Hackbarth CJ. Effect of NaCl and nafcillin on penicillin-binding protein 2a and heterogeneous expression of methicillin resistance in Staphylococcus aureus. Antimicrob Agents Chemother **1987**; 31:1982-8.

7. Sabath LD, Wallace SJ. The problems of drug-resistant pathogenic bacteria. Factors influencing methicillin resistance in staphylococci. Annals of the New York Academy of Sciences **1971**; 182:258-66.

8. Griffiths JM, O'Neill AJ. Loss of function of the gdpP protein leads to joint beta-lactam/glycopeptide tolerance in Staphylococcus aureus. Antimicrobial agents and chemotherapy **2012**; 56:579-81.

9. Dengler V, McCallum N, Kiefer P, et al. Mutation in the C-di-AMP cyclase dacA affects fitness and resistance of methicillin resistant Staphylococcus aureus. PloS one **2013**; 8:e73512.

10. Geiger T, Goerke C, Fritz M, et al. Role of the (p)ppGpp synthase RSH, a RelA/SpoT homolog, in stringent response and virulence of Staphylococcus aureus. Infection and immunity **2010**; 78:1873-83.

11. Geiger T, Kastle B, Gratani FL, Goerke C, Wolz C. Two small (p)ppGpp synthases in Staphylococcus aureus mediate tolerance against cell envelope stress conditions. J Bacteriol **2014**; 196:894-902.

12. Mwangi MM, Kim C, Chung M, et al. Whole-genome sequencing reveals a link between beta-lactam resistance and synthetases of the alarmone (p)ppGpp in Staphylococcus aureus. Microb Drug Resist **2013**; 19:153-9.

13. Fey PD, Endres JL, Yajjala VK, et al. A genetic resource for rapid and comprehensive phenotype screening of nonessential Staphylococcus aureus genes. MBio **2013**; 4:e00537-12.

14. Mlynek KD, Callahan MT, Shimkevitch AV, et al. Effects of Low-Dose Amoxicillin on Staphylococcus aureus USA300 Biofilms. Antimicrobial agents and chemotherapy **2016**; 60:2639-51.

15. Desjardins CA, Cohen KA, Munsamy V, et al. Genomic and functional analyses of Mycobacterium tuberculosis strains implicate ald in D-cycloserine resistance. Nat Genet **2016**; 48:544-51.

16. Chen JM, Uplekar S, Gordon SV, Cole ST. A point mutation in cycA partially contributes to the D-cycloserine resistance trait of Mycobacterium bovis BCG vaccine strains. PLoS One **2012**; 7:e43467.

17. Sieradzki K, Tomasz A. Suppression of beta-lactam antibiotic resistance in a methicillin-resistant Staphylococcus aureus through synergic action of early cell wall inhibitors and some other antibiotics. J Antimicrob Chemother **1997**; 39 Suppl A:47-51.

18. Pozzi C, Waters EM, Rudkin JK, et al. Methicillin resistance alters the biofilm phenotype and attenuates virulence in Staphylococcus aureus device-associated infections. PLoS Pathog **2012**; 8:e1002626.

19. Soper TS, Manning JM. Synergy in the antimicrobial action of penicillin and beta-chloro-D-alanine in vitro. Antimicrobial agents and chemotherapy **1976**; 9:347-9.

20. Plata KB, Riosa S, Singh CR, Rosato RR, Rosato AE. Targeting of PBP1 by beta-lactams determines recA/SOS response activation in heterogeneous MRSA clinical strains. PLoS One **2013**; 8:e61083.

21. Conlon KM, Humphreys H, O'Gara JP. icaR encodes a transcriptional repressor involved in
environmental regulation of ica operon expression and biofilm formation in Staphylococcus
epidermidis. J Bacteriol **2002**; 184:4400-8.

22. Halsey CR, Lei S, Wax JK, et al. Amino Acid Catabolism in Staphylococcus aureus and the Function
of Carbon Catabolite Repression. MBio **2017**; 8.

23. Tabuchi F, Matsumoto Y, Ishii M, et al. D-cycloserine increases the effectiveness of vancomycin
against vancomycin-highly resistant Staphylococcus aureus. J Antibiot (Tokyo) **2017**; 70:907-10.

24. Fujihira T, Kanematsu S, Umino A, Yamamoto N, Nishikawa T. Selective increase in the extracellular
D-serine contents by D-cycloserine in the rat medial frontal cortex. Neurochem Int **2007**; 51:233-6.

25. Batson S, de Chiara C, Majce V, et al. Inhibition of D-Ala:D-Ala ligase through a phosphorylated
form of the antibiotic D-cycloserine. Nat Commun **2017**; 8:1939.

26. Baisa G, Stabo NJ, Welch RA. Characterization of Escherichia coli D-cycloserine transport and
resistant mutants. J Bacteriol **2013**; 195:1389-99.

27. Curtiss R, 3rd, Charamella LJ, Berg CM, Harris PE. Kinetic and genetic analyses of D-cycloserine
inhibition and resistance in Escherichia coli. Journal of bacteriology **1965**; 90:1238-50.

28. Russell RR. Mapping of a D-cycloserine resistance locus in escherichia coli K-12. Journal of
bacteriology **1972**; 111:622-4.

29. Wargel RJ, Hadur CA, Neuhaus FC. Mechanism of D-cycloserine action: transport mutants for D-
alanine, D-cycloserine, and glycine. Journal of bacteriology **1971**; 105:1028-35.

30. Feher T, Cseh B, Umenhoffer K, Karcagi I, Posfai G. Characterization of cycA mutants of Escherichia
coli. An assay for measuring in vivo mutation rates. Mutat Res **2006**; 595:184-90.

31. Prosser GA, Rodenburg A, Khoury H, et al. Glutamate Racemase Is the Primary Target of beta-
Chloro-d-Alanine in Mycobacterium tuberculosis. Antimicrob Agents Chemother **2016**; 60:6091-9.

32. Sobral RG, Ludovice AM, Gardete S, Tabei K, De Lencastre H, Tomasz A. Normally functioning murF
is essential for the optimal expression of methicillin resistance in Staphylococcus aureus. Microb Drug
Resist **2003**; 9:231-41.

33. Sobral RG, Ludovice AM, de Lencastre H, Tomasz A. Role of murF in cell wall biosynthesis: isolation
and characterization of a murF conditional mutant of Staphylococcus aureus. J Bacteriol **2006**;
188:2543-53.

34. de Jonge BL, Chang YS, Gage D, Tomasz A. Peptidoglycan composition of a highly methicillin-
resistant Staphylococcus aureus strain. The role of penicillin binding protein 2A. J Biol Chem **1992**;
267:11248-54.

35. Li C, Sun F, Cho H, et al. CcpA mediates proline auxotrophy and is required for Staphylococcus
aureus pathogenesis. J Bacteriol **2010**; 192:3883-92.

36. Thurlow LR, Joshi GS, Richardson AR. Peroxisome Proliferator-Activated Receptor gamma Is
Essential for the Resolution of Staphylococcus aureus Skin Infections. Cell Host Microbe **2018**; 24:261-
70 e4.

37. Lehman MK, Nuxoll AS, Yamada KJ, Kielian T, Carson SD, Fey PD. Protease-Mediated Growth of
Staphylococcus aureus on Host Proteins Is opp3 Dependent. MBio **2019**; 10.

38. Hussain M, Hastings JG, White PJ. A chemically defined medium for slime production by coagulase-
negative staphylococci. Journal of medical microbiology **1991**; 34:143-7.

39. Alvarez L, Hernandez SB, de Pedro MA, Cava F. Ultra-Sensitive, High-Resolution Liquid
Chromatography Methods for the High-Throughput Quantitative Analysis of Bacterial Cell Wall
Chemistry and Structure. Methods Mol Biol **2016**; 1440:11-27.

40. De Jonge BL, Gage D, Xu N. The carboxyl terminus of peptidoglycan stem peptides is a determinant
for methicillin resistance in Staphylococcus aureus. Antimicrob Agents Chemother **2002**; 46:3151-5.

41. Kuhner D, Stahl M, Demircioglu DD, Bertsche U. From cells to muropeptide structures in 24 h:
peptidoglycan mapping by UPLC-MS. Sci Rep **2014**; 4:7494.

42. Boneca IG, Xu N, Gage DA, de Jonge BL, Tomasz A. Structural characterization of an abnormally
cross-linked muropeptide dimer that is accumulated in the peptidoglycan of methicillin- and
cefotaxime-resistant mutants of Staphylococcus aureus. J Biol Chem **1997**; 272:29053-9.

Table 1. Antibacterial activity (minimum inhibitory concentrations, MIC) and drug synergy (fractional inhibitory concentration indices, ΣFIC) of D-cycloserine (DCS) and several β-lactam antibiotics with different PBP specificity, namely oxacillin (OX; PBP1, 2, 3), nafcillin (NAF; PBP1), cefoxitin (FOX; PBP4), cefaclor (CEC; PBP3), cefotaxime (CTX; PBP2), piperacillin-tazobactin (TZP; PBP3/β-lactamase inhibitor) and imipenem (IMP; PBP1), alone and in β-lactam combinations, against fourteen *S. aureus* strains and *S. epidermidis* RP62A.

Antibiotic → ↓Strain	OX*	NAF*	DCS*	FOX*	DCS/FOX** (ΣFIC)***	CEC*	DCS/CEC** (ΣFIC)***	CTX*	DCS/CTX** (ΣFIC)***	TZP*	DCS/TZP** (ΣFIC)***	IPM*	DCS/IPM** (ΣFIC)***
JE2	64	32	32	64	8/8 (0.37)	64	8/2 (0.28)	64	8/4 (0.31)	32	8/0.5 (0.26)	1	ND
NE810 (*cycA*)	0.25	0.5	2	8	ND	4	ND	8	ND	2	ND	0.125	ND
USA300	64	32	32	64	8/8 (0.37)	64	8/8 (0.37)	64	8/8 (0.37)	64	8/2 (0.28)	1	ND
USA300 *cycA*	0.25	1	2	8	ND	4	ND	8	ND	4	ND	0.125	ND
DAR173	128	128	32	256	4/32 (0.25)	128	8/4 (0.28)	512	8/4 (0.25)	128	4/4 (0.15)	64	4/2 (0.15)
DAR173 *cycA*	0.5	8	4	32	ND	16	ND	16	ND	4	ND	1	ND
DAR22	128	128	32	128	8/8 (0.31)	128	8/8 (0.31)	512	8/4 (0.25)	128	4/4 (0.15)	128	4/1 (0.13)
DAR22 *cycA*	0.5	8	0.5	16	ND	16	ND	16	ND	4	ND	0.5	ND
DAR169	32	32	32	32	8/2 (0.31)	128	4/32 (0.37)	64	4/8 (0.25)	8	4/1 (0.25)	2	ND
DAR169 *cycA*	16	4	0.5	16	ND	64	ND	16	ND	2	ND	0.5	ND
COL	512	256	64	512	8/128 (0.37)	128	16/32 (0.5)	>2048	ND	128	32/0.5 (0.5)	256	8/64 (0.37)
DAR113	128	64	32	128	8/8 (0.3)	64	8/4 (0.3)	256	8/8 (0.2)	128	8/4 (0.2)	16	8/0.5 (0.2)
BH1CC	256	512	32	256	8/32 (0.3)	128	8/8 (0.3)	1028	8/16 (0.2)	256	8/4 (0.2)	64	8/1 (0.2)
BH14B(04)	128	64	16	128	4/32 (0.5)	128	4/32 (0.5)	512	4/32 (0.31)	64	4/1 (0.26)	64	2/8 (0.25)
BH8(03)	128	128	32	128	8/32 (0.5)	256	8/64 (0.5)	128	8/16 (0.25)	128	8/8 (0.3)	64	8/0.5 (0.1)
BH6	128	128	32	128	8/16 (0.37)	256	8/64 (0.5)	512	8/32 (0.31)	128	8/4 (0.28)	32	8/0.5 (0.26)
DAR202	64	64	32	128	4/32 (0.37)	128	4/32 (0.37)	64	8/16 (0.5)	64	8/2 (0.28)	4	8/1 (0.5)
DAR45	2	0.5	32	4	ND	32	8/2 (0.31)	4	ND	2	ND	0.5	ND
DAR13	128	32	32	128	4/32 (0.37)	128	8/4 (0.28)	128	4/32 (0.37)	32	8/1 (0.28)	8	4/2 (0.37)
RP62A	128	2	32	64	6/16 (0.5)	64	6/16 (0.5)	32	8/8 (0.5)	8	8/2 (0.5)	32	8/1 (0.2)

* MIC values for each antibiotic when measured individually; μg/ml

** MIC values for each antibiotic when measured in combination, also known as the fractional inhibitory concentration (FIC); μg/ml

*** FIC indices (ΣFIC) for the antibiotic combination. ΣFIC = FIC A + FIC B, where FIC A is the MIC of DCS in combination with the β-lactam/MIC of DCS alone, and FIC B is the MIC of the β-lactam in combination with DCS/MIC of the β-lactam alone. The combination is considered synergistic when the ΣFIC is ≤0.5, indifferent when the ΣFIC is >0.5 to <2.

‡ND. Not determined if strain is susceptible (or hyper-resistant) to β-lactam antibiotic or for *cycA* mutants with reduced DCS & β-lactam MICs.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

SUPPLEMENTARY DATA

Impaired alanine transport or exposure to D-cycloserine increases the susceptibility of MRSA to β-lactam antibiotics

Laura A. Gallagher[1], Rebecca K. Shears[2], Claire Fingleton[1], Laura Alvarez[3], Elaine M. Waters[1,2], Jenny Clarke[2], Laura Bricio-Moreno[2], Christopher Campbell[1], Akhilesh K. Yadav[3], Fareha Razvi[4], Eoghan O'Neill[5], Alex J. O'Neill[6], Felipe Cava[3], Paul D. Fey[4], Aras Kadioglu[2] and James P. O'Gara[1]*

Supplementary Tables

Table S1. Antibacterial activity (minimum inhibitory concentrations, MIC) and drug synergy (fractional inhibitory concentration indices, ΣFIC) of D-cycloserine (DCS), and clindamycin (CLI), tobramycin (TOB), trimethoprim (TMP), mupirocin (MUP), ciprofloxacin (CIP), kanamycin (KAN) and spectinomycin (SPT) alone and in DCS/antibiotic combinations, against fourteen *S. aureus* strains and *S. epidermidis* RP62A.

Antibiotic → ↓ Strain	DCS*	CLI*	DCS/CLI** (ΣFIC)***	TOB	DCS/TOB (ΣFIC)	TMP	DCS/TMP (ΣFIC)	CIP	DCS/CIP (ΣFIC)	KAN	DCS/KAN (ΣFIC)	SPT	DCS/SPT (ΣFIC)	MUP	DCS/MUP (ΣFIC)
JE2	32	0.25	ND	1	ND	1	ND	32	16/16 (1)	2	ND	64	16/32 (1)	0.25	ND
USA300	32	2048	32/2048 (2)	0.5	ND	1	ND	32	16/16 (1)	2	ND	64	16/32 (1)	2048	32/2048 (2)
DAR173	32	0.5	ND	512	32/512 (2)	0.5	ND	256	16/128 (1)	512	16/256 (1)	64	16/32 (1)	0.5	ND
DAR22	32	0.5	ND	512	16/256 (1)	0.5	ND	256	16/128 (1)	256	16/128 (1)	64	16/32 (1)	0.5	ND
DAR169	32	0.5	ND	0.5	ND	0.5	4/8 (0.25)	1	ND	2	ND	64	16/32 (1)	0.5	ND
COL	64	0.5	ND	0.5	ND	0.5	ND	1	ND	0.5	ND	64	16/32 (1)	0.5	ND
DAR113	32	0.25	ND	0.5	ND	1	ND	0.5	ND	2	ND	64	16/32 (1)	0.25	ND
BH1CC	32	1024	32/1024 (2)	32	32/32 (2)	256	32/256 (2)	32	16/16 (1)	>1024	ND	>1024	ND	32	16/8 (0.75)
BH14B(04)	16	0.5	ND	0.5	ND	0.5	ND	128	16/128 (1.5)	2	ND	64	8/32 (1)	0.5	ND
BH8(03)	32	1024	32/1024 (2)	0.5	ND	0.5	ND	256	16/128 (1)	8	32/4 (2)	128	16/64 (1)	4	32/4 (2)
BH6	32	0.5	ND	0.5	ND	0.5	ND	>512	16/256 (1)	1	ND	128	16/64 (1)	0.5	ND
DAR202	32	0.5	ND	128	16/64 (1)	0.5	ND	2	ND	>1024	ND	>1024	ND	0.5	ND
DAR45	32	1024	32/1024 (2)	512	16/256 (1)	0.5	ND	128	16/64 (1)	128	16/64 (1)	512	32/1024 (2)	0.5	ND
DAR13	32	0.5	ND	0.5	ND	32	16/16 (1)	32	16/16 (1)	2048	ND	>1024	ND	0.5	ND
RP62A	32	2048	32/2048 (2)	16	16/8 (1)	256	32/256 (2)	0.25	ND	>1024	ND	>1024	ND	0.25	ND

* MIC values for each antibiotic when measured individually; μg/ml

** MIC values for each antibiotic when measured in combination, also known as the fractional inhibitory concentration (FIC); μg/ml

*** FIC indices (ΣFIC) for the antibiotic combination. ΣFIC = FIC A + FIC B, where FIC A is the MIC of DCS in combination with the antibiotic/MIC of DCS alone, and FIC B is the MIC of the antibiotic in combination with DCS/MIC of the antibiotic alone. The combination is considered synergistic when the ΣFIC is ≤0.5, indifferent when the ΣFIC is >0.5 to <2.

[‡]ND. Not determined if strain is susceptible (or hyper-resistant) to the antibiotic or for *cycA* mutants with reduced DCS & antibiotic MICs.

Table S2. Bacterial strains and plasmids used in this study

Strains/plasmids	Relevant Details
RN4220	Restriction-deficient laboratory *S. aureus*
USA300 FPR3757	Community associated MRSA isolate of the USA300 lineage [43]. SCCmec type IV. CC8.
JE2	USA300 cured of p01 & p03. Parent of Nebraska Transposon Mutant Library (NTML).
13C	USA300 derivative cured of plasmid p03. Erm^s.
NE810	JE2 NTML *cycA* (SAUSA300_1642) mutation. Erm^r.
USA300 *cycA*	USA300 FPR3757 *cycA*. Constructed by transduction of *cycA*::Tn allele from NE810.
13C *cycA*	13C *cycA* mutant (*cycA*::Tn allele from NE810).
NE1868	JE2 NTML *mecA* mutation.
NE1713	JE2 NTML *alr* (SAUSA300_2027) mutation.
NE799	JE2 NTML *alr2* (SAUSA300_1292) mutation.
NE923	JE2 NTML *ynfA* (SAUSA300_2286) mutation.
NE1025	JE2 NTML *thrE* (SAUSA300_0729) mutation.
BH1CC	MRSA clinical isolate; SCC*mec* type II; CC8 [44]
COL	MRSA reference strain; SCC*mec* type I; CC8 [45]
BH14(04)	MRSA clinical isolate; SCC*mec* type IV; CC22 [44]
BH8(03)	MRSA clinical isolate; SCC*mec* type IV; CC22 [44]
BH6(03)	MRSA clinical isolate; SCC*mec* type II; CC8 [44]
DAR113	MRSA reference isolate; SCC*mec* type IV; CC22 [44, 46]
DAR13	MRSA reference isolate; SCC*mec* type IV; CC8 [44, 46]
DAR45	MRSA reference isolate; SCC*mec* type II; CC30 [44, 46]
DAR202	MRSA reference isolate; SCC*mec* type III; CC239 [44, 46]
DAR173	MRSA reference isolate; SCC*mec* type IV; CC5 [44, 46]
DAR173 *cycA*	DAR173 *cycA* mutant (*cycA*::Tn allele from NE810)
DAR22	MRSA reference isolate; SCC*mec* type III; CC5 [44, 46]
DAR22 *cycA*	DAR22 *cycA* mutant (*cycA*::Tn allele from NE810)
DAR169	MRSA Reference strain; SCCmec type I; CC8 [44, 46]
DAR169 *cycA*	DAR169 *cycA* mutant (*cycA*::Tn allele from NE810)
8325-4	NCTC 8325 derivative cured of prophages [47], methicillin susceptible, CC8.
8325-4 *cycA*	8325-4 *cycA* mutant (*cycA*::Tn allele from NE810).
ATCC 29213	Methicillin susceptible *S. aureus* strain for antibiotic susceptibility testing.
ATCC 29213 *cycA*	ATCC 29213 *cycA* mutant (*cycA*::Tn allele from NE810).
ATCC 25923	Methicillin susceptible *S. aureus* strain for antibiotic susceptibility testing.
S. epidermidis RP62A	ATCC 35984. Methicillin resistant, biofilm positive. [21]
E. coli	*E. coli* HST08
Plasmids	
pLI50	*E. coli-Staphylococcus* shuttle vector. Ap^r (*E. coli*), Cm^r. (*Staphylococcus*)
p*cycA*	pLI50 carrying *cycA* from JE2

Table S3. Oligonucleotide primers used in this study

Target Gene	Primer Name	Primer Sequence (5'-3')
cycA	NE810_Fwd	ACAGAATAGCCACAAATAGCACC
	NE810_Rev	ACAGAATAGCCACAAATAGCACC
cycA	NE810F1_Fwd	GTCTTCAAGAATTCGGCCACAAATAGCACCATTAA
	NE810F1_Rev	CGACTCTAGAGGATCATGTCCCAAGCCCTAAAAC
mecA	mecA1_Fwd	TGCTCAATATAAAATTAAAACAAACTACGGTAAC
	mecA1_Rev	GAATAATGACGCTATGATCCCAA
gyrB	gyrB_Fwd	CCAGGTAAATTAGCCGATTGC
	gyrB_Rev	AAATCGCCTGCGTTCTAGAG

Supplementary Figures

Figure S1. Mutation of *cycA* increases the susceptibility of MRSA to β-lactam antibiotics and D-cycloserine. A. Chromosomal location of *cycA* and neighbouring genes *icd* (isocitrate dehydrogenase), *gltA* (citrate synthase), *pyk* (pyruvate kinase) and *pfkA* (6-phosphofructokinase). The locations of transposon insertions in NE810, NE491, NE594 and NE1497 mutants from the Nebraska library are indicated. **B.** E-test measurement of oxacillin minimum inhibitory concentrations (MICs) in JE2 (wild type), NE810 (*cycA*::Tn), NE810 carrying pLI50 (control) and p*cycA*. **C.** Comparison of zones of inhibition around D-cycloserine 30μg disks on lawns of JE2, NE810 (*cycA*::Tn), NE810 pLI50 (control) and NE810 p*cycA* grown on Mueller Hinton (MH) agar.

Figure S2. Amino acid consumption by JE2 and NE810 grown aerobically in chemically defined media containing 14mM of glucose (CDMG). Residual amino acids were measured in spent media after 0, 4, 6 and 8 h growth.

Figure S3. Preparation of cell suspensions for peptidoglycan extraction and structural analysis by UPLC-MS. 50 ml flask cultures were inoculated into fresh BHI media from overnight cultures at a starting cell density of A_{600}=0.05 and incubated at 37°C. The number of CFU/ml was enumerated every 1-2 h for 6 h and again after 24 hours. For JE2 cultures being dosed with DCS, the antibiotic was added after approximated 2 h (A_{600}≈0.5) and the cells collected after a further 100 mins. Cells from untreated JE2 and NE810 control cultures were collected at the same time point.

Figure S4. *In vitro* kill curves for D-cycloserine (DCS), oxacillin and cefoxitin with methicillin resistant *S. epidermidis strain* RP62A. Antibiotics at the concentrations indicated (equivalent to 0.5×MIC) were added to suspensions of overnight bacterial cultures adjusted to 10^7CFU/ml in BHI, incubated at 37°C and the number of CFU/ml enumerated at 0, 2, 4, 8 and 24 h. The data presented are the mean of three independent experiments. Antibiotic synergism was defined as a ≥2 $\log^{10}$ decrease in the number of CFU/ml in cell suspensions exposed to DCS/β-lactam combinations compared to the most effective individual antibiotic alone.

Figure S5. Combination therapy with D-cycloserine and oxacillin significantly reduces the bacterial burden in the spleen of mice infected with MRSA. The number of colony-forming units (CFU) recovered from the spleens of mice infected by tail vein injection with 5×10^6 JE2 or NE810 (CycA) and left untreated or treated with 75mg of oxacillin (Ox)/kg, 30mg of DCS/kg or a combination of both Ox and DCS delivered subcutaneously every 12 hours for 5 days. The first antibiotic dose was given 16 hours after infection. Significant differences determined using one-way ANOVA with Kruskal-Wallis test followed by Dunn's multiple comparisons test are denoted using asterisks (*p≤0.05). The limit of detection (50 colonies) is indicated with a hashed red line.

Figure S6. Amino acid consumption by JE2 and NE810 grown aerobically in chemically defined media lacking glucose(CDM). Residual amino acids were measured in spent media after 0, 4, 6 and 8 h growth.