A web-based platform of nucleotide sequence alignments of plants

Chiara Santos[1,¶], João Carneiro[1,¶] and Filipe Pereira[2*]

[1] Interdisciplinary Centre of Marine and Environmental Research (CIIMAR), University of Porto,

Terminal de Cruzeiros do Porto de Leixões; Avenida General Norton de Matos, S/N 4450-208

Matosinhos – Portugal

[2] University of Porto, Praça Gomes Teixeira 4099-002 Porto, Portugal

[¶] These authors contributed equally to this work.

[*]Corresponding author:

F. Pereira; E-mail fpereirapt@gmail.com

ORCID Filipe Pereira 0000-0001-8950-1036

Abstract

In recent years, a large number of nucleotide sequences have become available for plant

species by the advent of massive parallel sequencing. The use of genomic data has been

important for agriculture, food science, medicine or ecology. Despite the increasing amount of

data, nucleotide sequences are usually available in public databases as isolated records with

some descriptive information. Researchers interested in studying a wide range of specific

plant families are forced to do multiple searches, sequence downloads, data curation and

sequence alignments. In order to help researches overcoming these problems, we have built a

comprehensive on-line resource of curated nucleotide sequence alignments for plant research,

named PlantAligDB (available at http://plantaligdb.portugene.com). The latest release

incorporates 514 alignments with a total of 66,052 sequences from six important genomic

regions: *atp*F-*atp*H, *psb*A-*trn*H, *trn*L, *rbc*L, *mat*K and ITS. The alignments represent 223

plant families from a variety of taxonomic groups. The users can quickly search the database,

download and visualize the curated alignments and phylogenetic trees using dynamic

browser-based applications. Different measures of genetic diversity are also available for each

plant family. We also provide the workflow script that allows the user to do the curation

process, explaining the steps involved. Overall, the PlantAligDB provides a complete, quality

checked and regularly updated collection of alignments that can be used in taxonomic, DNA

barcoding, molecular genetics, phylogenetic and evolutionary studies.

Keywords

DNA sequences, Multiple sequence alignments, Plant families

Introduction

The recent development of high-throughput sequencing technologies has significantly increased the number of nucleotide sequences available in public databases (Egan *et al.* 2012; Feuillet *et al.* 2011). Complete genome sequences are now accessible in public databases (e.g., EnsemblPlants) (https://plants.ensembl.org/index.html) for the analysis and visualisation of genomic data for an ever-growing number of plants, such as *Beta vulgaris*, *Prunus persica* and *Citrus sinensis*, among many others. Sequences from individual genes or gene regions have also been deposit in public databases as a result of international initiatives. For instance, the DNA barcoding project has released thousands of sequences aiming at species identification and taxonomic classification of plants, mostly from the chloroplast DNA (cpDNA) protein-coding genes *rbc*L and *mat*K [CBOL Plant Working Group' - (Group *et al.* 2009; Hollingsworth *et al.* 2011)]. The plastid *trn*L (UAA) intron is another good example of a cpDNA region highly represented in sequence databanks (Taberlet *et al.* 2007).

Several web-based databases are available for plant genome sequences, usually dedicated to a single species or a genomic feature [e.g., (Lai *et al.* 2012; Meyer *et al.* 2005; Numa & Itoh 2014; Sakai *et al.* 2013)]. However, most nucleotide sequences are accessible in public databases as isolated records with simple descriptive information (taxonomy, geography, publications, etc.). For instance, the NCBI Entrez Nucleotide database (http://www.ncbi.nlm. nih.gov) and the BOLD - The Barcode of Life Data System (www.barcodinglife.org) (Ratnasingham & Hebert 2007) are useful repositories with descriptive information for sequence or species. The TreeBASE (https://www.treebase.org) is a repository of phylogenetic information with user-submitted phylogenetic trees and the data used to generate them. Nevertheless, researchers interested in studying a large number of plant families are forced to do multiple searches and sequence downloads of genetic data for their investigations. Moreover, the available sequences are not aligned and curated. The

multiple sequence alignment step is critical because it determines the accuracy of the

subsequence analyses, such as phylogenetic inference, identification of conserved motifs,

function prediction, etc. Building accurate sequence alignments involves many steps,

including sequence file conversion, run of alignment algorithms in local computers or

webservers, selection of best alignment parameters, and manual fine-tuning of the alignment.

This process is laborious and requires costly computational resources, which are not always

available.

We describe an on-line database (PlantAligDB, available at

http://plantaligdb.portugene.com) with a comprehensive, automatically curated and regularly

updated collection of alignments from diverse plant families (Figure 1), and respective

phylogenetic inference of each alignment. The PlantAligDB is a consistent repository of

curated alignments and phylogenetic trees that are generated by the same workflow. For

example, the Orchidaceae family is frequently referred to as a critical group, whose species

are difficult to identify. Because of their ecological importance, ongoing studies continue to

be made to reduce their extinction risk and maintain their diversity (Li *et al.* 2018). The

PlantAligDB can help researchers designing accurate methods for plant identification (*mat*K

and *rbc*L are used in DNA barcoding projects) whether by identification of conserved motives

that enable the design of primers. The PlantAligDB alignments can be used as a reference

database for phylogenetic studies, allowing the construction of reference phylogenetic trees of

the different regions [e.g., genomic regions *atp*F-*atp*H (Domenech *et al.* 2014), *psb*A-*trn*H

(Dong *et al.* 2012) and ITS (Karehed *et al.* 2008)]. Moreover, it provides useful data to

understand the genetic diversity of the selected genomic regions.

Materials and Methods

Data curation

We retrieved all nucleotide sequences of different genomic regions from the NCBI

Entrez Nucleotide database (http://www.ncbi.nlm.nih.gov) using the Geneious software

(Drummond 2009). Different combinations of search terms (e.g. '*gene name*'; viridiplantae';

'chloroplast'; 'gene'; 'complete') and a maximum limit of 5000 bp as sequence length were

used in searches to retrieve the largest number of sequences. The selection was done by

eliminating sequences showing one or more of the following features: a) ambiguous name

description; b) sequences with high number of nucleotide ambiguities (>50%); c) sequences

with large stretches of the region missing (>50%). After a preliminary curation of the data, we

selected five cpDNA regions and one nuclear DNA region, which were the most represented

in the NCBI database, commonly used in phylogenetic studies for being relevant and

informative. The six genomic regions were named according to the gene regions where they

are located: *atp*F-*atp*H *(ATPase I subunit – ATPase III subunit)*, *psb*A-*trn*H [*Photosystem II*

32 kDa protein – tRNA-His (GUG)], *trn*L [*tRNA*-Leu (UAA)], *rbc*L *(rubisco large subunit)*,

*mat*K *(maturase* K*)* and ITS (internal transcribed spacer). We then removed from the datasets

all redundant sequences belonging to the same species and sequences without a clear species

assignment. We also reverse complement the sequences that were found in the opposite

direction. The sequence orientation for each region is that of the most commonly found in the

NCBI database. Therefore, the orientation of the *trn*L *(UAA)*, *atp*F-*atp*H and *rbc*L regions are

the same of that used in the reference cpDNA sequence of *Nicotiana tabacum*

(NC_001879.2), while the opposite orientation is used for regions *psb*A-*trn*H and *mat*K. The

target region named ITS in our database includes the internal transcribed spacer 1, 5.8S rRNA

and internal transcribed spacer 2 section of the nuclear ribosomal DNA.

Because a high number of sequences were detected for the *trn*L (UAA) region (more

than 50,000 hits), we used the external regions named "C" and "D" and the internal regions

named "G" and "H" by (Taberlet *et al.* 2007) as queries in the NCBI Basic Local Alignment

Search Tool (BLAST; http://blast.ncbi.nlm.nih.gov/). The search was made against the

nucleotide collection (nr/nt) of Tracheophyta (vascular plants) using the Biopython package

(www.biopython.org) with an expected threshold of 1000 and a minimum word size of 16.

Therefore, our database includes two datasets for the *trn*L (UAA) genomic region: the '*trn*L

CD' target region with a length of 577 bp in *N. tabacum,* and the '*trn*L GH' with a length of

78 bp in *N. tabacum*, located inside the *trn*L CD region. All information regarding the selected

target regions can be found in the *Genomic Regions* section of the PlantAligDB.

The nucleotide sequences of the six regions were organized by family according to the

NCBI taxonomy and were aligned (each region and family in separated alignments) using the

default parameters of the MUSCLE software (Edgar 2004) running in the Geneious software.

The alignment was repeated in some families after excluding sequences that do not cover the

entire region of interest and that had large stretches of nucleotide ambiguities. We only used

alignments with ten or more species per family to build the PlantAligDB. Some species

sequences were lost in this process filter. The neighbor joining phylogenetic tree of each

region-family were calculated using Tamura-Nei model using Geneious Tree Builder. The

methodology was built in Armadillo Workflow (http://www.bioinfo.uqam.ca/armadillo/) to

automate the update process of the database. The latest release update of June 2018

incorporates 514 alignments and phylogenetic trees, from 223 plant families.

Conservation measures

The database includes two measures of sequence conservation for each alignment:

percentage of identical sites (PIS), calculated by dividing the number of identical positions in

the alignment for an oligonucleotide by its length and the *percentage of pairwise identity*

(PPI), calculated by counting the average number of pairwise matches across the positions of

the alignment, divided by the total number of pairwise comparisons.

Results and Discussion

Database organization

Basic structure

The PlantAligDB database is divided in nine sections (Figure 1): 1) *Home*, provides a

brief description of what can be done in the database; 2) *Genomic regions*, describes the

regions used in the database; 3) *Taxonomic groups*, the table containing the plant

family/region alignments and phylogenetic trees; 4) *BLAST*, search the database using a query

sequence by means of the BLASTN algorithm (Altschul *et al.* 1990); 5) *Genetic diversity*,

describes the percentage of identical sites (PIS) and pairwise identity values (PPI) for each

alignment; 6) *Download*, provides hyperlinks to download the curated alignments; 7)

Tutorials, contains information about how the database was built, and how to use it; 8)

Citations; 9) *Contacts*.

Taxonomic groups

The database is being regularly updated by our team and currently includes 514

alignments and phylogenetic trees from seven target regions: *atp*F-*atp*H, *psb*A-*trn*H, *trn*L CD,

*trn*L GH, *rbc*L, *mat*K and ITS (Table 1). Sequence alignments are provided for 223 different

plant families. Currently, the *trn*L GH region has the largest number of sequences (n =

34,674). The *Fabaceae* family has the largest number of aligned species in a target region,

with 2599 sequences for the *trn*L GH region. When considering all regions together, the

Fabaceae (n = 4714), *Poaceae* (n = 4494) and *Asteraceae* (n = 4459) families are those with

the highest number of sequences. The alignments for each plant family can be accessed

through a dynamic table in the *Taxonomic groups* section of the database by following a

hyperlink   with   the   number   of   species   included   in   each   alignment

(http://plantaligdb.portugene.com/cgi-bin/PlantAligDB_taxonomicgroups.cgi). The users are

able to quickly search and locate a queried feature, order each column using the ascendant or

descendent mode, filter the information, download the curated datasets, among other features.

The multiple sequence alignment and phylogenetic tree can be visualized by clicking in the

number of species present in the alignment.

Genetic diversity

The PIS values in our current dataset vary from 0.16% to 99.07% (Table 2).

Nevertheless, the PIS and PPI sequence conservation measures are not intended to be used for

comparison of different families and/or regions, since the number of sequences in each

alignment can be very different. The results must be interpreted at the light of the number of

sequences and the representativeness of the sequences included in the alignment file. The

*mat*K was the region with the lowest PIS value (0.16%) [Figure 2. f)], while the *trn*L GH was

the region with the highest PIS value (99.07%), as can be seen in Figure 2 d) and Table 2. The

*rbc*L was the most conserved region [Figure 2 e)] with an average of 78.48%, while the ITS

was less conserved with an average of 28.84% (Table 2). Our results are in accordance with

earlier studies where *atp*F-*atp*H and *psbA-trn*H were found to be more variables than *mat*K

(Lahaye 2008). The *trn*L CD regions showed values slightly more conserved than *atp*F-*atp*H

[Figure 2 c) and a)]. The lowest PPI value (69.96%) was found in *psb*A-*trn*H region [Figure 2

b)] and the highest was 100% in *trn*L GH, as shown in Figure 2 d) and Table 2. The ITS

region was less conserved with an average of 87.21% [Table 3 and Figure 2 g)], while the

*trn*L GH was the most conserved with an average of 97.09% [Table 3 d)].

Sequence alignments and phylogenetic trees

The alignments stored in the database can be visualized using a dynamic browser-based

application named Wasabi (http://wasabiapp.org/) (Veidenberg *et al.* 2016) with multiple

options for the visualization and analysis of sequence data and phylogenetic trees. This

resource is particularly useful to help researchers selecting the most appropriated genomic

regions for their investigations. The users can zoom in and out the selected regions of the

alignments, collapse regions with gaps, alternate between column and row selection, remove

or add sequences, realign sequences, and export the sequence data in the FASTA format. If an

Wasabi account is created, the user can re-align specific alignments with PAGAN (Loytynoja

et al. 2012) and PRANK (Loytynoja 2014). The user can merge different alignments from

identical regions (Hollingsworth et al. 2009) and different families using the PAGAN

application. The download of the complete database of curated alignments is accessible in the

Download section of the database.

How to use the PlantAligDB

To explore a genomic region, a researcher must start by accessing the 'Genomic

Regions' section in the menu bar. For example, by selecting *psb*A-*trn*H, the user will find a

brief description of the genomic region and the name, size and position of that region in the

reference genome, and the families with available alignments. The user should select the

'Taxonomic Groups' section in the menu bar to search for a specific plant family. Then,

through the search tool on the right side of the page, the user can type the name of the family

and a dynamic table will be displayed with the number of sequences for each region. The user

can also access a hyperlink with the description of the family and taxonomic tree using the

resource Tree of Life Web Project (http://tolweb.org). Clicking on the number of sequences,

the database is redirected to the Wasabi tool. The user can create a Wasabi account by

providing an e-mail or choosing a temporary account, which allows to realign sequences with

PAGAN or PRANK. The user can merge a PAGAN realignment with alignments of other

families on the same region, by selecting a file on his local computer in the "alignment

extension" option.

Availability and design

The PlantAligDB is freely available at http://plantaligdb.portugene.com and is optimized for

the major web browsers (Internet Explorer, Firefox, Safari, and Chrome). The SQLite local

database is used for data storage and runs on an Apache web server. The dynamic HTML

pages were implemented using CGI-Perl and JavaScript and the dataset table views were

generated using the JQuery plugin DataTables v1.9.4 (http://datatables.net/). The PlantAligDB

visualization tables are generated automatically. The process of database update is optimized

for large datasets. There are no access restrictions for academic and commercial use.

Acknowledgments

This work was supported by the Portuguese Foundation for Science and Technology (FCT),

European Regional Development Fund (ERDF) and Operational Human Potential Program

[IF/01356/2012 to F.P. and SFRH/BPD/100912/2014 to J.C.]. C.S. was supported by the

"Programa Ciências Sem Fronteiras" from the Conselho Nacional de Desenvolvimento

Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível

Superior (CAPES), Brazil. CIIMAR was partially supported by the Strategic Funding

UID/Multi/04423/2019 through national funds provided by FCT and ERDF in the framework

of the programme PT2020.

Author Contributions

242 CS, JC and FP contributed equally to designed and performed research, analysed data and wrote

243 the paper.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol*
Biol **215**, 403-410.
Domenech B, Asmussen-Lange CB, Baker WJ*, et al.* (2014) A phylogenetic analysis of palm subtribe
Archontophoenicinae (Arecaceae) based on 14 DNA regions. *Botanical Journal of the*
Linnean Society **175**, 469-481.
Dong W, Liu J, Yu J, Wang L, Zhou S (2012) Highly variable chloroplast markers for evaluating plant
phylogeny at low taxonomic levels and for DNA barcoding. *PLoS One* **7**, e35071.
Drummond AB, Cheung M, Heled J, Kearse M, Moir R, Stones-Havas S, Thierer T, Wilson A (2009)
Geneious Pro 4.8.2.
Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res **32**, 1792-1797.
Egan AN, Schlueter J, Spooner DM (2012) Applications of next-generation sequencing in plant
biology. *Am J Bot* **99**, 175-185.
Feuillet C, Leach JE, Rogers J, Schnable PS, Eversole K (2011) Crop genome sequencing: lessons and
rationales. *Trends Plant Sci* **16**, 77-88.
Group CPW (2009) A DNA barcode for land plants. *Proc Natl Acad Sci U S A* **106**, 12794-12797.
Group CPW, Hollingsworth PM, Forrest LL*, et al.* (2009) A DNA barcode for land plants. *Proceedings*
of the National Academy of Sciences **106**, 12794-12797.
Hollingsworth ML, Andra Clark A, Forrest LL*, et al.* (2009) Selecting barcoding loci for plants:
evaluation of seven candidate loci with species-level sampling in three divergent groups of
land plants. *Mol Ecol Resour* **9**, 439-457.
Hollingsworth PM, Graham SW, Little DP (2011) Choosing and using a plant DNA barcode. *PLoS*
One **6**, e19254.
Karehed J, Groeninckx I, Dessein S, Motley TJ, Bremer B (2008) The phylogenetic utility of
chloroplast and nuclear DNA markers and the phylogeny of the Rubiaceae tribe
Spermacoceae. *Mol Phylogenet Evol* **49**, 843-866.
Lahaye RS, V.; Duthoit, S.; Maurin, O. and van der Bank, M. (2008) A test of psbK-psbI and atpF-
atpH as potential plant DNA barcodes using the flora of the Kruger National Park as a model
system (South Africa). *Nature Precedings*, 21.
Lai K, Berkman PJ, Lorenc MT*, et al.* (2012) WheatGenome.info: an integrated database and portal
for wheat genome information. *Plant Cell Physiol* **53**, e2.
Li J, Gale SW, Kumar P, Zhang J, Fischer G (2018) Prioritizing the orchids of a biodiversity hotspot
for conservation based on phylogenetic history and extinction risk. *Botanical Journal of the*
Linnean Society, box084-box084.
Loytynoja A (2014) Phylogeny-aware alignment with PRANK. *Methods Mol Biol* **1079**, 155-170.
Loytynoja A, Vilella AJ, Goldman N (2012) Accurate extension of multiple sequence alignments using
a phylogeny-aware graph algorithm. *Bioinformatics* **28**, 1684-1691.
Meyer S, Nagel A, Gebhardt C (2005) PoMaMo--a comprehensive database for potato genome data.
Nucleic Acids Res **33**, D666-670.
Numa H, Itoh T (2014) MEGANTE: a web-based system for integrated plant genome annotation.
Plant Cell Physiol **55**, e2.

Ratnasingham S, Hebert PD (2007) bold: The Barcode of Life Data System
(http://www.barcodinglife.org). *Mol Ecol Notes* **7**, 355-364.
Sakai H, Lee SS, Tanaka T*, et al.* (2013) Rice Annotation Project Database (RAP-DB): an integrative
and interactive database for rice genomics. *Plant Cell Physiol* **54**, e6.
Taberlet P, Coissac E, Pompanon F*, et al.* (2007) Power and limitations of the chloroplast trnL (UAA)
intron for plant DNA barcoding. *Nucleic Acids Res* **35**, e14.
Veidenberg A, Medlar A, Loytynoja A (2016) Wasabi: An Integrated Platform for Evolutionary
Sequence Analysis and Data Visualization. *Mol Biol Evol* **33**, 1126-1130.
295

Tables

Table 1: Summary of data currently available in the PlantAligDB.

Target region	Genome	Type	Length (bp) in *Nicotiana tabacum*	Number of alignments	Number of sequences
*atp*F-*atp*H	cpDNA	Inter-genic spacer	502	31	1025
*psb*A-*trn*H	cpDNA	Inter-genic spacer	509	79	4852
*trn*L CD	cpDNA	Intron	577	44	2527
*trn*L GH	cpDNA	Intron	78	173	34674
*rbc*L	cpDNA	Protein-coding gene	1434	39	1748
*mat*K	cpDNA	Protein-coding gene	1530	113	11341
ITS	nuDNA	Transcribed spacers and 5.8S gene	678	35	9885
			Total	514	66052

Table 2: Average percentage of identical sites (PIS) values in all plant families organized by genomic region.

PIS	*atp*F-*atp*H	*psb*A-*trn*H	*trn*L CD	*trn*L GH	*rbc*L	*mat*K	ITS
Mean	55.05	39.1	61.13	58.62	78.48	65	28.84
Max	85.95	97.42	96.46	99.07	95.24	97.3	78.45
Min	15.19	2.3	21	6.43	27.47	0.16	1.13

Table 3: Average percentage of pairwise identity (PPI) values in all plant families organized by genomic region.

PPI	*atp*F-*atp*H	*psb*A-*trn*H	*trn*L CD	*trn*L GH	*rbc*L	*mat*K	ITS
Mean	95.49	94.02	96.34	97.09	96.94	95.23	87.21
Max	99.37	99.94	99.6	100	99.48	99.62	97.75
Min	87.59	69.96	90.54	88.54	81.18	78.72	70.79

314 **Legends to figures**
315
316 Figure 1: Workflow used to generate the curated alignments, phylogenetic trees, and genetic
317 conservation values stored in PlantAligDB.
318
319 Figure 2: Graphic representation of the measures of sequence conservation PPI and PIS for each
320 region-family alignment: a) *atp*F-*atp*H region b), *psb*A-*trn*H region, c) *trn*L CD region, d) *trn*L GH
321 region, e) *rbc*L region, f) *mat*K region and g) ITS region.

Selection of target genomic regions

↓

DNA sequences obtained in public databases

↓

Manual curation | removal of duplicated sequences from the same species
removal of sequences without a clear species assignment
reverse complement of sequences found in the opposite direction

↓

Organization of sequences by family according to the NCBI taxonomy

↓

Selection of families represented by ten or more species

↓

Alignment of sequences in each family

↓↑

Manual curation | removal of sequences that do not cover the entire target region
removal of sequences with large stretches of nucleotide ambiguities

↓

Final sequence alignment

Figure 1