# The knockout of the HMG- Box Domain of the porcine SRY-gene by CRISPR/Cas RNP microinjection causes sex reversal in gene-edited pigs

**Stefanie Kurtz[1],** Antje Frenzel[1], Andrea Lucas-Hahn[1], Petra Hassel[1], Roswitha Becker[1], Maren Ziegler[1], Monika Nowak-Imialek[1], Brigitte Schlegelberger[2], Gudrun Göhring[2], Heiner Niemann[3], Björn Petersen[1*]

[1] Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut, Mariensee, Neustadt am Rübenberge, Germany
[2] Institute of Human Genetics, Hannover Medical School, Hannover, Germany
[3] Twincore, Hannover Medical School, Hannover, Germany

*Corresponding author: bjoern.petersen@fli.de; Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut, Höltystraße 10, 31535 Neustadt am Rübenberge, Germany

## Abstract

The sex-determining region on Y chromosome (SRY) is presumed as a main genetic switch of male sex development. Mutations within the SRY gene are associated with a male-to-female sex reversal syndrome in humans and other mammalian species such as mice and rabbits. However, the underlying mechanism causing this syndrome is not definitely known. To understand the genetic pathway of SRY, the investigation and analysis of morphological changes are essential. For the first time, we successfully generated a knockout of the SRY gene in pigs using microinjection of two Clustered regularly interspaced short palindromic repeats (CRISPR) – associated protein - 9 nuclease (Cas9) ribonucleoprotein (RNP) complexes targeting the centrally located "high mobility group"- box domain (HMG) of the SRY gene. Mutation within the porcine HMG- box domain resulted in the development of complete external and internal female genitalia in genetically male piglets. Uteri and oviducts displayed equal morphology compared to female wildtype controls, only the ovaries showed smaller size in sex reversal piglets. In contrast to the HMG- box domain deletion, a deletion within the 5' flanking region of the HMG domain was not associated with sex reversal. In summary, this study demonstrates for the first time the main role of the HMG- box domain of the SRY gene triggering the male sex determination in pigs.

## Keywords

Porcine SRY gene; sex reversal; CRISPR/Cas9 RNPs; HMG domain

## 1 Introduction

In mammals, the male and female sex are determined by the presence or absence of the Y chromosome [1]. The sex-determining region on Y chromosome (SRY) located on the short arm of the Y chromosome is mainly involved in the gender determination during embryogenesis [2, 3]. It is expressed in the male genital ridge at the time of sex determination causing the formation of primary precursor cells of tubuli seminiferi, leading to the development of testicle from undifferentiated gonads [4]. Mutations within the SRY gene are associated with male-to-female sex reversal syndrome in humans (Swyer syndrome)[5]. Not all cases of sex reversal can be explained by alterations of the SRY gene, however 15 % of humans with male-to-female sex reversal syndrome show mutations or dysfunction of the SRY gene. Moreover, most of the mutations are detectable in the HMG- box domain of the SRY gene [6, 7]. The centrally located "high mobility group"- box domain (HMG) of the SRY protein

is responsible for DNA binding [8, 9] and thought to act as the main functional domain of the SRY gene [10]. Furthermore, it displays high similarity between different species [11, 12]. In previous studies in mice [13] and rabbits [14], the SRY gene was knocked out using different targets of the SRY sequence. However, the direct knockout of the HMG- box domain to elucidate the function of the HMG- box domain was not achieved. The pig is a perfect human disease animal model due to its high degree of physiological, genetic and anatomical similarity [15]. Therefore, to provide a basis for diagnosis of the sex reversal syndrome in humans, we target either the 5' flanking region of the HMG-box domain of the porcine SRY gene or directly the HMG- box domain to investigate its central function.

The porcine SRY gene consists of a single exon, similar to the human SRY gene, with an open reading frame of 624 bp representing 206 amino acids encoding the testis-determining transcription factor (TDF). It is expressed specifically in the male genital ridge at the time of sex determination. No extra gonadal expression of the porcine SRY gene is detectable [16]. First expression of the porcine SRY gene can be measured on day 21 postcoitum (p.c.) with its highest expression rate during day 21 and 23 p.c. Testis formation can be histologically observed by day 24 to 27 p.c. shortly after beginning of the SRY expression [17, 18]. According to this, the SRY gene is presumed to serve as the master gene causing formation of testicle from undifferentiated gonads. Furthermore, it is still unknown whether the SRY gene is the only sex-determining gene on the Y-chromosome or if other genes such as SOX9 [19-21] and SOX3 [8] are involved as well. In an effort to investigate the molecular mechanism of the sex determination in pigs, we envisaged to knockout different target sites on the porcine SRY gene by cell transfection followed by somatic cell nuclear transfer or intracytoplasmic microinjection of two CRISPR/Cas9 RNPs(Fig. 1).

**2 Materials and Methods**

**Animal Welfare.** Animals were maintained and handled according to the German guidelines for animal welfare and the German law regarding genetically modified organisms. The animal experiments were approved by an external animal welfare committee (Niedersaechsische Landesamt fuer Verbraucherschutz und Lebensmittelsicherheit, LAVES, AZ: 33.9-42502-04-17/2541).

**Fig. 1** Schematic illustration of the experimental design to generate SRY-KO piglets (XY-) by either
intracytoplasmic microinjection of two CRISPR/Cas9 RNP complexes into IVF-produced zygotes or SCNT
of SRY-knockout cells. Embryos were surgically transferred into hormonally synchronized recipients
and the offspring were analyzed pheno- and genotypically.

**Transfection of gRNAs.** The CRISPR/Cas9 system was employed to induce defined deletions
within the SRY gene (Ensembl transcript: ENSSSCG00000037443). Guide RNAs (gRNAs)
targeting either the 5' flanking region of the HMG- box domain of the SRY gene (SRY_1 and
SRY_2) or encompassing the HMG- box domain (SRY_1 and SRY_3) were designed using the
web-based design tool *CRISPOR* (http://crispor.tefor.net/) (Fig.2). Target sequences were
further analyzed via BLAST to reduce the off-target probability. The gRNA oligos with a BbsI
overhang were cloned into the linearized CRISPR/Cas9 vector pX330 (addgene #42230).
Afterwards, two CRISPR/Cas9 plasmids were co-transfected (with a final concentration of
5µg/µl) into male porcine fibroblasts by electroporation (Neon™ Transfection System,
ThermoFisher Scientific) to test the efficacy of the plasmids to induce double-stranded breaks
at the targeted locus. The electroporation conditions were as following: 1350 V, 20 mm, 2
pulses. After lysis of transfected cells, the cell lysate was analyzed using SRY specific primer
(SRY-F: 5'-TGAAAGCGGACGATTACAGC and SRY-R: 5'-GGCTTTCTGTTCCTGAGCAC-3'). The
purified PCR product (10ng/µl) (Invisorb® Fragment CleanUp – Startec) was sanger sequenced
to detect mutations at the target site (Supplements Fig. 1/4/5).

A

106
107

108    B

Exon SRY gene

601        1225

HMG-box SRY gene

C T C C T C A C T A T T T C C G A C G G A C A A T C A ... A T A A A T A C C G A C C T C G T C G C A A G G G A G A

*C C T* C A C T A T T T C C G A C G G A C A A T     A A A T A C C G A C C T C G T C G C A A *G G G*

PAM     sgRNA (SRY_1)       sgRNA (SRY_3)     PAM

**Fig. 2 a** Schematic illustration showing the guide RNAs (yellow underlined) targeting an approx. 72 bp
segment in the 5' flanking region of the HMG- box domain (red box) of the SRY gene. Primer
amplifying the SRY exon are indicated with green arrows. **b** Location of two sgRNA target sites
(yellow underlined) flanking the HMG- box domain (red box) of the SRY gene.

**In-Vitro-Fertilization and In-Vitro-Maturation.** Fresh boar semen was thawn for 30 sec. in a
water bath (37 °C). Motility of sperm was analyzed using microscopy. After washing with
Androhep® and centrifugation for 6 minutes at 600 g, approx. 75 to 100 sperm per oocyte
(depending on semen capacity) were used for fertilization (no sexed sperm was utilized for
fertilization). After four hours of incubation, the fertilized oocytes were cultured in porcine-
zygote-medium (PZM).

**Somatic cell nuclear transfer.** SCNT was performed as previously described [22]. Fetal
fibroblasts transfected with gRNA SRY_1 and SRY_2 targeting the flanking region of the HMG-
box domain of the SRY gene were used as donor cells (Supplement Fig 1). Eighty-two and
eighty-six one- to two-cell embryos were surgically transferred into two hormonally
synchronized (20mg/day/gilt Altrenogest for 12 days, followed by an injection of 1500 IU
PMSG on day 13 and ovulation was induced by intramuscular injection of 500 IU hCG 78 h
after PMSG administration) German Landrace gilts (7-9 months of age) (Supplements Table 1).

**Preparation of RNP complexes for microinjection.** The Alt-R CRISPR/Cas9 system (IDT)
consists of two CRISPR RNA components (crRNA and tracrRNA). The crRNA was individually
designed to target the HMG- box domain of the SRY gene (SRY_3: 5' –
AAATACCGACCTCGTCGCAA – 3'). To generate an active gRNA, both components (crRNA and
tracRNA) were annealed (95 °C for 5 min and then ramp down to 25 °C at 5 °C/min) in a ratio
of 1:1 to reach a final concentration of 1 μg/μl. Afterwards, the gRNA complex was mixed with
Alt-R S.p. Cas9 nuclease 3NLS and incubated for 10 minutes at room temperature to form an
active RNP complex with a final concentration of 20 ng/μl. The second RNP complex was
prepared using the individually designed synthetic single-guide RNA (SRY_1: 5' –
ATTGTCCGTCGGAAATAGTG – 3') from Synthego. The sgRNA was mixed with purified 2NLS-
Cas9 nuclease using a ratio of approximately 1:1,5 (0,84 μl sgRNA [25pmols] and 1,25μl Cas9
protein [25pmols]) and incubated for 10 minutes at room temperature. After a centrifugation
step at 10.000 rpm for 10 minutes (4°C), the supernatant was transferred into a new tube.
Both RNP complexes were mixed in a ratio of 1 (SRY_1) to 1,7 (SRY_3) and directly used for
microinjection.

**Microinjection.** The RNPs targeting the SRY gene were intracytoplasmatically co-injected into
IVF-produced zygotes obtained from slaughterhouse ovaries. Therefore, approx. 10 pL of the
RNP solution was injected with a pressure of 1 bar into the IVF-produced zygotes (Eppendorf
transjector 5246). The injected oocytes were cultured in PZM-3-Medium (porcine-zygote-

medium) at 37°C, 5% $CO_2$ and 5% $O_2$. At day 6 when embryos reached the blastocyst stage, 31 or 32 embryos were surgically transferred into three recipients (Table 1).

**Establishing cell cultures from SRY-KO piglets.** Porcine fibroblasts were isolated from ear tissue of the piglets and cultured in Dulbecco´s modified Eagle's medium (DMEM) with 2% penicillin/streptomycin, 1% non-essential amino acids and sodium pyruvate and 30% fetal calf serum (FCS). When the cells reached confluency, they were lysed and genomic DNA was used for further analysis (PCR and karyotyping).

**PCR-based genotyping.** Genomic DNA of the piglets was extracted from tail tips. Cells were isolated from ear tissue. The DNA concentration was determined by a NanoDrop photometer (ThermoScientific, Waltham, USA). For genotyping of the piglets, polymerase chain reaction (PCR) was employed using specific primer (SRY-F: 5'-TGAAAGCGGACGATTACAGC-3' and SRY-R: 5'-GGCTTTCTGTTCCTGAGCAC-3') flanking a 498 bp segment of the SRY gene (Fig. 2). PCR amplification was performed in a total volume of 50 µl : 20 ng DNA, 0,6 µM reverse and forward primer, 1,5 mM $MgCl_2$, 0,2 mM dNTPs and 1,25 U *Taq* Polymerase (Program conditions for 32 cycles with denaturation at 94°C for 30 sec, annealing at 59 or 60 °C for 45, extension at 72°C for 30 sec and a final extension at 72°C for 5 minutes). Standard conditions for electrophorese were 80 V, 400 mA and 60 min using 1 % agarose gel. The PCR-product was purified (Invisorb®Fragment CleanUp-Kit, Startec) and sanger sequenced (Fig. 4/5 and Supplement Fig 2/3). To further analyze the genotype of the piglets Y-chromosome-specific genes such as KDM6A, DDX3Y, CUL4BY, UTY, UBA1Y and TXLINGY were amplified with specific primer sets (Supplement Table 2).

**Karyotyping of the cells.** Karyotyping was accomplished on porcine fibroblasts isolated from ear tissue of the piglets. After treatment of cells for 30 minutes with colcemide (Invitrogen), cells were trypsinised and metaphases were prepared according to standard procedures. Fluorescence R-banding using chromomycin A3 and methyl green was performed as previously described in detail [23]. At least 15 metaphases were analysed per offspring. Karyotypes were described according to Gustavsson, 1988 and the International System for Human Cytogenetic Nomenclature (ISCN).

**Off-target analysis.** The top ten off-target effects were selected from the gRNA design tool *CRISPOR* (http://crispor.tefor.net/). PCR primers used for amplifying the PCR product are listed in Supplements Table 3 for SRY_1 and Table 4 for SRY_3. The PCR product was purified (Invisorb®Fragment CleanUp-Kit, Startec, Germany) and analyzed via Sanger sequencing.

**3 Results**

**Investigation of the 5' flanking region of the HMG domain.** In order to generate SRY-KO pigs, a deletion in the 5' flanking region of the HMG- box domain of the SRY gene was introduced. One recipient went to term and delivered two healthy genetically male offspring (704/1 and 2) with a deletion of approx. 70bp (Supplement Table 1, Fig. 2). Sequencing of the target site detected a genetic modification of -72bp and -73bp in both piglets (Supplement Fig. 3). The piglets showed male external and internal genitalia. No sex reversal did occur. Therefore, we conclude, that mutations within the 5' flanking region of the HMG- box domain does not affect

196 the formation of the SRY protein. Moreover, these results give evidence of an assumed
197 duplication of the porcine SRY gene.
198
**Production of SRY-KO pigs.** In the next step, we induced a deletion of approx. 300bp
encompassing the HMG- box domain of the SRY gene (Fig. 2b). Two recipients went to term
and delivered in total twelve healthy piglets with a female phenotype (Tab. 1/Fig. 3). No sexed
sperm was used for fertilization. Three of the piglets showed a deletion of approx. 300bp
within the HMG- box domain motif of the SRY gene (Fig 4). Sequencing of the target region
revealed in-frame mutations of -292bp in piglet 715/2 and -267bp in piglet 715/7. In piglet
714/1 two different mutations with a deletion of 298 bp and an indel formation including a
deletion of 298 and an insertion of 1 bp were detected (Fig. 5). Furthermore, six different Y-
chromosome-specific genes (KDM6A, TXLINGY, DDX3Y, CUL4BY, UBA1Y, UTY) were analyzed.
Detection of these genes approved a male genotype and sex reversal in these three piglets
(Supplements Fig. 6). To further confirm the male genotype of the piglets, isolated cells from
ear tissue were karyotyped. Standard karyotype of the pig includes 38 chromosomes[24]. The
results of the karyotyping confirmed the male genotype of the piglets by detection of the Y
chromosome (Fig. 6, Supplements Fig 7/8). No chromosomal abnormalities were detected in
the sex-reversed pigs, except for pig 714/1, in which an inversion of chromosome 7 was
detected. If this inversion occurred through the use of CRISPR/Cas or naturally occurred
remains unclear. Further analysis is required to answer this question.
In total, 34 putative off-target sites within the porcine genome were determined
(http://crispor.tefor.net/). We designed primers for the top ten off-target events for each
gRNA (Supplements Tab. 2/3). The analyzes of the off-target sites is still in process.
Our SRY-KO study underline the role of the SRY gene in the male sex development during
porcine embryogenesis.
221

| Recipients | Transferred Embryos | Pregnancy | Piglets | Phenotype | Sex reversal |
|---|---|---|---|---|---|
| 8018 | 32 | - | - | | - |
| 714 | 32 | + | 1 | female | 1 |
| 715 | 31 | + | 11 | female | 2 |

222
223 **Table 1** Results of the embryo transfer of microinjected zygotes into recipients. Three of twelve piglets
224 showed sex reversal with a female phenotype and a male genotype.
225

**Fig. 3** Twelve healthy piglets were born after cytoplasmic microinjection of two CRISPR/Cas9 RNP complexes into IVF-produced zygotes and surgical embryo transfer. Three of the piglets showed complete female external genitalia. The deletion of the SRY gene had no effect on growth rate compared to wildtype. All piglet developed normal.

**Fig. 4** PCR detection of the mutated SRY gene in piglets generated via microinjection of CRISPR/Cas9 RNP complexes. Three piglets (715/2, 715/7 and 714/1, indicated with white asterisk) showed deletions of approx. 300 bp within the SRY gene compared to a male wildtype control (WT 578 F7). The male WT control showed an expected band of approx. 500 bp. Female WT control (WT 578 F4) is negative as expected for SRY amplification.

**Fig. 5** Sanger Sequencing of the purified PCR product of SRY-KO piglets (715/2, 715/7 and 714/1) showing genetic modification of the SRY locus. Piglet 715/2 with a deletion of -292bp and in piglet 715/7 a deletion of -267bp. Piglet 714/1 show two different mutations with a deletion of -298bp and an indel formation of -298bp and +1bp .

**Fig. 6** Karyotyping of cells from SRY-KO piglet 715/2 confirming the male genotype of this piglet. The karyotypes of piglet 715/7 and 714/1 are shown in the Supplement Fig. 6.

**External and internal genitalia of SRY-KO pigs.** We explored the external and internal genitalia of the SRY-KO piglets. Wildtype (WT) females from conventional artificial insemination and age-matched WT females generated at the same time as the KO pigs using microinjection served as controls. The external genitalia of the SRY-KO piglets at the age of a few days and one month were equal to the external genitalia of WT females from microinjection and conventional artificial insemination. To further investigate the internal genitalia, the ovaries, oviducts and uteri of the SRY-KO piglets and female controls were prepared at the age of 34 days. As shown in Fig. 7 the SRY-KO piglet had complete female internal genitalia including ovaries, oviducts and uterus. Interestingly, the ovaries were approx. 2-fold smaller compared to the ovaries of the female WT control. There were no differences in size and morphology in oviducts and uteri between SRY-KO and WT females. These results approved that the SRY-KO causes a sex reversal in genetically male pigs with development of complete female external and internal genitalia.

**A**

**Piglet 715/2**  **WT Control (artificial insemination)**

277
278

279
**B**

280
**Fig. 7** The uteri and ovaries of the SRY-KO, XY piglet (715/2) and the WT,XX piglet (control from artificial
insemination) at day 34. **a** No differences were shown in size of oviduct and uteri. **b** Solely the ovaries
in SRY-KO, XY piglets are approx. 2-fold smaller than the ovaries of the WT, XX piglet (B).

**4 Discussion**

A promising candidate gene presumably involved in the male sex development is the sex-
determining region on Y chromosome (SRY) [17]. Nevertheless, the molecular function and

the question if the SRY gene is the main switch for male sex development in all mammals is
still unknown. Previous studies in mice [13] and rabbits [14] were performed to investigate
the potential role of the sex-determining region. The murine SRY gene was knocked out by
introducing a two base pair insertion in the 5' part of the ORF (open reading frame) of the SRY
gene causing a frameshift mutation of the sequence. In one genetically male offspring the
mutation led to a female phenotype [25]. While, in rabbits a disruption of the Sp1-binding site
in the 5' flanking region of the SRY gene resulted in a sex reversal [14]. Interestingly, in our
experimental setting in pigs a mutation upstream of the HMG- box domain of the SRY gene
via SCNT did not result in the generation of genetically male offspring with a female
phenotype. It is thought that the porcine SRY gene is present in two palindromic head-to-head
copies[26], as in rabbits [27]. But it is still unknown, if both copies of the porcine SRY gene are
active and required for male sex development and if there is the need of reaching a certain
threshold to induce the development of a male gender as it was previously described in mice
[28-30]. In both studies, the HMG- box domain as the main functional domain was not
examined. Here, we report for the first time the successful knockout of the HMG- box domain
of the porcine SRY gene using intracytoplasmic microinjection of two CRISPR/Cas9 RNP
complexes resulting in genetically male piglets with a female phenotype. CRISPR/Cas9 RNP
components exist only temporary in the cells limiting the guide RNA and Cas9 expression to a
short time frame. Therefore, the use of CRISPR/Cas9 RNP complexes enable efficient genome
editing while significantly reducing the risk of possible off-targets and undesired random
integration of DNA segments into the host genome compared to DNA plasmids. Furthermore,
mosaicism can be reduced to a minimum by using RNPs in microinjection technique [31-33].
The CRISPR/Cas9 system has become the genome editing technology of choice for most
application due to its ease of use, cost-effectivity and highly specific way to introduce
mutations at the targeted loci [34, 35]. Nevertheless, there is the possibility of off-target
cleavages at undesired genomic sites. Therefore, it is necessary to increase the specificity of
the CRISPR/Cas system regarding gRNA design [36].
In the present study, healthy SRY-KO piglets showing normal development and growth rate
were born. Moreover, the knockout of the SRY HMG- box domain resulted in piglets with a
female phenotype including complete female external and internal genitalia. Only the ovaries
were smaller compared to the female WT controls.  A previous study in rabbits reported that
SRY-KO rabbits had a dramatically reduced number of follicles. Although, a normal copulatory
behavior existed, no pregnancy was established by mating of an SRY-KO rabbit and a wildtype
male rabbit. However, transfer of embryos at the blastocyst stage from wildtype female
rabbits into pseudo pregnant SRY-KO rabbits results in a successfully pregnancy with the birth
of twelve kitten. Therefore, they assume that the abnormal development and reduced
number of follicles are responsible for the subfertility in the sex-reversal rabbits [14]. We
suggest that SRY-KO pigs could also develop a reduced fertility because of the smaller size of
ovaries. Nevertheless, in regard of the fertility rate and copulatory behavior more studies are
needed after the piglets reach sexual maturity. In conclusion, these results described here,
provide first evidence that the centrally located HMG- box domain on SRY gene is the
functional domain of the porcine SRY protein responsible for male sex determination. The pig
serve as a suitable animal model to investigate the male-to-female sex reversal syndrome in
humans due to its physiological and anatomical closer relation to human as mice. In addition
to its importance for sex reversal studies, the pre-determination of sex using CRISPR/Cas9
system in pigs could be a great benefit due to its impact on economic profit. The pig
production would extremely benefit from a shift to generate more female offspring as the
male-specific boar taint remains a big obstacle. Our results could pave the way to generating

a boar, which produces only female offspring by integrating a CRISPR/Cas9 vector targeting the HMG- box domain on SRY gene into the porcine genome.

**5 Conflict of interest statement**

The authors disclose any financial and personal relationships with other people or organizations that could inappropriately influence this work.

**6 Acknowledgements**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

**8 References**

[1] Waters, P.D., M.C. Wallis and J.A.M. Graves. *Mammalian sex—origin and evolution of the Y chromosome and SRY*. in *Seminars in cell & developmental biology*. 2007. Elsevier, doi: https://doi.org/10.1016/j.semcdb.2007.02.007.

[2] Parma, P. and O. Radi, *Molecular mechanisms of sexual development.* Sex Dev, 2012. **6**(1-3): p. 7-17 10.1159/000332209.

[3] Gubbay, J., J. Collignon, P. Koopman, B. Capel, A. Economou, A. Münsterberg, et al., *A gene mapping to the sex-determining region of the mouse Y chromosome is a member of a novel family of embryonically expressed genes.* Nature, 1990. **346**(6281): p. 245-250

[4] Brennan, J. and B. Capel, *One tissue, two fates: molecular genetic events that underlie testis versus ovary development.* Nature Reviews Genetics, 2004. **5**(7): p. 509

[5] Battiloro, E., B. Angeletti, M.C. Tozzi, L. Bruni, S. Tondini, P. Vignetti, et al., *A novel double nucleotide substitution in the HMG- box domain of the SRY gene associated with Swyer syndrome.* Human genetics, 1997. **100**(5-6): p. 585-587

[6] Berta, P., J.B. Hawkins, A.H. Sinclair, A. Taylor, B.L. Griffiths, P.N. Goodfellow, et al., *Genetic evidence equating SRY and the testis-determining factor.* Nature, 1990. **348**(6300): p. 448

[7] Hawkins, J.R., *Mutational analysis of SRY in XY females.* Human mutation, 1993. **2**(5): p. 347-350

[8] Bergstrom, D.E., M. Young, K.H. Albrecht and E.M. Eicher, *Related function of mouse SOX3, SOX9, and SRY HMG domains assayed by male sex determination.* Genesis (New York, NY: 2000), 2000. **28**(3-4): p. 111 DOI:10.1002/1526-968X(200011/12)28:3/4<111::AID-GENE40>3.0.CO;2-5.

[9] Sekido, R., *SRY: A transcriptional activator of mammalian testis determination.* The international journal of biochemistry & cell biology, 2010. **42**(3): p. 417-420

[10] Harley, V.R., M.J. Clarkson and A. Argentaro, *The molecular action and regulation of the testis-determining factors, SRY (sex-determining region on the Y chromosome) and SOX9 [SRY-related high-mobility group (HMG) box 9].* Endocrine reviews, 2003. **24**(4): p. 466-487

[11] Whitfield, L.S., R. Lovell-Badge and P.N. Goodfellow, *Rapid sequence evolution of the mammalian sex-determining gene SRY.* Nature, 1993. **364**(6439): p. 713

[12] R. Harley, V. and P. N. Goodfellow, *The biochemical role of SRY in sex determination.* Molecular reproduction and development, 1994. **39**(2): p. 184-193

[13] Kato, T., K. Miyata, M. Sonobe, S. Yamashita, M. Tamano, K. Miura, et al., *Production of Sry knockout mouse using TALEN via oocyte injection.* Sci Rep, 2013. **3**: p. 3136 10.1038/srep03136.

| 384 | [14] | Song, Y., T. Liu, Y. Wang, J. Deng, M. Chen, L. Yuan, et al., *Mutation of the Sp1 binding site in the 5' flanking region of SRY causes sex reversal in rabbits.* Oncotarget, 2017. **8**(24): p. 38176 doi: 10.18632/oncotarget.16979. |

384    [14]    Song, Y., T. Liu, Y. Wang, J. Deng, M. Chen, L. Yuan, et al., *Mutation of the Sp1 binding site in*
385            *the 5' flanking region of SRY causes sex reversal in rabbits.* Oncotarget, 2017. **8**(24): p. 38176
386            doi: 10.18632/oncotarget.16979.
387    [15]    Flisikowska, T., A. Kind and A. Schnieke, *Genetically modified pigs to model human diseases.*
388            Journal of applied genetics, 2014. **55**(1): p. 53-64
389    [16]    Parma, P., E. Pailhoux and C. Cotinot, *Reverse transcription-polymerase chain reaction analysis*
390            *of genes involved in gonadal differentiation in pigs.* Biology of reproduction, 1999. **61**(3): p.
391            741-748 https://doi.org/10.1095/biolreprod61.3.741.
392    [17]    Daneau, I., J. Ethier, J. Lussier and D. Silversides, *Porcine SRY gene locus and genital ridge*
393            *expression.*    Biology    of    reproduction,    1996.    **55**(1):    p.    47-53
394            https://doi.org/10.1095/biolreprod55.1.47.
395    [18]    Silversides, D., N. Pilon, R. Behdjani, A. Boyer, I. Daneau and J. Lussier, *Genetic manipulation*
396            *of sex differentiation and phenotype in domestic animals.* Theriogenology, 2001. **55**(1): p. 51-
397            63 https://doi.org/10.1016/S0093-691X(00)00445-3.
398    [19]    Stachowiak, M., I. Szczerbal, J. Nowacka-Woszuk, H. Jackowiak, P. Sledzinski, P. Iskrzak, et al.,
399            *Polymorphisms in the SOX9 region and testicular disorder of sex development (38, XX; SRY-*
400            *negative)    in    pigs.*    Livestock    Science,    2017.    **203**:    p.    48-53
401            https://doi.org/10.1016/j.livsci.2017.07.002.
402    [20]    Bishop, C.E., D.J. Whitworth, Y. Qin, A.I. Agoulnik, I.U. Agoulnik, W.R. Harrison, et al., *A*
403            *transgenic insertion upstream of sox9 is associated with dominant XX sex reversal in the mouse.*
404            Nature genetics, 2000. **26**(4): p. 490
405    [21]    Huang, B., S. Wang, Y. Ning, A.N. Lamb and J. Bartley, *Autosomal XX sex reversal caused by*
406            *duplication of SOX9.* American journal of medical genetics, 1999. **87**(4): p. 349-353
407    [22]    Hölker, M., B. Petersen, P. Hassel, W.A. Kues, E. Lemme, A. Lucas-Hahn, et al., *Duration of in*
408            *vitro maturation of recipient oocytes affects blastocyst development of cloned porcine*
409            *embryos.* Cloning and stem cells, 2005. **7**(1): p. 35-44
410    [23]    Schlegelberger, B., S. Metzke, S. Harder, R. Zühlke-Jenisch, Y. Zhang and R. Siebert, *Diagnostic*
411            *cytogenetics*. Classical and molecular cytogenetics of tumor cells., ed. R. Wegner. 1999:
412            Springer. 35.
413    [24]    Gustavsson, I., *Standard karyotype of the domestic pig: Committee for the Standardized*
414            *Karyotype    of    the    Domestic    Pig.*    Hereditas,    1988.    **109**(2):    p.    151-157
415            https://doi.org/10.1111/j.1601-5223.1988.tb00351.x.
416    [25]    Kato, T., K. Miyata, M. Sonobe, S. Yamashita, M. Tamano, K. Miura, et al., *Production of Sry*
417            *knockout mouse using TALEN via oocyte injection.* Scientific Reports, 2013. **3**: p. 3136, doi: DOI:
418            10.1038/srep03136.
419    [26]    Skinner, B.M., C.A. Sargent, C. Churcher, T. Hunt, J. Herrero, J.E. Loveland, et al., *The pig X and*
420            *Y Chromosomes: structure, sequence, and evolution.* Genome research, 2016. **26**(1): p. 130-
421            139 doi: 10.1101/gr.188839.114
422    [27]    Geraldes, A., T. Rambo, R.A. Wing, N. Ferrand and M.W. Nachman, *Extensive gene conversion*
423            *drives the concerted evolution of paralogous copies of the SRY gene in European rabbits.*
424            Molecular    biology    and    evolution,    2010.    **27**(11):    p.    2437-2440
425            https://doi.org/10.1093/molbev/msq139.
426    [28]    Washburn, L.L., K.H. Albrecht and E.M. Eicher, *C57BL/6J-T-associated sex reversal in mice is*
427            *caused by reduced expression of a Mus domesticus Sry allele.* Genetics, 2001. **158**(4): p. 1675-
428            1681 https://mouseion.jax.org/stfb2000_2009/232
429    [29]    Wu, N., A.-B. Yu, H.-B. Zhu and X.-K. Lin, *Effective silencing of Sry gene with RNA interference*
430            *in developing mouse embryos resulted in feminization of XY gonad.* BioMed Research
431            International, 2012. **2012** doi: 10.1155/2012/343891.
432    [30]    Nagamine, C.M., K.-i. Morohashi, C. Carlisle and D.K. Chang, *Sex reversal caused by Mus*
433            *musculus domesticus Y chromosomes linked to variant expression of the testis-determining*
434            *gene    Sry.*    Developmental    biology,    1999.    **216**(1):    p.    182-194
435            https://doi.org/10.1006/dbio.1999.9436.

[31] Svitashev, S., C. Schwartz, B. Lenderts, J.K. Young and A.M. Cigan, *Genome editing in maize*
*directed by CRISPR–Cas9 ribonucleoprotein complexes.* Nature communications, 2016. **7**: p.
13274
[32] Kim, S., D. Kim, S.W. Cho, J. Kim and J.-S. Kim, *Highly efficient RNA-guided genome editing in*
*human cells via delivery of purified Cas9 ribonucleoproteins.* Genome research, 2014. **24**(6): p.
1012-1019
[33] Farboud, B., E. Jarvis, T.L. Roth, J. Shin, J.E. Corn, A. Marson, et al., *Enhanced genome editing*
*with Cas9 ribonucleoprotein in diverse cells and organisms.* Journal of visualized experiments:
JoVE, 2018(135)
[34] Petersen, B., *Basics of genome editing technology and its application in livestock species.*
Reproduction in Domestic Animals, 2017. **52**(S3): p. 4-13
[35] Petersen, B. and H. Niemann, *Molecular scissors and their application in genetically modified*
*farm animals.* Transgenic research, 2015. **24**(3): p. 381-396 https://doi.org/10.1007/s11248-
015-9862-z.
[36] Tasan, I. and H. Zhao, *Targeting Specificity of the CRISPR/Cas9 System.* ACS Synth Biol, 2017.
**6**(9): p. 1609-1613 10.1021/acssynbio.7b00270.

**9 Supplements**
454

455
456 **Fig. 1** PCR after co-transfection of two plasmids (SRY_1 and SRY_2) in male fetal fibroblasts and
457 selection of the edited cells via single cell dilution (lower band). The edited cells revealed a mutation
458 of approx. 70bp (indicated with a white asterisk) compared to WT control. The male WT control (WT
459 578 F7) showed an expected band of approx. 500bp. The edited cells were further employed for
460 SCNT.
461
462

| Recipients | Transferred Embryos | Pregnancy | Piglets | Sex reversal |
|---|---|---|---|---|
| 704 | 82 | + | 2 | - |
| 7266 | 86 | - | - | - |

**Table 1** Transfer of embryos generated by SCNT into recipients. Two piglets were born. Both of them showed a male phenotype and no sex reversal.

**Fig. 2** PCR to detect genetic mutations in the SRY gene of the two piglet generated via SCNT. Both piglets showed a mutation of approx. 70bp compared to the male WT control (WT 578 F7). The WT control displayed an expected band of approx. 500bp.

```
T C C T C A C T A T T T C C G A C G G A C A A T C A … G G C C A G G A T C G T G T C A A G C G A C C C A T
C C T C A C T A T T T C C G A C G G A C A A T          C C A G G A T C G T G T C A A G C G A C C C A
T C C T C A C T – – – – – – – – – – – – – – – – … – – – – – – – – T C G T G T C A A G C G A C C C A T  -72bp
T C C T C A C – – – – – – – – – – – – – – – – – … – – – – – – – – T C G T G T C A A G C G A C C C A T  -73bp
                                    46bp
```

**Fig. 3** Both piglets (704/1 and 704/2) generated via SCNT of two gRNAs (SRY_1 and SRY_2) targeting the 5' flanking region of the HMG- box domain of the SRY gene showed two different mutation including a deletion of -72bp and -73bp.

477
478 **Fig. 4** PCR of cells after co-transfection of two gRNAs targeting the porcine SRY gene using
479 electroporation (Neon ™ Transfection System, ThermoFisher Scientific). Male WT control showed an
480 expected band of approx. 500bp. The additional lower band indicate a deletion of approx. 300 bp
481 caused by the CRISPR/Cas9 system at the SRY locus. For further experiments the combination of
482 SRY_1 and SRY_3 were used.

```
C T C C T C A C T A T T T C C G A C G G A C A A T C A … A T A A A T A C C G A C C T C G T C G C A A G G G A G A
   C C T C A C T A T T T C C G A C G G A C A A T          A A A T A C C G A C C T C G T C G C A A G G G
C T C C T C A C T – – – – – – – – – – – – – – – – – … – – – – – – – – – – – – – – – – – – – C A A G G G A G A
                                                    260bp
```

484
485 **Fig. 5** Sequencing of the PCR product from co-transfection of SRY_1 and SRY_3 revealed a deletion of
486 298 bp compared to the wildtype in fetal fibroblasts.

| Genloci | Primer | Sequence (5´- 3´) | Annealing Temp. (°C) | Length (bp) |
|---------|--------|-------------------|----------------------|-------------|
| KDM6A | KDM6A_1f | AAATCAAGGTATTACTTCACTATCCTG | 60 | 442 |
| | KDM6A_1r | ACCATGGAGATTTGACATCACCA | | |
| CUL4BY | CUL4BY_1f | AGCCAGCCGGATAGAAAGTT | 60 | 353 |
| | CUL4BY_1r | ACAAAAGACACTCAGTTAAAACTTACC | | |
| DDX3Y | DDX3Y_1f | ACTGGATTCTGTGTTCTTTGGA | 60 | 291 |
| | DDX3Y_2r | TTGGGGTGTTCTGTGCATGA | | |
| TXLINGY | TXLINGY_1f | CCTCTAGCTTGTGGTTGGCA | 60 | 249 |
| | TXLINGY_1r | GTGGCCCTCTGAATCTTGCT | | |
| UTY | UTY_1f | TCGTAAAGTGCTAAGTGGAGAAGA | 60 | 345 |
| | UTY_2r | TCAGGAACACACTGACGCTC | | |
| UBA1Y | UBA1Y_1f | GCTGACACACTCACTGACCA | 59 | 438 |
| | UBA1Y_1r | AGCCATCAGAATCGTGTGGG | | |

488
489 **Table 2** Primer set for six different Y-chromosome-specific genes are listed. They can be used for Y
490 chromosome detection.

| Genloci | Primer | Sequence (5´- 3´) |
| --- | --- | --- |
| intergenic:XM_021091945.1 | 101rev1_f | AGGGCTTTACATGGCTTCTCC |
| | 101rev1_r | TGGAGGCGCAAGTCATAGG |
| intron:XM_021075830.1 | 101rev2_f | GTCCTCATGCTCTCGGCTAC |
| | 101rev2_r | CCAACTGGACCCCTAGCTTG |
| intron:XM_005663347.3 | 101rev3_f | GTAGGTGTGGCCTTCGTCTT |
| | 101rev3_r | CCATCAGGGGGCTATCACAC |
| intron:XM_021091957.1 | 101rev4_f | CTCTGACTTGCACCCTGCTT |
| | 101rev4_r | ACTTCTCAATCCGCCCTATGC |
| exon:XM_021086306.1 | 101rev5_f | AACAACATGCGTCCAAACCG |
| | 101rev5_r | GCATCAGCACTCACCTGGAT |
| exon:XM_021069624.1 | 101rev6_f | AGTGACTGGGTTTGGGGTTG |
| | 101rev6_r | CGCCAGAGTCCCATACACTC |
| intergenic:XM_021090127.1 | 101rev7_f | TCCATAAGGAGTGCACGACG |
| | 101rev7_r | TTTGCCCAGGTTCTGTCTGG |
| intergenic:XM_021082147.1 | 101rev8_f | GGGTGAATCCTGACCACGTC |
| | 101rev8_r | AGTGTCCTTAGGGGGCAGAA |
| intergenic:XM_021101305.1 | 101rev9_f | TAGCACCCCCAGAAACTCCT |
| | 101rev9_r | GGTGGATACTGTCAGCTGGG |
| intergenic:NM_001315615.1 | 101rev10_f | GCCTGCGCTTCTCATTTAGC |
| | 101rev10_r | ATCCTGGACCTGGGCCTTAT |

492
493 **Table 3** Top ten off-target sites for gRNA SRY_1 on the porcine genome. Primer pair for sequencing of
494 the off-target loci are listed.
495

| Genloci | Primer | Sequence (5´- 3´) |
|---|---|---|
| intergenic:XM_021073683.1 | 408fw1_f | CTCCCCACAGCTGCTCTTTT |
| | 408fw1_r | GAATTGGGCACTTGCTGGAC |
| intergenic:XM_003358165.4 | 408fw2_f | CTGTGCTACGTCCCCCAAAT |
| | 408fw2_r | GCTTGTGGTCTTTGCCAGTG |
| intergenic:XM_021065302.1 | 408fw3_f | AGAGGAATGGGCAAACAGCA |
| | 408fw3_r | CCGTGGTTAGCACCTGGTAG |
| intergenic:XM_021093786.1 | 408fw4_f | TTGGTCTGGGGAAGAAACCG |
| | 408fw4_r | TCTGCCCTACCACTTCAGGA |
| intergenic:XM_003134254.5 | 408fw5_f | AGCCTTATCCAATGAGGCCG |
| | 408fw5_r | CTAATGCCAGGGCAGTTTGC |
| intron:XM_021070776.1 | 408fw6_f | CCTTACCTGGTCCAGTTGGC |
| | 408fw6_r | GCATTCATTTCCCCACGCAT |
| intron:XM_021069725.1 | 408fw7_f | ACAAAGTCCCTAGCTCACGC |
| | 408fw7_r | GCTCCGCATCCTCAGATGTT |
| intergenic:XM_013983698.2 | 408fw8_f | GTCACCTGCCACTCTCTCAC |
| | 408fw8_r | GGTCTGGAGTTCCTGTCGTG |
| intron:XM_021090110.1 | 408fw9_f | CTTCTCTGGTAACTGGCCCC |
| | 408fw9_r | TCCCGCAGATCCATTCCAAC |
| intergenic:NM_001204768.1 | 408fw10_f | GACAGTGTTGGCGTATGCAC |
| | 408fw10_r | GGAAAGTCAGTAGGGCCGAG |

496
497 **Table 4** Top ten off-target sites for gRNA SRY_3 on the porcine genome. Primer pair for sequencing
498 the off-target loci are listed.
499

503 **Fig. 6** PCR of six different Y-chromosome-specific genes (KDM6A, CUL4BY, DDX3Y, TXLINGY, UTY and
504 UBA1Y) for detection of Y chromosome in SRY-KO piglets (715/2, 715/7 and 714/1, indicated by a
505 white asterisk) compared to female wildtype controls (715/1, 715/3 and 715/4) from same litter
506 generated by microinjection. Moreover, a male (WT 578/F7) and a female (WT 578/F4) DNA sample
507 was used as control.
508

509
**Fig. 7** Karyotyping of the SRY-KO piglet 715/7 confirming the male genotype by analysis of the sex
chromosomes.
512

513
514 **Fig. 8** Karyotyping of the SRY-KO piglet 714/1 confirming the male genotype by analysis of the sex
515 chromosomes. In this piglet, a clonal aberration (inversion) could be found in chromosome 7.
516