Incursion of killer sponge *Terpios hoshinota* on the coral reefs of

Lakshadweep archipelago

Rocktim Ramen Das[1,2]., Sreeraj, C.R[1]., Gopi, M[1]., Abhilash, K.R[1]., Deepak Samuel.V[1]*.,

Purvaja, R[1]. and Ramesh, R[1].

[1]*National Centre for Sustainable Coastal Management, Ministry of Environment, Forest and*

Climate Change, Chennai, India

[2]*Graduate School of Engineering and Science, University of the Ryukyus, Nishihara, Okinawa,*

Japan

*Email: deepakocean@gmail.com

Abstract

Our Study documents the outbreak of killer sponge *Terpios hoshinota* on the coral reefs of

Lakshadweep archipelago, Arabian Sea.

Introduction

Coral killing sponges have the potential to overgrow live corals, eventually killing the coral

polyps, and thus leading to an epidemic[1]. The cyanobacteriosponge *Terpios hoshinota* Rützler & Muzik,

1993 first reported from Guam[1] and later described from the coral reefs of the Ryukyu archipelago

(Japan)[2], is identified by its gray to blackish encrustations. Since its first occurrence, it has been observed

in several coral reef localities around the globe viz. the Great Barrier Reef[3], Papua New Guinea[4],

Taiwan[5], Philippines[6], Indonesia[7], South China Sea[8,9], Thailand[6], Palk Bay (PB)/Gulf of Mannar (GOM)

(India)[10-12], Maldives[13], Mauritius[14] and our present observation, confirms that the species has further

extended its habitat into the pristine atolls of Lakshadweep (Figure 1) (Indian Ocean) and requires urgent

attention.

Results and Discussion

During the coral reef surveys conducted at Lakshadweep in November 2016, *Terpios hoshinota*

was observed overgrowing on several colonies of *Acropora formosa, Acropora palifera, Cyphastrea* sp.,

Favia lizardensis and *Porites lutea* (Figure 2 and 3) in the atoll encircling Bangaram and Thinnakara

Islands. Out of 33 stations surveyed, 6 stations exhibited the presence of *T.hoshinota* incrustations (Figure

1). The coral colonies in the atoll were patchy and the depth varied between 2 to 12 meters. As depth

increased, (i.e. >5 m) large boulder corals were observed whereas the shallow regions (<5 m) had greater

coral diversity. Certain areas consisting of large *Acropora* beds, rocks, rubbles and dead reef were also

observed. The affected corals displayed grayish/blackish encrustations of *T.hoshinota* forming a mat-like

layer on live corals taking the shape of the coral in all cases. The osculum in the sponge, a primary

character with a radiating network was clearly visible, the thickness of the mat was less than 1mm (Figure

1a). It was observed that the encrustations were propagating laterally and infecting the other live coral

colonies. Further, in some colonies along with *T.hoshinota*, algal presence was also noted (Figure 3a) but

the sponge was absent in the colonies which were completely covered with turf (Figure 3b). Other

associated communities such as ascidians and clams remain unaffected but interestingly the calcareous

serpulid tubes, though overgrew by the *Terpios* like sponge, the animal was unharmed[15] (Figure 2d).

Environmental parameters assessed in the atoll revealed that the area was unpolluted with an optimum

level of dissolved oxygen (5.04~8.21 mg/l), and low turbidity (0.3 to 0.8 NTU). Sea surface temperature

(SST) during the survey was 28.2°~30.1°C. It is important to note that, Bangaram and Thinnakara is one

of the few atolls in Lakshadweep where tourism is permitted, as a result, limited amounts of diving and

other water-related recreational activities can be seen in the area.

Previous studies[2,10] suspected that the outbreak of *T. hoshinota* is related to increased water

turbidity. Polluted seawater following anthropogenic impacts favours the development of *T.hoshinota* as

seen in Guam[2,6] and Green Island[16]. Our findings contradict these above statements and is more in line

with the findings of Shi et al.[8], who observed *T.hoshinota* outbreak in unpolluted areas of Yongxing

Island (South China Sea). Prevalence of *T.hoshinota* in the southeastern reefs of India (~800km from

Lakshadweep) is due to high anthropogenic stress and its close proximity to mainland[10-12], however, a

similar conclusion cannot be applied in the case of Lakshadweep because of its isolated geography[17] and

with comparatively less anthropogenic activities. In terms of host selectivity, the killer sponge has

affected several coral species in different parts of the world[1,10,12,14] and in the reefs of Palk Bay (PB), it

affected all genus surveyed[10]. In Vaan Island (GOM) the dominant genus *Montipora* was the most

susceptible[12]. Our observation though could not reveal any specific host coral selectivity, we can

speculate that the dense *Acropora* (ACB) beds in station 3, 5 and 6 were more easily overgrown because

the killer sponge prefers branching corals as reported from Mauritius[14]. We would further conclude that

the coral composition in any specific location may play an important role in determining its host.

Terpios hoshinota is a belligerent contender for space[6] and is known to overgrow corals from its

base where it interacts with turf algae[14]. In stations 3, 5 and 6 (Figure 3a), we observed ACB colonies

with both algae (e.g. *Ceratodicyton repens*? *Dictyota* sp.) and killer sponge, however, the observation of a

massive turf algal cover dominating an area of ~0.35km of *Acropora* bed in *T.hoshinota* occurrence site

station 5 and 6 indicates towards a much complex ecological scenario. Such complexity between sponges,

corals and algae can be only understood through long term monitoring. Gonzalez-Rivero et al.[18] stated

that sponges can act as a potential group that can facilitate and influence coral-algal shifts by acting as a

"*third antagonist*" as observed in Glover's atoll (Belize). Based on our knowledge of the life history of

T.hoshinota we can just hypothesize the scenario in station 5 and 6 as follows: - (1) *T.hoshinota* invades

and overgrows the *Acropora* beds → (2) The coral dies which is followed by the death of the killer

sponge → (3) Turf algae takes over. (Figure 3a, b). Moreover, reports of turf algae being a dominant

component in the atolls[17] might indicate towards a faster transition. Globally Elevated SST is a major

threat to coral reefs[19], and the reefs of India[20,21,22] including the atolls[23] are no different. With reports

indicating that elevated SST has already depleted the coral ecosystem of Lakshadweep which was evident

during 1998[17], 2010[23] and 2016[19] mass bleaching events, it can provide an opportunity for sponges to

invade[24]. Dynamics of waterflow[17] may also play a crucial role in this regard.

Conclusion

Our findings confirm that the infestation of *T.hoshinota* on the coral colonies of Lakshadweep is currently

limited to only Bangaram and Thinnakara atoll as it was not observed in other islands surveyed. There is

although a possibility that the killer sponge could invade nearby atolls as seen in other regions[1,25]. Our

observation is indeed the first documentation of *T.hoshinota* on the reefs of Lakshadweep and can be

regarded as a baseline for subsequent studies in the islands. Further, to protect the reefs of Lakshadweep,

coral health monitoring is required which will allow us to understand the nature of occurrence,

distribution, the impact and the causative factors of the killer sponge as well as other coral diseases as per

which various coral reef management plans can be initiated.

Acknowledgement

This study was undertaken as part of the grant-in-aid project of "Mapping of Ecologically Sensitive Areas

(ESAs) and Critically Vulnerable Coastal Areas (CVCAs) along the Coast of India", "[F.No. 22-29/2008-

WBICZM-IA-III; 19 June 2014]". The authors acknowledge the financial and technical support provided

by MoEF&CC, Government of India, and the World Bank under the India ICZM Project and are grateful

to Dr. Shao-Lun Liu (Tunghai University) for identifying few algae species.

References

1. Bryan, P., *Micronesica*, 1973, **9**:237–242.

2. Rützler, K. and Muzik, K., *Scientia Marina*, 1993, **57**(4), 395-403.

3. Fujii, T. *et al., Coral Reefs*, 2011, **30**: 483.

4. Ekins, M. *et al., Mem. Queensl. Mus*, 2017, **60**: 174-175.

5. Liao, M. H. *et al., Zool. Stud.*, 2007, **46**(4): 520.

6. Plucer-Rosario, G. *Coral Reefs*, 1987. **5**(4), 197-200.

7. De Voogd, N. J. *et al., Coral Reefs*, 2013, **32**(3), 755.

8. Shi, Q. *et al., AMBIO*, 2012, DOI 10.1007/s13280-011-0245-2

9. Hoeksema, B. W. *et al., Bull. Mar. Sci.,* 2014 **90**(4): 989-990.

10. Thinesh, T. *et al., Curr. Sci.,* 2015*,* **109**(6): 1030-1032.

11. Thinesh, T. *et al., Dis. Aquat. Org.,* 2017, **126**: 63-70.

12. Diraviya Raj, K. *et al., Curr. Sci.,* 2018, DOI: 10.18520/cs/v114/i05/1117-1119

13. Montano, S. *et al., Bull. Mar. Sci.*, 2015, **91**(1), 000-000.

14. Elliott, J. *et al., Coral Reefs*, 2016, 1-13; DOI 10.1007/s00338-016-1434-z

15. García-Hernández, J. E. and Hoeksema, B. W., *Coral Reefs*, 2017, DOI 10.1007/s00338-017-1617-2

16. Chen, S. L. *et al., Journal of National Park*, 2009, **19**: 33–45.

17. Arthur, R. *et al., Curr. Sci.,* 2005, **89**(4), 694-699.

18. Gonzalez-Rivero, M. *et al., Ecol. Modell.,* 2011, **222**, 1847–1853.

19. Hughes, T. P. *et al., Science*, 2018, **359**, 80-83.

20. Krishnan, P. *et al.*, *Curr. Sci.,* 2018, DOI: 10.18520/cs/v114/i03/679-685

21. Diraviya Raj, K. *et al., Curr. Sci.,* 2018, DOI: 10.18520/cs/v114/i02/384-387

22. Patterson Edward, J.K. *et al., Curr. Sci,* 2018, DOI: 10.18520/cs/v114/i09/1967-1972

23. Vinoth, R. *et al., J. Ocean Univ. China*, 2012, **11**(1), 105-110.

24. Bell, J. J. *et al., Global. Change. Biol.,* 2013, **19**, 2613-2624.

25. Reimer, J. D. *et al., Galaxea*, 2011, **13**, 35–44.

129
130

Figure 1. Bangaram & Thinnakara Atoll (Red star); killer sponge infested areas (1-6) (QGIS 3.0.1)

Figure 2. Incrustations of *T.hoshinota* (A) *Acropora formosa*, A1. *T.hoshinota* exhibiting osculum with radiating networks
(B) Incrustation on *A.palifera*,B1. *T.hoshinota* mat, B2. Bleached ring, B3. Live coral (C) *T.hoshinota* taking shape of a
Coral (*Cyphastrea* sp.) (D) *Terpios hoshinota* overgrowing calcareous serpulid tubes, animal unaffected.

Figure 3. *Acropora* colonies (Station 3) A1. *T.hoshinota* A2. Algae (B) *Acropora* colonies (Station 5) completely over grown
by turf algae, killer sponge absent.