1

**Evaluation of biodiversity metrics through environmental DNA metabarcoding**

**outperforms visual and capturing surveys**

Hideyuki Doi[1,*,†], Ryutei Inui[2,*,†], Shunsuke Matsuoka[1, †], Yoshihisa Akamatsu[2], Masuji

Goto[2], and Takanori Kono[2]

[1] Graduate School of Simulation Studies, University of Hyogo, 7-1-28 Minatojima

Minami-machi, Chuo-ku, Kobe, 650-0047, Japan

[2] Graduate School of Science and Engineering, Yamaguchi University, 2-16-1

Tokiwadai, Ube, Yamaguchi, 755-8611, Japan

[†]These authors equally contributed to this study.

[*]Corresponding authors:

Hideyuki Doi (hideyuki.doi@icloud.com)

Ryutei Inui (inuiryutei@gmail.com)

**Abstract**

Information on alpha (local), beta (between habitats), and gamma (regional) diversity is

fundamental to conserving biodiversity and the functions and stability of ecosystem

processes. Robust methods like environmental DNA (eDNA) metabarcoding are

currently considered useful to investigate biodiversity. However, the performance of

eDNA methods in evaluating diversity has not been tested quantitatively. We compared

the performance of eDNA metabarcoding and visual and capturing surveys in

estimating alpha, beta, and gamma diversity in river fish communities, particularly

considering community nestedness and turnover. In five rivers across west Japan, when

compared with visual and capturing surveys, eDNA metabarcoding detected higher

alpha and gamma diversity in local habitats, and indicated differences in beta diversity

more clearly; this suggests the superiority of eDNA metabarcoding over

visual/capturing surveys in estimating diversity. The statistical frameworks, particularly

nestedness and turnover, can provide quantitative evidences needed to assess diversity

component estimation by new survey methods.

The maintenance of biodiversity underpins the stability of ecosystem processes in constantly changing environments[1,2,3]. Moreover, biodiversity loss affects ecosystem functions and services and consequently human society[1,2,3]. Ecologists have made efforts to conserve biodiversity based on essential biodiversity data, e.g., species richness and distribution[1,2,3,4]. Biodiversity can be evaluated in different ways: viz., by estimating alpha (local), beta (between habitats), and gamma (regional), diversity. To conserve local communities, ecologists incorporated these diversity measurements into management decision-making[1,2,5]. For example, beta diversity can quantify biodiversity loss and inform the placement of protected areas and the management of biological invasions and landscapes[5]. Thus, robust methods for monitoring biodiversity are fundamental for biodiversity and environmental management.

Since recent advances in molecular techniques, environmental DNA (eDNA) analysis has been considered a useful tool to investigate the distribution and richness of aquatic and terrestrial organisms[6,7,8,9,10,11,12,13,14]. High-throughput sequencing (HTS) derived from eDNA, called "eDNA metabarcoding", became a useful and powerful tool for biodiversity surveys[13,16,17,18,19,20]. eDNA metabarcoding has recently been applied in fish community surveys, e.g., Miya et al.[11] (2015) developed the universal PCR primers for fish community (MiFish primers). To confirm the usefulness of eDNA metabarcoding for community assessment, many studies conducted eDNA metabarcoding comparing the observed species list with traditional direct surveys including visual and capturing surveys[13,16,20,21,22]. However, evaluating the performance of eDNA metabarcoding estimating alpha and beta diversity was still limited quantitively and statistically, especially when evaluating beta diversity and comparing to traditional surveys.

Beta diversity is a fundamental aspect for communities, and it is important for

evaluating community responses to environmental gradients and the spatial locations

across the different habitats and ecosystems[5,23,24,25]. Beta diversity itself and the trends

along environmental gradients such as productivity and disturbance can be affected by

alpha diversity[24]. Furthermore, beta diversity can decline even if alpha and gamma

diversity remain unchanged or even increase in the area, mainly due to biological

homogenization[26,27,28]. Therefore, beta diversity evaluation of communities would be

considerably different among the survey methods when alpha and gamma diversity

evaluations were different between the survey methods, i.e., eDNA metabarcoding and

traditional surveys.

Beta diversity is considered to reflect two different components: nestedness

and spatial turnover[29,30,31]. Nestedness of species assemblages in the communities

occurs when the community at the sites with less species are subsets of the community

at the sites with higher species richness[32,33]. Nestedness generally reflects a non-random

process of species loss as a consequence of any factor that promotes the orderly

disaggregation of assemblages[34]. Contrastingly, spatial turnover implies the replacement

of some species by others because of environmental sorting or spatial/historical

constraints[31]. Baselga[31] (2010) developed statistical separation methods to evaluate beta

diversity considering nestedness and spatial turnover and applied them to beta-diversity

evaluation in various systems[31,35]. However, the method has never been applied to

evaluate the performance of eDNA metabarcoding estimating beta diversity. Moreover,

Baselga's framework[31] (2010) can be applied to comparing the performance among

methods when evaluating alpha diversity via nestedness and species turnover.

Using statistical methods, we can quantitatively and statistically compare the

performance of eDNA metabarcoding and traditional surveys in community evaluation

using alpha, beta, and gamma diversity. Here, we tested the performance of eDNA

metabarcoding in five river systems in different regions with various fish species. We

conducted eDNA metabarcoding using the MiFish universal primer for fish and also

identified the fish by visual snorkeling and hand-net capturing surveys. We evaluated

the performance of eDNA metabarcoding by comparing the obtained fish community

structure to that evaluated by visual/capturing survey with special regard to nestedness

and spatial turnover. We finally validated the ability of eDNA metabarcoding to

estimate biodiversity.

**Results**

**Overview.** We detected 53 fish taxa, almost all identified to species or genus level, by

eDNA metabarcoding in five rivers (Table S1, and S2) and visually observed 38 fish

taxa in total. A MiSeq paired-end sequencing for the 40 libraries (30 samples plus 10

negative controls) yielded a total of 1,601,816 reads (53,351 ± 17,639; mean ± S. D. for

each sample, Table S2). We confirmed very low reads from negative controls (Table

S2).

A**lpha, beta, and gamma diversity between the methods.** We found significant

differences in fish richness between eDNA metabarcoding and visual/capturing surveys

(Fig. 1, GLMM, t = –5.45, P = 0.000018). Richness was not significantly different

among river segments (t = –5.85, P = 0.000004) but not among rivers (t = 1.737, P =

0.0942), indicating higher alpha and gamma diversity estimated by eDNA

metabarcoding than by visual/capturing surveys.

We found differences in community structures between the two methods by

NMDS ordination (Fig. 2). PERMANOVA results for the ordination suggested there

were differences in community composition evaluated by each method, eDNA

metabarcoding and visual/capturing survey (P = 0.003, Table S3 for full statistical

results). Moreover, communities from the combined results of eDNA metabarcoding

and visual/capturing survey were significantly different among rivers and segments (P =

0.001, Table S3). We found different patterns in ordinated river sites for each method

(Fig. 2). The PERMANOVA results determined that communities were significantly

different among the rivers by eDNA metabarcoding (P = 0.011) but not by

visual/capturing survey (P = 0.12, Table S4 for full statistical results). Conversely, the

communities were significantly different among river segments by visual/capturing

survey (P = 0.011) but marginally not significantly different by eDNA metabarcoding

(P = 0.061, Table S4). The differences in PERMANOVA results suggested that

differences in beta-diversity among rivers across regions can be detected by eDNA

metabarcoding but not by visual/capturing survey.

Indicator taxa analysis comparing the communities estimated by both

methods, eDNA metabarcoding and visual/capturing survey, detected significantly

different method preferences of several taxa, including Japanese eel (*Anguilla*

*japonica*), salmon (e.g., *Oncorhynchus keta)*, and Amur catfish (*Silurus asotus*),

between both methods ($p < 0.05$, Table 1 for statistically significant taxa, Table S5 for

all taxa).

**Nestedness and turnover**. We compared the pairwise standardized effect size (pSES)

between eDNA metabarcoding and visual/capturing survey in nestedness and species

turnover (Fig. 3). Nestedness pSES was significantly positive without overlapping the

zero-pSES 95% confidence interval, indicating that the visual/capturing survey

community was nested in that detected by eDNA metabarcoding. The significantly

negative pSES in species turnover indicated that the taxa did not significantly turnover

in the community evaluated by eDNA metabarcoding and visual/capturing survey

comparing with random communities.

We compared the longitudinal nested structure (upstream to downstream) of

the fish community in the study rivers using nested pSES (Fig. 4a). The longitudinal

nested structure was not significantly different between both methods (GLMM, $P =$

0.302). The positive pSES indicated that downstream community significantly nested

the upstream ones. The species per segment nested as downstream > mid-stream >

upstream in both methods (Fig. S4, 5, NOFD, $P < 0.001$). We also compared the

longitudinal spatial turnover of fish community in the study rivers using turnover pSES

(Fig. 4b). The longitudinal spatial turnover was not significantly different between both

methods (GLMM, $P = 0.280$).

**Discussion**

We found that river fish communities estimated by eDNA metabarcoding significantly

nested the communities estimated by visual/capturing survey. Furthermore, the species

turnover in the communities was very weak between methods. Nestedness of species

assemblages occurs when the communities obtained by the method estimating lower

number of species are subsets of the communities estimated by other methods with

higher species richness. Although it is difficult to assume the effort of sampling was

enough to evaluate the species compositions, we assumed the efforts for the visual and

capturing survey are enough to confirm the species composition in the study sites

because we combined two high-detection surveys, i.e. visual observation and capturing

surveys[12,13]. Therefore, the performance of eDNA metabarcoding in estimating alpha

and gamma diversity outperformed that of visual/capturing survey for river fish

community. In fact, the fish local and regional richness (alpha and gamma diversity)

evaluated by eDNA metabarcoding was significantly higher, including almost all taxa

evaluated by visual/capturing survey.

168           eDNA metabarcoding has been reported to perform better than traditional

methods in evaluating species richness[13,16,19,20,21,22]. Nakagawa et al.[21] (2018)

investigated freshwater fish community in 100 rivers and confirmed that the community

detected by eDNA metabarcoding was similar to the species list observed in

governmental-authorized monitoring. Furthermore, several eDNA metabarcoding

studies on fish communities have been performed in other river systems[20], marine

habitats[22] , and freshwater lakes[13,19]. Deiner et al.[16] (2016) showed that river eDNA

metabarcoding can reflect the community in a watershed, indicating that eDNA

metabarcoding has high-performance gamma diversity evaluation. These studies

indicated the great potential of eDNA metabarcoding as a useful tool for alpha and

gamma diversity assessment by simply comparing the community data obtained from

eDNA metabarcoding and traditional surveys. However, previous studies did not

evaluate the performance by nestedness and species turnover between eDNA

metabarcoding and other community data. Here, we supported previous literature by

showing nestedness, species turnover, and species preferences between the communities

detected by eDNA metabarcoding and traditional methods, and we found the nestedness

structure eDNA metabarcoding > visual/capturing survey, with scarce species turnover

in the community.

We especially focused on beta-diversity for assessing the performance of

eDNA metabarcoding compared to the visual/capturing survey. Higher beta diversity of

the rivers were statistically detected by eDNA metabarcoding than by visual/capturing

surveys. Furthermore, the spatial nestedness and turnover were not significantly

different between eDNA metabarcoding and traditional methods. These results

suggested that eDNA metabarcoding also outperformed visual/capturing survey in

evaluating river fish community beta diversity, probably due to the lower alpha

diversity detected by visual/capturing survey, especially in the upstream segments.

Moreover, beta diversity between segments could be detected by visual/capturing

surveys but not by eDNA metabarcoding. These differences in beta diversity evaluation

may lead us to interpret beta diversity using the results from both survey methods,

eDNA and traditional survey. Deiner et al.[16] (2016) suggested that eDNA sampling

allows an estimate of catchment-level diversity and integrates this information across

space due to downstream transport of eDNA. We agree with the suggestion for estimate

of catchment-level diversity and further suggest that the catchment-level diversity

metrics including alpha, beta, and gamma diversity, can be evaluated similarly to the

direct survey.

We further compared the indicator taxa for the communities obtained from

both eDNA metabarcoding and visual/capturing survey and concluded that several taxa,

including eel, salmon, and catfish, were significantly better detected by eDNA

metabarcoding, whereas non-indicator taxa were detected by visual/capturing surveys.

These results indicated that eDNA metabarcoding outperformed visual/capturing

surveys in fish taxa detection. The community structures estimated by eDNA

metabarcoding and visual/capturing survey were slightly different, as reported in

previous studies[13,19], probably because of the differences in taxa-detection

performances. Discriminated taxa in this analysis included eel, salmon, and catfish,

which mostly had larger body size and lower abundances in these rivers[36,37]. In fact, the

Japanese eel *Anguilla japonica*, was difficult to find by visual observation, probably due

to its hiding behavior[38]. Such endangered species would be important as top-predator[37].

eDNA metabarcoding can evaluate the distribution of such rare and important taxa in

fish communities better than traditional surveys.

To evaluate the comparing among the local sites and the rivers, we examined

the performances of eDNA metabarcoding at three sites of five rivers with eDNA

sampling from only two habitats. Thus, our understanding of details in the fish

community spatial structure in the rivers and the performance of community evaluation

in local habitats, such as backwater, was still limited. In fact, Bylemans et al.[20] (2018)

found that river morphology in the habitats influenced the optimal sampling strategy for

eDNA metabarcoding. Moreover, in backwater lakes, the performance of eDNA

metabarcoding varied with different lake morphology[13]. However, testing the usefulness

of eDNA metabarcoding for assessing river fish community biodiversity has been

limited. Further research is needed to evaluate fish community spatial structure in

rivers.

In conclusion, eDNA metabarcoding outperformed the visual/capturing

survey in evaluating alpha, beta, and gamma diversity of fish communities in five rivers

across the west Japan region. Biodiversity testing using statistical frameworks,

especially community nestedness and turnover, provided the quantitative evidence

needed to support the advantages of eDNA metabarcoding against traditional surveys.

eDNA methods for biodiversity assessment may provide more information to us, as

shown here, but we should also pay attention to the unknown characteristics of eDNA,

such as the origins, degradation, and transport of eDNA in the water, which are still

unknown in various habitats[39,40]. To routinely use eDNA methods to assess biodiversity,

we encourage testing the effect of eDNA degradation and transport on the performance

of eDNA methods for biodiversity evaluation.

**Methods**

**Study sites.** In 2016, we conducted field surveys in five river systems across Japan

(river map in Fig. S1): the Kyuragi River on 10th October, the Koishiwara River on 21st

October, the Yato River on 25th October, the Hazuki River on 2nd November, and the

Oze River on 6th November. The survey sites were set at three sites in each of three

segments (upstream, mid-stream, and downstream, the internal distances ranged from

4.5 to 25.8 km, Fig. S1) for each river. Each site was set so that the length in the up-

down direction is approximately 100 m and with a riffle at the downstream end (e. g.,

Fig. S2).

**Water collection for eDNA survey.** In each site, we collected 1 L of surface water in

bleached bottles at two points, the stream near the downstream end and the shore with

static or semi-static water (Fig. S2). The bottles and filtering equipment removed the

DNA using 10% commercial bleach (approximately 0.6% hypochlorous acid) and

washed it using DNA-free distilled water. One milliliter of benzalkonium chloride

(BAC, 10% w/v) was added per liter of water sample to avoid a decrease in eDNA

concentration in the samples[41]. During transport, samples were stored in a cooler with

ice packs. The 'cooler blank' contained 1 L DNA-free water, which we brought to the

field and was treated identically to the other water samples, except that it was not

opened at the field sites.

**Visual observation and capturing methods.** After water sampling, the fish fauna

survey was conducted by visual observation with snorkel and collection with hand net.

For visual observation, we observed and recorded fish species by snorkeling in 100-m

area (1 person × 1 h, Fig. S2). We observed at various micro habitats including riffle,

pool, and shore bank from downstream end to upstream end. We also conducted the

hand-net capturing survey (1 person × 1 h) using a D-flame net (net opening: 0.16 $m^2$)

in the various habitats in the river including riffle, pool, and shore bank. In this study,

Because of the impossibility of sampling some habitats such as behind stones, we did

not use electrofishing to observe species. Fishes were identified according to Nakabo et

al.[37] (2013) at the survey site. We used the combined taxa list from both surveys to

compare with that from eDNA metabarcoding. In order to prevent contamination of

DNA, the investigator who collected and identified the fish and the investigator who

sampled water were different.

**Water filtering and eDNA extraction from filter samples.** Bottled water samples

were vacuum-filtered in the laboratory into 47 mm GF/F glass filters (pore size: 0.7 μm,

GE Healthcare, Little Chalfont, UK) within 24 h after sampling. After filtration, all

filters were stored at -20 °C before eDNA extraction. The cooler blank was also

processed in the same manner. A liter of Milli-Q water was used as the equipment

control to monitor contamination during filtering in each site and during subsequent

DNA extraction.

To extract the DNA from the filters, we followed the methods described in

Uchii, Doi & Minamoto[42] (2016). We incubated the filter by submerging the mixed

buffer of 400 μL of Buffer AL (Qiagen, Hilden, Germany) and 40 μL of Proteinase K

(Qiagen, Hilden, Germany), using a Salivette tube (Sarstedt, Nümbrecht, Germany) at

56 °C for 30 min. The Salivette tube with filters was centrifuged at 5000 × g for 5 min

and then we added 220 µL of TE buffer (10 mM Tris-HCl and 1 mM EDTA, pH: 8.0)

onto the filter and again centrifuged at 5000 × g for 5 min. The DNA in the eluted

solution was purified using a DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany)

following the manufacturer's protocol. The final volume of the extracted sample was

200 µL with Buffer AE of DNeasy Blood & Tissue Kit. Samples were stored at -20 °C

until qPCR assay.

**Library preparation and MiSeq sequencing.** The procedures used for molecular

methods are described in Fujii et al.[13] (2019). A two-step PCR-procedure was used for

library preparation of Illumina MiSeq sequencing. As the first step, a fragment of the

mitochondrial 12S rRNA gene was amplified using the MiFish-U-F and MiFish-U-R

primers[11] (Miya et al. 2015) which were designed to contain Illumina sequencing primer

regions and 6-mer Ns;

Forward: 5′-*ACACTCTTTCCCTACACGACGCTCTTCCGATCT* NNNNNN

GTCGGTAAAACTCGTGCCAGC-3′,

Reverse: 5′-*GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT* NNNNNN

CATAGTGGGGTATCTAATCCCAGTTTG-3′

The italicized and normal letters represent MiSeq sequencing primers and

MiFish primers, respectively, and the six random bases (N) were used to enhance

cluster separation on the flow cells during initial base call calibrations on MiSeq. We

used a KOD FX Neo polymerase (Toyobo, Osaka, Japan) for the first PCR to facilitate

amplifications of DNA from crude extracts. The first PCR was performed with a 12 µL

reaction volume containing 1× PCR Buffer for KOD FX Neo, 0.4 mM dNTP mix, 0.24

U KOD FX Neo polymerase, 0.3 µM of each primer, and 2 µL template (Sato et al.

2017). The thermal cycles of this step were as follows: initial denaturation at 94 °C for

2 min, followed by 35 cycles of denaturation at 98 °C for 10 s, annealing at 65 °C for 30

s, and elongation at 68 °C for 30 s, followed by final elongation at 68 °C for 5 min. The

first PCRs were performed using eight replicates to mitigate false negatives (PCR

dropouts). Thereafter, individual first PCR replicates were pooled. Each PCR product

(30 µL) was purified using AMPure XP (Beckman Coulter, Brea CA, USA) and eluted

with 30 µL of sterilized water. The purified first PCR products were used as templates

for the second PCR. The Illumina sequencing adaptors and the eight bp identifier

indices (XXXXXXXX) were added to the subsequent PCR process using a forward and

reverse fusion primer:

Forward: 5´-*AATGATACGGCGACCACCGAGATCTACA* XXXXXXXX

ACACTCTTTCCCTACACGACGCTCTTCCGATCT-3´

Reverse: 5´-*CAAGCAGAAGACGGCATACGAGAT* XXXXXXXX

GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT-3´

The italicized and normal letters represent MiSeq P5/P7 adapter and

sequencing primers, respectively. The 8X bases represent dual-index sequences inserted

to identify different samples[43] (Hamady 2008). The second PCR was conducted with 12

cycles of a 12 µL reaction volume containing 1× KAPA HiFi HotStart ReadyMix, 0.3

µM of each primer, and 1.0 µL template from the first PCR production. The thermal

cycle profile after an initial 3 min denaturation at 95 °C was as follows: denaturation at

98 °C for 20 s, annealing, and extension combined at 72 °C (shuttle PCR) for 15 s, with

the final extension at the same temperature for 5 min. The second PCR products were

pooled in equal volumes and purified using AMPure XP as in the first PCR.

The purified PCR products were loaded on a 2% E-Gel SizeSelect (Thermo

Fisher Scientific, Waltham, MA, USA) and the target size of the libraries

(approximately 370 bp) was collected. The library concentration and quality were

estimated by a Qubit dsDNA HS assay kit and a Qubit 2.0 (Thermo Fisher Scientific).

The amplicon libraries were sequenced by $2 \times 250$ bp paired-end sequencing on the

MiSeq platform using the MiSeq v2 Reagent Kit according to the manufacturer's

instructions. Note that the sequencing run contained a total of 339 libraries including 40

of our libraries (30 samples plus ten negative controls) and 299 libraries from other

research projects. The MiSeq sequencing was conducted in the Department of

Environmental Solution Technology, Faculty of Science and Technology, Ryukoku

University. Sequence data were deposited in the Sequence Read Archive of DNA Data

Bank of Japan (DRA, Accession number: DRA008090).

**Bioinformatic analysis for MiSeq sequencing.**  The procedures used for

bioinformatics analysis are described in Fujii et al.[13] (2019). The processing formality

of the MiSeq reads was evaluated by the FASTQC program

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc). After confirming a lack of

technical errors in the MiSeq sequencing, low-quality tails were trimmed from each

read using "DynamicTrim.pl" in the SOLEXAQA software package[44] (Cox et al. 2010)

with a cut-off threshold set at a Phred score of 10. The trimmed paired-end reads (reads

1 and 2) were then merged. The assembled 1,823,446 reads were further filtered by

custom Perl scripts to remove reads with either ambiguous sites (Ns) or those exhibiting

unusual lengths with reference to the expected size of the PCR amplicons ($297 \pm 25$ bp).

The software TagCleaner[45] (Schmieder et al. 2010) was used to remove primer

sequences with a maximum of three-base mismatches and transform the FASTQ format

into FASTA. The pre-processed reads with an identical sequence (i.e., 100% sequence

similarity) were assembled using UCLUST[46] (Edgar 2010) and the number of identical

reads was added to the header line of the FASTA formatted data file. The sequences

represented by 10 or more identical reads were subjected to the downstream analyses.

Processed reads were subjected to local BLASTN searches on the previously

established comprehensive reference database of fish species[11] (Miya et al. 2015).

The top BLAST hit with a sequence identity ≥97% and the E-value threshold of $10^{-5}$

was applied to species detection of each sequence but the species were mostly identified

with ≥99% match. From the BLAST results, we identified the species using methods

previously described[19] (Sato et al. 2017).

**Statistical analyses**. All statistical analyses and graphics were conducted in R ver.

3.4.4[47] (R Core Team, 2018). All statistics were set at the significance level α = 0.05. To

compare between eDNA metabarcoding and visual survey data, the taxonomic levels in

the species list from visual survey were adjusted to the lists from eDNA metabarcoding

(Table S1, S2) in reference to previous studies using MiFish primer[13,19] (Sato et al.

2017, Fujii et al. 2019). Before statistical analysis, we confirmed that the sequencing

depth was sufficient to detect alpha diversity in the samples by "iNEXT" and

"ggiNEXT" functions in "iNEXT" package (Fig. S3). We merged the community data

from two points, the stream near the downstream end and the shore, to compare with

visual/capturing surveys.

We tested the differences in fish richness of sites, segments, and rivers

between both methods by general linear mixed model (GLMM) with "lmer" function in

"lme4" package. In the GLMM models, the method was treated as a fixed effect, and

the rivers and segments were treated as the random effects.

The differences in community compositions were visualized using nonmetric

multidimensional scaling (NMDS) with 500 separate runs of real data. For NMDS, the

community dissimilarity was calculated based on incidence-based Jaccard indices. We

evaluated the differences in community structures between methods and sites using

Permutational multivariate analysis of variance (PERMANOVA). For PERMANOVA,

we used Jaccard similarity matrix and calculated the statistical values with 999

permutations. We used "metaMDS" and "adonis" functions in vegan package for

NMDS ordination and PERMANOVA, respectively.

For the communities evaluated by both methods, indicator taxa analysis[48]

(Cáceres & Legendre, 2009) was performed to determine which taxa had significantly

different frequencies between both methods. The analysis was performed using the

"signassoc" function in the "indicspecies" package on the present/absence data. We

used mode = 1 (group-based) and calculated the p-values with 999 permutations after

Sidak's correction of the multiple testing.

To compare community composition (i.e., beta diversity) focusing on nested

and turnover structures between eDNA metabarcoding and visual/capturing survey, we

calculated the pairwise indices for nestedness and turnover[31,49] (Baselga 2010; Baselga

2012) using the "beta.pair" function in the "betapart" package. Standardized effective

size (SES) of nestedness and turnover indices were calculated to show the degree of

nestedness and turnover structure. The significance was defined by deviation from zero,

the expectation of random assembly (a null model) estimated with 999 random

sampling replicates. The SES was defined as: $(\beta_{obs} - \beta_{null}) / \beta_{sd}$, where $\beta_{obs}$ is the

observed beta diversity, $\beta_{null}$ is the mean of the null distribution of beta diversity, and $\beta_{sd}$

is the standard deviation of the null distribution. SES values greater than zero indicate

statistically stronger nestedness or turnover structure than expected under a random

model of community assembly, while negative values indicate weaker nestedness or

turnover than expected. The randomized community data were generated with

independent swap algorithm[50] (Gotelli 2000) using "randomizeMatrix" function in

"picante" package. Then, to evaluate the fish community longitudinal nestedness and

turnover structure along with river flow (i.e., upstream to downstream) by each method,

a nestedness metric[51] (NODF, Almeida-Neto et al., 2008) and pairwise indices were

calculated. First, to test whether the communities show significant longitudinal

nestedness, SES value of NODF and their SES were calculated with 999 randomizations

using "oecosimu" function in "vegan" package. Then, the SES of pairwise nestedness

and turnover were calculated for each sample pair within the river. Finally, we tested

the differences in SES of pairwise indices between the survey methods (eDNA

metabarcoding vs. visual/capturing survey) by general linear mixed model (GLMM)

with "lmer" function in "lme4" package. In the GLMM models, the SES of pairwise

indices was treated as a fixed effect, and the rivers-pairs and segment-pairs (i.e., three

pairs in each river) were treated as the random effects.

For the map graphics in Supplemental Materials, we used the stream and coast line data

from the National Land Information Division, Ministry of Land, Infrastructure,

Transport and Tourism of Japan (http://nlftp.mlit.go.jp/ksj/).

**Data availability**

All data of the MiSeq sequencing was shared in DRA (Accession number:

DRA008090), and the observed species data was shared in Table S2 in Supplemental

Materials.

**References**

1. Primack, R. B. (1993). Essentials of conservation biology (Vol. 23). Sunderland,

Massachusetts: Sinauer Associates.

2. Margules, C. R. & Pressey, R. L. Systematic conservation planning. *Nature*, **405**, 243

(2000).

3. Pecl, G. T., et al. Biodiversity redistribution under climate change: Impacts on

ecosystems and human well-being. *Science*, **355**, 9214 (2017).

4. Doi, H. & Takahara, T. Global patterns of conservation research importance in

different countries of the world. *PeerJ*, **4**, e2173 (2016).

5. Socolar, J. B., Gilroy, J. J., Kunin, W. E. & Edwards, D. P. How should beta-

diversity inform biodiversity conservation? *Trends Ecol. Evol.*, **31**(1), 67-80 (2016).

6. Takahara, T., Minamoto, T. & Doi, H. Using environmental DNA to estimate the

distribution of an invasive fish species in ponds. *PLoS ONE*, **8**, e56584 (2013).

doi: 10.1371/journal.pone.0056584.

7. Takahara, T., Minamoto, T. & Doi, H. Using environmental DNA to estimate the

distribution of an invasive fish species in ponds. *PLoS ONE,* **8**, e56584 (2013).

8. Goldberg, C. S., Strickler, K. M. & Pilliod, D. S. Moving environmental DNA

methods from concept to practice for monitoring aquatic macroorganisms. *Biol*

*Conserv.*, **183**, 1-3 (2015). doi: 10.1016/j.biocon.2014.11.040.

9. Rees, H. C., Maddison, B. C., Middleditch, D. J., Patmore, J. R. & Gough, K. C.

Review: the detection of aquatic animal species using environmental DNA — a

review of eDNA as a survey tool in ecology. *J. Appl. Ecol.*, **51**, 1450-1459

(2014). doi: 10.1111/1365-2664.12306.

10. Thomsen, P. F., Willerslev, E. Environmental DNA–an emerging tool in

conservation for monitoring past and present biodiversity. *Biol Conserv.*, **183**, 4-

18 (2015). doi: 10.1016/j.biocon.2014.11.019.

11. Miya, M., et al. MiFish, a set of universal PCR primers for metabarcoding

environmental DNA from fishes: detection of more than 230 subtropical marine

species. *R. Soc. Open Sci.*, **2**, 150088 (2015).

12. Doi, H., et al. Environmental DNA analysis for estimating the abundance and

biomass of stream fish. *Freshw. Biol.*, **6**, 30-39 (2017).

13. Fujii, K., Doi, H., Matsuoka, S., Nagano, S., Sato, H. & Yamanaka, H.

Environmental DNA metabarcoding for fish community analysis in backwater

lakes: A comparison of capture methods. *PLoS ONE*, **14**, e0210357 (2019).

14. Doi, H., et al. Evaluation of detection probabilities at the water-filtering and initial

PCR steps in environmental DNA metabarcoding using a multispecies site

occupancy model. *Sci. Rep.*, **9**, 3581(2019).

15. Bista, I., et al. Annual time-series analysis of aqueous eDNA reveals ecologically

relevant dynamics of lake ecosystem biodiversity. *Nat. Commun.*, **8**, 14087

(2017). doi: 10.1038/ncomms14087.

16.Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J. C. & Altermatt, F.

Environmental DNA reveals that rivers are conveyer belts of biodiversity

information. *Nat. Commun.*, **7**, 12544 (2016).

17.Deiner K, et al. Environmental DNA metabarcoding: Transforming how we survey

animal and plant communities. *Mol. Ecol.*, **26**, 5872-5895 (2017). doi:

10.1111/mec.14350.

18.Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C. & Willerslev, E. Towards

next-generation biodiversity assessment using DNA metabarcoding. *Mol. Ecol.*,

**21**(8), 2045-2050 (2012).

19.Sato, H., Sogo, Y., Doi, H. & Yamanaka, H. Usefulness and limitations of sample

pooling for environmental DNA metabarcoding of freshwater fish communities. *Sci.*

*Rep.*, **7**, 14860 (2017). doi: 10.1038/s41598-017-14978-6.

20. Bylemans, J., et al. Monitoring riverine fish communities through eDNA

metabarcoding: determining optimal sampling strategies along an altitudinal and

biodiversity gradient. *Metabarcoding Metagenom.*, **2**, e30457 (2018).

21. Nakagawa, H., et al. Comparing local- and regional-scale estimations of the

diversity of stream fish using eDNA metabarcoding and conventional

observation methods. *Freshw. Biol.*, **63**, 569-580 (2018).

22. Yamamoto, S., et al. Environmental DNA metabarcoding reveals local fish

communities in a species-rich coastal sea. *Sci. Rep.*, **7,** 40368 (2017).

23. Chase, J. M. & Ryberg, W. A. Connectivity, scale-dependence, and the

productivity–diversity relationship. *Ecol. Lett.*, **7**(8), 676-683 (2004).

24. Chase, J.M. Stochastic community assembly causes higher biodiversity in more

productive environments. *Science*, **328**, 1388–1391 (2010).

25. McCune, J. L. & Vellend, M. Gains in native species promote biotic

homogenization over four decades in a human-dominated landscape. *J. Ecol.*,

**101**, 1542-1551 (2013).

26. Olden, J. D. & Poff, N. L. Toward a mechanistic understanding and pre-diction of

biotic homogenization. *Am. Nat.*, **162**, 442–460 (2003).

27. Van Calster, H., et al. Management driven changes (1967–2005) in soil acidity and

the understorey plant community following conversion of a coppic-with-

standards forest. *For. Ecol. Manage.*, **241**, 258–271 (2007).

28. Keith, S. A., Newton, A. C., Morecroft, M. D., Bealey, C. E. & Bullock, J. M.

Taxonomic homogenization of woodland plant communities over70 years. *Proc.*

*R. Soc. B. Biol. Sci.*, **276**, 3539–3544 (2009).

29. Harrison, S. et al. Beta-diversity on geographic gradients in Britain. *J. Anim. Ecol.*,

**61**, 151-158 (1992).

30. Baselga, A., Jimenez-Valverde, A. & Niccolini, G. A multiple-site similarity

measure independent of richness. *Biol. Lett.*, **3**, 642–645 (2007).

31. Baselga, A. Multiplicative partition of true diversity yields independent alpha and

beta components; additive partition does not. *Ecology*, **91**(7), 1974-1981 (2010).

32. Wright, D. H. & Reeves, J. H. On the meaning and measurement of nestedness of

species assemblages. *Oecologia*, **92**(3), 416-428 (1992).

33. Ulrich, W. & Gotelli, N. J. Null model analysis of species nestedness patterns.

*Ecology*, **88**(7), 1824-1831 (2007).

34. Gaston, K. J. & T. M. Blackburn. Pattern and process in macroecology. Blackwell

Scientific, Oxford (2000).

35. Baselga, A., Gómez-Rodríguez, C. & Lobo, J. M. Historical legacies in world

amphibian diversity revealed by the turnover and nestedness components of beta

diversity. *PloS ONE*, **7**(2), e32341 (2012).

36. Kawanabe, H., Mizuno, N. & K. Hosoya (eds.) Freshwater Fishes of Japan Yama-

Kei, Tokyo (2001) (in Japanese).

37. Nakabo, T. Fishes of Japan with pictorial keys to the species, 3rd ed. Tokai

University Press (2013) (in Japanese).

38. Itakura H, et al. Environmental DNA analysis reveals the spatial distribution,

abundance, and biomass of Japanese eels at the river-basin scale. *Aquat.*

*Conserv.*, 1–13 (2019).

39. Barnes, M. A. & Turner, C. R. The ecology of environmental DNA and implications

for conservation genetics. *Conserv. Genet.*, **17**(1), 1–17 (2016).

40. Seymour, M., et al. Acidity promotes degradation of multi-species environmental

DNA in lotic mesocosms. *Commun. Biol.*, **1**(1), 4 (2018).

41. Yamanaka, H., et al. On-site filtration of water samples for environmental DNA

analysis to avoid DNA degradation during transportation. *Ecol. Res.,* **31**(6), 963-

967 (2016).

42. Uchii, K., Doi, H. & Minamoto, T. A novel environmental DNA approach to

quantify the cryptic invasion of non-native genotypes. *Mol. Ecol. Res*, **16**, 415-422

(2016). doi: 10.1111/1755-0998.12460.

43. Hamady, M., Walker, J. J., Harris, J. K., Gold, N. J. & Knight, R. Error-correcting

barcoded primers for pyrosequencing hundreds of samples in multiplex. *Nat.*

*Methods*, **5**, 235-237 (2008).

44. Cox, M. P., Peterson, D. A., Biggs, P. J. SolexaQA: at-a-glance quality assessment

of Illumina second-generation sequencing data. BMC *Bioinform.*, **11**, 485 (2010).

doi: 10.1186/1471-2105-11-485.

45. Schmieder, R., Lim, Y. W., Rohwer, F. & Edwards, R. TagCleaner: Identification

and removal of tag sequences from genomic and metagenomic datasets. *BMC*

*Bioinform.*, **11**, 341 (2010). doi: 10.1186/1471-2105-11-341.

46. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST.

*Bioinformatics*, **26**, 2460-2461 (2010). doi: 10.1093/bioinformatics/btq461.

47. R Development Core Team. R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria (2018).

https://www.R-project.org/.

48. Cáceres, M. D. & Legendre, P. Associations between species and groups of sites:

indices and statistical inference. *Ecology*, **90**(12), 3566-3574 (2009).

49. Baselga, A. The relationship between species replacement, dissimilarity derived

from nestedness, and nestedness. *Glob. Ecol. Biogeogr.*, **21**, 1223-1232 (2012).

50. Gotelli, N. J. Null model analysis of species co-occurrence patterns. *Ecology*, **81**,

2606-2621 (2000).

51. Almeida-Neto, M., Guimaraes, P., Guimaraes, P. R., Loyola, R. D. & Ulrich, W. A

consistent metric for nestedness analysis in ecological systems: reconciling concept

and measurement. *Oikos*, **117**, 1227-1239 (2008).

**Acknowledgement**

MiSeq sequencing was conducted in the Department of Environmental Solution

Technology, Faculty of Science and Technology, Ryukoku University, and we thank to

H. Yamanaka and H. Sato for supporting the experiments involving MiSeq sequencing.

This study was supported by the Environment Research and Technology Development

Fund (4-1602) of Environmental Restoration and Conservation Agency, Japan and JST-

CREST (JPMJCR13A2).

**Author contributions**. HD, RI, and YA designed the study, RI, MG, TK, and YA

contributed to field survey and sampling. SM, HD, RI, and MG contributed to

molecular experiments. SM and HD analyzed the data and interpreted the results. HD,

SM, and RI wrote the initial draft of the manuscript. All other authors critically

reviewed the manuscript.

**Competing financial interests.** The authors declare no competing financial or non-

financial interests.

**Materials & Correspondence.** H. D. (hideyuki.doi@icloud.com) and R. I.

(inuiryutei@gmail.com)

**Figure legends**

Figure 1. Venn diagrams for the number of detected taxa. eDNA metabarcoding

(eDNA, blue) and visual/capturing methods (V/C, red) in each site of the five study

river systems. The numbers mean the number of taxa.

Figure 2. Nonmetric multidimensional scaling (NMDS) ordination. Fish communities

evaluated by eDNA metabarcoding (blue) and visual/capturing methods (red) in each

site of the river. MDS stress was 0.158.

Figure 3. Pairwise effect size (SES, standard effect size). Comparison between

community nestedness and spatial turnover between the communities detected by

eDNA metabarcoding and visual/capturing methods. Error bars indicate 95%

confidence interval. The horizontal dotted line represents SES = 0, indicating non-

significant effect.

Figure 4. Effect size (SES, standard effect size). a) community nestedness and b) spatial

turnover SES among the segment communities evaluated by eDNA metabarcoding and

visual/capturing methods. The error bars indicate 95% confidence interval. The

horizontal dotted line represents SES = 0, indicating non-significant effect.

611    Figure 1

614     Figure 2

617 Figure 3

621    Figure 4