Estimation of biodiversity metrics by environmental DNA metabarcoding

compared with visual and capture surveys of river fish communities

Hideyuki Doi[1,*,†], Ryutei Inui[2,3*,†], Shunsuke Matsuoka[1, †], Yoshihisa Akamatsu[2],

Masuji Goto[2,4], and Takanori Kono[2,5]

[1] Graduate School of Simulation Studies, University of Hyogo, 7-1-28 Minatojima

Minami-machi, Chuo-ku, Kobe, 650-0047, Japan

[2] Graduate School of Science and Engineering, Yamaguchi University, 2-16-1

Tokiwadai, Ube, Yamaguchi, 755-8611, Japan

[3] Faculty of Socio-Environmental Studies, Fukuoka Institute of Technology, 3-30-1

Wajiro-higashi, Higashi-ku, Fukuoka, 811-0295, Japan

[4] Nippon Koei Research and Development Center, 2304 Inarihara, Tsukuba City, Ibaraki

300-1259, JAPAN

[5] Aqua Restoration Research Center, Public Works Research Institute National Research

and Development Agency, Kawashima, Kasada-mathi, Kakamigahara-City, Gifu, 501-

6021, Japan

[†] These authors equally contributed to this study.

[*] Corresponding authors:

Hideyuki Doi (hideyuki.doi@icloud.com)

Ryutei Inui (inuiryutei@gmail.com)

Running head: eDNA metabarcoding for fish diversity

Abstract

1. Information on alpha (local), beta (between habitats), and gamma (regional) diversity

is fundamental to understanding biodiversity as well as the function and stability of

community dynamics. Methods like environmental DNA (eDNA) metabarcoding are

currently considered useful to investigate biodiversity.

2. We compared the performance of eDNA metabarcoding with visual and capture

surveys for estimating alpha and gamma diversity of river fish communities, and

nestedness and turnover in particular.

3. In five rivers across west Japan, by comparison to visual/capture surveys, eDNA

metabarcoding detected more species in the study sites (i.e., alpha diversity).

Consequently the overall number of species in the region (i.e., gamma diversity) was

higher. In particular, the species found by visual/capture surveys were encompassed by

those detected by eDNA metabarcoding.

4. Estimates of community diversity within rivers differed between survey methods.

Although we found that the methods show similar levels of community nestedness and

turnover within the rivers, visual/capture surveys showed more distinct community

differences from upstream to downstream. Our results suggest that eDNA

metabarcoding may be a suitable method for community assemblage analysis,

especially for understanding regional community patterns, for fish monitoring in rivers.

Key words: eDNA, community, alpha and gamma diversity, nestedness, turnover

Introduction

The maintenance of biodiversity underpins the stability of ecosystem processes in

constantly changing environments (Primack, 1993; Margules & Pressey, 2000; Pecl et

al., 2017). Moreover, biodiversity loss affects ecosystem functions and services and,

consequently, human society (Primack 1993; Margules & Pressey, 2000, Pecl et al.

2017). Ecologists have made efforts to conserve biodiversity based on essential

biodiversity survey methods, especially for species richness and distribution (Primack,

1993; Margules & Pressey, 2000, Doi & Takahara, 2016, Pecl et al., 2017). Biodiversity

can be evaluated in different levels: e.g., by estimating alpha (local), beta (between

habitats), and gamma (regional) diversity and the variation of the community

assemblages. To conserve local communities, ecologists incorporate these diversity

metrics into management decisions (Primack 1993; Margules & Pressey, 2000, Socolar

et al., 2016). For example, measuring variation among community assemblages can

identify biodiversity loss and inform the placement of protected areas and the

management of biological invasions and landscapes (Socolar et al., 2016). Thus, robust

methods for monitoring biodiversity are fundamental for biodiversity and environmental

management.

70   Environmental DNA (eDNA) analysis is considered a useful tool to

investigate the distribution and richness of aquatic and terrestrial organisms (Takahara

et al., 2012, 2013; Rees et al., 2014; Goldberg et al., 2015; Miya et al., 2015; Thomsen

& Willerslev, 2015; Doi et al., 2017; Doi et al., 2019; Fujii et al., 2019). High-

throughput sequencing derived from eDNA, called "eDNA metabarcoding", is an

exceptionally useful and powerful tool for community biodiversity surveys (Taberlet et

al., 2012; Deiner et al., 2016, 2017; Sato et al., 2017; Bylemans et al., 2018; Fujii et al.,

2019). eDNA metabarcoding has recently been applied in fish community surveys, e.g.

Miya et al. (2015) designed and applied universal PCR primers (the MiFish primers) to

survey marine fish communities. To confirm the usefulness of eDNA metabarcoding for

community assessment, many studies have compared eDNA metabarcoding to a species

list generated by traditional surveys including visual and capture methods (Deiner et al.,

2016; Bylemans et al., 2018; Nakagawa et al., 2018; Yamamoto et al., 2018; Fujii et al.,

2019). However, comparisons of eDNA and traditional survey methods for estimating

regional (gamma) diversity or differences in community assemblage composition

among locations are more limited (but see Drummond et al., 2015; Deiner et al., 2016;

Staehr et al. 2016, Maechler et al. 2019).

Variation in community assemblage composition reflects two components,

nestedness and species turnover for considering regional structure of community

assemblage (Harrison et al., 1992; Baselga et al., 2007; Baselga, 2010). Nestedness

occurs when the community at the sites with fewer species are subsets of the

community at the sites with higher species richness (Wright & Reeves, 1992; Ulrich &

Gotelli, 2007) and generally reflects a non-random process of species loss (Gaston &

Blackburn, 2000). Contrastingly, species turnover implies the replacement of some

species by others because of environmental sorting or spatial/historical constraints

(Baselga, 2010). Statistical separation methods for nestedness and species turnover were

applied for evaluating variation of the community assemblages in various systems

(Baselga, 2010; Baselga et al., 2012). Moreover, Baselga's (2010) framework can be

applied to compare the performance among methods when evaluating beta diversity via

nestedness and species turnover. Nestedness and species turnover can also occur due to

differences in sampling methods, even when the surveys are conducted on the same

community. For example, differing biases in species detection between two methods

can result in apparent species turnover. Evaluating nestedness and species turnover

would allow us to reveal the potential bias of community analysis of eDNA

metabarcoding compared to traditional surveys. Therefore, we can quantitatively

compare the performance of eDNA metabarcoding and traditional surveys for

alpha/gamma diversity evaluation of biological communities and the variation of the

community assemblages among the study sites.

108   Using statistical methods, we tested the performance of eDNA metabarcoding

in five river systems in different regions with various fish species. We conducted eDNA

metabarcoding using universal MiFish primers that target fish and identified the fish by

visual snorkeling and hand-net capture surveys. We evaluated the performance of

eDNA metabarcoding by comparing the obtained fish community structure to that

evaluated by visual/capture survey with special regard to nestedness and species

turnover.

Methods

Site description

In 2016, we conducted field surveys in five river systems across Japan (river map in

Fig. S1): the Kyuragi River on October 10, the Koishiwara River on October 21, the

Yato River on October 25, the Hazuki River on November 2, and the Oze River on

November 6. The survey sites were set at a site at each of three river segments

(upstream, mid-stream, and downstream, the internal distances ranged from 4.5 to 25.8

km, Fig. S1) for each river. Each site was set so that the length in the up-down direction

was approximately 100 m with a riffle at the downstream end (e.g., Fig. S2).

Water collection for eDNA survey

In each site, we collected 1 L of surface water in bleached bottles at two points, in

flowing water near the downstream end of the site, and in stagnant or nearly stagnant

water near the shore (Fig. S2) immediately before visual and capture surveys. Prior to

sample collection, eDNA was removed from the bottles and filtering equipment using

10% commercial bleach (ca. 0.6% hypochlorus acid) and washing with DNA-free

distilled water. One milliliter of benzalkonium chloride (BAC, 10% w/v) was added per

liter of water sample to avoid a decrease in eDNA concentration in the samples

(Yamanaka et al., 2016). During transport, samples were stored in a cooler with ice

packs. The 'cooler blank' contained 1 L DNA-free water, which we brought to the field

and treated identically to the other water samples, except that it was not opened at the

field sites.

Visual observation and capture methods

After water sampling, the fish fauna survey was conducted by visual observation with

snorkel and collection with hand net. For visual observation, we observed and recorded

fish species by snorkeling in a 100-m transect (snorkeling by 1 person for 1 h, Fig. S2).

We observed at various micro habitats, including the riffle, pool, and shore bank from

the downstream end to upstream end. We also conducted a hand-net capture survey (1

person × 1 h) using a D-frame net (2 mm mesh, net opening: 0.16 m^2) in the various

habitats in the river, including the riffle, pool, and shore bank. Fishes were identified

according to Nakabo et al. (2013) at the survey site. We used the combined taxa list

from both traditional surveys to compare to that of eDNA metabarcoding. In order to

prevent contamination of eDNA samples, the investigator who collected and identified

the fish (i.e., handled the fish) and the investigator who sampled the water were

different.

eDNA collection, extraction and measurements

Collected water samples were vacuum-filtered into GF/F glass filters (47 mm diameter,

pore size: 0.7 μm, GE Healthcare, Little Chalfont, UK) in the laboratory within 24 h of

sampling. After filtration, all filters were stored at -20 °C before eDNA extraction. The

cooler blank was also processed in the same manner. A liter of Milli-Q water was used

as the filtering blank to monitor contamination during filtering in each site and during

subsequent DNA extraction.

To extract the DNA from the filters, we followed the methods described in

Uchii, Doi, & Minamoto (2016). We incubated the filter by submerging the mixed

buffer of 400 μL of Buffer AL in DNeasy Blood & Tissue Kit (Qiagen, Hilden,

Germany) and 40 μL of Proteinase K (Qiagen, Hilden, Germany), using a Salivette tube

(Sarstedt, Nümbrecht, Germany) at 56 °C for 30 min. The Salivette tube with filters was

centrifuged at $5000 \times g$ for 5 min. Then, we added 220 μL of TE buffer (pH: 8.0) onto

the filter and again centrifuged at $5000 \times g$ for 5 min. The DNA was purified using a

DNeasy Blood & Tissue Kit with extracted the DNA in 200 μL in Buffer AE. Samples

were stored at -20 °C until the 1st-PCR assay.

Library preparation and MiSeq sequencing

The detailed molecular methods are described in Fujii et al. (2019) with a two-step

PCR-procedure for Illumina MiSeq sequencing. Briefly, we performed 1st-PCR

withMiFish-U-F and MiFish-U-R primers (Miya et al., 2015), which were designed to

contain Illumina sequencing primer regions and 6-mer Ns;

Forward: 5'-*ACACTCTTTCCCTACACGACGCTCTTCCGATCT* NNNNNN

GTCGGTAAAACTCGTGCCAGC-3′ ,

Reverse: 5′ -*GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT* NNNNNN

CATAGTGGGGTATCTAATCCCAGTTTG-3′

The italicized and non-italicized letters represent MiSeq sequencing primers

and MiFish primers, respectively, and the six random bases (N) were used to enhance

cluster separation on the flow cells during initial base call calibrations on the MiSeq

(Miya et al. 2015, Doi et al. 2019).

We performed the 1st-PCR with a 12 μL reaction volume containing 1× PCR

Buffer for KOD FX Neo (Toyobo, Osaka, Japan), 0.4 mM dNTP mix, 0.24 U KOD FX

Neo polymerase, 0.3 μM of each primer, and 2 μL template. The thermocycling

conditions for this step were as follows: initial denaturation at 94 °C for 2 min, followed

by 35 cycles of denaturation at 98 °C for 10 s, annealing at 65 °C for 30 s, and

elongation at 68 °C for 30 s, followed by final elongation at 68 °C for 5 min. The first

PCRs were performed using eight replicates (Doi et al. 2019) and individual first PCR

replicates were pooled and purified using AMPure XP (Beckman Coulter, Brea CA,

USA) as templates for the 2nd-PCR. The Illumina sequencing adaptors and the eight bp

identifier indices (XXXXXXXX) were added to the subsequent PCR process using a

forward and reverse fusion primer:

Forward: 5′ -*AATGATACGGCGACCACCGAGATCTACA* XXXXXXXX

ACACTCTTTCCCTACACGACGCTCTTCCGATCT-3′

Reverse: 5′ -*CAAGCAGAAGACGGCATACGAGAT* XXXXXXXX

GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT-3′

The italicized and non-italicized letters represent MiSeq P5/P7 adapter and

sequencing primers, respectively. The 8X bases represent dual-index sequences inserted

to identify different samples (Hamady et al. 2008). We performed the 2nd-PCR with 12

cycles of a 12 µL reaction volume containing 1× KAPA HiFi HotStart ReadyMix, 0.3

µM of each primer, and 1.0 µL of the first PCR production. The thermocycling

conditions profile after an initial 3 min denaturation at 95 °C was as follows:

denaturation at 98 °C for 20 s, annealing, and extension combined at 72 °C (shuttle

PCR) for 15 s, with the final extension at the same temperature for 5 min. We

confirmed the positive bands of the targeted 1st-PCR amplicons by electrophoresis. The

2nd-PCR products were pooled in equal volumes and purified using AMPure XP.

The purified PCR products were loaded on a 2% E-Gel SizeSelect (Thermo

Fisher Scientific, Waltham, MA, USA) and the target size of the amplified DNA

(approximately 370 bp) was collected. The samples concentration and quality were

estimated by a Qubit dsDNA HS assay kit and a Qubit 2.0 (Thermo Fisher Scientific).

The amplicons were sequenced by 2 × 250 bp paired-end sequencing on the MiSeq

platform using the MiSeq v2 Reagent Kit. Note that the sequencing run contained a total

of 339 samples including 40 of our samples (30 samples plus five cooler and five filter

negative controls) and 299 samples from other research projects. The MiSeq sequencing

was conducted in the Department of Environmental Solution Technology, Faculty of

Science and Technology, Ryukoku University. All sequence data were deposited in

DNA Data Bank of Japan (DRA, Accession number: DRA008090).

Bioinformatic analysis for MiSeq sequencing

The detailed procedures used for bioinformatics analysis are described in Fujii et al.

(2019). First, low-quality tails were trimmed from each read and paired-end reads were

then merged. For the obtained 1,823,446 reads, primer sequences were removed and

identical sequences (i.e., 100% sequence similarity) were merged using UCLUST

(usearch 7.0.1001, Edgar, 2010). The sequences with 10 or more identical reads were

subjected to the downstream processes. To annotate the taxonomy, local BLASTN

search using BLAST 2.2.29 was conducted with the reference database of fish species

for processed reads (Miya et al., 2015). The top BLAST hit with a sequence identity ≥

97% was applied to species detection of each sequence. Note that the species were

mostly identified with a ≥ 99% match. From the BLAST results, we identified the

species using methods previously described (Sato et al., 2017). Also, we detect most of

the fish species known to occur in the rivers of the region, according to Kawanabe et al.

(2001).

Statistical analyses

All statistical analyses and graphics were conducted in R ver. 3.4.4 (R Core Team,

2018). All statistics were set at the significance level $\alpha = 0.05$. To compare between

eDNA metabarcoding and visual survey data, the taxonomic levels in the species list

from visual survey were compared to the lists from eDNA metabarcoding (Table S1,

S2) in reference to previous studies using the MiFish primer (Sato et al., 2017; Fujii et

al., 2019). Before statistical analysis, we confirmed that the sequencing depth was

sufficient to detect alpha diversity in the samples by "iNEXT" and "ggiNEXT"

functions in the "iNEXT" ver. 2.0.19 package (Chao et al. 2014, Fig. S3). We merged

the community data from two points, the stream near the downstream end and the shore,

to compare with visual/capture surveys.

We tested the differences in fish richness of sites, segments, and rivers

between both methods using generalized linear mixed models (GLMMs) with the

"lmer" function in the "lme4" ver. 1.1-21 package (Bates et al. 2015). In the GLMM

models, the method was treated as a fixed effect with Poisson distribution, and the

rivers and segments were treated as random effects.

The differences in community compositions were visualized using nonmetric

multidimensional scaling (NMDS) with 500 separate runs of real data. For NMDS, the

two data sets (eDNA metabarcoding and visual/capture survey) were combined into one

ordination (but showed divided into two panels by "facet_grid" function in ggplot2).

The community dissimilarity was calculated based on incidence-based Jaccard indices.

We evaluated the differences in community structures between methods and sites using

permutational multivariate analysis of variance (PERMANOVA). For PERMANOVA,

we used Jaccard and Raup-Crick dissimilarity indices and calculated the statistical

values with 999 permutations. The Raup-Crick index estimates the probability that two

samples have greater differences in species composition than would be expected by

chance draws from the regional species pool, and accounts for differences in alpha

diversity between samples. We used "metaMDS" and "adonis" functions in the "vegan"

ver. 2.5-6 package (https://github.com/vegandevs/vegan) for NMDS ordination and

PERMANOVA, respectively.

Indicator taxa analysis (Cáceres & Legendre, 2009) was performed to

determine which taxa had significantly different frequencies between eDNA and

visual/capture methods. The analysis was performed using the "signassoc" function in

the "indicspecies" ver. 1.7.8 package on the presence/absence data for the testing with

regarding the package description and Cáceres & Legendre (2009). The "signassoc"

function can calculate the index with both presence/absence and abundance data. We

used the presence/absence data with mode = 1 (group-based) and calculated the P-

values with 999 permutations after Sidak's correction of the multiple testing.

To compare community nestedness and turnover (Baselga, 2010; Baselga,

2012) as estimated by eDNA and visual/capture surveys, we used the "beta.pair"

function in the "betapart" ver. 1.5.1 package (Baselga and Orme 2012).   Standardized

effect sizes (SESs) were calculated to show the degree of nestedness and turnover

structure. The significance was defined by deviation from zero, and the expectation of

random assembly (a null model) was estimated with 999 random sampling replicates.

The SES was defined as follows: $(\beta_{obs} - \beta_{null})/\beta_{sd}$, where β_{obs} is the observed beta

diversity (here, the variation of the community assemblages among the sites), β_{null} is the

mean of the null distribution of beta diversity, and β_{sd} is the standard deviation of the

null distribution. SES values greater than zero indicate statistically stronger nestedness

or turnover structure than expected under a random model of community assembly,

while negative values indicate weaker nestedness or turnover than expected. The

randomized community data were generated with independent swap algorithm (Gotelli,

2000) using "randomizeMatrix" function in the "picante" ver. 1.8 package (Kembel et

al. 2010). First, to evaluate the differences in nestedness and species turnover between

the survey methods (eDNA metabarcoding vs. visual/capture survey) at the same

segments, the SES of pairwise nestedness and turnover were calculated for each sample

pair within the river. Then, the fish community longitudinal nestedness and turnover

structure along with river flow (i.e., upstream to downstream) by each method was

evaluated with NODF and pairwise indices of nestedness. First, a nestedness metric

(NODF, Almeida-Neto et al., 2008) and their SES value were calculated with 999

randomizations using the "nestednodf" and the "oecosimu" function in the "vegan"

package. Then, we tested the differences in SES of pairwise indices between the survey

methods (eDNA metabarcoding vs. visual/capture survey) by generalized linear mixed

model (GLMM, with Gaussian distribution) with the "lmer" function in the "lme4"

package. In the GLMM models for the SES of pairwise indices, the survey method was

treated as a fixed effect, and the rivers-pairs and segment-pairs (i.e., three pairs in each

river) were treated as random effects.

Results

Overview

We detected 53 fish taxa, almost all identified to the species or genus level, by eDNA

metabarcoding in five rivers (Table S1, and S2) and visually observed 38 fish taxa in

total. MiSeq paired-end sequencing for the library (30 samples plus five cooler and five

filter negative controls) yielded a total of 1,601,816 reads (53,351 ± 17,639; mean ± S.

D. for each sample, Table S2). We confirmed very low reads from negative controls

(Table S2) with only one detection of a fish species, *Tridentiger* sp. in the blank of the

Yato River, probably because of the cross-contamination among the samples. The reads

of *Tridentiger* sp. in water samples (4783-9818 reads) were higher than in the blank

sample (1286 reads), and so we retained this species in the analysis.

Diversity indices between methods

We consistently detected more taxa at a site (i.e., higher alpha diversity) with eDNA

metabarcoding than with visual/capture surveys (Fig. 1; GLMM, $t = -5.45$, $P =$

0.000018). Estimated taxa richness generally increased from upstream to downstream

(Fig. 1, $t = -5.85$, $P = 0.000004$, random effect for variation among segments: $26.879 \pm$

5.185, SD). The random effect for "river" was not significantly different from zero ($t =$

1.737, $P = 0.0942$, random effect for variation among river: 3.896 ± 1.974).

We ordinated the differences in community structure between the two

methods by NMDS ordination by Jaccard index (Fig. 2) as well as Raup-Crick (Fig.

S4). The patterns of differences in the both Jaccard and Raup-Crick indices were

similar. The ordination showed the differences in community structure between the two

methods, probably due to the differences in species detection in the communities. The

PERMANOVA results with Jaccard and Raup-Crick indices for the ordination

suggested there were differences in community composition evaluated by each method,

eDNA metabarcoding and visual/capture survey ($P < 0.012$, Table S3). Moreover,

communities from the combined results of eDNA metabarcoding and visual/capture

survey were significantly different among rivers and segments ($P < 0.029$, Table S3).

We found different patterns in ordinated river sites for each method (Fig. 2 for Jaccard

index, Fig. S4 for Raup-Crick index).

The PERMANOVA results with Jaccard index determined that communities

were significantly different among the five rivers by eDNA metabarcoding ($P = 0.001$)

but not by visual/capture survey ($P = 0.12$, Table S4). Conversely, the communities

were significantly different among river segments (from downstream to upstream) by

visual/capture survey ($P = 0.011$) but not significantly different (albeit marginally) by

eDNA metabarcoding ($P = 0.061$, Table S4). The differences in PERMANOVA results

with Jaccard index suggested that differences in the variation of the community

assemblages among rivers across regions can be detected by eDNA metabarcoding but

not by visual/capture survey. While, the PERMANOVA results with Raup-Crick index

determined that communities were significantly different among the rivers by both

eDNA metabarcoding ($P = 0.001$) and visual/capture survey ($P = 0.018$, Table S4).

Indicator taxa analysis comparing the communities estimated by both methods

showed that eDNA metabarcoding more frequently detected several species than

visual/capture survey, including Japanese eel (*Anguilla japonica*), salmon (e.g.,

Oncorhynchus masou), and Amur catfish (*Silurus asotus*) ($P < 0.05$, Table 1 for

statistically significant taxa, Table S5 for all taxa).

Nestedness and species turnover

We evaluated nestedness and turnover between communities as assessed using eDNA

compared with visual/capture surveys using pairwise standardized effect size (pSES)

calculated for all sites. Nestedness pSES was significantly positive without overlapping

the zero-pSES 95% confidence interval (Fig. 3). The significantly negative pSES in

species turnover indicated that taxa were not replaced in communities as assessed by

eDNA compared to visual/capture surveys any more than would be expected to occur

with random sampling.

We compared the longitudinal nested structure and turnover strength

(upstream to downstream) of the fish communities as assessed by eDNA metabarcoding

and visual/capture surveys (Fig. 4). The longitudinal nested structure was not

significantly different between both methods. The pSES values were significantly

positive for both eDNA metabarcoding and visual/capture survey, indicating significant

nested structures between segments, regardless of sampling method. Moreover, the

pSES values were not significantly different between the eDNA metabarcoding and

visual/capture survey, so the strength of the nested structures did not differ between the

two methods (GLMM, $P = 0.302$). The species per segment were nested as downstream

> mid-stream > upstream according to both methods (Fig. S5 for all species and S6 for

each river, NOFD, $P < 0.001$). Longitudinal species turnover pSES values were not

different between sampling methods (GLMM, p=0.28) and were significantly negative

(Fig. 4b) indicating a lack of turnover whether communities were sampled by eDNA or

visual/capture survey.

Discussion

We here examined the nestedness and species turnover patterns of fish communities

evaluated by eDNA metabarcoding in five rivers to evaluate the performance of eDNA

metabarcoding. Our results suggested that the nestedness and species turnover from

upstream to downstream did not significantly differ within a river. Therefore, we found

that the both methods show similar levels of community nestedness and turnover within

the rivers. In general, nestedness of species assemblages occurs when the communities

obtained by the method estimating lower number of species are subsets of the

communities estimated by other methods with higher species richness (Baselga 2010).

Furthermore, the species turnover in the communities was very weak between the both

methods.

eDNA metabarcoding has been reported to perform better than traditional

methods in evaluating species richness (Deiner et al., 2016; Sato et al., 2017; Bylemans

et al., 2018; Nakagawa et al., 2018; Yamamoto et al., 2018; Fujii et al., 2019).

Nakagawa et al. (2018) investigated freshwater fish communities in 100 rivers and

confirmed that the communities detected by eDNA metabarcoding were similar to the

species lists observed in government-authorized monitoring. Furthermore, several

eDNA metabarcoding studies on fish communities have been performed in other river

systems (Bylemans et al., 2018), marine habitats (Yamamoto et al., 2018), and

freshwater lakes (Sato et al., 2017; Fujii et al., 2019). Deiner et al. (2016) showed that

river eDNA metabarcoding can reflect the regional community in a watershed,

indicating that eDNA metabarcoding has high performance for gamma diversity

evaluation. These studies indicated the great potential of eDNA metabarcoding as a

useful tool for alpha and gamma diversity assessment by simply comparing the

communities detected by eDNA metabarcoding and traditional surveys. Here, we

support the previous literature (e.g., Nakagawa et al. 2018, Maechler et al. 2019) by

showing patterns similar to those observed in alpha and gamma diversity in river

macroinvertebrates. However, previous studies, excluding Maechler et al. (2019), did

not evaluate performance in terms of nestedness and species turnover between eDNA

metabarcoding and other community data. We also estimated nestedness, species

turnover, and the capture bias of communities detected by eDNA metabarcoding and

traditional methods in segment scale. We found that communities detected by

traditional methods were nested within those detected by eDNA metabarcoding in a

given river locale, with scarce species turnover between methods.

We especially focused on the variation of the community assemblages

evaluated by eDNA metabarcoding compared to visual/capture survey. The two survey

methods showed different sources of community variation. That is, eDNA

metabarcoding detected significant differences between the rivers, while visual/capture

survey detected significant differences between the segments rather than between the

rivers. These differences may lead us to interpret the variation in community

assemblages using the results from both survey methods. We recommend using eDNA

for screening surveys because the method performs similarly to visual/capture surveys

for estimating alpha and gamma diversity, although eDNA may fail to detect some

species that can be captured or observed using traditional methods.

Using Jaccard index, a higher variation of the river fish communities was

statistically detected by eDNA metabarcoding than by visual/capture survey and the

variation of the community assemblages between segments could be significantly

detected by visual/capture surveys but not by eDNA metabarcoding. While the results

of Raup-Crick index, considering the alpha diversity of sites, showed the same results of

the both methods. We further compared the indicator taxa for the communities obtained

from both eDNA metabarcoding and visual/capture survey and concluded that several

taxa, including eel, salmon, and catfish, were significantly better detected by eDNA

metabarcoding, whereas non-indicator taxa were detected by visual/capture surveys.

These results indicated that eDNA metabarcoding had higher detection frequency than

visual/capture surveys in fish taxa detection. The community structures estimated by

eDNA metabarcoding and visual/capture survey were slightly different, as reported in

previous studies (e.g., Sato et al., 2017; Fujii et al., 2019), probably because of the

differences in taxa-detection performances.

Discriminated taxa in this analysis included eel, salmon, and catfish, which

mostly had larger body size and lower abundances in these rivers (Kawanabe, 2001;

Nakabo, 2013), although we here could not separately examine the detected capture bias

and actual pattern differences in species detection. In fact, the Japanese eel *Anguilla*

japonica, was difficult to find by visual observation, probably due to its hiding behavior

(Itakura et al., 2019). Such endangered species would be important as top predators

(Nakabo, 2013). eDNA metabarcoding can evaluate the distribution of such rare and

important taxa in fish communities better than traditional surveys. While we did not

detect any indicator taxa by visual/capture surveys, two species *Lepomis macrochirus*

and *Biwia zezera* were only detected by this method. *Lepomis macrochirus* often inhabit

lentic systems and are rarely observed in rivers (Nakabo, 2013) and *Biwia zezera* are not

widely distributed in this region (Hosokawa et al., 2007). In fact, we only observed *L.*

macrochirus and *B. zezera* in downstream segments of the Kyuragi and Oze river,

respectively.

We sampled eDNA from only two habitats at each study locale and examined

the performances of eDNA metabarcoding. Thus, our understanding of some aspects of

the fish community spatial structure in the rivers and the performance of community

evaluation in local habitats, such as backwater, was still limited. In fact, Bylemans et al.

(2018) found that river morphology in these habitats influenced the optimal sampling

strategy for eDNA metabarcoding. Moreover, in backwater lakes, the performance of

eDNA metabarcoding varied with different lake morphologies (Fujii et al. 2019).

However, testing the usefulness of the eDNA metabarcoding for assessing river fish

community biodiversity has been limited. Further research is needed to evaluate fish

community spatial structure in rivers. In addition, we should consider whether eDNA

recovered from a water sample came from an individual in the survey area. Previous

studies have suggested the eDNA came from upstream (Deiner, K., & Altermatt 2014;

Deiner et al. 2016). Therefore, the comparisons between eDNA metabarcoding and

visual/capture methods could be using community data with different spatial scales. For

example, visual/capture methods obtained community data from a 100-m reach but

eDNA potentially detected the community in larger area than surveyed. However, the

visual/capture survey also does not cover all species due to limited sampling time and

efforts.

Community assessment with eDNA metabarcoding required much less effort

in the field and detected the community in broader area than visual/capture survey.

Therefore, eDNA metabarcoding may be a suitable method, especially for regional

community patterns. Biodiversity testing using statistical frameworks, especially

community nestedness and turnover, provided the quantitative evidence to compare the

performance of eDNA metabarcoding and traditional surveys. eDNA methods for

biodiversity assessment may provide more information to us, as shown here, but we

should also pay attention to the unknown characteristics of eDNA, such as the origins,

degradation, and transport of eDNA in water which are still unknown in various

habitats (Barnes & Turner, 2016; Seymour et al., 2018).

Acknowledgments

MiSeq sequencing was conducted in the Department of Environmental Solution

Technology, Faculty of Science and Technology, Ryukoku University, and we thank to

H. Yamanaka and H. Sato for supporting the experiments involving MiSeq sequencing.

Fish visual/capture surveys were permitted by the prefectures (No. 38 for Fukuoka, No.

3024 for Yamaguchi prefecture) for the survey period; 23 Sep. to 30 Nov. 2016. We

conducted the fish survey under the guidance of Fisheries Adjustment Rules of the

prefectures and Yamaguchi University. This study was supported by the Environment

Research and Technology Development Fund (4-1602) of Environmental Restoration

and Conservation Agency, Japan and JST-CREST (JPMJCR13A2).

Data availability

All data of the MiSeq sequencing was shared in DRA (Accession number:

DRA008090). Data was available in the supplementary material Table S1 and S2.

Author contributions

HD, RI, and YA designed the study, RI, MG, TK, and YA contributed to field survey

and sampling. SM, HD, RI, and MG contributed to molecular experiments. SM and HD

analyzed the data and interpreted the results. HD, SM, and RI wrote the initial draft of

the manuscript. All other authors critically reviewed the manuscript.

Conflict of Interest

The authors declare no competing financial interests.

References

Almeida-Neto, M., Guimaraes, P., Guimaraes, P. R., Loyola, R. D., & Ulrich, W.

(2008). A consistent metric for nestedness analysis in ecological systems:

reconciling concept and measurement. Oikos, 117, 1227–1239.

Barnes, M. A., & Turner, C. R. (2016). The ecology of environmental DNA and

implications for conservation genetics. Conservation Genetics, 17(1), 1–17.

Baselga, A., Jimenez-Valverde, A., & Niccolini, G. (2007). A multiple-site similarity

measure independent of richness. Biology Letters, 3, 642–645.

Baselga, A. (2010). Multiplicative partition of true diversity yields independent alpha

and beta components; additive partition does not. Ecology, 91(7), 1974–1981.

Baselga, A. (2012). The relationship between species replacement, dissimilarity derived

from nestedness, and nestedness. Global Ecology and Biogeography, 21, 1223–

1232.

Baselga, A., Gómez-Rodríguez, C., & Lobo, J. M. (2012). Historical legacies in world

amphibian diversity revealed by the turnover and nestedness components of beta

diversity. PloS ONE, 7(2), e32341.

Baselga, A., & Orme, C. D. L. (2012). betapart: an R package for the study of beta

diversity. Methods in Ecology and Evolution, 3, 808-812.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects

Models Using lme4. Journal of Statistical Software, 67, 1-48.

Bylemans, J., Gleeson, D. M., Lintermans, M., Hardy, C. M., Beitzel, M., … Furlan, E.

M. (2018). Monitoring riverine fish communities through eDNA metabarcoding:

determining optimal sampling strategies along an altitudinal and biodiversity

gradient. Metabarcoding and Metagenomics, 2, e30457.

Cáceres, M. D., & Legendre, P. (2009). Associations between species and groups of

sites: indices and statistical inference. Ecology, 90(12), 3566–3574.

Chao, A., Gotelli, N. J., Hsieh, T. C., Sander, E. L., Ma, K. H., Colwell, R. K., &

Ellison, A. M. (2014). Rarefaction and extrapolation with Hill numbers: a

framework for sampling and estimation in species diversity studies. Ecological

Monographs, 84, 45-67.

Deiner, K., & Altermatt, F. (2014). Transport distance of invertebrate environmental

DNA in a natural river. PLOS ONE, 9, e88786.

Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J. C., & Altermatt, F. (2016).

Environmental DNA reveals that rivers are conveyer belts of biodiversity

information. Nature Communications, 7, 12544.

Deiner K, Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt,

F., ... Pfrender, M. E. (2017). Environmental DNA metabarcoding:

Transforming how we survey animal and plant communities. Molecular

Ecology, 26, 5872–5895. doi: 10.1111/mec.14350.

Doi, H., Fukaya, K., Oka, S. I., Sato, K., Kondoh, M., & Miya, M. (2019). Evaluation

of detection probabilities at the water-filtering and initial PCR steps in

environmental DNA metabarcoding using a multispecies site occupancy model.

Scientific Reports, 9, 3581.

Doi, H., Inui, R., Akamatsu, Y., Kanno, K., Yamanaka, H., Takahara, T., & Minamoto,

T. (2017). Environmental DNA analysis for estimating the abundance and

biomass of stream fish. Freshwater Biology, 6, 30–39.

Doi, H., & Takahara, T. (2016). Global patterns of conservation research importance in

different countries of the world. PeerJ, 4, e2173.

Drummond, A. J., Newcomb, R. D., Buckley, T. R., Xie, D., Dopheide, A., Potter, B.

C., Heled, J., Ross, H. A., Tooman, L., Grosser, S. & Park, D. (2015).

Evaluating a multigene environmental DNA approach for biodiversity

assessment. GigaScience, 4:s13742-015.

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST.

Bioinformatics, 26, 2460–2461. doi: 10.1093/bioinformatics/btq461.

Fujii, K., Doi, H., Matsuoka, S., Nagano, S., Sato, H., & Yamanaka, H. (2019).

Environmental DNA metabarcoding for fish community analysis in backwater

lakes: A comparison of capture methods. PLoS ONE, 14, e0210357.

Gaston, K. J., & T. M. Blackburn. (2000). Pattern and process in macroecology.

Blackwell Scientific, Oxford.

Goldberg, C. S., Strickler, K. M., & Pilliod, D. S. (2015). Moving environmental DNA

methods from concept to practice for monitoring aquatic macroorganisms.

Biological Conservation, 183, 1–3. doi: 10.1016/j.biocon.2014.11.040.

Gotelli, N. J. (2000). Null model analysis of species co-occurrence patterns. Ecology,

81, 2606–2621.

Hamady, M., Walker, J. J., Harris, J. K., Gold, N. J., & Knight, R. (2008). Error-

correcting barcoded primers for pyrosequencing hundreds of samples in multiplex.

Nature Methods, 5, 235–237.

Harrison, S., Ross, S. J., & Lawton, J. H. (1992). Beta-diversity on geographic gradients

in Britain. Journal of Animal Ecology, 61, 151–158.

Horikawa, M., Nakajima, J. and Mukai, T. (2007). Distribution of indigenous and non-

indigenous mtDNA haplotypes of *Biwia zezera* (Cyprinidae) in northern

Kyushu. Japanese Journal of Ichthyology, 54, 149-159.

Itakura, H., Wakiya, R., Yamamoto, S., Kaifu, K., Sato, T., & Minamoto, T. (2019).

Environmental DNA analysis reveals the spatial distribution, abundance, and

biomass of Japanese eels at the river-basin scale. Aquatic Conservation: Marine

and Freshwater Ecosystems, 29(3), 361–373.

Kawanabe, H., Mizuno, N. & K. Hosoya (eds.) (2001) Freshwater Fishes of Japan

Yama-Kei, Tokyo (in Japanese).

Kembel, S. W., Cowan, P. D., Helmus, M. R., Cornwell, W. K., Morlon, H., Ackerly,

D. D., et al. (2010). Picante: R tools for integrating phylogenies and ecology.

Bioinformatics, 26, 1463-1464.

Margules, C. R., & Pressey, R. L. (2000). Systematic conservation planning. Nature,

405, 243.

Maechler, E., Little, C.J., Wuethrich, R., Alther, R., Fronhofer, E.A., Gounand, I.,

Harvey, E., Huerlemann, S., Walser, J.-C. & Altermatt, F. (2019). Assessing

different components of diversity across a river network using eDNA.

Environmental DNA, 1, 290-301.

Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J. Y., Sato, K., ... Kondoh, M.

(2015). MiFish, a set of universal PCR primers for metabarcoding environmental

DNA from fishes: detection of more than 230 subtropical marine species. Royal

Society Open Science, 2, 150088.

Nakabo, T. (2013). Fishes of Japan with pictorial keys to the species, 3rd ed. Tokai

University Press (in Japanese).

Nakagawa, H., Yamamoto, S., Sato, Y., Sado, T., Minamoto, T., & Miya, M. (2018).

Comparing local- and regional-scale estimations of the diversity of stream fish

using eDNA metabarcoding and conventional observation methods. Freshwater

Biology, 63, 569–580.

Pecl, G. T., Araújo, M. B., Bell, J. D., Blanchard, J., Bonebrake, T. C., Chen, I. C., ...

Falconi, L. (2017). Biodiversity redistribution under climate change: Impacts on

ecosystems and human well-being. Science, 355, 9214.

Primack, R. B. (1993). Essentials of conservation biology (Vol. 23). Sunderland,

Massachusetts: Sinauer Associates.

R Development Core Team. (2018). R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

https://www.R-project.org/.

Rees, H. C., Maddison, B. C., Middleditch, D. J., Patmore, J. R., & Gough, K. C.

(2014). Review: the detection of aquatic animal species using environmental DNA

— a review of eDNA as a survey tool in ecology. Journal of Applied Ecology, 51,

1450–1459. doi: 10.1111/1365-2664.12306.

Sato, H., Sogo, Y., Doi, H., & Yamanaka, H. (2017). Usefulness and limitations of

sample pooling for environmental DNA metabarcoding of freshwater fish

communities. Scientific Reports, 7, 14860. doi: 10.1038/s41598-017-14978-6.

Seymour, M., Durance, I., Cosby, B. J., Ransom-Jones, E., Deiner, K., Ormerod, S.

J., ... Edwards, F. (2018). Acidity promotes degradation of multi-species

environmental DNA in lotic mesocosms. Communications Biology, 1(1), 4.

Socolar, J. B., Gilroy, J. J., Kunin, W. E., & Edwards, D. P. (2016). How should beta-

diversity inform biodiversity conservation? Trends in Ecology & Evolution, 31(1),

67–80.

Stæhr, P. A., Santos, S., Hansen, L. H., Lundsteen, S., Haraguchi, L., Dahl, K., Ávila,

M. P., Winding, A. & Boutrup, S. (2016). Comparison of eDNA and conventional

techniques for monitoring species diversity of boulder reefs in Danish waters. DCE-

Danish Centre for Environment and Energy. Aarhus University, p.22.

Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., & Willerslev, E. (2012).

Towards next-generation biodiversity assessment using DNA metabarcoding.

Molecular Ecology, 21, 2045–2050.

Takahara T., Minamoto T., Yamanaka H., Doi H. & Kawabata Z. (2012). Estimation of

fish biomass using environmental DNA. PLOS ONE, 7, e35868

Takahara, T., Minamoto, T., & Doi, H. (2013). Using environmental DNA to estimate

the distribution of an invasive fish species in ponds. PLoS ONE, 8, e56584

Thomsen, P. F., Willerslev, E. (2015). Environmental DNA–an emerging tool in

conservation for monitoring past and present biodiversity. Biological

Conservation, 183, 4–18. doi: 10.1016/j.biocon.2014.11.019.

Uchii, K., Doi, H., & Minamoto, T. (2016). A novel environmental DNA approach to

quantify the cryptic invasion of non-native genotypes. Molecular Ecology

Resources, 16, 415–422. doi: 10.1111/1755-0998.12460.

Ulrich, W., & Gotelli, N. J. (2007). Null model analysis of species nestedness patterns.

Ecology, 88(7), 1824–1831.

Wright, D. H., & Reeves, J. H. (1992). On the meaning and measurement of nestedness

of species assemblages. Oecologia, 92(3), 416–428.

Yamamoto, S., Masuda, R., Sato, Y., Sado, T., Araki, H., Kondoh, M., ... Miya, M.

(2017). Environmental DNA metabarcoding reveals local fish communities in a

species-rich coastal sea. Scientific Reports, 7, 40368.

Yamanaka, H., Motozawa, H., Tsuji, S., Miyazawa, R. C., Takahara, T., & Minamoto,

T. (2016). On-site filtration of water samples for environmental DNA analysis to

avoid DNA degradation during transportation. Ecological Research, 31(6), 963–

967.

Table 1 Indicator taxa analysis for the taxa which had significantly different detection

frequencies between eDNA metabarcoding and visual/capture methods (P < 0.05). Best

means preferred methods. P-values was calculated with 999 permutations after Sidak's

correction of the multiple testing.

Taxa	P-value for eDNA metabarcoding	P-value for visual and capturing survey	best	P-value for multiple testing
Anguilla japonica	0.001	1.000	eDNA metabarcoding	0.002
Oncorhynchus masou	0.001	1.000	eDNA metabarcoding	0.002
Oncorhynchus mykiss	0.016	1.000	eDNA metabarcoding	0.032
Cyprinus carpio	0.001	1.000	eDNA metabarcoding	0.002
Silurus asotus	0.003	1.000	eDNA metabarcoding	0.006
Misgurnus anguillicaudatus	0.009	1.000	eDNA metabarcoding	0.018

Figure legends

Figure 1. Numbers of taxa detected by eDNA metabarcoding (red), visual/capture

surveys (blue), or both methods (white) at individual sites and for sites combined (ALL)

within each of five river systems in Japan. Total taxa detected by eDNA metabarcoding

is the sum of values included within the larger circle. Total taxa detected by

visual/capture surveys is the sum of values in blue and white areas.

Figure 2. Nonmetric multidimensional scaling (NMDS) ordination of fish communities

evaluated by study river (shapes) and segment (colors), using Jaccard index (ordination

with Raup-Crick index shown in Fig. S4). A single ordination is divided into two panels

to display differences in communities as measured by sampling methods. MDS stress

was 0.158.

Figure 3. Species nestedness and turnover between the communities detected by eDNA

metabarcoding and visual/capture methods, expressed as pairwise standardized effect

sizes (SES). Error bars indicate 95% confidence intervals. The horizontal line represents

SES=0, indicating a non-significant effect.

Figure 4. Effect sizes for (a) community nestedness and (b) species turnover among

river segments as assessed by eDNA metabarcoding and visual/capture methods. The

error bars indicate 95% confidence intervals

Figure 1

Figure 2

Figure 3

Figure 4