A neural system for encoding odor valence which influences approach behaviors

Marie A. Gadziola[1,2,3,#], Lucas A. Stetzik[1,2#], Adrianna J. Milton[3], Keiko Arakawa[3], Daniel W. Wesson[1,2,3,*]

[1]Department of Pharmacology & Therapeutics
[2]Center for Smell & Taste
University of Florida
1200 Newell Dr.
Gainesville, FL, 32610. U.S.A.

[3]Department of Neurosciences
Case Western Reserve University
2109 Adelbert Rd.
Cleveland, OH, 44106. U.S.A.

[#]Equal contributing first-authors

[*]corresponding author: danielwesson@ufl.edu (D.W.)

Running head: Valence and approach in olfaction.

Abstract

Odors may elicit strong emotional and behavioral responses which are known to be strengthened through learning. Presumably subserving this, olfactory structures are well-known for their experience-dependent plasticity, yet the specific cellular systems involved and the actual influence of these systems on behavior are less understood. Here we investigated the transformation of odor information throughout the association of odors with learned-rewarded outcomes and also sought to link those neural systems with displays of reinforcement-based approach behaviors. First, we investigated the representation of odor-reward associations, also known as valence, within two areas recipient of dense olfactory bulb input, the posterior piriform cortex (pPCX) and olfactory tubercle (OT), using simultaneous multi-site electrophysiological recordings from mice engaged in a reward-based olfactory learning task. As expected, neurons in both regions represented the reward-associated valence of odors, yet both the proportion of neurons recruited by conditioned rewarded odors and the magnitudes of their recruitment were distinctly pronounced in the OT. Using fiber photometry we found that odor valence coding in the OT occurs among D1- but not D2-type dopamine receptor expressing neurons. In both the recordings and imaging, statistically meaningful changes in activity occurred prior to instrumental behavior responses. Finally, using optogenetics we show that OT D1-receptor expressing neurons strongly influence reward-motivated approach behavior. Together our results contribute to a model whereby OT D1 neurons support olfactory valence and ultimately hedonic-based behavioral responses.

Introduction

Upon learning, stimuli may acquire meaning, also known as valence, which is considered integral in guiding behavioral choice (Berridge and Aldridge, 2008; Chikazoe et al., 2014; Gottfried, 2009; Kumar et al., 2012; Pessoa and Adolphs, 2010; Veldhuizen et al., 2009). Learning to associate a stimulus with an outcome shapes our actions in profound manners – including impacting the choices of foods we eat and can lead to maladaptive outcomes such as those resulting from engaging in compulsive behaviors. This process enables stimuli to hold 'valence' (Lewin, 1935) which is of essence in the execution of behaviors which support satiation of basic needs (Lewin, 1935; Tolman, 1932). Despite holding fundamental importance for our day-to-day actions, the neural systems underlying the generation of valence within the brain are unresolved.

Neural signatures of valence can be observed in the form of flexible expressions of neural activity in response to stimuli throughout reward- or emotion-based associative learning (*e.g.,* (Berridge, 2019; Gore et al., 2015; Hickey and Peelen, 2015; Isosaka et al., 2015; Morrison and Salzman, 2009; Schoenbaum et al., 1998)). This has been well-established in the orbitofrontal cortex and basolateral amygdala, for instance, wherein neurons in these associative structures display differential firing following reinforcement learning for stimuli conditioned to predict a salient outcome (Roesch et al., 2007; Schoenbaum and Eichenbaum, 1995; Schoenbaum et al., 2003).

Behavioral displays linked to the valence of odors have been observed in all animals studied to date and involve a variety of brain systems (Knaden and Hansson, 2014; Li and Liberles, 2015). Olfactory system structures display forms of plasticity (Barnes et al., 2008; Chapuis and Wilson, 2011; Dias and Ressler, 2014; Doucette et al., 2011; Kass et al., 2013; Lebel et al., 2001; Li et al., 2008; Mandairon and Linster, 2009; Murata et al., 2015; Ross and Fletcher, 2018; Schoenbaum et al., 2000), yet in many of these cases the specific cellular systems involved and/or whether those systems specifically influence behavior are less understood. Among these, olfactory cortical structures including both the piriform cortex (PCX) and olfactory tubercle (OT) encode the valence of odors. Specifically, work from several groups including ours (Gadziola et al., 2015; Gire et al., 2013; Roesch et al., 2007; Schoenbaum and Eichenbaum, 1995), have demonstrated that PCX and OT neurons are more greatly recruited by odors conditioned to predict rewards versus those which are unreinforced. These results have indicated that olfactory cortices encode learned odor valence by a combination of recruiting neurons into an ensemble and also divergent firing for rewarded odors.

The representation of odor valence by both PCX and OT neurons leads to the assumption that valence is a distributed, global property of the olfactory cortex. Indeed, that both PCX and OT neurons encode odor valence with divergent firing (Gadziola et al., 2015; Gire et al., 2013;

Roesch et al., 2007; Schoenbaum and Eichenbaum, 1995) suggests a possible substrate wherein

information within these systems is more-or-less equally relayed into downstream structures

important for learning and memory (PCX; (Gottfried, 2010; Wilson and Sullivan, 2011)) or those

integral for motivated and affective behaviors (OT; (Wesson and Wilson, 2011; Zhang et al.,

2017a)). However, this assumption is not warranted since presently no information exists

regarding whether either of these structures are more specialized in their capacity to encode odor

valence. As a specific void, no investigations of these structures simultaneously during valence

learning exist. Thus, an alternative system may be in place: that olfactory cortices differentially

represent odor valence in manners suggesting that one structure is specialized in valence

assignment functions. Determining whether valence is a global property distributed among the

PCX and OT is an important question in the overall goal of resolving the neural circuitry underlying

olfactory valence.

Based upon the representation of learned odor valence in the OT (Gadziola et al., 2015;

Murata et al., 2015), and the positioning of the OT within the ventral striatum (Heimer et al., 1982;

De Olmos and Heimer, 1999) wherein it receives dense innervation from dopaminergic neurons

in the ventral tegmental area (Ikemoto, 2007; Voorn et al., 1986), we predicted that the OT is a

central figure in assigning odors with valence. Infusion of psychoactive substances into the OT is

reinforcing, and rodents more robustly seek infusions into the OT than even into the nucleus

accumbens (Ikemoto, 2003) – highlighting the likely importance of OT dopamine in guiding

approach behavior, as also indicated by a recent optogenetic study (Zhang et al., 2017b). An

additional feature of the OT which supports this hypothesis is its cellular composition. The OT is

primarily composed of medium spiny neurons expressing either the D1- or D2-type dopamine

receptor (Murata et al., 2015). D1 MSNs in many striatal structures are activated by appetitive

stimuli and are often considered integral for promoting stimulus reward and approach, whereas

at least some reports indicate D2-type MSNs are less shaped by reinforcement (for reviews (Lobo

and Nestler, 2011; Soares-Cunha et al., 2016). Therefore, it is possible that previous reports of

neuromodulation driving reinforcement through the OT (Ikemoto, 2003) is due to direct actions

upon OT D1-receptor expressing neurons. Supporting this hypothesis, c-Fos expression is

elevated in OT D1-type neurons following associative learning (Murata et al., 2015). Based upon

the above, we predicted that OT D1-receptor expressing neurons may underlie odor valence

representations in the OT and separately, should therefore also be capable of promoting

reinforcer-motivated approach behaviors, similar to that needed to subserve engagement with

odors.

Here we tested the above hypotheses through a combination of multi-site single-unit

recordings, cell type-specific fiber photometry, and cell type-specific optogenetic studies, all in

mice engaged in reinforcer-motivated operant tasks. All together our results contribute to a model

whereby the OT, and specifically OT D1-receptor expressing neurons support olfactory hedonics

and ultimately hedonic-based behavioral responses.

Results

Compared to pPCX, neurons in the OT robustly encode valence.

We began by examining whether valence is a global property distributed among the pPCX

and OT. We simultaneously monitored OT and posterior PCX (pPCX) single-unit activity from 11

mice implanted with chronic electrode arrays (**Fig 1 & S1A**) while they engaged in a head-fixed

odor discrimination task requiring them to lick a spout for a palatable reinforcer available at the

offset of conditioned-rewarded but not unrewarded odors (**Fig 1B&C**). This head-fixed paradigm

allows odor-guided behaviors while maintaining precision in odor-delivery (Verhagen et al., 2007).

To optimize the data contributed from each animal, all mice were shaped to discriminate between

2-4 odor pairs (Methods). Further, by excluding trials wherein animals prematurely licked during

the odor we can reduce influences of motor planning or execution on the neural analyses.

Baseline and odor-evoked activity were measured for isolated single neurons. The baseline firing

rates of OT (median rate: 3.1 Hz; range = 0 - 56.9 Hz) and pPCX neurons (median rate: 2.1 Hz;

range = 0 - 35.6 Hz) were low across the sampled population (**Fig S1B**, measured 2s prior to odor

onset). In both structures, odor-evoked firing was observed in response to the odors and occurred

soon after odor onset (0.35 ± 0.31s [OT], 0.57 ± 0.48s [pPCX], **Fig 1D**).

After mice learned to discriminate between odor pairs (**Fig S1C**) we observed that the

majority of OT neurons were significantly modulated in their firing during the task window (65%,

52/80 OT neurons 'task-responsive' within 7s of odor onset). This was comparable to the

population of pPCX neurons modulated in the same time (X^2(df=1, n=87) = 3.32, p>0.05; 44%,

35/79 neurons). To characterize responses across these populations, we measured the temporal

response profile of each neuron-odor pair during correct trials by quantifying changes in firing rate

from baseline using a sliding receiver operating characteristic (ROC) analysis (Cohen et al., 2012;

Gadziola et al., 2015; Veit and Nieder, 2013). Area under the ROC curve (auROC) values greater

than 0.5 indicate increases (excitation) in firing rate compared to baseline, with values under 0.5

indicating decreases (suppression). This analysis revealed that 91% (189/208) of OT neuron-odor

pairs and 78% (110/140) of pPCX neuron-odor pairs displayed significant stimulus-evoked activity

during odor presentation, with significant changes in firing rate observed shortly (<500 ms) after

odor onset. Of these responses, 71% of OT and 59% of pPCX neurons increased their firing rate

relative to baseline, with the remaining neurons suppressing their firing rates relative to baseline.

Across the population of neurons from both regions, we observed that OT neurons

displayed more divergent responses to the conditioned odors than pPCX neurons, with more

robust and longer-lasting responses to conditioned rewarded than unrewarded odors (**Fig 2**). To

test whether the populations of OT and pPCX neurons preferentially encode one conditioned odor

over the other, we averaged the auROC values computed across the first 500-ms following odor

onset. OT neurons preferentially responded to rewarded odor with higher firing than the

unrewarded odors (0.59 ± 0.02 vs 0.55 ± 0.01, respectively; paired t test: $t_{(103)}$ = -6.05, p <

0.0001), whereas pPCX neurons did not respond differentially to rewarded and unrewarded odors

across the population (0.53 ± 0.01 vs 0.53 ± 0.01, respectively; paired t test: $t_{(69)}$ = -0.73, p =

0.47). Furthermore, average auROC values in response to rewarded odors were greater for the

population of OT neurons when compared to pPCX neurons (0.59 ± 0.02 vs 0.53 ± 0.01,

respectively; two-sample t test: $t_{(172)}$ = -3.49, p < 0.001). There were no discernable differences in

the temporal response pattern of neurons that increased vs decreased firing rates, suggesting

that these neurons are similarly driven by conditioned odors (**Fig S2**). Next, we compared the

duration of significant auROC responses in the first 500ms after odor onset across the population

of neurons in both regions (including units with ≥ 1 significant bin). OT responses were

significantly longer in duration compared to pPCX responses (209 ± 97 vs 166 ± 91 ms,

respectively; two-sample t test: $t_{(290)}$ = -3.64, p < 0.001). In addition, OT response durations to

rewarded odors were longer than unrewarded odors (225 ± 94 vs 193 ± 98 ms, respectively;

paired t test: $t_{(96)}$ = -12.01, p < 0.0001), and longer than pPCX responses to rewarded odors (225

± 94 vs 167 ± 84 ms, respectively; two-sample t test: $t_{(141)}$ = -3.51, p < 0.0001). These results

support the conclusion that the representation of odor valence is not equally distributed among

the OT and pPCX – with the OT dominating in assigning valence.

D1 receptor expressing OT neurons display divergent responses to conditioned rewarded odors.

Which neurons are responsible for the pronounced display of valence in the OT? We

addressed this question by performing fiber photometry-based imaging (*e.g.,* (Gunaydin et al.,

2014; Kudo et al., 1992)) of GCaMP6f (Chen et al., 2013) from D1 and D2 receptor-expressing

neurons in the OT. Mice expressing Cre-recombinase in either neurons expressing the D1
receptor (D1-Cre) or D2 receptor (D2-Cre) (Gong et al., 2003) were injected with floxed GCaMP6f
into their OT and later implanted with optical fibers (see Methods) before being water deprived
and shaped in the same lick/no-lick odor discrimination task used above. During imaging (**Figs**
3A-C), the GCaMP6f emission spectrum was collected as a measure of aggregate OT D1 or D2
neuron activity. Simultaneously, from all recordings, we also collected endogenous UV emission
spectrum as a control measure for movement artifacts. The resultant GCaMP6f signal was
subtracted from the UV signal for a single output, which is considered closely reflective of
aggregate neural activity. In separate experiments we confirmed that GCaMP6f expression
robustly colocalizes with D1 and D2 receptor expressing neurons in these Cre lines by crossing
the Ai9 TdTomato reporter line (Madisen et al., 2012) with the D1-Cre and D2-Cre mice. The
resultant D1-Cre;Ai9 and D2-Cre;Ai9 mice were injected with floxed GCaMP6f into their OT and
allowed to recover 2 weeks prior to tissue collection. High levels of colocalization were found in
the OTs of both lines (**Fig 3A**; D1-Cre;Ai9: 95.01 ± 0.96 [inter-animal range 94.0-95.9]; D2-
Cre;Ai9: 95.52 ± 1.7 [inter-animal range 93.6-97.0] mean ± sd), thus confirming the targeted
specificity of GCaMP to the desired cell populations.

We imaged 5 D1- and 5 D2-Cre mice with fiber implants and GCaMP6f expression both
confirmed in the OT (**Fig S3**) throughout learning and performance in the lick/no-lick odor
discrimination task. To enhance each animal's contributions for a more rigorous data set, we
shaped each mouse on several odor pairs over differing daily sessions, yet only one pair of odors
in a given session. This allowed each animal to contribute odor-modulated GCaMP6f signals to
2-3 rewarded and unrewarded odors (on different sessions) at or above criterion performance
(mean rewarded odor trials / session = 44.5 [24-75 min-max range]; mean unrewarded odor trials
/ session = 91.3 [52-133 min-max range]). Across all rewarded and unrewarded odors, we
observed modulations in both D1 and D2 neuron activity following odor onset. This was evident
both looking within individual mice at subsets of trials (**Fig 3C & D**), and averaged across mice
over several sessions (**Fig 3E & F**). No discernable differences were observed in the time prior
to odor onset, or before 0.5s following odor onset between any stimulus type in either D1 or D2
mice (**Fig 3E & F**). Later during the odor period however (0.5-0.75s), across all D1 mice,
conditioned rewarded and unrewarded odors elicited differing amounts of activity ($p<0.001$; **Fig**
3E). Whereas D1 neurons were largely unmodulated by the conditioned unrewarded odor,
conditioned rewarded odors elicited large increases in activity during the odor. In contrast, no
differential representation of conditioned rewarded versus unrewarded odors was observed in D2

mice, with both conditioned stimuli eliciting similar increases in activity during odor (p=0.483; **Fig
3F**).

The responses of D1 receptor-expressing OT neurons are flexible and encode the associated
outcomes of odors.

Having imaged D1 and D2-type receptor expressing OT neurons and observing that they
differently encode odor-reward associations, we were next able to ask whether this differential
representation of conditioned rewarded odors observed among the D1 neuron population is
flexible. In other words, does the representation of conditioned-rewarded odors flexibly follow as
odors are associated to predict new outcomes? To address this, in a separate behavioral session
we used reversal learning to reverse the odor-outcome contingencies so that a previously
rewarded odor is no longer rewarded and a previously unrewarded odor is now rewarded. Most
mice were shaped upon at least 2 odor-reversals (differing odor pairs, range 2-3). If a population
of neurons is 'identity' encoding they should exhibit the same odor selectivity before and after
reversal learning, whereas a population encoding the associative valence of an odor (rewarded
or not) should reverse its odor preference (Gire et al., 2013; Roesch et al., 2007). Animals took
several sessions to learn the reversal to criterion in this task requiring them to withhold licking
until odor offset, and comparisons were made only from sessions following task acquisition, during
blocks consisting of criterion-level performance, and correct responses during each trial.

On average across animals, the evoked GCaMP6f signals in D1-Cre mice tracked the
behavioral contingencies – with the original rewarded odor and the newly rewarded odor eliciting
evoked responses with similar amplitudes (p=0.202; **Fig. 3G**). Likewise, the original unrewarded
odors elicited similar responses as the new unrewarded odors (p=0.177). In contrast, no effect of
reversal on either the rewarded (p=0.785) or unrewarded odors (p=0.522) were observed in the
D2-Cre mice (**Fig. 3H**). D2 neurons maintained somewhat positive responses for all odors,
regardless of valence, indicating that while these neurons are modulated by odor, they do not
represent odor valence. These data support our hypothesis that specifically D1 receptor-
expressing neurons are responsible for the profound representation of odor valence in the OT.

OT D1 receptor-expressing neurons drive robust approach behaviors.

Having observed reward-associated odor valence coding in OT D1, but not D2 receptor-
expressing neurons, we next asked whether the activity of these neurons directly influences the
occurrence of approach behaviors. Indeed, the latency and robustness of OT D1 neuron encoding
of conditioned rewarded odors may serve as an effective way whereby to activate approach
behaviors. To test this, D1-Cre mice injected with a floxed AAV encoding channelrhodopsin
(ChR2) [n = 7] (or floxed AAV encoding a reporter fluorophore [n = 5]) into their ventral striatum
were later implanted with an unilateral optical fiber into their OT (**Fig 4A**). Mice were shaped in
an optical intra-cranial self-stimulation task, adapted from (Carlezon and Chartoff, 2007; Ilango et
al., 2014; Vicente et al., 2016), allowing for assaying of stimulation-seeking goal-directed
behaviors (**Fig 4B**). Shaping in this instrumental task occurred over several phases (See
Methods). In the first phase, mice could poke their snout into either of two ports in exchange for
blue light stimulation into their OT (25Hz, for 2s, fixed ratio 1). All but one of the D1-Cre mice
injected with AAV.ChR2 reached a set level of criterion performance (40 pokes into either port
during a 1 hr session)(**Fig 4C**), indicating that they readily learned to poke in exchange for blue
light stimulation. No control AAV injected mice reached criterion performance (**Fig 4C**). The mice
reaching criterion on Phase 1 then progressed onto Phase 2 wherein only one of the two ports
delivered light. The port receiving the greater number of pokes during Phase 1 was selected as
the 'active' port for Phase 2. The vast majority of AAV.ChR2 mice (5/6) reached criterion on Phase
2, with some poking over 500 times in the single hour of testing (**Fig 4Di & ii**). To ensure that
these mice were indeed seeking light stimulation, on the subsequent Phase 3, the location of the
active port was reversed and despite this, all the mice who reached criterion on Phase 2 redirected
their responding to criterion levels for the new active port. Compared to the first session of reversal
phase testing, there were significantly more pokes for the active port during both the last session
of phase 2 testing (p=0.0002), as well as during the last session of reversal (p=0.0004). These
results indicate that OT neurons expressing the D1-type dopamine receptor support behavioral
approach, specifically by signaling reinforcement.

Discussion

The olfactory system holds a remarkable capacity for experience-based plasticity and informing
a wide-range of critical behaviors. Olfactory structures, including those more associative in nature
(*e.g.,* the orbitofrontal cortex), are well-known for their plasticity (Barnes et al., 2008; Chapuis and
Wilson, 2011; Dias and Ressler, 2014; Doucette et al., 2011; Kass et al., 2013; Lebel et al., 2001;

Li et al., 2008; Mandairon and Linster, 2009; Murata et al., 2015; Ross and Fletcher, 2018;
Schoenbaum et al., 2000), yet in many of these cases the specific cellular systems involved and/or
whether those systems specifically influence behavior is less understood. Here we build off of
previous studies which have uncovered odor-reward association-based plasticity among neurons
in both the PCX and OT (in separate studies; (Calu et al., 2007; Gadziola et al., 2015; Gire et al.,
2013)) to interrogate the respective roles of each of these structures in the representation of odor
valence by means of simultaneous single-unit recordings. After identifying robust odor valence
coding among neurons in the OT significantly more-so than those in the pPCX, we employed fiber
photometry in D1-Cre and D2-Cre mice to uncover evidence indicating that OT D1-, but not D2-
type dopamine receptor expressing neurons encode odor valence. Finally, we linked the activity
of OT D1 neurons to the display of reward-driven behaviors, specifically approach behaviors.

This work adds to a growing body of literature seeking to understand the brain systems
governing the generation of stimulus hedonics and preferences (for review (Berridge and
Kringelbach, 2015)). Attraction to odors is well established and may employ innate (unlearned) or
acquired (learned) mechanisms. Since the olfactory bulb is the starting-point for odor perception,
it is not surprising that the olfactory bulb holds importance for both innate and learned odor
hedonics (*e.g.,* (Doucette et al., 2011; Kermen et al., 2016; Kobayakawa et al., 2007; Wilson et
al., 1987). Indeed, recent work uncovered evidence that a specific zone within the main olfactory
bulb is necessary for generating odor hedonics (Kermen et al., 2016). The hedonic information of
odors in the OB next may be distributed into the OT and PCX. Both PCX and OT neurons may
be recruited to encode odor valence and do so with divergent firing to conditioned rewarded
versus unrewarded odors (Gadziola et al., 2015; Gire et al., 2013; Roesch et al., 2007;
Schoenbaum and Eichenbaum, 1995). A key prediction in this work was that OT units would more
greatly represent odor valence than those in the PCX. That said, it is quite surprising the
magnitude at which we found this occurs. OT units far surpassed the size of the population of
pPCX units encoding valence and also exceeded pPCX units in their magnitude of firing for
conditioned rewarded odors. This inter-regional difference could not simply be the result of
behavioral, affective, or other state-dependent influences since recordings were sampled from
both the pPCX and OT simultaneously. Thus, this work uncovers that systems downstream from
the OB are specialized in their representation of odor valence. This outcome is consistent with
our hypothesis and also that of other groups (Kermen et al., 2016; Murata et al., 2015).

Specific cell populations in several brain regions have been linked to stimulus valence or
to approach responses (Betley et al., 2015; Kim et al., 2019; Li and Liberles, 2015; Li et al., 2019;
Miller et al., 2019; Soares-Cunha et al., 2016). This work adds OT D1 receptor expressing
neurons, in an underappreciated ventral striatum structure, the OT, to this list and suggests a
model wherein OT D1-MSNs may signal to downstream systems which also influence approach
like the amygdala or hypothalamic nuclei. This model, as also supported by immediate early gene
mapping experiments following olfactory learning (Murata et al., 2015), would be potentially
powerful in cases wherein odors associated with positive outcomes, which as we show here are
represented in OT D1 neurons, could influence downstream limbic systems to strengthen
approach responding through their coordinated output.

All together these results support a role for OT D1 neurons, not unlike D1 neurons in the
nucleus accumbens (Lobo and Nestler, 2011), in behaviors integral for valence -- reinforcement
and stimulus-seeking. To come to this conclusion, we used a straight-forward instrumental
responding paradigm wherein mice were allowed to nose-poke in exchange for OT D1 neuron
stimulation. This paradigm has several advantages over real-time place preference testing or
other tests which serve to assay approach behaviors since it allows for measures of discrete,
stimulus (*i.e.*, light) seeking behaviors (nose-pokes). The finding that some animals responded at
the rate they did, indeed up to 500 pokes in an hour, provides a compelling demonstration that
these neurons influence motivation states, or tap into reward circuitry, in a manner which
promotes approach behavior. This test was done in the absence of any experimentally-delivered
odors which further reinforces the notion that optical stimulation of OT D1 neurons engages
important internal circuitry for guiding behavior, versus that which may get paired with odor
information entering the OT from the olfactory bulb. OT D1 MSNs send efferents throughout the
brain's affective and sensory centers (Zhang et al., 2017a). What specific output targets of OT D1
cells are responsible for approach responses will be an important question for future research to
address.

Taken together, our results contribute to a model whereby the OT is a specialized center
for encoding odor valence. This along with the finding that OT D1 neurons encode olfactory
valence and robustly motivate reinforcement-seeking in a nose-poke based task which emulates
snout-directed investigation behaviors like those used by rodents to sample odors, leads us to
predict that OT D1-type MSNs are a fundamental component of brain systems needed to inform
and respond to the hedonics of odors.

Acknowledgements: This work was supported by NIH NIDCD grants R01DC014443 and
R01DC016519 to D.W. Some mouse strains used in this research were obtained from the Mutant
Mouse Regional Resource Center (MMRRC), an NIH funded strain repository, and were donated
to the MMRRC by the MMRRC facility at the University of California, Davis.

Author Contributions: Conceptualization: M.A.G., L.A.S., D.W.W.; Methodology: M.A.G.,
L.A.S., D.W.W.; Investigation: M.A.G., L.A.S., A.J.M., K.A., D.W.W.; Writing – Original Draft:
M.A.G., L.A.S., D.W.W.; Writing – Review & Editing: M.A.G., L.A.S., D.W.W.; Visualization:
M.A.G., L.A.S., D.W.W.; Supervision: D.W.W.; Funding Acquisition: D.W.W.

Declaration of Interests: The authors declare no competing interests.

References

Barnes, D.C., Hofacer, R.D., Zaman, A.R., Rennaker, R.L., and Wilson, D.A. (2008). Olfactory
perceptual stability and discrimination. Nat Neurosci *11*, 1378–1380.

Berridge, K.C. (2019). Affective valence in the brain: modules or modes? Nat. Rev. Neurosci.
20, 225–234.

Berridge, K.C., and Aldridge, J.W. (2008). Decision utility, the brain, and pursuit of hedonic
goals. Soc. Cogn. *26*, 621–646.

Berridge, K.C., and Kringelbach, M.L. (2015). Pleasure systems in the brain. Neuron *86*, 646–
664.

Betley, J.N., Xu, S., Cao, Z.F.H., Gong, R., Magnus, C.J., Yu, Y., and Sternson, S.M. (2015).
Neurons for hunger and thirst transmit a negative-valence teaching signal. Nature *521*, 180.

Calu, D.J., Roesch, M.R., Stalnaker, T.A., and Schoenbaum, G. (2007). Associative Encoding in
Posterior Piriform Cortex during Odor Discrimination and Reversal Learning. Cereb Cortex *17*,
1342–1349.

Carlezon, W.A., and Chartoff, E.H. (2007). Intracranial self-stimulation (ICSS) in rodents to
study the neurobiology of motivation. Nat. Protoc. *2*, 2987–2995.

Chapuis, J., and Wilson, D.A. (2011). Bidirectional plasticity of cortical pattern recognition and
behavioral sensory acuity. Nat Neurosci *15*, 155–161.

Chen, T.-W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R.,
Kerr, R.A., Orger, M.B., Jayaraman, V., et al. (2013). Ultrasensitive fluorescent proteins for
imaging neuronal activity. Nature *499*, 295–300.

Chikazoe, J., Lee, D.H., Kriegeskorte, N., and Anderson, A.K. (2014). Population coding of
affect across stimuli, modalities and individuals. Nat Neurosci *17*, 1114–1122.

Cohen, J.Y., Haesler, S., Vong, L., Lowell, B.B., and Uchida, N. (2012). Neuron-type-specific
signals for reward and punishment in the ventral tegmental area. Nature *482*, 85–88.

Dias, B.G., and Ressler, K.J. (2014). Parental olfactory experience influences behavior and
neural structure in subsequent generations. Nat Neurosci *17*, 89–96.

Doucette, W., Gire, D.H., Whitesell, J., Carmean, V., Lucero, M.T., and Restrepo, D. (2011).
Associative Cortex Features in the First Olfactory Brain Relay Station. Neuron *69*, 1176–1187.

Gadziola, M.A., and Wesson, D.W. (2016). The Neural Representation of Goal-Directed Actions
and Outcomes in the Ventral Striatum's Olfactory Tubercle. J. Neurosci. *36*, 548–560.

Gadziola, M.A., Tylicki, K.A., Christian, D.L., and Wesson, D.W. (2015). The Olfactory Tubercle
Encodes Odor Valence in Behaving Mice. J. Neurosci. *35*, 4515–4527.

Gire, D.H., Whitesell, J.D., Doucette, W., and Restrepo, D. (2013). Information for decision-
making and stimulus identification is multiplexed in sensory cortex. Nat Neurosci *16*, 991–993.

Gong, S., Zheng, C., Doughty, M.L., Losos, K., Didkovsky, N., Schambra, U.B., Nowak, N.J.,
Joyner, A., Leblanc, G., Hatten, M.E., et al. (2003). A gene expression atlas of the central
nervous system based on bacterial artificial chromosomes. Nature *425*, 917–925.

Gore, F., Schwartz, E.C., Brangers, B.C., Aladi, S., Stujenske, J.M., Likhtik, E., Russo, M.J.,
Gordon, J.A., Salzman, C.D., and Axel, R. (2015). Neural Representations of Unconditioned
Stimuli in Basolateral Amygdala Mediate Innate and Learned Responses. Cell *162*, 134–145.

Gottfried, J.A. (2009). Olfaction and Its Pleasures: Human Neuroimaging Perspectives. In
Pleasures of the Brain, M.L. Kringelbach, and K.C. Berridge, eds. (Oxford University Press), pp.
125–145.

Gottfried, J.A. (2010). Central mechanisms of odour object perception. Nat Rev Neurosci *11*,
628–641.

Green, D.M., and Swets, J.A. (1966). Signal detection theory and psychophysics (New York:
Wiley).

Gunaydin, L.A., Grosenick, L., Finkelstein, J.C., Kauvar, I. V., Fenno, L.E., Adhikari, A.,
Lammel, S., Mirzabekov, J.J., Airan, R.D., Zalocusky, K.A., et al. (2014). Natural Neural
Projection Dynamics Underlying Social Behavior. Cell *157*, 1535–1551.

Heimer, L., Switzer, R.D., and Van Hoesen, G.W. (1982). Ventral striatum and ventral pallidum:
Components of the motor system? Trends Neurosci. *5*, 83–87.

Hickey, C., and Peelen, M.V. (2015). Neural Mechanisms of Incentive Salience in Naturalistic
Human Vision. Neuron *85*, 512–518.

Ikemoto, S. (2003). Involvement of the Olfactory Tubercle in Cocaine Reward: Intracranial Self-
Administration Studies. J. Neurosci. *23*, 9305–9311.

Ikemoto, S. (2007). Dopamine reward circuitry: Two projection systems from the ventral
midbrain to the nucleus accumbens-olfactory tubercle complex. Brain Res Rev *56*, 27–78.

Ilango, A., Kesner, A.J., Keller, K.L., Stuber, G.D., Bonci, A., and Ikemoto, S. (2014). Similar
Roles of Substantia Nigra and Ventral Tegmental Dopamine Neurons in Reward and Aversion.
J. Neurosci. *34*, 817–822.

Isosaka, T., Matsuo, T., Yamaguchi, T., Funabiki, K., Nakanishi, S., Kobayakawa, R., and
Kobayakawa, K. (2015). Htr2a-Expressing Cells in the Central Amygdala Control the Hierarchy
between Innate and Learned Fear. Cell *163*, 1153–1164.

Kass, M.D., Rosenthal, M.C., Pottackal, J., and McGann, J.P. (2013). Fear Learning Enhances
Neural Responses to Threat-Predictive Sensory Stimuli. Science (80-.). *342*, 1389–1392.

Kermen, F., Midroit, M., Kuczewski, N., Forest, J., Thevenet, M., Sacquet, J., Benetollo, C.,
Richard, M., Didier, A., and Mandairon, N. (2016). Topographical representation of odor
hedonics in the olfactory bulb. Nat Neurosci *19*, 876–878.

Kim, J., Lee, S., Fang, Y.-Y., Shin, A., Park, S., Hashikawa, K., Bhat, S., Kim, D., Sohn, J.-W.,
Lin, D., et al. (2019). Rapid, biphasic CRF neuronal responses encode positive and negative
valence. Nat. Neurosci. *22*, 576–585.

Knaden, M., and Hansson, B.S. (2014). Mapping odor valence in the brain of flies and mice.
Curr. Opin. Neurobiol. *24*, 34–38.

Kobayakawa, K., Kobayakawa, R., Matsumoto, H., Oka, Y., Imai, T., Ikawa, M., Okabe, M.,
Ikeda, T., Itohara, S., Kikusui, T., et al. (2007). Innate versus learned odour processing in the
mouse olfactory bulb. Nature *450*, 503–508.

Kudo, Y., Akita, K., Nakamura, T., Ogura, A., Makino, T., Tamagawa, A., Ozaki, K., and
Miyakawa, A. (1992). A single optical fiber fluorometric device for measurement of intracellular
Ca2+ concentration: Its application to hippocampal neurons in vitro and in vivo. Neuroscience
50, 619–625.

Kumar, S., von Kriegstein, K., Friston, K., and Griffiths, T.D. (2012). Features versus Feelings:

Dissociable Representations of the Acoustic Features and Valence of Aversive Sounds. J.
Neurosci. *32*, 14184–14192.

Lebel, D., Grossman, Y., and Barkai, E. (2001). Olfactory learning modifies predisposition for
long-term potentiation and long-term depression induction in the rat piriform (olfactory) cortex.
Cereb. Cortex *11*, 485–489.

Lewin, K. (1935). A Dynamic Theory of Personality (McGraw-Hill Book Company, Inc.).

Li, Q., and Liberles, S.D. (2015). Aversion and attraction through olfaction. Curr. Biol. *25*, R120–
R129.

Li, H., Pullmann, D., Cho, J.Y., Eid, M., and Jhou, T.C. (2019). Generality and opponency of
rostromedial tegmental (RMTg) roles in valence processing. Elife *8*, e41542.

Li, W., Howard, J.D., Parrish, T.B., and Gottfried, J.A. (2008). Aversive learning enhances
perceptual and cortical discrimination of indiscriminable odor cues. Science (80-.). *319*, 1842–
1845.

Lobo, M.K., and Nestler, E.J. (2011). The Striatal Balancing Act in Drug Addiction: Distinct
Roles of Direct and Indirect Pathway Medium Spiny Neurons. Front. Neuroanat. *5*, 41.

Madisen, L., Mao, T., Koch, H., Zhuo, J., Berenyi, A., Fujisawa, S., Hsu, Y.-W.A., Garcia, A.J.,
Gu, X., Zanella, S., et al. (2012). A toolbox of Cre-dependent optogenetic transgenic mice for
light-induced activation and silencing. Nat Neurosci *15*, 793–802.

Mandairon, N., and Linster, C. (2009). Odor perception and olfactory bulb plasticity in adult
mammals. J. Neurophysiol. *101*, 2204–2209.

Miller, S.M., Marcotulli, D., Shen, A., and Zweifel, L.S. (2019). Divergent medial amygdala
projections regulate approach–avoidance conflict behavior. Nat. Neurosci. *22*, 565–575.

Morrison, S.E., and Salzman, C.D. (2009). The Convergence of Information about Rewarding
and Aversive Stimuli in Single Neurons. J. Neurosci. *29*, 11471–11483.

Murata, K., Kanno, M., Ieki, N., Mori, K., and Yamaguchi, M. (2015). Mapping of Learned Odor-
Induced Motivated Behaviors in the Mouse Olfactory Tubercle. J. Neurosci. *35*, 10581–10599.

De Olmos, J.S., and Heimer, L. (1999). The Concepts of the Ventral Striatopallidal System and

Extended Amygdala. Ann. N. Y. Acad. Sci. *877*, 1–32.

Paxinos, G., and Franklin, K. (2000). The Mouse Brain in Stereotaxic Coordinates (San Diego:
Academic Press).

Pessoa, L., and Adolphs, R. (2010). Emotion processing and the amygdala: from a "low road" to
"many roads" of evaluating biological significance. Nat Rev Neurosci *11*, 773–783.

Roesch, M.R., Stalnaker, T.A., and Schoenbaum, G. (2007). Associative encoding in anterior
piriform cortex versus orbitofrontal cortex during odor discrimination and reversal learning.
Cereb Cortex *17*, 643–652.

Ross, J.M., and Fletcher, M.L. (2018). Learning-Dependent and -Independent Enhancement of
Mitral/Tufted Cell Glomerular Odor Responses Following Olfactory Fear Conditioning in Awake
Mice. J. Neurosci.

Schoenbaum, G., and Eichenbaum, H. (1995). Information coding in the rodent prefrontal
cortex. I. Single-neuron activity in orbitofrontal cortex compared with that in pyriform cortex. J
Neurophysiol *74*, 733–750.

Schoenbaum, G., Chiba, A.A., and Gallagher, M. (1998). Orbitofrontal cortex and basolateral
amygdala encode expected outcomes during learning. Nat Neurosci *1*, 155–159.

Schoenbaum, G., Chiba, A.A., and Gallagher, M. (2000). Changes in functional connectivity in
orbitofrontal cortex and basolateral amygdala during learning and reversal training. J Neurosci
20, 5179–5189.

Schoenbaum, G., Setlow, B., Saddoris, M.P., and Gallagher, M. (2003). Encoding predicted
outcome and acquired value in orbitofrontal cortex during cue sampling depends upon input
from basolateral amygdala. Neuron *39*, 855–867.

Soares-Cunha, C., Coimbra, B., Sousa, N., and Rodrigues, A.J. (2016). Reappraising striatal
D1- and D2-neurons in reward and aversion. Neurosci. Biobehav. Rev. *68*, 370–386.

Thompson, K.G., Hanes, D.P., Bichot, N.P., and Schall, J.D. (1996). Perceptual and motor
processing stages identified in the activity of macaque frontal eye field neurons during visual
search. J. Neurophysiol. *76*, 4040–4055.

Tolman, E. (1932). Purposive Behavior (New York: Appleton-Century).

Veit, L., and Nieder, A. (2013). Abstract rule neurons in the endbrain support intelligent
behaviour in corvid songbirds. Nat Commun *4*, 2878.

Veldhuizen, M.G., Rudenga, K.J., and Small, D.M. (2009). The pleasure of taste, flavor, and
food. In Pleasures of the Brain, M.L. Kringelbach, and K.C. Berridge, eds. (Oxford University
Press), pp. 146–168.

Verhagen, J. V, Wesson, D.W., Netoff, T.I., White, J.A., and Wachowiak, M. (2007). Sniffing
controls an adaptive filter of sensory input to the olfactory bulb. Nat Neurosci *10*, 631–639.

Vicente, A.M., Galvão-Ferreira, P., Tecuapetla, F., and Costa, R.M. (2016). Direct and indirect
dorsolateral striatum pathways reinforce different action strategies. Curr. Biol. *26*, R267–R269.

Voorn, P., Jorritsma-Byham, B., Van Dijk, C., and Buijs, R.M. (1986). The dopaminergic
innervation of the ventral striatum in the rat: A light- and electron-microscopical study with
antibodies against dopamine. J. Comp. Neurol. *251*, 84–99.

Wesson, D.W., and Wilson, D.A. (2011). Sniffing out the contributions of the olfactory tubercle to
the sense of smell: hedonics, sensory integration, and more? Neurosci Biobehav Rev *35*, 655–
668.

Wilson, D.A., and Sullivan, R.M. (2011). Cortical processing of odor objects. Neuron *72*, 506–
519.

Wilson, D.A., Sullivan, R.M., and Leon, M. (1987). Single-unit analysis of postnatal olfactory
learning: modified olfactory bulb output response patterns to learned attractive odors. J
Neurosci *7*, 3154–3162.

Zhang, Z., Zhang, H., Wen, P., Zhu, X., Wang, L., Liu, Q., Wang, J., He, X., Wang, H., and Xu,
F. (2017a). Whole-Brain Mapping of the Inputs and Outputs of the Medial Part of the Olfactory
Tubercle. Front. Neural Circuits *11*, 52.

Zhang, Z., Liu, Q., Wen, P., Zhang, J., Rao, X., Zhou, Z., Zhang, H., He, X., Li, J., Zhou, Z., et
al. (2017b). Activation of the dopaminergic pathway from VTA to the medial olfactory tubercle
generates odor-preference and reward. Elife *6*.

Figure Legends

Figure 1. Paradigm for investigating the distribution of odor valence encoding among olfactory cortices. A) Recording schematic indicating simultaneous monitoring of OT and pPCX activity with multi-channel arrays in mice (8 wires/region). pPCX and OT activity were recorded on opposite hemispheres within mice. **B)** Schematic of head-fixed behavioral set-up and **C)** experimental trial outline. Water-deprived mice implanted with chronic multi-site arrays into their OT and pPCX (as in A) and a head-bar were acclimated to head restraint and shaped in a lick/no-lick odor discrimination task wherein following a variable inter-trial interval (ITI) one odor signaled the availability of a small palatable fluid reward from a reward spout following odor offset (during the 'lick reward available' epoch). Alternative (unreinforced) odors were conditioned to not signal reward availability. OT and pPCX unit activity was acquired throughout performance in the task. **D)** Example OT and pPCX unit activity indicating odor valence encoding by neurons in both structures. Multiunit activity (MUA) was spike sorted off-line to identify single-unit activity (SUA) using template matching and cluster cutting with principal component analyses. Vertical lines on the lick trace indicate timing of individual licks (detected with an infrared photo-beam in front of the lick spout). Vertical scale bar applies to both OT and pPCX MUA channels. Data are from two consecutive trials.

Figure 2. OT neurons display greater divergent responses to conditioned odors than pPCX. A) Time course of odor-evoked changes in firing rate relative to baseline as indicated by auROC in response to unrewarded and rewarded odors in the pPCX (left; n=70 neurons) and OT (right; n=104 neurons). Each row represents one single responsive neuron, and neurons are sorted by baseline rates during rewarded odor trials (top row = lowest rate). **B)** Population average auROC values for all responsive neurons in response to rewarded and unrewarded odors compared against a shuffled control distribution. Both regions show an increase in auROC values >0.5 shortly after odor onset, indicating increases in firing relative to baseline. Across the population, OT neurons displayed more divergent responses to conditioned odors than pPCX neurons, with the rewarded odor eliciting larger and more persistent changes to firing rate relative to the unrewarded odor. Population values represent the mean ± SEM.

Figure 3. D1 receptor expressing neurons underlie odor valance coding in the OT. Ai) Example images of GCaMP6f expression in D1- and D2-Cre mice crossed with the rosa Ai9.TdTomato reporter line which were later injected with GCaMP6f. In both Cre lines, GCaMP6f colocalized robustly with the targeted cells (**Aii**). Data points in Aii = means of individual mice (4-

5 sections/mouse). Dashed lines define approximate OT cell layers and its ventral boundary. **B)**
Example localization of an optical fiber implant into a mouse injected with GCaMP6f (green). The
optical fiber was positioned to terminate immediately dorsal to the OT. **C)** Schematic of the fiber
photometry system for use in head-fixed behaving mice. 465nm and 405nm LEDs excitation
wavelengths passed through dichroic mirrors prior to being sent into the implanted optical fiber
via a patch cable. GCaMP6f and UF emission were then both amplified by femtowatt
photoreceivers prior to being digitized simultaneously with behavioral data and stimulus
presentation events. Please see Fig 1 for a description of the head-fixed paradigm. Example
GCaMP6f responses to the original rewarded and unrewarded odors in a single D1-Cre **(D)** and
D2-Cre mouse **(E)** averaged over 10 trials of correct behavioral responses during blocks of
criterion behavioral performance. These example averaged traces indicate divergent responses
to the original rewarded odor vs the original unrewarded odor in D1-Cre but not D2-Cre mice,
which is maintained at the population level when averaging across 5 mice, throughout 2-3
sessions during criterion behavioral performance (**F & G**). The encoding of odor valence in D1-
Cre mice (as in **F**) was confirmed in experiments wherein the odor-reward contingencies were
reversed (original rewarded = no unrewarded, and vice versa) (**H & I**). Insets in **H & I** display the
mean delta f/f during 0.5-0.75sec of odor in for all four stimulus types. ***=p<0.001. ns = not
significant (p>0.05). No significance was found in any planned comparison in the D2-Cre mice (**I**).

Figure 4. **OT neurons expressing the D1 receptor promote reinforcement and approach**
behaviors. (A) Optical fiber implant locations from mice contributing optical-intracranial self-
stimulation data (opto-ICSS), segregated based upon those mice which learned versus failed to
acquire phase 1 of the task. **(B)** Opto-ICSS task schematic wherein mice were required to nose
poke in exchange for blue-light mediated optogenetic excitation of OT D1 neurons. At least 2
weeks following injection with either a floxed AAV encoding ChR2 and a reporter fluorophore
(AAV.ChR2) or a floxed AAV solely encoding a reporter fluorophore (AAV), mice were shaped on
the opto-ICSS task. Please see text for a description of the four task phases. **(C)** Pie chart
indicating that all but one AAV.ChR2 mouse reached criterion performance on phase 1 of the
task. **(Di)** Performance of two example mice during phases 2-4 of the opto-ICSS task. The mice
displayed diversity in their number of pokes into the active port, and in their % of pokes displayed
for the active (blue-light emitting) versus inactive ports. Both of these mice reached criterion on
phase 2 and phase 3 wherein they had to learn to redirect their poking for a new active port
location (reversal learning). Shown also is the behavior of these two mice during extinction,
wherein the port lights were on (*viz.,* both ports were visually 'active') but no optogenetic

stimulation was available regardless of poking. **(Dii)** A win-loss plot for mouse 127 from session
2-5. **(E)** Quantification of opto-ICSS data indicating that ChR2 mediated stimulation of OT D1
neurons promotes reinforcement and approach behaviors. The first phase of reversal learning
resulted in a significant reduction in % pokes for the new active port, which with experience was
restored on the last session of the reversal phase.

Supplementary Figure Legends

Supplementary Figure 1. A) Histological verification of pPCX and OT recording sites. **B)**
Background firing rate distribution of all pPCX and OT single-units used for analysis. **C)** Summary
of learning and performance in the lick/no-lick odor discrimination task for mice contributing single-
unit data. Red dashed horizontal line = criterion behavioral performance level.

Supplementary Figure 2. Time course of odor-evoked changes in firing rate separated by
response type. Population average auROC values for all responsive neurons in pPCX and OT
(same neurons as in Fig. 2), separated by whether the response was excited or suppressed
relative to baseline. Population values represent the mean ± SEM.

Supplementary Figure 3. A) Histological verification of optical fiber implants from all mice
contributing fiber photometry data. Summary of learning and performance in the lick/no-lick odor
discrimination task for D1-Cre **(B)** and D2-Cre mice **(C)** contributing fiber photometry data. Red
dashed horizontal line = criterion behavioral performance level.

Methods

Animals. Eleven 2-4 months of age male C57bl/6 mice originating from Harlan Laboratories (Indianapolis, IN) were used for single-unit recordings. For fiber photometry, we utilized male bacterial artificial chromosome transgenic mice expressing Cre-recombinase under control of the *drd1a* (D1-Cre; *n* = 7; obtained from the UC Davis Mutant Mouse Regional Resource Center, strain EY262Gsat/Mmucd) or *drd2* gene (D2-Cre; *n* = 7; obtained from the UC Davis Mutant Mouse Regional Resource Center, strain ER44Gsat)(Gong et al., 2003). The offspring of homozygous tdTomato Ai9 reporter mice (Madisen et al., 2012), crossed with D1- or D2-Cre heterozygotes were used to confirm specificity of GCaMP expression. Genotyping was performed following standard protocols using tail tissue DNA. Mice were housed on a 12:12 h light-dark cycle with food and water available *ad libitum*, except when water was restricted for behavioral training (see below). Up to 5 mice were co-housed in a cage before experimentation, but all post-surgical animals were housed individually. All experimental procedures were conducted in accordance with the guidelines of the National Institutes of Health and were approved by the Case Western Reserve University and University of Florida Institutional Animal Care and Use Committees.

Methods for multi-site single-unit recordings during lick/no-lick odor discriminations.
Surgical procedures were conducted as described previously (Gadziola et al., 2015; Gadziola & Wesson, 2016). Briefly, mice were anesthetized with Isoflurane (2-4% in oxygen, Abbott Laboratories, Green Oaks, IL), and mounted in a stereotaxic frame with a water-filled heating pad (38°C) beneath to maintain body temperature. An injection of a local anesthetic (0.05% marcaine, 0.1 ml s.c.) was administered before exposing the dorsal skull. A craniotomy was made to access the OT and pPCX, each contralateral from another (see **Fig S1A** for distribution). An 8-channel micro-wire electrode array (PFA-insulated tungsten wire, with four electrode wires encased together in a 254 μm diameter polyimide tube (Gadziola et al., 2015)) was implanted within the OT (4.9 mm ventral) and posterior PCX (4.2 mm ventral) (**Fig 1A and S1A**), and cemented in place, along with a headbar for later head fixation. The hemisphere receiving an array into each structure was not held constant across animals but instead varied (**Fig S1A**). Additional craniotomies were drilled over a single neocortex for placement of a stainless steel ground wire. During a three-day recovery period, animals received a daily injection of carprofen (5 mg/kg, s.c., Pfizer Animal Health, New York, NY) or meloxicam (5mg/kg; Putney, Inc., Portland, ME) and *ad libitum* access to food and water. Animals were allowed 5-7 days for post-op recovery before beginning water restriction.

Olfactory lick/no-lick odor discrimination and reversal learning behavioral task.

Mice were mildly water-restricted for three days prior to behavioral training. Bodyweight was monitored daily and maintained at 85% of their original weight. Mice were trained in cohorts of three to four. All behavioral procedures were performed during the light hours, in a dim room. Head-fixed mice were trained in a lick/no-lick odor discrimination task (**Fig 1C**) across multiple 1 hour recording sessions in which the mice obtained a fluid reward for licking a spout positioned in front of their snouts in trials with reinforced odors (Gadziola et al., 2015), adapted from (Verhagen et al., 2007).

Licking was measured by a pair of infrared photo-beams positioned to cross in front of the lick spout by ~2 mm. Mice were first trained to lick the water spout for reward, with a progressively increasing inter-trial interval (ITI) (3 ± 1 s, Phase 1). In Phase 2, odor presentation began with a 10 ± 2 s ITI, and mice were only rewarded for licking during a 2 s period after odor offset (FR1). Licking during the odor was discouraged by moving the lick spout further away from the snout. In Phase 3, trials were randomized between rewarded and unrewarded trials (17 ± 2 sec ITI); for unrewarded odor trials, mice were presented with a "blank" stimulus (mineral oil) and had to learn to withhold licking during these trials. Finally, in Phase 4, trials (17 ± 2 s ITI) were randomized between the rewarded and unrewarded odors. In go trials, mice would receive a reward for licking a spout within 2sec following the offset of the rewarded odor (hit); not licking would be considered a miss. In no-go trials, mice were presented with an unreinforced odor and did not receive a water reward regardless of whether they licked (false alarm) or correctly withheld licking for the total odor duration of the trial (correct reject). Next, mice continued on Phase 5 to learn a novel odor pair. Phase 5 consisted of pseudo-random trials (17 ± 2 s ITI) among which all four odors were separately presented, with the two reinforced odors both resulting in a low-value reward. Finally, in Phase 6, the four odors were presented (17 ± 2 s ITI), with one rewarded odor being assign to the high value saccharine and the other to the low value saccharine. Mice discriminated among the four odors, two of which predicted presentation of reward (either low or high value saccharine), and two unrewarded odors. These sessions include the original training odor set, as well as novel odors presented on a different experimental day (each mouse was shaped on 2 novel sets of four odors on different sessions). No major effects were observed between high versus low value rewarded odor evoked responses nor behavior and therefore herein any odors paired with reinforcers are classified as conditioned rewarded odors, whereas those not paired are termed conditioned unrewarded odors. Throughout all phases, behavioral performance was evaluated in

blocks of 20 trials, and mice were required to achieve a performance criterion ≥80% correct for
two consecutive blocks in order to advance to the next phase. Neural activity was recorded
throughout all training phases, but analysis of odor-evoked activity was restricted to post-training
sessions.

Stimulus Delivery.
Odors were presented through a custom air-dilution olfactometer with independent stimulus lines
up to the point of entry into the odor port. In addition to a blank stimulus (mineral oil), odors
included ethyl butyrate, 1,7-octadiene, isopentyl acetate, heptanal, 2-heptanone, (+)-limonene,
ethyl propionate, (-)-limonene, methanol, methyl valerate, 2-butanone, 1,4-cineole, butanal,
propyl acetate, allylbenzene, allyl bromide, isobutyl propionate, and 2-methylbutyraldehyde
(Sigma Aldrich, St. Louis, MO; all >97% purity). These molecularly diverse odors were diluted in
their liquid state to 1 Torr (133.32 Pa) in mineral oil and were then further diluted to 10% (vol/vol)
by mixing 100 ml odor vaporized N_2 with 900 ml medical grade N_2 (Airgas, Radnor, PA). Thus,
stimuli were delivered at a total flow rate of 1 L/min. Not all animals were tested with all odors.
The rewarded and unrewarded odor pairs were pseudo-randomly assigned to each cohort prior
to training (neurons were not initially screened for odor responsiveness). The experimenter was
not blind to odor assignment, but all stimulus presentation was automated. Presentation of
rewarded and unrewarded odors were pseudo-randomized within each block, delivered for 2 s
duration with a 17 ± 2 s ITI through a Teflon odor-port (9 mm diameter opening) directed towards
the animal's snout at a distance of 1 cm. Odor was continuously flowing to the odor-port but was
removed by a vacuum before exiting towards the animal. Recordings with a photoionization
detector (miniPID, Aurora Scientific, Ontario, Canada) were used to confirm the temporal
dynamics of the odor presentation in this design. While the dynamics may vary slightly across
odors, they confirm the precision and stability of the odor presentation methods used in this study.

Reward Delivery.
Reward fluids were delivered through a custom 3D-printed polylactic acid lick spout, as reported
previously (Gadziola and Wesson, 2016). Independent stimulus lines terminated onto 20G
blunted needles that passed through one of seven 1-mm holes and extended to the tip of the
spout. In the current task, two adjacent holes on the lick spout were used for reward delivery,
three were connected to a vacuum line, and the last two unused holes were blocked. Animals
were reinforced with a 4µL drop of 2-20mM saccharin (Sigma Aldrich, St. Louis, MO; dissolved in
water). Reward volumes were calibrated for the individual reward valves.

In vivo Electrophysiology.

The outputs of the electrode arrays were amplified, digitized at 24.4 kHz, filtered (bandpass 300-5000 Hz), and monitored (Tucker-Davis Technologies, Alachua, FL), along with licking (300 Hz sampling rate), and odor and reward presentation events. One electrode wire was selected to serve as a local reference for each headstage. Our electrode arrays were fixed in place and no attempt was made to record from unique populations of neurons on different sessions. To compensate for the possibility that the same neurons were recorded across multiple days, two different behavioral tasks were employed and statistical comparisons are only made within each task type. Sessions of the same task and odor pair were run for 1-3 consecutive days to achieve adequate behavioral performance and/or capture the dynamics of newly identified neurons. On average, 3.5 ± 1.9 single neurons were identified per mouse per session (range: 1-7 neurons), with an average of 1.4 ± 0.6 neurons recorded per viable electrode wire per mouse per session (range: 1-3 neurons). Numbers were comparable when separated by region (± 0.1 single units / region difference).

Analysis of odor-evoked electrophysiology data.
Single neurons were sorted offline in Spike2 (Cambridge Electronic Design, Cambridge, England), using a combination of template matching and cluster cutting based on principle component analysis. Single neurons were further defined as having <2% of the spikes occurring within a refractory period of 2 ms. Spike times associated with each trial were extracted and exported to MATLAB (Mathworks, Natick, MA) for further analysis. To examine modulations in firing rate within a single trial, spike density functions were calculated by convolving spike trains with a function resembling a postsynaptic potential (Thompson et al., 1996). Mean firing rates across trials were measured in 50 ms bins, along with the 95% confidence interval. Mean baseline firing rate for each neuron was averaged across a 2 s period prior to odor onset.

Animals were typically presented with at least 40 trials of each rewarded and unrewarded odor set per session. To ensure that animals were engaged in the task, only blocks possessing high behavioral performance were analyzed. On some trials, mice may have been licking during odor presentation. Any trials in which the animal licked during the first 1.5s period after odor onset were removed. This ensured that any odor-evoked activity observed during the first several hundred ms was not due to licking-related activity. Of the 11 animals with bilateral implants, 10

792 were confirmed to reside within the OT and 8 within the pPCX. One of the 10 OT-confirmed
793 animals did not contribute data since they did not yield well-isolated neurons. From all mice
794 contributing data, 7 yielded simultaneous OT and pPCX single-units.
795
ROC Analysis.
The area under the receiver operating characteristic (auROC) is a nonparametric measure of the
discriminability of two distributions (Green and Swets, 1966). To normalize activity across
neurons, we used an auROC method that quantifies stimulus-related changes in firing rate to the
baseline activity on a 0-1 scale (for more details see (Cohen et al., 2012)). A value of 0.5 indicates
completely overlapping distributions, whereas values of 0 or 1 signal perfect discriminability. We
calculated the auROC at each 50-ms time bin over a 4 s period centered on stimulus onset for
each neuron. Values greater than 0.5 indicate the probability that firing rates were increased
relative to the pre-stimulus background (excitation), whereas values less than 0.5 indicate the
probability that firing rates were decreased relative to the pre-stimulus background (inhibition).
Similar trial numbers have been used for calculating auROC (Veit and Nieder, 2013; Gadziola et
al., 2015, Gadziola & Wesson, 2016). To obtain mean auROC values, the auROC values of
individual neurons were averaged at each time bin. In some cases mean auROC values were
computed separately for all excitatory and inhibitory neurons.
810

Methods for fiber photometric-based imaging of D1- and D2-type dopamine receptor expressing
OT neurons during lick/no-lick odor discrimination and reversal learning.

Virus injection and optical fiber Implantation. To achieve cell-type specific GCaMP6f (Chen et al.,
2013) expression, D1- and D2-Cre mice were injected with AAV5-EF1a-DIO-GCaMP6f-WPRE
(5.43^{e13} GC/ml, obtained from the Penn Vector Core in the Gene Therapy Program of the Univ of
Pennsylvania). Stereotaxic surgery was performed as described above with exceptions as noted
below. 1µl of virus was infused into the OT at a rate of 0.1nl/sec and allowed to diffuse for 10mins
before needle was slowly withdrawn over the subsequent 20mins. This injection paradigm was
not intended to infect solely OT neurons, but to yield a bolus of infected cells which could be later
targeted selectively for imaging by careful localization of the optical fiber. Following, the
craniotomy was sealed with wax and the wound margin closed. Animals were allowed 2 weeks
for the virus to transduce and then a 400µm core, 0.48NA optical fiber, threaded through metal
ferrules were chronically implanted to terminate within the OT. Implants were cemented in place

along with a headbar for later head fixation. As with the single-unit recordings, animals were

allowed 5-7 days for post-op recovery before beginning water restriction. D1- and D2-Cre mice

were tested in mixed cohorts.

Imaging. 465nm (GCaMP excitation wavelength, driven at 210 Hz) and 405nm (UV excitation

wavelength, driven at 330Hz; control channel) light emitting diodes were coupled to a 5 port

fluorescence mini cube (FMC5, Doric Lenses) using 400μm core, 0.48NA, 2.5mm FCM optical

fiber patchcords. Excitation and emission light were directed through a single optical fiber

patchcord connected to the animal via a 2.5mm metal ferrule. Emission light was directed through

the 5-port mini cube, and subsequently coupled to a pair of femtowatt photoreceivers (model

2151, Newport) for monitoring of the GFP and UV fluorescence. Photometry data were digitized

and pulse demodulated at 1kHz, along with the timing of behavioral events (licking) and odor and

reward delivery using a Tucker Davis Technologies digital processor (TDT). Parameters used for

imaging and acquisition of photometry data were held constant across all mice.

Lick/no-lick odor discrimination and reversal learning during photometry. The methods for

operant olfactory behavior employed for fiber photometry were mostly identical to those used

for the single-unit recordings (see above) with two exceptions. First, during the fiber photometry

imaging, mice were only asked to discriminate between two odors in any given session (one

conditioned rewarded and one conditioned unrewarded). Second, mice were shaped in a

reversal learning paradigm wherein following reaching successful criterion performance in the

original odor discrimination, the contingencies were reversed so that the original rewarded odor

no longer signals reinforcement availability (now the 'new unrewarded odor') and the original

unrewarded odor now signals reward availability (now the 'new rewarded odor').

Processing and analysis of photometry data. Off-line, the UV signal was subtracted from that of

the GFP, filtered (2nd order, 25Hz low-pass), smoothed, and down sampled to 200Hz using a

custom script written in Spike2 (Cambridge Electronic Design, Inc). Photometry recordings were

normalized across trials off-line, using the mean signal across 3 sec before odor delivery as a trial

specific baseline for the $\Delta F/F$ values. Of the 7 D1-Cre animals with bilateral implants, 1 was

confirmed to have poor/weak viral expression and one have a fiber tip residing outside of the OT.

Of the 7 D2-Cre mice, 2 were confirmed to have fiber tips residing outside of the OT. Thus, in

total, 5 mice per genotype, which had 1) criterion-level behavioral performance, 2) on target OT

fiber tips, and 3) quality viral expression contributed data.

Methods for optogenetic stimulation of OT D1 expressing neurons and intra-cranial self-stimulation.

The surgical methods to allow for expression of ChR2 in OT D1-expressing neurons are similar to those used for GCaMP6f as described above. Two surgical procedures were performed on each mouse. First, mice were intracranially injected with one of two floxed AAV vectors to allow for expression of AAV5.EF1a.DIO.hChR2(H134R).eYFP or mCherry (to yield optogenetic stimulation) or AAV5.EF1a.DIO.eYFP or mCherry (as control). AAVs were obtained from the University of North Carolina viral vector core. Second, mice were intracranially implanted with a unilateral optic fiber (300um, 0.39NA) into their OT which was attached to a ferrule to allow light transmittance into the brain. AAV injections and ferrule implants occurred in the left hemisphere or right hemisphere to prevent biasing results towards effects which may be lateralized. Using a light meter (Thorlabs, Inc., Newton, New Jersey), all fiber optic implants were verified prior to implantation to deliver 6.5-7.5mW[3] of light energy when using the same LED and LED driver as used during Opto-ICSS testing (see below).

Opto-ICSS apparatus and light stimulation. We used a dual nose-poke port apparatus wherein poke into each port is monitored by 880nm infrared photobeams and digitized for later analysis. The floor and three walls of the apparatus were made of black acrylonitrile butadiene styrene (ABS) plastic (30cm tall walls), consisting of a 15 x 15cm square floor and a guillotine style ABS door to allow gentle insertion and removal of the mouse from the testing apparatus. One wall of the apparatus was constructed of clear acrylic to allow visual monitoring of the mouse by an experimenter when needed. The ceiling was 'open' to allow for tethering of the animal to an optical rotary joint (Doric Lenses, Inc., Quebec, Canada) which was suspended above the apparatus. The nose-poke ports were 3D-printed out of polylactic acid plastic and were T-shaped with a 2cm opening which faced into the behavioral apparatus to allow nose-entry and monitoring of infrared beam breaks. Photobeam status (open or closed) was acquired continuously and relayed to a PC for data collection using an Arduino microcontroller board running custom code written in the Arduino open source language (Arduino, https://www.arduino.cc). Immediately next to each port was a 3mm white LED, also controlled by the Arduino, which could signal to the mouse active port status (on = active and optical stimulation available, off = inactive and no available optical stimulation).

Optogenetic stimulation (25Hz train, 15ms pulse width, 2 sec duration [for opto-ICSS the 2

sec stimulation occurred regardless of the nose-poke duration]) was accomplished by means of

a 447.5nm LED (Luxeon Rebel ES, Luxeon Stars, Lethbridge, Alberta) driven by a Thorlabs T-

Cube LED driver (Thorlabs) and controlled by the digital output of an Arduino microcontroller. This

stimulation paradigm was selected following the results of physiological and behavioral pilot

studies. An SMA adapter was adhered atop of the LED using optical epoxy to allow for butt

coupling of an SMA terminated multimode fiber to the LED. The fiber used to connect the LED to

the optic rotary joint as well as the fiber used to connect the rotary joint to the mouse were both

300μm core multimode fibers with a 0.39NA (same as the fiber used for the optical implants),

encased in 2mm furcation tubing.

Opto-ICSS shaping and testing. All behavioral procedures were performed during the light hours,

in a dim room. At the start of all sessions, the mice were first gently connected to the fiber optic

tether. The mice were then placed into the apparatus and the door closed. The stages of shaping

and testing in the opto-ICSS task are illustrated in **Figure 4** and occurred on separate daily 60min

sessions. In phase 1, both ports were active (both active port LEDs were on) and nose-poke into

either port triggered blue light stimulation. After performing ≥ 40 pokes into either or both ports in

a given session, for at least two consecutive sessions, mice were then transitioned into phase 2

of the task. In phase 2, the port wherein the mouse displayed the greatest number of pokes on

subsequent days, was set to be active (port LED on, blue light stimulation available) and the

alternative port was inactive (port LED off, blue light stimulation unavailable). Only nose poke into

the active port triggered light stimulation. After mice reached criterion performance in phase 2

(≥40 pokes, with ≥80% into the active port), they were switched to phase 3 wherein the active and

inactive ports were reversed so that the previously inactive port is now the only port which would

trigger light stimulation. Phase 3 is thus a test of reversal learning and ensures that the animal's

tendency to poke is indeed motivated by light-mediated ChR2 stimulation. Those mice which

reached criterion performance on phase 3 (see above), were advanced onto phase 4. In all of the

previous phases stimulation was available on an FR1 (fixed ratio of 1) schedule. In phase 4, both

ports were rendered inactive (both port LEDs were off, blue light off) in a test of extinction.

Transitions between phases occurred on subsequent daily testing sessions. Mice which did not

reach criterion performance on a given phase by the end of the 6[th] session of that phase were

eliminated from further testing.

Histology

Following the end of experiments, mice were overdosed with Fatal-plus (0.01mL/g; Vortech
Pharmaceutical, Dearborn, MI) and perfused with 10mL of cold saline followed by 15mL of cold
10% phosphate buffered formalin. Brains were stored in 10% formalin/30% sucrose (4°C) prior to
sectioning at 40μm thickness on a Leica sliding microtome. Tissue was later mounted on slides
using Fluoromount-G containing DAPI (for fluorescence analysis; 4',6-diamidino-2-phenylindole;
Invitrogen, Carlsbad, CA) or stained with 1% cresyl-violet (for tissue contributing single-unit data).
Microscopic examination of tissue was conducted on a Nikon Ti2e inverted fluorescent
microscope equipped with 5MP color and 15MP monochrome cameras. Histological definition of
brain regions was aided by use of a brain atlas (Paxinos and Franklin, 2000). Quantification of
fluorophore colocalization (*i.e.,* as in the case of D1-Cre;Ai9 and D2-Cre;Ai9 GCaMP6f injections)
was performed using semi-automated methods in NIS Elements (Nikon). Briefly, background
fluorescence was decreased to enhance contrast, cells were detected based upon their
fluorescence intensity (via a threshold) and their size, and detected objects next post-processed
based upon their area to eliminate fluorescent objects which were not cell bodies (*e.g.,* brightly
labeled fibers). Counts of tdTomato (Ai9), GFP (GCaMP), and colocalized OT cells (including all
OT cell layers) were next automatically exported following manual setting of OT boundaries.

Overall statistical methods
All performed tests were two-sided and met assumptions of normality (Kolmogorov-Smirnov test).
Sample sizes are consistent with numbers reported in the field and no statistical method was used
to predetermine these. Statistical analyses were performed in SPSS 22.0 (SPSS Inc., Chicago,
IL) or MATLAB. All data are reported as mean ± SD unless otherwise noted. Specific details
regarding some analytical methods (*e.g.,* ROC) can be found in above sections.
945

Figure 1

Figure 2

Figure 3

Figure 4

A

+2 mm Bregma

OT

+0.5 mm Bregma

● learned phase 1
● failed phase 1

B

AAV injection & implant optical fiber

phase 1 — **phase 2** — **phase 3** (reversal) — **phase 4** (extinction)

○ **Inactive port** (LED off)
● **Active port** (LED on + intracranial blue light delivery available)

C

■ learned
□ failed

1/7
6/7

Di

mouse 124

phase 2 | phase 3 | phase 4

Poke # / % pokes for active port

session # 1 2 3 4 5 6 7 8 9 10 11 12

mouse 127

phase 2 | phase 3 | phase 4

Poke # / % pokes for active port

session # 1 2 3 4 5 6 7

■ active poke
■ inactive poke

Dii *session #*

reversal →

session	total # pokes
2	437
3	244
4	514
5	378

E

% pokes for active port ± SEM

n=6 mice

*** phase 2 (last session)
reversal (first session)
*** reversal (last session)

Figure S1

A

pPCX

OT

-0.5 mm Bregma

+2 mm Bregma

pPCX

OT

-2 mm Bregma

+0.5 mm Bregma

right hemisphere
left hemisphere

C

Phase 1   Phase 2   Phase 3   Phase 4

% Correct

relative block within training phase

number of sessions

Phase

B

pPCX
n=109 units

units

OT
n=98 units

units

Background firing Hz (2-sec pre odor)

Figure S2

rewarded (odor suppressed)
rewarded (odor excited)
unrewarded (odor suppressed)
unrewarded (odor excited)
shuffled control

Figure S3

A

+2 mm Bregma

OT

D1-Cre
D2-Cre

+0.5 mm Bregma

B D1- Cre

Phase 1 Phase 2 Phase 3 Phase 4

% Correct

1st   last 1st   last 1st   last 1st   last

Relative Block Within Training Phase

number of sessions

1 2 3 4
Phase

C D2- Cre

Phase 1 Phase 2 Phase 3 Phase 4

% Correct

1st   last 1st   last 1st   last 1st   last

Relative Block Within Training Phase

number of sessions

1 2 3 4
Phase