A developmental framework linking neurogenesis and circuit formation in the Drosophila CNS

Brandon Mark[1], Sen-Lin Lai[1], Aref Arzan Zarin[1,2], Laurina Manning[1], Ashok Litwin-Kumar[3], Albert Cardona[4], James W. Truman[4,5], and Chris Q. Doe[1]*

[1]Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon, Eugene, OR 97403
[2]Current address: Institute of Neuroscience, Texas A&M University, College Station, TX 77843
[3]Mortimer B Zuckerman Mind Brain Behavior Institute, Department of Neuroscience, Columbia University, New York, NY
[4]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA 20147
[5]Friday Harbor Laboratories, University of Washington. Friday Harbor, WA 98250

* Author for correspondence at cdoe@uoregon.edu

Key words: neuroblast, hemilineage, temporal identity, pathfinding, cell lineage, neural circuits

Abstract

The mechanisms specifying neuronal diversity are well-characterized, yet it remains unclear how or if these mechanisms regulate neural circuit assembly. To address this, we mapped the developmental origin of 160 interneurons from seven bilateral neural progenitors (neuroblasts), and identify them in a synapse-scale TEM reconstruction of the *Drosophila* larval CNS. We find that lineages concurrently build the sensory and motor neuropils by generating sensory and motor hemilineages in a Notch-dependent manner. Neurons in a hemilineage share common synaptic targeting within the neuropil, which is further refined based on neuronal temporal identity. Connectome analysis shows that hemilineage-temporal cohorts share common connectivity. Finally, we show that proximity alone cannot explain the observed connectivity structure, suggesting hemilineage/temporal identity confers an added layer of specificity. Thus, we demonstrate that the mechanisms specifying neuronal diversity also govern circuit formation and function, and that these principles are broadly applicable throughout the nervous system.

Introduction

Tremendous progress has been made in understanding the molecular mechanisms generating neuronal diversity in both vertebrate and invertebrate model systems. In mammals, spatial cues generate distinct pools of progenitors which generate a diversity of neurons and glia appropriate for each spatial domain (Sagner and Briscoe, 2019). The same process occurs in invertebrates like *Drosophila*, but with a smaller number of cells, and this process is particularly well-understood. The first step occurs when spatial patterning genes act combinatorially to establish single, unique progenitor (neuroblast) identities (Skeath and Thor, 2003). These patterning genes endow each neuroblast with a unique spatial identity (Figure 1A, left).

The second step is temporal patterning – the specification of neuronal identity based on birth-order – an evolutionarily-conserved mechanism for generating neuronal diversity (Kohwi and Doe, 2013; Rossi et al., 2017). Here we focus on *Drosophila* embryonic neuroblasts, which undergo a cascade of temporal transcription factors: Hunchback (Hb), Krüppel (Kr), Pdm, and Castor (Cas) (Isshiki et al., 2001). Each temporal transcription factor is inherited by ganglion mother cells (GMCs) born during each expression window. The combination of spatial and temporal factors endows each GMC with a unique identity (Figure 1A, middle).

The third step is hemilineage specification, which was initially characterized in *Drosophila* larval and adult neurogenesis (Lee et al., 2020; Truman et al., 2010), and may also be used in vertebrate neurogenesis (Peng et al., 2007). Hemilineages are formed by GMC asymmetric division into a pair of post-mitotic neurons; during this division, the Notch inhibitor Numb (Nb) is partitioned into one neuron (Notch[OFF] neuron) whereas the other sibling neuron receives active Notch signaling (Notch[ON] neuron), thereby establishing Notch[ON] and Notch[OFF] hemilineages (Figure 1A, right). In summary, three mechanisms generate neuronal diversity within the embryonic central nervous system (CNS): neuroblast spatial identity, GMC temporal identity, and neuronal hemilineage identity.

A great deal of progress has also been made in understanding neural circuit formation in both vertebrates and invertebrate model systems, revealing a multi-step mechanism. Neurons initially target their axons to broad regions (e.g. thalamus/cortex), followed by targeting to a neuropil domain (glomeruli/layer), and finally forming highly specific synapses within the targeted domain (Kolodkin and Tessier-Lavigne, 2011).

Despite the progress in understanding the generation of neuronal diversity and the mechanisms governing axon guidance and neuropil targeting, how these two developmental processes are coordinated remains mostly unknown. While it is accepted that the identity of a neuron is linked to its connectivity, the developmental mechanisms involved remain unclear. For example, do clonally-related neurons target similar regions of the neuropil due to the expression of similar guidance cues? Do temporal cohorts born at similar times show preferential connectivity? Here we address the question of whether any of the three developmental mechanisms (spatial, temporal, hemilineage identity) are correlated with any of the three circuit-wiring mechanisms (neurite targeting, synapse localization, connectivity). We map the developmental origin for 80 bilateral pairs of interneurons in abdominal segment 1 (A1) by identifying and reconstructing these neurons within a full CNS TEM volume (Ohyama et al., 2015); this is over a quarter of the ~300 neurons per hemisegment. We make the unexpected observation that hemilineage identity determines neuronal projection to sensory or motor neuropils; thus, neuroblast lineages coordinately produce sensory and motor circuitry. In addition, we show that neurons with shared hemilineage-temporal identity target pre- and postsynapse localization within the neuropil, and that hemilineage-temporal cohorts have greater shared connectivity than that produced by neuropil proximity alone. Thus, temporal and hemilineage identity play essential roles in establishing neuronal connectivity.

Results

Mapping neuronal developmental origin in a TEM reconstruction

To relate developmental mechanisms to circuit establishment mechanisms, we first needed to identify the developmental origin of neurons within a TEM reconstruction of the larval CNS (Ohyama et al., 2015), allowing us to quantify neuronal projections, synapse localization, and connectivity. We took three

approaches. First, we generated GFP-marked clones for seven different neuroblasts, representing different
spatial positions in the neuroblast array (Figure 1A, left). We imaged each clone by light microscopy in newly-
hatched larvae – the same stage used for the TEM reconstruction – so we could match clonal morphology at
the light and TEM levels (Figure 1B,C). All assayed neuroblast clones had a reproducible clonal morphology
including the number of fascicles entering the neuropil, cell body position, and axon/dendrite morphology
(Figure 1B; and data not shown). We determined the identity of the neuroblast generating for each clone by
comparing clonal morphology to embryonic DiI single neuroblast clones (Bossing et al., 1996; Schmid et al.,
1999; Schmidt et al., 1997), larval neuroblast clones (Birkholz et al., 2015; Lacin and Truman, 2016), and cell
body position.
To identify each of the seven genetically labeled neuroblast clones in the TEM volume, we matched
lineage-specific features present in both light and TEM analyses. We identified neurons that had clustered cell
bodies, clone morphology matching that seen by light microscopy (Figure 1C), and one or two fascicles
entering the neuropil (Figure 1C,D). The similarity in overall clone morphology between genetically marked
clones and TEM reconstructed clones was striking (compare Figures 1B,C). We validated these assignments
using two methods. First, we used neuroblast-specific Gal4 lines (Lacin and Truman, 2016; Seroka and Doe,
2019) and multicolor flip out (MCFO) (Nern et al., 2015) to label individual neurons within each lineage, and
found that in each case we could match the morphology of an MCFO-labeled single neuron from a known
neuroblast to an identical single neuron in the same neuroblast clone within the TEM reconstruction (data
not shown). Second, we reconstructed the same seven lineages in a second hemisegment, A1R. We observed
similar neuron numbers, similar fascicles per clone, and similar clonal morphology (Figure 1E; data not
shown). Thus, neuroblast lineages are highly stereotyped in left and right hemisegments.
We found that, while lineages have stereotyped and distinct morphology, they all had broad projections
within the neuropil. We mapped the distribution of pre- and postsynapses for each neuroblast clone and
found that, consistent with neuronal morphology, synapses were distributed widely across the neuropil
(Figure 1F). We conclude that clonally-related neurons project widely, and localize synapses widely (Figure
1G,H).
In total, we have mapped 14 neuroblast clones in the TEM dataset (seven in A1L, seven in A1R). These
lineages contain 160 interneurons (80 each in A1l and A1r), containing 7794 presynapses and 19468
postsynapses. We also include the previously traced sensory afferents and dendrites from all 32 motor
neurons (Jovanic et al., 2016; Ohyama et al., 2015; Zarin et al., 2019). We note that some of the earliest born
neurons are not included either because their cell bodies are in contact with the neuropil and they do not
fasciculate with clonal fascicles, precluding assignment to a specific neuroblast lineage, or they do not
maintain marker expression at larval hatching. The morphology and function of the earliest born neurons will
be described elsewhere. Nevertheless, this dataset is sufficiently comprehensive for mapping developmental
mechanisms to circuit assembly mechanisms.
In the sections below we first analyze the relationship of developmental mechanisms to neuronal
projections and synapse localization within the neuropil; we conclude by exploring the relationship between
developmental mechanisms and neuronal connectivity.
125
Individual lineages generate two distinct morphological classes of neurons
Perhaps the most important neuropil subdivision in both mammalian spinal cord and *Drosophila* ventral
nerve cord is the segregation of motor and sensory processing domains. In *Drosophila*, motor dendrites target
a dorsal region of the neuropil, while sensory neurons target a ventral region of the neuropil (Landgraf et al.,
2003; Mauss et al., 2009; Zarin et al., 2019; Zlatic et al., 2009). We extend these findings to show that
premotor neurons also target pre- and postsynapses to the dorsal neuropil, and targets of sensory afferents
target pre- and postsynapses to the ventral neuropil (Figure 2A-C). Thus, the neuropil has an important
functional subdivision along the dorsoventral axis.
Upon examination of each lineage, we found that nearly all neuroblasts made two morphologically
distinct types of neurons (except NB2-1; Figure 2–figure supplement 1). We used NBLAST (Costa et al.,
2016) to quantify the morphological similarity of neurons within each neuroblast lineage, and found that most
lineages made two distinct morphological classes of neurons (or one class of neurons plus glia). Strikingly,
each neuronal class projected to the dorsal/motor neuropil or the ventral/sensory neuropil (Figure 2D-H). In

the case of NB3-3, where only a single class of neurons was observed, all neurons were ventral-projecting
(Figure 2I). These classes were more distinct from one another than from neurons in other lineages (Figure
2–figure supplement 2). We conclude that most lineages do not generate neurons that share common
projections, but rather generate two distinct classes of neurons that target either the motor or sensory
neuropil.
144
145 **Hemilineages produce sensory and motor processing neurons via a Notch-dependent mechanism**
146
Recent work has shown that within the larval and adult CNS, most neuroblast lineages generate Notch^{ON}
neurons with a similar clonal morphology (called the Notch^{ON} hemilineage), and Notch^{OFF} hemilineage with a
different morphology (Harris et al., 2015; Lacin and Truman, 2016; Lee et al., 2020; Truman et al., 2010). We
hypothesized that the observed morphological differences within our seven embryonic neuroblast lineages
may be due to hemilineage identity. This hypothesis was strengthened by the fact that the sole neuroblast
lineage that generated a single morphological class, NB3-3, had previously been shown to make only one
"Notch^{OFF}" hemilineage via at type 0 lineage mode (Baumgardt et al., 2014; Wreden et al., 2017). Based on
these data, we hypothesize that each neuroblast makes a Notch^{ON} hemilineage projecting to the dorsal motor
neuropil (or glia), and a Notch^{OFF} hemilineage projecting to the ventral sensory neuropil (Figure 2G).
To test this hypothesis, we took two approaches: first, we generated a lineage-specific reporter for
Notch^{ON} neurons to see if Notch^{ON} neurons in specific neuroblast linages always project to the dorsal
neuropil; and second, we tested whether misexpression of constitutively active Notch can redirect ventral
projections into the dorsal neuropil. To generate a neuroblast lineage-specific Notch^{ON} reporter, we CRISPR
engineered the Notch target gene *hey*, placing a *T2A:FLP* exon in frame with the terminal *hey* exon at the
endogenous *hey* locus. Combining this fly with a neuroblast-specific Gal4 line (*NB5-2-gal4* or *NB7-1-gal4*) plus
a FLP-dependent reporter (*UAS-FRT-stop-FRT-myr:sfGdP:HA*) resulted in myr:GdP:HA expression
specifically in Notch^{ON} neurons within a neuroblast lineage. This resulted in specific labeling of dorsal
projecting neurons within the NB5-2 lineage (Figure 3A,B) or the NB7-1 lineage (Figure 3C,D). We conclude
that Notch^{ON} neurons specifically project to the dorsal motor neuropil.
We next asked whether Notch activity determines dorsal/ventral neuropil projections. We used Gal4
lines specifically expressed in single neuroblast lineages (NB1-2, NB5-2, NB7-1, or NB7-4)(Lacin and
Truman, 2016; Seroka and Doe, 2019) to misexpress a constitutively active form of Notch (Notch^{intra}) in
individual neuroblast lineages. Notch expression throughout the NB1-2, NB5-2, or NB7-1 lineages led to a
loss of ventral projections and a concomitant increase in dorsal projections (Figure 3E-I). Notch expression
throughout the NB7-4 lineage led to a loss of ventral projecting neurons and an increase in glia (Figure 3E-I).
In addition, we note that ascending and descending projection neurons are normally generated by ventral
Notch^{OFF} hemilineages (Figure 3–figure supplement 1). Importantly, these ascending and descending
projection neurons were completely lost following Notch^{intra} expression (Figure 3G,H insets). Our
observation that Notch^{OFF} hemilineages make more complex and lengthy neurons than the Notch^{ON}
hemilineages is similar to that observed for larval brain lineages (Lee et al., 2020), suggesting that this is a
conserved mechaism. The Notch^{intra} phenotypes we observed are likely due to a Notch^{OFF} to Notch^{ON}
hemilineage transformation, rather than death of ventral projecting neurons, as we observed no change in
neuron number following Notch^{intra} misexpression (e.g. NB5-2 control: 20.3, n=16 hemisegments; NB5-2
Notch^{intra}: 19.3, n=26). In conclusion, we show that NBLAST can be used to accurately identify neuroblast
hemilineages; that Notch^{ON}/Notch^{OFF} hemilineages project to motor/sensory neuropil domains, respectively;
and most importantly, that hemilineage identity determines neuronal targeting to the motor or sensory
neuropil. Thus, each neuroblast coordinately generates similar numbers of sensory and motor processing
neurons throughout its embryonic lineage.
185
186 **Hemilineages target synapses to subdomains of motor or sensory neuropil**
187
188 To identify a relationship between hemilineage identity and synapse localization, we mapped the pre- and
189 postsynapse localization for 12 bilateral hemilineages (24 total) in segment A1. Note we show synapses from
190 both A1L and A1R neuroblast lineages, which highlights the similarity and stereotypy of synapse localization

between left and right lineages (Figure 4A). We found that the neurons in dorsal hemilineages localized both pre- and postsynaptic sites to the dorsal/motor neuropil, whereas neurons in ventral hemilineages localized both pre- and postsynaptic sites to the ventral/sensory neuropil (Figure 4A). Consistent with these observations, we found that the vast majority of sensory output was to ventral hemilineages, and the vast majority of input to motor neurons was from dorsal hemilineages (Figure 4B). We conclude that, at least for the assayed hemilineages, Notch[ON] hemilineages target projections and pre- and postsynapses to the motor neuropil, whereas Notch[OFF] hemilineages target projections and pre- and postsynapses to the sensory neuropil (Figure 4H).

After showing that hemilineages target synapses to dorsal or ventral neuropil, we asked if individual hemilineages target distinct regions of the neuropil or if they have overlapping territories. We mapped the pre- and postsynaptic position for both dorsal and ventral hemilineages (excluding the NB7-4 glial hemilineage). We found that presynapses were localized to distinct regions of the neuropil (Figure 4C,D). Similarly, postsynapses were localized to distinct but slightly more overlapping regions of the neuropil (Figure 4E,F). We quantified synapse similarity (a measure of the average distance between synapses of two neurons) using published methods (Schlegel et al., 2016). We found that synapses from lineages in A1L and A1R had similar relative positions in the neuropil (Figure 4–figure supplement 1A-D), highlighting the stereotypy of synapse localization. Importantly, neurons in a hemilineage showed greater similarity in synaptic positions (pre or post) than unrelated neurons (Figure 4G, Figure 4–figure supplement 1E,F). We conclude that each hemilineage targets its presynapses (and to lesser extent postsynapses) to small domains of the sensory or motor neuropil (Figure 4H), strongly suggesting that the developmental information needed for neuropil targeting is shared by neurons in a hemilineage, but not by all neurons in a complete neuroblast lineage (see Discussion).

Mapping temporal identity in a TEM reconstruction: radial position is a proxy for neuronal birth-order

To investigate the role of temporal identity in determining neuronal projections, synapse localization, or connectivity we needed to identify the temporal identity of all 160 interneurons within our dataset. We used two methods. First, we confirmed that temporal transcription factors (Hb, Kr, Pdm, and Cas) have a radial distribution in the embryonic CNS, with early-born Hb+ neurons positioned in a deep layer adjacent to the developing neuropil, and late-born Cas+ neurons are at the most superficial position within the CNS (Isshiki et al., 2001; Kambadur et al., 1998) (Figure 5A). Importantly, we show that this distribution persists for more stable Hb and Cas reporters in the newly hatched larval CNS (Figure 5B,C), the stage of the TEM reconstruction. Thus, radial position can be used as a proxy for temporal identity in both embryos and newly hatched larvae (Figure 5D). Second, we used MCFO to identify additional single Hb+ or Cas+ neurons and matched them to the morphologically identical neuron in the TEM volume (Figure 5E,F; Figure 5–figure supplement 1; and data not shown). In total, we identified 54 Hb+ neurons and 44 Cas+ neurons within the TEM volume. We measured their cortex neurite length (neurite length from the cell body to the neuropil entry point; i.e. the length of the neurite as it traverses the cellular cortex), and found that experimentally verified Hb+ neurons were closer to the neuropil, whereas Cas+ neuron were further from the neuropil (Figure 5G). We conclude that neuronal radial position can be used as a proxy for neuronal temporal identity.

To determine the temporal identity of neurons within the seven bilateral neuroblast lineages, we measured cortex neurite length for each neuron. Importantly, the same neuron in A1L and A1R had similar cortex neurite lengths (Figure 5H), showing that cell body radial position was reproducible. We assigned 70 interneurons in our dataset to one of four temporal cohorts (early, mid-early, mid-late and late born) based on radial position (Figure 5I). Note that we excluded the NB1-2 dorsal hemilineage from radial analysis as we found it to be an unreliable measure of birth order for that hemilineage (see Figure 5–figure supplement 2). We conclude that cortex neurite length can be used as a proxy for temporal identity, that it is reproducible across at least two hemisegments, and that it can be used to approximate the temporal identity of any neuron in the TEM reconstruction. We use these temporal cohorts to explore the relationship between temporal identity and neuronal projections, synapse localization, and connectivity in the following sections.

After determining the temporal identity for 140 neurons in our dataset, we asked whether neurons with a shared temporal identity had similar axon/dendrite projections or synapse localization. We found that

²⁴³ neurons in each temporal class had broad neuronal projections, and no greater than random synapse
²⁴⁴ similarity (Figure 5–figure supplement 3). We conclude that temporal identity, on its own, does not confer
²⁴⁵ shared neuronal projections or synapse localization.
²⁴⁶
²⁴⁷ **Hemilineage-temporal cohorts have distinct synapse localization within the neuropil**
²⁴⁸
²⁴⁹ Although temporal identity alone did not correlate with axon projections or synapse localization, it still could
²⁵⁰ play a role in restricting synapse localization within individual hemilineages. Here we analyze neurons in a
²⁵¹ hemilineage that share a temporal identity, subsequently called hemilineage-temporal cohorts (Supplemental
²⁵² Table 1). We mapped presynaptic and postsynaptic localization for 20 temporal-hemilineage cohorts from
²⁵³ seven neuroblasts, as illustrated for the NB5-2 dorsal hemilineage and the NB3-3 ventral hemilineage (Figure
²⁵⁴ 6A-B) and the remainder of the hemilineage-temporal cohorts (Figure 6–figure supplements 1,2). We found
²⁵⁵ that NB5-2 dorsal hemilineage-temporal cohorts target their <u>presynapses</u> to mostly non-overlapping neuropil
²⁵⁶ domains (Figure 6A), whereas the NB3-3 hemilineage-temporal cohorts target their <u>postsynapses</u> to mostly
²⁵⁷ non-overlapping neuropil domains (Figure 6B). When we expand this analysis to all hemilineages, we found
²⁵⁸ that (a) hemilineage-temporal cohorts had more similar synaptic positions than hemilineage alone, and (b)
²⁵⁹ dorsal hemilineage-temporal cohorts preferentially clustered presynapses, whereas ventral hemilineage-
²⁶⁰ temporal cohorts preferentially clustered postsynapses (Figure 6C-F). This latter observation may reflect the
²⁶¹ need to precisely target premotor output to specific regions of motor neuron dendrites, and the need to
²⁶² precisely receive sensory input from distinct sensory modalities (see Discussion). Our results also suggest that
²⁶³ dorsal and ventral hemilineages utilize temporal identity in different ways to specify the targeting of either
²⁶⁴ presynapses or postsynapses. We conclude that temporal identity subdivides hemilineages into smaller
²⁶⁵ populations of neurons that target both projections and synapses to distinct sub-domains within the larger
²⁶⁶ hemilineage targeting domain (Figure 6G). Thus, hemilineage identity provides coarse targeting within
²⁶⁷ neuropil, and temporal identity refines targeting to smaller sub-domains.
²⁶⁸
²⁶⁹ **Hemilineage-temporal cohorts have shared connectivity**
²⁷⁰
²⁷¹ Work from the Heckscher lab has shown that early-born and late-born temporal cohorts in the NB3-3 lineage
²⁷² have unique connectivity and participate in escape or proprioception locomotor circuits, respectively (Wreden
²⁷³ et al., 2017). This has led them to hypothesize that other hemilineage-temporal cohorts may have shared
²⁷⁴ connectivity (Meng and Heckscher, 2020). Below, we test this hypothesis for 20 hemilineage-temporal
²⁷⁵ cohorts from seven different neuroblasts (Supplemental Table 1). Here we compare connectivity of
²⁷⁶ hemilineage-temporal cohorts to unrelated neurons, neurons sharing a temporal identity, and neurons sharing
²⁷⁷ a hemilineage identity. This allows us to test the hypothesis that hemilineage-temporal cohorts have more
²⁷⁸ shared connectivity other developmental groupings. First, we analyzed the connectome of 12 hemilineages
²⁷⁹ plus the motor and sensory neurons in segment A1. In total, we analyzed 160 interneurons, 56 motor
²⁸⁰ neurons, and 86 sensory neurons, which corresponded to approximately 25% of all inputs and 14% of all
²⁸¹ outputs for the 12 hemilineages. We found similar connectivity for the same neurons on the left and right
²⁸² side of the segment (Figure 7A-C), validating the accuracy of the neuronal reconstructions as well as the
²⁸³ stereotypy of neurogenesis. Consistent with anatomical observations, we found that neurons in a hemilineage
²⁸⁴ were clustered together in network space, suggesting hemilineage-related neurons are highly interconnected
²⁸⁵ and functionally distinct (Figure 7D-F), similar to what has been suggested previously (Harris et al., 2015).
²⁸⁶   Next, to determine if temporal identity could provide additional specificity to the observed hemilineage
²⁸⁷ connectivity, we compared the connectivity of hemilineages to hemilineage-temporal cohorts. Here we focus
²⁸⁸ specifically on the 160 interneurons in our dataset, excluding motor and sensory neurons. We quantified the
²⁸⁹ average connectivity distance between neurons related by hemilineage or hemilineage-temporal cohort.
²⁹⁰ Neuron pairs that are directly connected have a network distance of 1 synapse; neurons that share a common
²⁹¹ input or output have a network distance of 2 synapses, up to an observed maximum of seven synapses.
²⁹² Importantly, neurons within a hemilineage had a lower network distance (i.e. greater connectivity) than
²⁹³ unrelated neurons, while neurons within a hemilineage-temporal cohort had a lower network distance than
²⁹⁴ those related by hemilineage alone (Figure 7G-J). Interestingly, we found that the largest difference between

hemilineage and hemilineage-temporal cohort connectivity was at the 2 synapse distance. While less than 20% of hemilineage-related or hemilineage-temporal cohort related neurons were directly connected, 75% of hemilineage-temporal cohort neurons were separated by two synapses or less (Figure 7I,J). We conclude that hemilineage-temporal cohorts share common connectivity, and provide increased partner specificity compared to connectivity of hemilineages alone (Figure 7K).

Temporal cohorts have greater shared connectivity with each other than predicted by neuropil proximity alone

Next, we next sought to determine how temporal identity maps to connectivity. Peter's rule proposes that connectivity is determined primarily by the overlap of axons and dendrites (Rees et al., 2017), raising the possibility that developmental mechanisms confer connectivity specificity simply by targeting sets of neurons to appropriate regions of the neuropil. To determine the role of neuropil proximity in connectivity, we calculated the spatial overlap presynapses and postsynapses for all pairwise neuronal combinations, which are expected to range from no overlap to high overlap (Figure 8A, schematic). This measurement is analogous to previous synapse similarity measures, but instead we compared the presynapse positions of one neuron to the postsynapse positions of another. The majority of neurons have very little spatial overlap between presynapses and postsynapses (Figure 8A'). Interestingly, even neurons with the highest observed levels of overlap were not always connected (Figure 8A"). Thus, proximity alone can't explain the observed connectivity, consistent with a role for hemilineage-temporal cohorts providing increased synaptic specificity.
    To test the hypothesis that hemilineage-temporal identity confers synaptic specificity, we shuffled the connectivity by preserving the output degree of each neuron while setting the connection probability as a function of pre/post synaptic overlap. We found that hemilineage-temporal cohort to hemilineage-temporal cohort connectivity (Figure 8B, red dashed line) was greater than random wiring (Figure 8B, black) or proximity based wiring (Figure 8B, cyan and magenta).
    Finally, we asked if there was greater connectivity hemilineage-temporal cohorts compared to unrelated neurons, neurons from the same temporal cohort, or neurons from the same hemilineage. We found that presynapses of a hemilineage-temporal cohort have greater shared connectivity than observed for unrelated neurons, temporal cohorts, or hemilineages (Figure 8C). In contrast, postsynapses of a hemilineage-temporal cohort have greater shared connectivity than observed for unrelated neurons, temporal cohorts, but not for hemilineages (Figure 8D). Moreover, pairwise comparisons revealed greater hemilineage-temporal cohort interconnectivity than hemilineages alone (Figure 8E, top). Furthermore, we detected greater connectivity between neurons within an individual hemilineage cohort than within an individual hemilineage (Figure 8E, bottom). These results suggest that within each hemilineage, temporal identity confers an added level of connectivity compared to proximity or hemilineage alone. In conclusion, we propose that neuroblast lineage, hemilineage, and temporal identity function to progressively refine neurite projections, synapse localization and connectivity (Figure 8F).

Hemilineage-temporal cohorts and circuit formation: the Eve proprioceptive circuit

We previously identified a proprioceptive motor circuit that is required to maintain left/right symmetry of body wall muscle contractions (Heckscher et al., 2015)(Figure 9A). There are three Even-skipped (Eve)+ interneurons at the core of the circuit, receiving strong input from the proprioceptive sensory neruons and three local interneurons (Jaam1-3), and strong output to three premotor neurons (Saaghi1-3) (Heckscher et al., 2015). Recent work has shown that the three Eve+ interneurons at the core of the circuit are from a single hemilineage-temporal cohort: late-born neurons from the NB3-3 hemilinage (Wreden et al., 2017). This has led to speculation that the input Jaam neurons and the output Saaghi neurons may also be hemilineage-temporal cohorts (Meng and Heckscher, 2020). Fortunately, the Eve+, Jaam and Saaghi neurons are all contained within our data set, allowing us to test this hypothesis. We confirm the results of Wreden et al., showing that the three Eve+ neurons (A08e1-3) are in a single late-born hemilineage-temporal cohort (Figure 9B; Supplemental Table 1). In addition, we find that the Jaam neurons are all derived from a single hemilineage-temporal cohort: late-born neuron in the NB5-2 ventral (Notch^{OFF}) hemilineage (Figure 9B;

347 Supplemental Table 1). Similarly, all three Saaghi neurons are derived from a single hemilineage-temporal
348 cohort: late-born neuron in the NB5-2 dorsal (Notch^{ON}) hemilineage (Figure 9B; Supplemental Table 1).
349 Thus, the Jaam-EL-Saaghi proprioceptive circuit is assembled from three distinct hemilineage-temporal
350 cohorts (Figure 9C). We propose that other motor circuits may also be assembled by preferential connectivity
351 between distinct hemilineage-temporal cohorts.
352
353 **Discussion**
354
Here we determine the relationship between developmental mechanisms (spatial, temporal, and hemilineage
identity) and circuit assembly mechanisms (projections, synapse localization, and connectivity). To do this we
map both developmental and circuit features for 160 neuronal progeny of 14 neuroblast lineages in a serial
section TEM reconstruction – this allows us to characterize neurons that share a developmental feature at
single synapse resolution. It is important to note that we chose the seven neuroblasts in this study based on
successful clone generation and availability of single neuroblast Gal4 lines, and thus there should be no bias
towards a particular pattern of neurite projections, synapse localization or connectivity. Our results show that
individual neuroblast lineages have unique but broad axon and dendrite projections to both motor and
sensory neuropil; hemilineages restrict projections and synapse localization to either motor or sensory
neuropil; and distinct temporal identities within hemilineages provide additional specificity in synapse
localization and connectivity. Thus, all three developmental mechanisms act sequentially to progressively
refine neurite projections, synapse localization and connectivity (Figure 8F).
In mammals, clonally-related neurons often have a similar location (Fekete et al., 1994; Mihalas and
Hevner, 2018), morphology (Mihalas and Hevner, 2018; Wong and Rapaport, 2009), and connectivity (Yu et
al., 2009). In contrast, we found that clonal progeny of each neuroblast project widely in the neuropil, to
both sensory and motor domains, and thus clones lack shared morphology, with some neurons projecting to
motor neuropil, and other neurons projecting to sensory neuropil. Perhaps as brain size expands to contain
an increasing number of progenitors, each clone takes on a more uniform structure and function. Yet the
observation that each neuroblast clone had highly stereotyped projections suggests that neuroblast identity
(determined by the spatial position of the neuroblast) determines neuroblast-specific projection patterns.
Testing this functionally would require manipulating spatial patterning cues to duplicate a neuroblast and
assay both duplicate lineages for similar projections and connectivity.
We found that hemilineages produce sensory and motor processing units via a Notch-dependent
mechanism. Pioneering work on *Drosophila* third instar larval neuroblast lineages showed that each neuroblast
lineage is composed of two hemilineages with different projection patterns and neurotransmitter expression
(Harris et al., 2015; Lacin and Truman, 2016; Truman et al., 2010). We extend these studies to embryonic
neuroblasts, and show that Notch signaling determines motor versus sensory neuropil projections in all
lineages examimed. Surprisingly, the Notch^{ON} hemilineage always projected to the dorsal/motor neuropil,
whereas the Notch^{OFF} hemilineage always projected to the ventral/sensory neuropil. The relationship
between the Notch^{ON} hemilineage projecting to motor neuropil may be a common feature of all 30 segmental
neuroblasts, or it could be that the Notch^{ON}/Notch^{OFF} provides a switch to allow each hemilineage to
respond differently to dorsoventral guidance cues, with some projecting dorsally and some projecting
ventrally. Analysis of additional neuroblast lineages will resolve this question.
Hemilineages have not been well described in vertebrate neurogenesis. Notch signaling within the Vsx1+
V2 progenitor lineage generates Notch^{OFF} V2a excitatory interneurons and Notch^{ON} V2b inhibitory
interneurons, which may be distinct hemilineages (Del Barrio et al., 2007; Francius et al., 2016; Peng et al.,
2007; Skaggs et al., 2011). Interestingly, both V2a and V2b putative hemilineages contain molecularly distinct
subclasses (Harris et al., 2019); our work raises the possibility that these subtypes arise from temporal
patterning within the V2 lineage. In addition, Notch^{ON}/Notch^{OFF} hemilineages may exist in the pineal
photoreceptor lineage, where Notch^{ON} and Notch^{OFF} populations specify cell type identity (Cau et al., 2019).
Elegant work has identified neuropil gradients of Slit and Netrin along the mediolateral axis (Zlatic et al.,
2009), Semaphorins along the dorsoventral axis (Zlatic et al., 2009), and Wnt5 along the anteroposterior axis
(Yoshikawa et al., 2003). Our finding that neurons in a hemilineage project to a common region of the
neuropil strongly suggests that all neurons within a hemilineage respond in the same way to these global

pathfinding cues. Conversely, our finding that neurons in different hemilineages target distinct regions of the neuropil suggests that each hemilineage expresses a different palette of guidance receptors, which enable them to respond differentially to the same global cues. For example, neurons in ventral hemilineages may express Plexin receptors to repel them from high Semaphorins in the dorsal neuropil.

Only recently have the role of hemilineages been tested for their functional properties. In adults, activation of each larval hemilineage from NB5-2 showed similar behavioral output, whereas each hemilineage from NB6-1 elicited different behaviors (Harris et al., 2015). Our previous work showed that the Eve+, Saaghi and Jaam neurons are part of a proprioceptive circuit (Heckscher et al., 2015); here we show that each class of neurons represents a hemilineage-temporal cohort. Note that the Jaam neurons process sensory input and are in a Notch^{OFF} hemilineage, supporting our conclusion that Notch^{OFF} hemilineages are devoted to sensory processing; the Saaghi premotor neurons are in a Notch^{ON} hemilineage consistent with their role in motor processing. Interestingly, both input and output neurons in this circuit arise from a common progenitor (NB5-2), which may generate late-born Jaam/Saaghi sibling neurons (Figure 9B). In the future, it would be interesting to determine if other sibling hemilineages are in a common circuit to generate a specific behavior.

Our hemilineage results have several implications. First, our results reveal that sensory and motor processing components of the neuropil are being built in parallel, with one half of every GMC division contributing to either sensory or motor networks. This would be an efficient mechanism to maintain sensory/motor balance as lineage lengths are modified over evolutionary time. Second, our results suggest that analyzing full neuroblast clones may be misleading due to the clone comprising two quite different hemilineages. For example, performing bulk RNAseq on all neurons in a neuroblast lineage is unlikely to reveal key regulators of pathfinding or synaptic connectivity, due to the mixture of disparate neurons from the two hemilineages.

We used the cortex neurite length of neurons as a proxy for birth-order and shared temporal identity. We feel this is a good approximation, but it clearly does not precisely identify neurons born during each of the Hb, Kr, Pdm, Cas temporal transcription factor windows. Nevertheless, we had sufficient resolution to observe that neurons with the same temporal identity clustered their pre- or postsynapses, rather than localizing them uniformly through the hemilineage neuropil domain (Figure 4H). Interestingly, the three-dimensional location of each hemilineage temporal cohort synaptic cluster is identical on the left and right side of A1 (data not shown), ruling out the mechanism of stochastic self-avoidance (Zipursky and Grueber, 2013). Other possible mechanisms include hemilineage-temporal cohorts expressing different levels of the spacing cue Sequoia (Kulkarni et al., 2016; Petrovic and Hummel, 2008), or hemilineage-temporal cohorts exhibiting different responses to global patterning cues. Testing the function of temporal identity factors in synaptic tiling will require hemilineage-specific alteration of temporal identity, followed by assaying synapse localization within the neuropil.

Our results strongly suggest that hemilineage identity and temporal identity act combinatorially to allow small pools of neurons to target pre- and postsynapses to highly precise regions of the neuropil, thereby restricting synaptic partner choice. Yet precise neuropil targeting is not sufficient to explain connectivity, as many similarly positioned axons and dendrites fail to form connections (Figure 8C). We favor the model that hemilineages direct gross neurite targeting to motor or sensory neuropil, whereas temporal identity acts combinatorially with each hemilineage to direct more precise neurite targeting and synaptic connectivity. Thus, the same temporal cue (e.g. Hb) could promote targeting of one pool of neurons in one hemilineage, and another pool of neurons in an adjacent hemilineage. This limits the number of regulatory mechanisms needed to generate precise neuropil targeting and connectivity for all ~600 neurons in a segment of the larval CNS.

In conclusion, we demonstrate how developmental information can be mapped into large scale connectomic datasets. We show that lineage information, hemilineage identity, and temporal identity can all be accurately predicted using morphological features (e.g. number of fascicles entering the neuropil for neuroblast clones, and radial position for temporal cohorts). This both greatly accelerates the ability to identify neurons in a large EM volume as well as sets up a framework in which to study development using datasets typically intended for studying connectivity and function. We have used this framework to relate developmental mechanism to neuronal projections, synapse localization, and connectivity. We find that

451   lineage, hemilineage, and temporal identity act sequentially to progressively refine neuronal projections,

452   synapse localization and connectivity, and our data supports a model where hemilineage-temporal cohorts are

453   units of connectivity for assemblying motor circuits.

Methods

Key Resources Table

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Species (*Drosophila melanogaster*)	R16A05[AD] R28H10[DBD]	(Lacin et al., 2019)		NB1-2 split Gal4
Species (*Drosophila melanogaster*)	R70D06[AD] R28H10[DBD]	(Lacin et al., 2019)		NB2-1 split Gal4
Species (*Drosophila melanogaster*)	Ac[AD] Gsb[DBD], 25A05[kz]	(Seroka and Doe, 2019)		NB7-1 split Gal4
Species (*Drosophila melanogaster*)	R19B03[AD] R18F07[DBD]	(Lacin et al., 2019)		NB7-4 split Gal4
Species (*Drosophila melanogaster*)	castor-gal4	Gift from Technau lab		Late-born neuron marker
Species (*Drosophila melanogaster*)	hsFlp.G5.PEST.Opt	BDSC	RRID: 77140	Heat-inducible Flp recombinase
Species (*Drosophila melanogaster*)	26XLexAop2-mCD8::GFP	BDSC	RRID: 32207	LexA reporter
Species (*Drosophila melanogaster*)	13XLexAop2-IVS-myr::GFP	BDSC	RRID: 32210	LexA reporter
Species (*Drosophila melanogaster*)	dpn(FRT.stop)LexA.p65	BDSC	RRID: 56162	Used with hsFlp and lexAop-GFP (above) to create and visualize clones in single Dpn+ neuroblasts
Species (*Drosophila melanogaster*)	13XLexAop2-IVS-myr::GFP	BDSC	RRID: 32210	LexA reporter
Species (*Drosophila melanogaster*)	w[1118] P{y[+t7.7] w[+mC]=R57C10-FLPG5.PEST}attP18; P{y[+t7.7] w[+mC]=10xUAS(FRT.stop)myr::smGdP-OLLAS}attP2 PBac{y[+mDint2] w[+mC]=10xUAS(FRT.stop)myr::smGdP-HA}VK00005 P{10xUAS(FRT.stop)myr::smGdP-V5-THS-	BDSC	RRID: 64091	Multicolor Flp Out stock (Nern et al., 2015)

	10xUAS(FRT.stop)myr::smGdP-FLAG}su(Hw)attP1			
Species (*Drosophila melanogaster*)	Sco/CyO; Dr/TM3,Sb	BDSC	RRID: 34516	
Species (*Drosophila melanogaster*)	LexA-T2A-Hb	This work		Endogenous hb locus CRISPR engineered to place LexAp65-T2A upstream and in frame with the first hb ORF
Species (*Drosophila melanogaster*)	hey:T2A:FLP, UAS-myr:GFP, UAS-FRT-stop-FRT-myr:sfGdP:HA	This work		Labels Hey+ (Notch[ON]) neurons within a Gal4+ neuronal population
Antibody, polyclonal	Rabbit anti-GFP	ThermoFisher, Waltham, MA		1:500
Antibody, polyclonal	Rabbit anti-Venus	ThermoFisher, Waltham, MA		1:500
Antibody, polyclonal	Chicken anti-GFP	Abcam, Eugene, OR	RRID: 13970	1:1000
Antibody, polyclonal	Camelid sdAB direct labeled with AbberiorStar635P	NanoTab Biotech., Gottingen, Germany		1:1000
Antibody, polyclonal	Rabbit anti-mCherry	Novus, Littleton, CO		1:1000
Antibody, polyclonal	Alexa Fluor 488-conjugated rabbit anti-GFP	Thermofisher (Eugene, OR)	RRID:AB_221477	1:1000
Antibody, monoclonal	Mouse anti-FasII	DSHB (Iowa City, IA)		1:100
Antibody, monoclonal	Mouse anti-HA	Cell signaling, Danvers, MA		1:200
Antibody, polyclonal	Rabbit anti-V5	Rockland, Atlanta, GA		1:400
Antibody, polyclonal	Rabbit anti-FLAG	Rockland, Atlanta, GA		1:200
Antibody, monoclonal	Mouse anti-Engrailed	DSHB (Iowa City, IA)	RRID:AB_528224	1:100
Antibody, polyclonal	Rabbit anti-Hb	Doe lab		1:400
Antibody, polyclonal	Alexa Fluor™ 405 Phalloidin	Thermofisher (Eugene, OR)		1:40
Antibody, polyclonal	Secondary antibodies	Thermofisher (Eugene, OR)		1:400
DNA	pHD-DsRed	Addgene	RRID:Addgene_51434	
DNA	pCFD5	Addgene	RRID:Addgene_73914	

Transgenic fly stocks

Transgenic lines were made by BestGene (Chino Hills, CA) or Genetivision (Houston, TX).

<u>NB Clone generation and lineage identification</u>

The NB clones were generated with the following flies: *hs-Flp.G5.PEST.Opt, dpn(FRT.stop)LexA.p65*,

26XLexAop2-mCD8::GFP. The embryos were collected in 25 °C for 3 hours and then incubated in 25 °C for

another 3 hours. The aged embryos were then submerged in 32 °C water bath for 5-minute heat shock, and then

incubated in 25 °C until larvae hatched. The CNS of newly hatched larvae was dissected and mounted as

previously described (Clark et al., 2016; Heckscher et al., 2014; Syed et al., 2017). The neuropil was stained with

mouse anti-engrailed (RRID:AB_528224) (DSHB, 4D9), and Alexa Fluor™ 647 Phalloidin (ThermoFisher) by

following manufacturer's protocol. The images were collected with Zeiss710 and processed with Imaris.

Lineages were identified in the EM volume by finding neurons with morphologies that matched the

clonal morphology, and then identifying their neuropil entry point. We then examined every neuron which

entered the neuropil in the same fascicle. In most cases, every neuron in the fascicle had a morphology that

matched the clonal morphology. In a small number of cases, the fascicles diverged slightly before the

neuropil entry point. We verified the number of neurons by looking at fasciculating cell populations from at

least two hemisegments (A1L and A1R). In some cases, we were able to identify a stereotyped number of

cells across as many as four hemisegments, suggesting that fasciculation is stereotyped and reliable.

<u>Hb+ single cell clone generation</u>

Hb+ single cell clones were generated with the following flies: *hs-Flp.G5.PEST.Opt, 13XLexAop2-IVS-*

myr::GFP, and *LexAp65-T2A-hb* (see below). The embryos were collected for 7 hours in 25 °C, submerged in 32

°C water bath for 10-min heat shock, and then follow the protocol as described above.

LexAp65-T2A-hb was generated by in-frame fusion of *(FRT.stop)::LexA.P65::T2A* to the N-terminus of

the hb open reading frame with CRISPR-Cas9 gene editing. The ds-DNA donor vector for homology-directed

repair was composed of left homologous arm (1000bp), LexA.P65 (Pfeiffer et al., 2008), T2A (Nern et al., 2015),

and the right homologous arm (1000bp); the fragments were amplified with PCR and then assembled in pHD-

DsRed (RRID:Addgene_51434) with NEBuilder (New England BioLabs). The gRNAs were generated from the

vector pCFD5 (RRID:Addgene_73914) (Port et al., 2014) containing target sequence

TGCATCTTGGCGGCTCTAGA and ACTACGAGCAGCACAACGCC. The ds-DNA donor vectors and

gRNA vectors were co-injected into *yw;nos-Cas9* (Kondo and Ueda, 2013) flies by BestGene. The selection marker

3xP3-DsRed was then removed in transgenic flies by *hs-Cre*.

<u>Immunostaining and imaging</u>

Standard confocal microscopy, immunocytochemistry and MCFO methods were performed as previously

described for larvae (Clark et al., 2016; Heckscher et al., 2014; Syed et al., 2017) or adults (Nern et al., 2015;

Pfeiffer et al., 2008). Secondary antibodies were from Jackson Immunoresearch (West Grove, PA) and used

according to manufacturer's instructions. Confocal image stacks were acquired on Zeiss 700, 710, or 800

microscopes. Images were processed in Fiji (https://imagej.net/Fiji), Adobe Photoshop (Adobe, San Jose,

CA), and Adobe Illustrator (Adobe, San Jose, CA). When adjustments to brightness and contrast were

needed, they were applied to the entire image uniformly. Mosaic images to show different focal planes were

assembled in Fiji or Photoshop.

<u>Morphological analysis of lineages</u>

Morphological analysis was done using NBLAST and the NAT package (Costa et al., 2016), and analysis and

figure generation was done using R. Neurons were preprocessed by pruning the most distal twigs (Strahler order

4), converting neurons to dot-props, and running an all-by-all NBLAST. For individual lineages, clusters were set
using a cutoff of 3.0. In the case of NB2-1, where nearly every neuron shares a very similar morphology, we first
confirmed the presence of a hemilineage using anti-Hey staining. After confirmation of a hemilineage, we next
removed A02o and A02l since we could not find any clones that contained either an anterior projection (A02o) or
a second contralateral projection (A02l). We reasoned that the hemilineages would represent the next largest
morphological division (Fig. S1).

<u>Synaptic distributions and density analysis</u>
Synapse distribution plots and density contours were generated using MATLAB. Neuron synaptic and
skeleton information was imported to MATLAB using pymaid (Schlegel et al., 2016). Cross sectional synapse
distribution plots were made by taking all synapse positions between the T3 and A2 segments as positional
information becomes lost due to changes in brain shape beyond these bounds. Synapse distribution plots are
1D kernel density estimates. Sensory and motor density maps were made by taking the synapse positions of
all sensory neurons entering the A1 nerve, and all motor neurons exiting the A1 nerve as well as all neurons
with at least 3 synapses connected to one of these neurons. For sensory and motor maps as well as individual
hemilineages, density plots are 2D kernel density estimates of all synapse positions across the neuropil. A
cutoff of 60% was used to set the outermost contour. For lineage maps (Figure 4A), we used 80% as a
cutoff. Polyadic synapses were counted as many times as they have targets. For synapse distribution plots,
polyadic synapses are scaled by their number of targets.

<u>Temporal Cohort assignment</u>
Cortex neurite length was calculated by converting the skeletonized neuronal arbor into a directed graph away
from the soma, and performing a depth-first-search of all vertices. The neuropil borders were defined by a
previously created neuropil volume object (Costa et al., 2016). The neuropil entry point was defined as the
first vertex within the neuropil volume object. Cortex neurite length was then the path length between the
soma and the neuropil entry point. Neurons were binned into 5 groups with 6μm edges to define temporal
cohorts.

<u>Synapse similarity measurements</u>
Synapse similarity was calculated as described previously (Schlegel et al., 2016):

534
$$f(is, jk) = e^{\frac{-d_{sk}^2}{2\sigma^2}} e^{\frac{|n_{is} - n_{jk}|}{n_{is} + n_{jk}}}$$

where $f(is,jk)$ is the mean synapse similarity between all synapses of neuron i and neuron j. d_{sk} is the Euclidean
distance between synapses s and k such that synapse k is the closest synapse of neuron j to synapse s of neuron i.
σ is a bandwidth term that determines what is considered close. n_{is} and n_{jk} are the fraction of synapses for neuron
i and neuron j that are within ω of synapse s and synapse k respectively. We used parameters ω= σ = 2000nm.
For presynaptic or postsynaptic comparisons, the score for a given pair of neurons was the average similarity
between the left and right sides. In order to calculate pre/post overlap, we applied the same measure, instead
using the presynapses of neuron i and the postsynapses of neuron j.

<u>Connectivity similarity measurement and random models</u>
Connectivity similarity was calculated as one minus the cosine distance between either the row vectors (outputs)
or column vectors (inputs) of the binarized connectivity matrix for all neurons. For models of random

connectivity, we used two methods to determine the individual connection probabilities. First, we used a standard degree based method in which the number of inputs for a given neuron was divided by the total number of inputs received by that neuron. Second, we used the pre-post synapse overlap scores. For a given neuron, its probability of connecting to another neuron was equal to the overlap score divided by the sum of all overlap scores for that neuron. Pairs with a score of 0 had a 0 probability of connecting. Thresholds were done by setting all values below the threshold to zero when determining connection probability.

Electron microscopy and CATMAID

We reconstructed neurons in CATMAID as previously described (Carreira-Rosario et al., 2018; Heckscher et al., 2015; Ohyama et al., 2015).

Figures

Figures were generated using Matlab, R, CATMAID, and FIJI, and edited in either Illustrator or Photoshop (Adobe, San Jose CA).

Statistical analysis

Statistical significance is denoted by asterisks: ****p<0.0001; ***p<0.001; **p<0.01; *p<0.05; n.s., not significant. All statistical analysis was done in MATLAB. When comparing two groups of quantitative data, an unpaired t-test was performed if data was normally distributed (determined using a one-sample Kolmogorov-Smirnov test) and Wilcoxon rank sum test if the data was not normally distributed. Two sample Kolmogorov-Smirnov tests were used on empirical distributions. Linear models were generated in MATLAB using lmfit.

Acknowledgements

We thank Haluk Lacin for unpublished fly lines. We thank Todd Laverty, Gerry Rubin, and Gerd Technau for fly stocks; Luis Sullivan, Emily Sales, John Tuthill and Tim Warren for comments on the manuscript; Avinash Khandelwal and Laura Herren for annotating neurons; Keiko Hirono for generating transgenic constructs; and Keiko Hirono, Rita Yazejian, Dalton Lee, Cooper Doe and Casey Doe for confocal imaging. Stocks were obtained from the Bloomington *Drosophila* Stock Center (NIH P40OD018537). Funding was provided by HHMI (CQD, BM, LM, AAZ), NIH HD27056 (CQD), and NIH T32HD007348-24 (BM).

Figure 1. **Mapping neurons with shared developmental origin in a TEM reconstruction: clonally-related neurons project widely and localize synapses to both sensory and motor neuropil**

(A) Three mechanisms specifying neuronal diversity. Neuroblasts characterized here are shown in dark gray and arise from all anteroposterior and mediolateral positions of the neuroectoderm. Nb, Numb; N, Notch; A1L, abdominal segment A1 left side.

(A') Schematic of newly hatched larval CNS in a dorsal view, anterior up (top) or in a posterior view, dorsal up (bottom). All images in the figures are shown in posterior view, dorsal up unless noted otherwise.

(B) Single neuroblast clones for the indicated neuroblasts (larval neuroblast names given in Supplemental Table 1) each generated in A1L with *dpn(FRT.stop)LexA.p65* and assayed in newly-hatched larvae. We recovered n>2 clones for each newly characterized lineage; NB4-1 was previously characterized (Lacin and Truman, 2016). Dashed lines, neuropil border; vertical dash, midline. Scale bar, 10 μm.

(C) The corresponding neurons traced in the TEM reconstruction. Dashed lines, neuropil border; vertical dash, midline.

(D) Each clone characteristically has either one or two fascicles at neuropil entry (yellow, arrows).

(E) There are a similar number of neurons per neuroblast clone in A1L and A1R.

(F) Presynaptic and postsynaptic density maps (75% threshold) for the each neuroblast lineage. Dashed lines, neuropil border; vertical dash, midline. All density maps are from neurons in A1L (cell bodies not shown).

(G) Seven bilateral neuroblast lineages in segment A1 left traced in the TEM reconstruction. Inset: same projections, lateral view, anterior up.

(H) Summary.

Figure 2. Each neuroblast lineage generates neurons projecting to dorsal/motor or ventral/sensory neuropil.

(A-C) Organization of motor and sensory domains withing the ventral nerve cord neuropil. (A) Motor neuron postsynapses (purple) and sensory neuron pre-synapses (green) showing dorsoventral segregation. Plots are 1D kernel density estimates for dorsoventral or mediolateral axes. Purple dots represent a single postsynaptic site. Green dots represent a single pre-synaptic site scaled by the number of outputs from that presynaptic site. (B) Premotor neuron post-synaptic sites (>3 synapses onto a motor neuron in segment A1), or post-sensory neuron presynaptic sites (pre >3 synapses with an A1 sensory neuron) show that connecting neurons are still restricted to dorsal or ventral neuropil domains. (C) 2D kernel density estimates of all presynaptic and postsynaptic sites for pre-motor and post-sensory neurons outlines the regions of sensory (green) and motor (magenta) processing in the ventral nerve cord.

(D-I) NBLAST clustering for the indicated neuroblast progeny typically reveals two morphological groups of neurons or glia (red/cyan) that project to dorsal or ventral neuropil; these are candidate hemilineages. Cluster cutoffs were set at 3.0 for all lineages.

(J) Superimposition of all dorsal candidate hemilineages (red) and all ventral candidate hemilineages (cyan). Data from NBs 1-2, 3-3, 4-1, 5-2, 7-1, and 7-4 (this figure) and NB2-1 (Figure 2–figure supplement 1).

Figure 2–figure supplement 2. Clonally related neurons can have different morphology.

Dendrogram showing NBLAST results clustering neurons based on similar morphology. Clustered neurons were all from hemisegment A1L. Colored bars denote lineage identity. Note that some lineages contain neurons with highly divergent morphology as indicated by their lack of clustering together in the dendrogram (e.g. NB4-1 and NB5-2 lineages).

Figure 2–figure supplement 1. NB2-1 has Notch[ON] and Notch[OFF] neurons, but all project to a similar neuropil domain.

(A) NBLAST clustering for the NB2-1 progeny reveals two morphological groups (red/cyan). The magenta and green neurons are outliers and may be from another lineage because we never identified a NB2-1 clone containing these two neurons; the two outlier neurons were removed prior to hemilineage analysis below. Cluster cutoffs were set at 3.0.

(B,B') Reconstruction of NB2-1 progeny; presumptive Notch[ON] neurons (red) are slightly more dorsal and lateral than the presumptive Notch[OFF] neurons (cyan).

 (C,C') The Notch target gene Hey is expressed in ~half of the neurons in the NB2-1 lineage, confirming that
there are Notch[ON] and Notch[OFF] hemilineages. Stage 16 embryo; scale bar, 5μm. Posterior view, dorsal up,
midline at center of panel. These hemilineages are likely to be the red/cyan neurons identified by NBLAST.

Figure 3. Hemilineage identity drives neuronal projections to motor or sensory neuropil
(A-D) A Notch reporter (*hey:T2A:FLP, UAS-myr:GFP, UAS-FRT-stop-FRT-myr:sfGdP:HA*) specifically labels
dorsal projections within the indicated neuroblast lineages. Myr:GFP labels the whole lineage (green),
myr:sfGdP:HA is a myristoylated (myr) superfolder green dead protein (sfGdP) fused to hemoagglutinin (HA),
which labels the Notch[ON] hemilineage (magenta). Vertical dashes, midline; dorsal, up.
(E) NBLAST clustering reveals two morphological classes of neurons in each indicated lineage. Cluster cutoffs
were set at 3.0.
(F) TEM reconstruction of neuronal projections for the indicated lineages; color coding to match NBLAST class.
Dashed lines, neuropil border; vertical dashes, midline; dorsal, up.
(G,H) Lineage-specific Notch[intra] expression transforms ventral projections to dorsal projections. (G) Control
neuroblast lineages project to both dorsal neuropil (red arrowhead) and ventral neuropil (cyan arrowhead). NB7-4
makes glia (red arrowheads) and ventral projections. (H) Notch mis-expression in the indicated neuroblast
lineages results in a loss of ventral projections and an increase in dorsal projections or glia. Cell numbers in
control and Notch mis-expression are similar (data not shown). n>3 for all experiments.
(I) Summary.

Figure 3–figure supplement 1. Ventral hemilineages have projection neurons
The indicated neuroblast lineages traced in catmaid showing the dorsal (red) and ventral (cyan) predicted
hemilineages. Note that the ventral (cyan) hemilineages contains significantly longer axons (ascending and
descending projection neurons) compared to dorsal (red) hemilineage neurons, consistent with what has been
observed in larva (Truman, 2010). P = .0034, via 2-sided Wilcoxon rank sum test.

Figure 4. Hemilineages target pre- and postsynapses to subdomains of dorsal or ventral neuropil
(A) Density maps for dorsal synapses (red) or ventral syapses (blue) within each lineage. Dorsal hemilineages
localize both presynapses and postsynapses to dorsal neuropil, whereas ventral hemilineages localize both
presynapses and postsynapses to ventral neuropil. Vertical dashes, midline. Synapses from both A1L and A1R are
shown to highlight left/right stereotypy.
(B) Connectivity diagram showing sensory neurons provide inputs to neurons in ventral hemilineages, while
motor neurons preferentially receive inputs from neurons in dorsal hemilineages. Edges represent fractions of
outputs for sensory neurons, and fraction of inputs for motor neurons.
(C,D) Presynaptic distributions of of the indicated dorsal or ventral hemilineages from A1L. Dots represent single
presynaptic sites with their size scaled by the number of associated (polyadic) postsynaptic sites. Circles, location
of cell bodies.
(E,F) Postsynaptic distributions of the indicated dorsal or ventral hemilineages from A1L. Dots represent single
postsynaptic sites. Circles, location of cell bodies.
(G) Mean inter- versus intra-hemilineage synapse similarity scores for dorsal and ventral hemilineages show intra-
hemilineage presynapse (blue) and postsynapse (orange) similarity is greater than inter-hemilineage (black)
similarity. In this case, intra-hemilineage similarity represents comparisons only to neurons targeting the same
region of the neuropil (dorsal-dorsal / ventral-ventral). Error bars represent SEM. P<.05 in all cases (Wilcoxon
rank sum test).
(H) Summary.

Figure 4–figure supplement 1. Neurons in a hemilineage cluster their synapses to specific regions of the neuropil
(A) Diagram of neuropil volume analyzed for synapse spatial positions.
(B) Pre- and postsynapse positions are highly reproducible for the same neuron in A1L and A1R.

(C,D) Presynapse positions are more stereotyped than postsynaptic positions. (C) Left/right variation between pairwise synapse similarity scores. (D) Distribution of per-neuron correlation coefficients for similarity scores shows presynaptic sites are more stereotyped. P<.05, 2 sample KS-test.

(E,F) Synapse similarity shows greater presynapse clustering (blue) than postsynapse clustering (orange) in both dorsal and ventral hemilineages. Postsynaptic clustering is greater in ventral hemilineages than dorsal hemilineages.

Figure 5. Mapping temporal identity in the TEM reconstruction: radial position is a proxy for neuronal birth-order

(A-C) Early-born Hb+ neurons are near the neuropil and late-born Cas+ neurons are far from the neuropil in the late embryo (A) and newly hatched larvae (B,C); larval reporters are recombineered *Hb:GFP* and *cas-gal4 UAS-RFP*.

(D) Schematic showing correlation between radial position and early or late born temporal identity.

(E,F) Examples of experimentally verified 'ground truth' early-born Hb+ neurons and late-born Cas+ neurons, identified using MCFO with *cas-gal4* or using a CRISPR engineered *hb* locus where LexA:T2A is in frame with start of the *hb* coding sequence (see methods); additional examples in (Figure 5–figure supplement 1).

(G) Fraction of experimentally validated Hb+ or Cas+ 'ground truth' neurons at the indicated distance from the neuropil (cortex neurite length). n = 47 Cas+ neurons, and 55 Hb+ neurons from segments T3-A2.

(H) Cortex neurite lengths between the same neuron in the left and right hemisegment, showing that radial position is highly stereotyped.

(I) Assignment of temporal identity based on radial position for neurons within the TEM reconstruction, calculated as the mean left/right distance for the same neuron in A1L and A1R. Note that not all lineages have all temporal cohorts, mirroring experimental observation that some neuroblast lineages do not express all temporal transcription factors (Benito-Sipos et al., 2010; Cui and Doe, 1992; Isshiki et al., 2001; Tsuji et al., 2008).

Figure 5–figure supplement 1. Ground truth Hb+ and Cas+ neurons used to define radial position as a proxy for temporal identity.

(A) Previously identified Hb+ or Cas+ neurons from the A1L TEM reconstruction. Hb neurons, cyan; Cas neurons, magenta. Dorsal view, anterior up; midline, dashes. Inset, posterior view.

(B) Newly discovered Hb+ or Cas+ neurons identified by MCFO using *hb-lexA* or *cas-gal4*. MCFO neurons, black panels; matching TEM reconstructed neurons, white panels. Midline, dashes.

(C) All reconstructed Hb+ and Cas+ neurons from A1L and A1R shown in posterior view, dorsal up.

Figure 5–figure supplement 2. NB1-2 dorsal hemilineage: radial proxy is unreliable for determining the temporal identity.

(A) Presumptive temporal cohorts for NB1-2 dorsal (NB1-2D) hemilineage show an early cohort with 6 neurons, while mid and late cohorts have only 1 or 2 neurons respectively.

(B) Synaptic positions of presumptive temporal cohorts for NB1-2D show no tiling, unlike other hemilineage-temporal (HL-T) cohorts.

(C) Mean presynapse similarity of NB1-2D early is ~70% less than the average synapse similarity observed in other HLTs.

(D) Radially determined NB1-2D early has more cells than any other temporal cohort. The only other HLT with a similar number is NB3-3, a late forming neuroblast with a known large Cas window (Baumgardt et al., 2014; Tsuji et al., 2008; Wreden et al., 2017).

(E) NB1-2 Cas+ neurons are observed both close and far from the neuropil in suggesting radial proxy is a poor measure of birth-order for this lineage. Scale bar, 2 µm.

Figure 5–figure supplement 3. Neurons with a common temporal identity project widely within the neuropil

(A) Temporal cohorts determined by radial proxy show broad projections throughout the neuropil. Posterior view, dorsal up; midline, vertical dashed line.

(B) Temporal cohorts do not have synapse similarity greater than random. Synapse similarity of temporal cohorts (across hemilineages) does not confer a common axon/dendrite projection pattern, except for a small but significant difference observed among ventral hemilineage presynapses. NS = not significant, * signifies p<.05, Wilcoxon rank sum test. In this case, inter-cohort comparisons represent a common (radially determined) birth-date, a common dorsal/ventral target, but unrelated by hemilineage.

Figure 6. Hemilineage-temporal cohorts show synaptic tiling within motor or sensory neuropil
(A-B) Hemilineages target their synapses within the neuropil. (A) NB5-2 dorsal hemilineage neurons segregate their presynapses to distinct regions of the neuropil. (B) NB3-3 ventral hemilineage neurons segregate their postsynapses to distinct regions of the neuropil. Circles, location of cell bodies.
(C-F) Hemilineage-temporal (HL-T) cohorts target their synapses within each hemilineage domain. (C,D) All dorsal hemilineages in A1L. Left: cumulative plots showing that hemilineages have greater synaptic clustering than random, but that HL-T cohorts have greater presynaptic clustering than hemilineages alone. Right: Histograms showing that hemilineages have greater presynaptic clustering than random, but that HL-T cohorts have more presynaptic clustering than hemilineages alone. Circles, location of cell bodies. (E,F) All ventral hemilineages in A1L. Left: cumulative plots showing that hemilineages have greater postsynaptic clustering than random, but that HL-T cohorts have greater postsynaptic clustering than hemilineages alone. Right: Histograms showing that hemilineages have greater postsynaptic clustering than random, but that HL-T cohorts have more postsynaptic clustering than hemilineages alone.
For all quantifications, random refers to hemilineage-unrelated neurons projecting to the same (dorsal or ventral) region of the neuropil. Hemilineage similarity refers to hemilineage related neurons from different temporal cohorts. *:P<.05 or *** P<.001 (Wilcoxon rank-sum test.). Error bars, SEM.
(G) Summary. Dorsal hemilineages tend to cluster presynapses, which provide input to motor neurons; ventral hemilineages tend to cluster postsynapses, which receive input from sensory neurons.

Figure 6–figure supplement 1. Hemilineage-temporal cohorts show synaptic tiling within motor or sensory neuropil
Presynaptic distributions across M/L, D/V, and A/P axes for each hemilineage-temporal cohort.

Figure 6–figure supplement 2 Hemilineage-temporal cohorts show synaptic tiling within motor or sensory neuropil
Postsynaptic distributions across M/L, D/V, and A/P axes for each hemilineage-temporal cohort.

Figure 7. Hemilineages and hemilineage-temporal cohorts have more shared connectivity than unrelated neurons.
(A-C) Connectivity is similar for hemilineages in A1L and A1R. (A,B) Fraction of inputs/outputs for each hemilineage; (C) heatmap showing connectivity between dorsal hemilineages (red), ventral hemilineages (blue), motor neuron dendrites (MN), and sensory neuron axons (SN).
(D-F) Dorsal and ventral hemilineages have distinct connectivity. (D) Force directed network graph of all 160 interneurons, together with sensory afferents and motor dendrites. Neurons with similar connectivity appear closer in network space. Red edges represent dorsal hemilineage connectivity; blue edges represent ventral hemilineage connectivity. (E) Force directed network graph highlighting the lack of shared connectivity between dorsal and ventral hemilineages in the NB5-2 progeny. (F) Force directed network graph highlighting the lack of shared connectivity between dorsal and ventral hemilineages in the NB7-1 progeny.
(G-J) Hemilineage-temporal cohorts within a hemilineage have shared connectivity. (G,H) Force directed network graphs highlighting the shared connectivity of hemilineage-temporal cohorts within NB5-2 or NB7-1 progeny. (I) Cumulative distribution of the number of synapses between temporal cohorts of hemilineage related neurons, hemilineage related neurons, or random neurons. Neurons that belonged to a temporal cohort with only one neuron were not analyzed (16 neurons). Random neurons were selected from the same hemisegment.
(J) Quantification of the number of directly connected pairs of neurons, neurons separated by 1 or 2 synapses. Black circles represent pairs of neurons connected by 1 synapse (top) or two synapses (bottom).
(K) Summary.

Figure 8: Hemilineage-temporal cohorts have greater shared synaptic connectivity than unrelated neurons or hemilineages.

(A) Axon-dendrite proximity alone does not predict connectivity. The distribution of presynaptic/postsynaptic overlap for all neurons in our dataset; higher "similarity" = smaller average 3D distance between presynapses and postsynapses. (Top) Most neuronal pairs have little overlap (0-0.1 score) in presynapse/postsynapse position. (Bottom) Presynaptic and postsynaptic neurons have increasing connectivity probability as the distance between presynapses and postsynapses decreases, but even neurons with the most presynapse/postsynapse overlap often fail to be connected (e.g. 0.7-0.9 similarity are <o.6 connectivity).

(B) HL-T to HL-T connectivity occurs more frequently than can be explained by proximity alone. Red dashed line, observed frequency that a pair of neurons in a HL-T cohort connect to one or more neuron in another HL-T cohort. Two synapse threshold used. Colored distributions represent datasets shuffled on the basis of proximity, while the black distribution is a dataset shuffled on the basis of input probability (see methods).

(C,D) Hemilineage-temporal (HL-T) cohorts share common connectivity. Left: violin plots; median, white circle. Right: cumulative plots. (A) HL-T cohort <u>presynapses</u> have greater shared connectivity than observed for unrelated neurons, temporal cohorts, or hemilineages. (B) In contrast, HL-T cohort <u>postsynapses</u> have greater shared connectivity than observed for unrelated neurons, temporal cohorts, but not for hemilineages. Connectivity similarity is equivalent to 1 minus the cosine distance between the presynapses (A, blue) or postsynapses (B, orange) vectors of the binarized connectivity matrix. **** $p < .0001$ in a Mann-Whitney test.

(E) Top: there is greater connectivity between pairs of HL-T cohorts (green) than between pairs of hemilineages (magenta). Bottom: there is greater connectivity within a single HL-T cohort (green) than within a single hemilineage (magenta).

(F) Summary.

Figure 9: Hemilineage-temporal cohorts assemble a proprioceptive circuit

(A) The EL proprioceptive circuit (Heckscher et al., 2015), including hemilineage-temporal cohort membership. Proprio, proprioceptive neurons (e.g. dbd); MNs, motor neurons.

(B) Average cortex-neuron length for the nine interneurons shown in (A) in the left and right hemisegments; all are in the late-born groups 3 and 4, colored orange and red respectively (Supplemental Table 1).

(C) Developmental origin of the Eve+ proprioceptive circuit. NB5-2 generates a $Notch^{OFF}$ hemilineage including late-bon Jaam1-3 neurons, and a $Notch^{ON}$ hemilineage including the late-born Saaghi1-3 neurons. NB3-3 undergoes a type 0 division pattern where the neuroblast progeny do not divide and remain $Notch^{OFF}$ (Baumgardt et al., 2014; Wreden et al., 2017), effectively creating a $Notch^{OFF}$ "hemilineage" that includes the late-born A08e1-3 Eve+ neurons. Thus, the proprioceptive circuit shown in (A) is comprised of three interconnected hemilineage-temporal cohorts.

References

Baumgardt M, Karlsson D, Salmani BY, Bivik C, MacDonald RB, Gunnar E, Thor S. 2014. Global programmed switch in neural daughter cell proliferation mode triggered by a temporal gene cascade. *Dev Cell* **30**:192–208. doi:10.1016/j.devcel.2014.06.021

Benito-Sipos J, Estacio-Gomez A, Moris-Sanz M, Baumgardt M, Thor S, Diaz-Benjumea FJ. 2010. A genetic cascade involving klumpfuss, nab and castor specifies the abdominal leucokinergic neurons in the Drosophila CNS. *Development* **137**:3327–36. doi:137/19/3327 [pii] 10.1242/dev.052233

Birkholz O, Rickert C, Nowak J, Coban IC, Technau GM. 2015. Bridging the gap between postembryonic cell lineages and identified embryonic neuroblasts in the ventral nerve cord of Drosophila melanogaster. *Biol Open* **4**:420–34. doi:10.1242/bio.201411072

Bossing T, Udolph G, Doe CQ, Technau GM. 1996. The embryonic central nervous system lineages of Drosophila melanogaster. I. Neuroblast lineages derived from the ventral half of the neuroectoderm. *Dev Biol* **179**:41–64. doi:S0012-1606(96)90240-7 [pii] 10.1006/dbio.1996.0240

Carreira-Rosario A, Zarin AA, Clark MQ, Manning L, Fetter RD, Cardona A, Doe CQ. 2018. MDN brain descending neurons coordinately activate backward and inhibit forward locomotion. *Elife* **7**. doi:10.7554/eLife.38554

Cau E, Ronsin B, Bessiere L, Blader P. 2019. A Notch-mediated, temporal asymmetry in BMP pathway activation promotes photoreceptor subtype diversification. *PLOS Biol* **17**:e2006250. doi:10.1371/journal.pbio.2006250

Clark MQ, McCumsey SJ, Lopez-Darwin S, Heckscher ES, Doe CQ. 2016. Functional Genetic Screen to Identify Interneurons Governing Behaviorally Distinct Aspects of Drosophila Larval Motor Programs. *G3 Bethesda*. doi:10.1534/g3.116.028472

Costa M, Manton JD, Ostrovsky AD, Prohaska S, Jefferis GS. 2016. NBLAST: Rapid, Sensitive Comparison of Neuronal Structure and Construction of Neuron Family Databases. *Neuron* **91**:293–311. doi:10.1016/j.neuron.2016.06.012

Cui X, Doe CQ. 1992. ming is expressed in neuroblast sublineages and regulates gene expression in the Drosophila central nervous system. *Development* **116**:943–52.

Del Barrio MG, Taveira-Marques R, Muroyama Y, Yuk DI, Li S, Wines-Samuelson M, Shen J, Smith HK, Xiang M, Rowitch D, Richardson WD. 2007. A regulatory network involving Foxn4, Mash1 and delta-like 4/Notch1 generates V2a and V2b spinal interneurons from a common progenitor pool. *Development* **134**:3427–36. doi:10.1242/dev.005868

Fekete DM, Perez-Miguelsanz J, Ryder EF, Cepko CL. 1994. Clonal analysis in the chicken retina reveals tangential dispersion of clonally related cells. *Dev Biol* **166**:666–82. doi:10.1006/dbio.1994.1346

Francius C, Hidalgo-Figueroa M, Debrulle S, Pelosi B, Rucchin V, Ronellenfitch K, Panayiotou E, Makrides N, Misra K, Harris A, Hassani H, Schakman O, Parras C, Xiang M, Malas S, Chow RL, Clotman F. 2016. Vsx1 Transiently Defines an Early Intermediate V2 Interneuron Precursor Compartment in the Mouse Developing Spinal Cord. *Front Mol Neurosci* **9**. doi:10.3389/fnmol.2016.00145

Harris A, Masgutova G, Collin A, Toch M, Hidalgo-Figueroa M, Jacob B, Corcoran LM, Francius C, Clotman F. 2019. Onecut Factors and Pou2f2 Regulate the Distribution of V2 Interneurons in the Mouse Developing Spinal Cord. *Front Cell Neurosci* **13**. doi:10.3389/fncel.2019.00184

Harris RM, Pfeiffer BD, Rubin GM, Truman JW. 2015. Neuron hemilineages provide the functional
ground plan for the Drosophila ventral nervous system. *Elife* **4**. doi:10.7554/eLife.04493
Heckscher ES, Long F, Layden MJ, Chuang CH, Manning L, Richart J, Pearson JC, Crews ST, Peng
H, Myers E, Doe CQ. 2014. Atlas-builder software and the eNeuro atlas: resources for
developmental biology and neuroscience. *Development* **141**:2524–32.
doi:10.1242/dev.108720
Heckscher ES, Zarin AA, Faumont S, Clark MQ, Manning L, Fushiki A, Schneider-Mizell CM,
Fetter RD, Truman JW, Zwart MF, Landgraf M, Cardona A, Lockery SR, Doe CQ. 2015.
Even-Skipped(+) Interneurons Are Core Components of a Sensorimotor Circuit that
Maintains Left-Right Symmetric Muscle Contraction Amplitude. *Neuron* **88**:314–29.
doi:10.1016/j.neuron.2015.09.009
Isshiki T, Pearson B, Holbrook S, Doe CQ. 2001. Drosophila neuroblasts sequentially express
transcription factors which specify the temporal identity of their neuronal progeny. *Cell*
106:511–21. doi:S0092-8674(01)00465-2 [pii]
Jovanic T, Schneider-Mizell CM, Shao M, Masson JB, Denisov G, Fetter RD, Mensh BD, Truman
JW, Cardona A, Zlatic M. 2016. Competitive Disinhibition Mediates Behavioral Choice and
Sequences in Drosophila. *Cell* **167**:858-870.e19. doi:10.1016/j.cell.2016.09.009
Kambadur R, Koizumi K, Stivers C, Nagle J, Poole SJ, Odenwald WF. 1998. Regulation of POU
genes by castor and hunchback establishes layered compartments in the Drosophila CNS.
Genes Dev **12**:246–60.
Kohwi M, Doe CQ. 2013. Temporal fate specification and neural progenitor competence during
development. *Nat Rev Neurosci* **14**:823–38.
Kolodkin AL, Tessier-Lavigne M. 2011. Mechanisms and molecules of neuronal wiring: a primer.
Cold Spring Harb Perspect Biol **3**. doi:10.1101/cshperspect.a001727
Kondo S, Ueda R. 2013. Highly improved gene targeting by germline-specific Cas9 expression in
Drosophila. *Genetics* **195**:715–21. doi:10.1534/genetics.113.156737
Kulkarni A, Ertekin D, Lee CH, Hummel T. 2016. Birth order dependent growth cone segregation
determines synaptic layer identity in the Drosophila visual system. *Elife* **5**:e13715.
doi:10.7554/eLife.13715
Lacin H, Chen H-M, Long X, Singer RH, Lee T, Truman JW. 2019. Neurotransmitter identity is
acquired in a lineage-restricted manner in the Drosophila CNS. *Elife* **8**:e43701.
doi:10.7554/eLife.43701
Lacin H, Truman JW. 2016. Lineage mapping identifies molecular and architectural similarities
between the larval and adult Drosophila central nervous system. *Elife* **5**:e13399.
doi:10.7554/eLife.13399
Landgraf M, Sanchez-Soriano N, Technau GM, Urban J, Prokop A. 2003. Charting the Drosophila
neuropile: a strategy for the standardised characterisation of genetically amenable neurites.
Dev Biol **260**:207–25.
Lee Y-J, Yang C-P, Miyares RL, Huang Y-F, He Y, Ren Q, Chen H-M, Kawase T, Ito M, Otsuna H,
Sugino K, Aso Y, Ito K, Lee T. 2020. Conservation and divergence of related neuronal
lineages in the Drosophila central brain. *eLife* **9**. doi:10.7554/eLife.53518
Mauss A, Tripodi M, Evers JF, Landgraf M. 2009. Midline signalling systems direct the formation
of a neural map by dendritic targeting in the Drosophila motor system. *PLOS Biol*
7:e1000200. doi:10.1371/journal.pbio.1000200
Meng JL, Heckscher ES. 2020. Development of motor circuits: From neuronal stem cells and
neuronal diversity to motor circuit assembly. *Curr Top Dev Biol* **in press**.

Mihalas AB, Hevner RF. 2018. Clonal analysis reveals laminar fate multipotency and daughter cell apoptosis of mouse cortical intermediate progenitors. *Development* **145**. doi:10.1242/dev.164335

Nern A, Pfeiffer BD, Rubin GM. 2015. Optimized tools for multicolor stochastic labeling reveal diverse stereotyped cell arrangements in the fly visual system. *Proc Natl Acad Sci U A* **112**:E2967-76. doi:10.1073/pnas.1506763112

Ohyama T, Schneider-Mizell CM, Fetter RD, Aleman JV, Franconville R, Rivera-Alba M, Mensh BD, Branson KM, Simpson JH, Truman JW, Cardona A, Zlatic M. 2015. A multilevel multimodal circuit enhances action selection in Drosophila. *Nature* **520**:633–9. doi:10.1038/nature14297

Peng CY, Yajima H, Burns CE, Zon LI, Sisodia SS, Pfaff SL, Sharma K. 2007. Notch and MAML signaling drives Scl-dependent interneuron diversity in the spinal cord. *Neuron* **53**:813–27. doi:10.1016/j.neuron.2007.02.019

Petrovic M, Hummel T. 2008. Temporal identity in axonal target layer recognition. *Nature* **456**:800–3. doi:10.1038/nature07407

Pfeiffer BD, Jenett A, Hammonds AS, Ngo TT, Misra S, Murphy C, Scully A, Carlson JW, Wan KH, Laverty TR, Mungall C, Svirskas R, Kadonaga JT, Doe CQ, Eisen MB, Celniker SE, Rubin GM. 2008. Tools for neuroanatomy and neurogenetics in Drosophila. *Proc Natl Acad Sci U A* **105**:9715–20. doi:0803697105 [pii] 10.1073/pnas.0803697105

Port F, Chen HM, Lee T, Bullock SL. 2014. Optimized CRISPR/Cas tools for efficient germline and somatic genome engineering in Drosophila. *Proc Natl Acad Sci U A* **111**:E2967-76. doi:10.1073/pnas.1405500111

Rees CL, Moradi K, Ascoli GA. 2017. Weighing the Evidence in Peters' Rule: Does Neuronal Morphology Predict Connectivity? *Trends Neurosci* **40**:63–71. doi:10.1016/j.tins.2016.11.007

Rossi AM, Fernandes VM, Desplan C. 2017. Timing temporal transitions during brain development. *Curr Opin Neurobiol* **42**:84–92. doi:10.1016/j.conb.2016.11.010

Sagner A, Briscoe J. 2019. Establishing neuronal diversity in the spinal cord: a time and a place. *Development* **146**. doi:10.1242/dev.182154

Schlegel P, Texada MJ, Miroschnikow A, Schoofs A, Huckesfeld S, Peters M, Schneider-Mizell CM, Lacin H, Li F, Fetter RD, Truman JW, Cardona A, Pankratz MJ. 2016. Synaptic transmission parallels neuromodulation in a central food-intake circuit. *Elife* **5**. doi:10.7554/eLife.16799

Schmid A, Chiba A, Doe CQ. 1999. Clonal analysis of Drosophila embryonic neuroblasts: neural cell types, axon projections and muscle targets. *Development* **126**:4653–89.

Schmidt H, Rickert C, Bossing T, Vef O, Urban J, Technau GM. 1997. The embryonic central nervous system lineages of Drosophila melanogaster. II. Neuroblast lineages derived from the dorsal part of the neuroectoderm. *Dev Biol* **189**:186–204. doi:S0012160697986607 [pii]

Seroka AQ, Doe CQ. 2019. The Hunchback temporal transcription factor determines motor neuron axon and dendrite targeting in Drosophila. *Development* **146**:dev175570. doi:10.1242/dev.175570

Skaggs K, Martin DM, Novitch BG. 2011. Regulation of spinal interneuron development by the Olig-related protein Bhlhb5 and Notch signaling. *Development* **138**:3199–211. doi:10.1242/dev.057281

Skeath JB, Thor S. 2003. Genetic control of Drosophila nerve cord development. *Curr Opin Neurobiol* **13**:8–15. doi:S0959438803000072 [pii]

Syed MH, Mark B, Doe CQ. 2017. Steroid hormone induction of temporal gene expression in Drosophila brain neuroblasts generates neuronal and glial diversity. *Elife* **6**. doi:10.7554/eLife.26287

Truman JW, Moats W, Altman J, Marin EC, Williams DW. 2010. Role of Notch signaling in establishing the hemilineages of secondary neurons in Drosophila melanogaster. *Development* **137**:53–61. doi:10.1242/dev.041749

Tsuji T, Hasegawa E, Isshiki T. 2008. Neuroblast entry into quiescence is regulated intrinsically by the combined action of spatial Hox proteins and temporal identity factors. *Development* **135**:3859–69. doi:10.1242/dev.025189

Wong LL, Rapaport DH. 2009. Defining retinal progenitor cell competence in Xenopus laevis by clonal analysis. *Development* **136**:1707–15. doi:136/10/1707 [pii] 10.1242/dev.027607

Wreden CC, Meng JL, Feng W, Chi W, Marshall ZD, Heckscher ES. 2017. Temporal Cohorts of Lineage-Related Neurons Perform Analogous Functions in Distinct Sensorimotor Circuits. *Curr Biol* **27**:1521-1528.e4. doi:10.1016/j.cub.2017.04.024

Yoshikawa S, McKinnon RD, Kokel M, Thomas JB. 2003. Wnt-mediated axon guidance via the Drosophila Derailed receptor. *Nature* **422**:583–8. doi:10.1038/nature01522

Yu YC, Bultje RS, Wang X, Shi SH. 2009. Specific synapses develop preferentially among sister excitatory neurons in the neocortex. *Nature* **458**:501–4. doi:10.1038/nature07722

Zarin AA, Mark B, Cardona A, Litwin-Kumar A, Doe CQ. 2019. A multilayer circuit architecture for the generation of distinct locomotor behaviors in Drosophila. *Elife* **8**. doi:10.7554/eLife.51781

Zipursky SL, Grueber WB. 2013. The molecular basis of self-avoidance. *Annu Rev Neurosci* **36**:547–68. doi:10.1146/annurev-neuro-062111-150414

Zlatic M, Li F, Strigini M, Grueber W, Bate M. 2009. Positional cues in the Drosophila nerve cord: semaphorins pattern the dorso-ventral axis. *PLOS Biol* **7**:e1000135. doi:10.1371/journal.pbio.1000135

Figure 1

A **Motor inputs** **Sensory outputs**

B **Pre-motor inputs** **Post-sensory outputs**

C **Combined Synaptic Density**
Pre-motor **Post-sensory**

D NBLAST Embryonic Hemilineages Summary
NB1-2 N^OFF N^ON

E NB7-1 N^OFF N^ON

F NB5-2

G NB4-1

H NBLAST Embryonic Hemilineages Summary
NB7-4 N^OFF N^ON
Glia

I NB3-3 N^OFF

J **Dorsal Hemilineages**
Ventral Hemilineages

Figure 2

A

Figure 2 - figure supplement 1

 Figure 2 - figure supplement 2

Figure 3

Figure 3 - figure supplement 1

Figure 4

Figure 4 - figure supplement 1

Figure 5

Figure 5 - figure supplement 1

Figure 5 - figure supplement 2

Figure 5 -- figure supplement 3

Figure 6 - supplement 1. Presynapse distribution

Figure 6 - supplement 2. Postsynapse distribution

Figure 7

Figure 8

Figure 9