Fitness-related traits are maximized in recently introduced, slow-growing populations of an invasive clam: is this a response to strong *r*-selection?

Leandro A. Hünicken[1,2]*, Francisco Sylvester[2,3], Nicolás Bonel[2,4,5]

[1] Museo Argentino de Ciencias Naturales 'Bernardino Rivadavia', Av. Angel Gallardo 470 (C1405DJR), Buenos Aires, Argentina. Present Address: Centro de Investigación Aplicada y Transferencia Tecnológica en Recursos Marinos ''Almirante Storni'', Güemes 1030, San Antonio Oeste, R8520CXV Río Negro, Argentina

[2] Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

[3] Instituto para el estudio de la Biodiversidad de Invertebrados, Facultad de Ciencias Naturales, Universidad Nacional de Salta, Av. Bolivia 5150 (4400), Salta, Argentina.

[4] Laboratorio de Zoología de Invertebrados I, Departamento de Biología, Bioquímica y Farmacia, Universidad Nacional del Sur (UNS), San Juan 670, (B8000ICN) Bahía Blanca, Argentina.

[5] Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175, CNRS—Université de Montpellier, Université Paul-Valéry Montpellier—Ecole Pratique des Hautes Etudes—IRD, 34293 Montpellier cedex 05, France.

* Corresponding author. E-mail: leandrohunicken@gmail.com

Key words: Asian clam, *Corbicula fluminea*, Invasive species, Individual growth rates, Age at first reproduction, Front-edge populations, *r*-selection.

Author Contributions: LAH, FS, and NB conceived the study. LAH, FS, and NB conducted the literature search and extracted the data. NB designed the methodological approach. LAH and NB performed the data analysis. NB led the writing.

ABSTRACT

Many species are shifting their ranges being forced to rapidly respond to novel stressful environmental conditions. Colonizing individuals experience strong selective forces that favor the expression of life history traits notably affecting dispersal and reproductive rates in newly invaded habitats. Limited information is currently available on trait variation within the invasive range despite being critical for understanding ecological and evolutionary factors that drive the process of range expansion of invasive species. Here we evaluated life history shifts of the widely introduced Asian clam *Corbicula* within its invaded range. Through an exhaustive literature search, we obtained data for 17 invasive *Corbicula* populations from different ecosystems worldwide to test the relationship between population and individual parameters relevant to the process of range expansion. Our main results show that (i) recently introduced *Corbicula* populations are characterized by low conspecific density and low rate of population increase, (ii) clams reproduce earlier in slow-growing populations, and (iii) density had no effect on exponential population increase. All invasive *Corbicula* populations analyzed in this study (Form A/R) are mostly fixed for one genotype. Our results therefore suggest that plasticity favored the expression of traits that maximize fitness in recently established populations facing stronger *r*-selective forces, thereby increasing their chances to overcome difficulties associated with low densities and low population increase in newly invaded areas.

INTRODUCTION

The ongoing increase in frequency of extreme climatic events, caused by anthropogenic climate change, and global transport of people and goods is altering ecosystems by triggering a wide array of irreversible consequences (Walther et al. 2002; Bossdorf et al. 2005; Hoegh-Guldberg and Bruno 2010; IPCC 2014). Changes in local conditions, in particular those driven by human-induced climate change, are predicted to affect ecological processes across several levels of organization, likely over the coming decades. Such changes are altering the performance of individual organisms, the dynamics of populations and the distribution of species, ultimately modifying ecosystem properties and function (Hoffmann and Sgró 2011; Lonhart et al. 2019).

Many species are forced to rapidly respond to novel stressful environmental conditions when shifting their distribution and colonizing new areas. If non-native species have difficulties in adapting to such conditions and fail to track abrupt environmental changes, populations become vulnerable to decline and extinction (Hill et al. 2011; Hoffmann and Sgró 2011). Phenotypic plasticity is the ability of a single genotype to express different phenotypes when exposed to different environmental conditions (Travis 1994; West-Eberhard 2003). In a new environment, plasticity rapidly favors the appearance of a novel phenotypic variant, promoting successful colonization and increasing the likelihood for the long-term persistence of populations facing new environmental stress brought about by range shifting (Pigliucci et al. 2006; Pfennig et al. 2010; Hill et al. 2011). Such stressful conditions can have, for instance, a strong effect on species life histories by favoring the expression of traits that optimize population growth and survival from offspring to maturity (e. g. reduced age of maturity and reproduction, increased growth rate, increased energy allocated to reproduction; Stearns, 1976, 1977). Thus, individuals at the expanding edge of the species distribution experience strong selective forces that can have profound impacts on the evolution of a species' life history traits affecting, for instance, dispersal and reproductive rates (Phillips 2009; Shine et al. 2011).

Colonization events through either continuous range expansion or jump dispersal are usually driven by a small number of individuals that are the first to arrive at a new point of the distribution, where they benefit from increased space open for colonization and

low conspecific density, below carrying capacity (Cole 1954; Sakai et al. 2001; Phillips

et al. 2010). Hence, the density gradient occurring at the expanding range (i. e. grading

from zero to carrying capacity) creates different selective environments where

individuals from a recently established population face stronger r-selection compared to

those from long established ones, which are exposed to selective K-type forces (e. g.

Phillips et al., 2010). According to theory, low densities favor the evolution of life-

history traits related with r-type selections (MacArthur and Wilson 1967; Phillips 2009;

Burton et al. 2010), leading to exponential (non-density regulated) population growth in

recently established populations while denser, long-established ones follow a density-

regulated population growth (Brook and Bradshaw 2006).

Recently established populations are characterized by low conspecific density and low

rate of increase, often leading to lag-times between initial colonization and the onset of

rapid population growth (Sakai et al. 2001). The duration of the lag phase is mediated,

among others, by a number of non-mutually exclusive ecological and evolutionary

mechanisms (e. g. low encounter rate for reproduction, density-dependent effects,

adaptation, and selection of new genotypes; Bossdorf et al., 2005; Davis, 2005;

Ricciardi, 2013). An effective mechanism that increases the rate of population growth is

early reproduction, which results from fast growing individuals (Cole 1954; Lewontin

1965; Roff 1993). Individuals that grow faster will attain reproductive sizes earlier and

bring down generation times, promoting a rapid build-up of population numbers (Cole

1954; Lewontin 1965; Roff 1993; Phillips 2009) and accelerating primary invasion and

secondary spread of invasive species (Skellam 1951; Phillips et al. 2006; Lockwood et

al. 2007). Several studies have compared life-history traits between native and invaded

ranges ('home-and-away' comparisons). Such comparisons could lead to erroneous

interpretations of the patterns observed because it would be difficult to disentangle what

were the processes (e.g. propagule bias and/or evolution in populations at spatial

(dis)equilibrium) causing the shifts in life-history traits (Phillips et al., 2010 and

references therein). Interestingly, only a few studies have looked for such trait changes

on introduced species (Gaston 2009) despite knowledge on life-history trait variation

within the invasive range being critical for better understanding the process of species

range expansion, largely induced by climate change and global transport.

Bivalves of the genus *Corbicula* (Megerle Von Mühlfeld 1811) are native to Southeast

Asia, the Middle East, Australia and Africa (Araujo et al. 1993) but they have colonized

a large part of the Americas and Europe (Mouthon 2001; Schmidlin and Baur 2007;

Crespo et al. 2015). The successful invasion of *Corbicula* clams has been mainly

attributed to their rapid growth and maturation, high fecundity, and high dispersal

making this bivalve genus one of the most relevant faunal non-indigenous groups in

aquatic ecosystems (reviewed in Sousa et al. 2008a). Invasive freshwater *Corbicula*

populations are constituted by a small number of lineages composed of hermaphroditic

individuals, which reproduce through androgenesis, where the most widespread and

abundant lineage is the so-called Form A/R, widely known as *C. fluminea* (Komaru et

al. 1998; Pigneur et al. 2012, 2014). The current taxonomic status of the invasive

Corbicula lineages in the Americas and Europe is still largely unresolved despite

several morphological and genetic studies (reviewed in Pigneur et al. 2014). In the

present paper, we deal with this form and refer to it as *Corbicula* to avoid entering

unsettled taxonomic debates.

Here we investigated how different life-history traits of the Asian clam *Corbicula*

responded to different selective pressures occurring at varying population contexts

within the introduced range: (i) newly- vs. long-established, (ii) low vs. high density,

and (iii) low vs. high rates of population increase; all of them being linked to the

geographic range shifting of species' distribution (i. e. front/core populations). To do so,

we searched for peer-reviewed articles providing enough information to estimate time

since population introduction as well as different population and individual processes

and characteristics including population density, individual growth rate, minimum age

at sexual maturity, and lifespan of global populations of this worldwide invader.

Different studies suggest that *Corbicula*'s population dynamics are chiefly regulated by

environmental conditions (e.g. Crespo et al., 2015, 2017; Gama et al., 2017), whereby

population declines are typically driven by periodic mass mortality due to extreme

hydro-meteorological events (Ilarri et al. 2011; McDowell et al. 2016; McDowell and

Sousa 2019). This strongly suggests that invasive *Corbicula* populations would

primarily follow a non-density regulated population increase and, hence, they might not

level off at carrying capacity as happens with many other species (e.g. Brook and

Bradshaw, 2006). We focused our predictions on life history strategies exhibited by

individuals from recently established populations, which typically occur at the invasion

front, and likely subjected to strong *r*-selective pressures with respect to those from

older, long-established ones. We tested the hypotheses that population density and

population increase are directly related with time since introduction. *Corbicula* is

widely known to exhibit rapid population growth, being able to reach extremely high

densities (reviewed in McMahon, 2002). We expected density to have no significant

effect on population increase even within a wide range of densities in nature, though

high densities are considered likely to have a negative effect on individual growth rates.

We also asked whether clams that grow faster and reproduce earlier are favored in slow-

growing populations at the expanding front. A related prediction is that clams that grow

rapidly would have a shorter lifespan with respect to individuals exhibiting slow

growth.

METHODS

Literature search

We exhaustively searched the literature for peer-reviewed articles on *Corbicula*

(*Corbicula fluminea sensu lato*) providing enough information to estimate the

individual growth rate, population density, rate of exponential population increase,

time since colonization, water temperature, and conductivity of global populations of

this clam. To do this, we first identified relevant studies from ISI Web of Knowledge

(Web of Science Core Collection, 1900 through 2017) by conducting a first search

using the field tag "topic (TS)" in the "Advanced search" option and the search string:

TS="Corbicula fluminea" OR TS="asia* clam" (search #1). We conducted a second

search including topics TS="population dynamic*" OR TS="population size*" OR

TS="population densit*" OR TS="population structure" (search #2). We ran a third

search using TS="growth" OR TS="individual growth" OR TS="population growth"

(search #3). Finally, we combined searches #2 and #3 into search #4 (#4= #3 OR #2),

and restricted the latter by search #1 to obtain a final set of candidate studies (search

#5= #4 AND #1).

This literature search was finished on 27 September 2018 and this protocol identified

215 candidate studies (database search: 215 candidates; gray literature and unpublished

data: 0 candidates). To identify papers potentially missing in this list, we subsequently

scanned the references in these papers and conducted a Google Scholar search using

keywords mentioned above and added 8 studies to this candidate pool (Fig. S1). A total

of 67 studies that belonged to research areas unrelated to the topic of the current study

(*e. g.*, applied chemistry, toxicology, paleontology) were excluded from this candidate

pool by refining the set based in 'Web of Science Categories'. The resulting studies (n =

156) were finally inspected individually in detail to assess their eligibility for our

analyses according to the following inclusion criteria: i) Studies that assessed invasive

populations of *Corbicula* clams (i. e. studies exclusively providing estimates for native

populations were excluded); ii) Studies that reported or provided data enough to extract

the population density and age/size distribution for >6-months (see below); iii) Studies

that assessed populations in reasonably "average" natural environments, such that

populations from heavily impacted sites (e. g. thermal plumes) were excluded. We

finally retained a sample of 16 studies, covering 17 invasive populations of *Corbicula*

from different ecosystems worldwide (n = 140 out of 156 were excluded; see Fig. S1 for

further details on screening and eligibility criteria). Whenever population data was only

graphically presented, we contacted directly the authors. However, if data were no

longer available or authors did not respond, we used the WebPlotDigitizer software

(http://arohatgi.info/WebPlotDigitizer/) to extract data and accurately estimate

population density and growth rates. This step-by-step criterion for the literature search

and data collection attempted to reduce any potential selection bias in the long-term

populations that might occur because old high-density populations (with higher growth

rates) could be more studied than young low-density populations (with low growth

rates), which would be more likely to be extirpated if population characteristics were

not changing with time.

Population density and exponential population increase

For each population selected from our literature search, we obtained the mean

population density for the entire study period or extracted temporal values to estimate

it whenever this value was missing in the text. *Corbicula* clams generally exhibit a

bivoltine juvenile release pattern (i.e. producing two generations a year; McMahon,

2002; Sousa et al., 2008a), leading to one or more high-density peaks related to

recruitment events yearly. However, density in the field fluctuates almost constantly

generating several peaks and valleys that do not necessarily indicate significant

recruitment events. *Corbicula* populations exhibit rapid growth (McMahon 2002;

Sousa et al. 2006, 2008a; Franco et al. 2012). Hence, the rate of exponential

population increase of *Corbicula* can be expressed by deriving the classical

exponential growth equation:

$$r = \frac{\ln\left(\frac{Nt}{N0}\right)}{t}$$

One of the goals of this study was to compare population growth rate during periods of

substantial growth across populations, and it is such capacity that is of interest to better

understand *Corbicula* invasiveness. Hence, to estimate the rate of exponential

population increase (*r*, hereafter population growth rate), we set a criterion to

determine relevant recruitment peaks in each study by using the average density

throughout the study period as a threshold. By doing so, we excluded time periods

where population growth was extremely low likely due to mass mortality due to

extreme hydro-meteorological events (e.g. a period of dramatic demographic decline),

usually characterized by a lag phase (where population growth can be negligible)

followed by an exponential increase (Ricciardi 2013). Likewise, this approach allows

for excluding large windows of the time period resulting from *Corbicula*'s inability to

reproduce year-round as they generally exhibit a uni- or bivoltine juvenile release. We

therefore considered the lowest density value (N_0), the highest peak above the average

density (N_t), and the time period elapsed between the valley and the peak (*t*).

Intermediate density values comprised in each period (i. e. defined by N_0 and N_t) were

used to fit the growth equation and obtain *r*. Population rates of increase were

estimated using GraphPad Prism version 7.00 for Windows, GraphPad Software, La

Jolla California USA, www.graphpad.com. Note that periods defined by non-relevant

peaks (i. e. below the average density threshold) were discarded, as were also periods

of progressive density decline between a relevant peak and the next N_0 value (see Fig.

S2 for further details on the step-by-step procedure to estimate population increase).

When populations exhibited more than one relevant density peak throughout the study

period, the population increase rate for such population was calculated as the average

of single *r* estimations. Then, we standardized estimates on an annual basis to make

them comparable.

Individual growth rates and parameters

For studies that only presented data on shell length-frequency distributions (shell

length being the greatest distance between the anterior and posterior margins of the

shell valve), we identified age cohorts using that information and applied the

Bhattacharya's method available in FISAT II software (Version 1.2.0, FAO-ICLARM

Fish Assessment Tools; (Gayanilo et al. 2002). To confirm each component of normal

distributions from the modal progression analysis, we used the NORMSEP method

also available in the FISAT II software (Pauly and Caddy 1985). Once we had cohorts

identified for all the studies, we calculated individual growth parameters using the

same equations for all the studies considered in order to make sure that results were

fully comparable across studies. For each cohort present in each of the populations

considered in this study, we fitted the von Bertalanffy-growth function with seasonal

oscillations (hereafter SVBGF) proposed by Pauly and Gaschutz (1979), Hoenig and

Hanumara (1982), and Somers (1988):

$$L_t = L_\infty \, (1 - exp^{-K\,[(t - t_0) + T_1 - T_2]}),$$

$$T_1 = C \sin \left(2\pi \, (t - t_s)\right) / \, 2\pi,$$

and

$$T_2 = C \sin \left(2\pi \, (t_0 - t_s)\right) / \, 2\pi,$$

where L_t is the predicted shell length at age t, L_∞ is the asymptotic shell length, K is

the growth constant of dimension time (year^{-1} in most seasonally oscillating growth

curves) expressing the rate at which L_∞ is approached, t_0 is the theoretical 'age' the

clam have at shell length zero, C expresses the relative amplitude of the seasonal

oscillation and varies between 0 and 1 (0 indicating lack of summer-winter differences

in growth, it was constrained to be less than or equal to 1 during model fitting), and t_s

is the starting point of the oscillation.

The parameters of the function were estimated by the modeling method available in

JMP statistical software (v9.0 SAS Institute). When preliminary results of the SVBGF

failed to converge to estimate the asymptotic shell length (L_∞), we used the maximum

shell length (*Lmax*) observed or reported at/for each study site to calculate the

asymptotic shell length following the equation suggested by Taylor (1958):

$$Lmax \, / \, 0.95 = L_\infty$$

These values were fixed when we performed a second fit.

Because of the negative correlation between growth parameters (K and L_∞) prevents

making comparisons based on individual parameters (Vakily 1992; Ramón et al. 2007),

we used the growth-performance index (*GPI*), which reflects the growth rate of a given

organism of unit length. In other words, *GPI* can be viewed as the (theoretical) value of

K that would occur in organisms with a L_∞ value of 1 unit of length (Munro and Pauly

1983), and it was defined by Pauly and Munro (1984) as:

$$GPI = 2 \log_{10} L_\infty + \log_{10} K$$

We calculated the *GPI* values for cohorts from each study site based on growth

parameters (L_∞ and *K*) obtained after fitting the SVBGF using the *Growth*

Performance Indices application of the FISAT II software (Gayanilo et al. 2002).

Minimum age at sexual maturity

The minimum age at sexual maturity was estimated as the time (months) elapsed until

clams reach the minimum size (shell length) at which invasive *Corbicula* clams become

sexually mature, which, on average, is 10.0 mm ± 0.8 SE (n = 13 reported sizes). This

value was obtained from shell sizes reported in the literature (Heinsohn 1958; Gardner

et al. 1976; Kraemer and Lott 1977; Aldridge and McMahon 1978; Eng 1979; Kraemer

1979; Ituarte 1985; Kennedy and Van Huekelem 1985; French and Schloesser 1991;

Rajagopal et al. 2000; McMahon 2002; Cao et al. 2017). We acknowledge that in some

conditions and populations, clams might mature at a different pace and thus introduce a

degree of error in such cases. While this error is unavoidable until more detailed data is

available, we are confident that the 10-mm estimate applies to many real-life situations

and thus our results based on it are robust.

Lifespan

As *t0* is the theoretical 'age' the clam have at shell length zero, *tmax* can be defined as

the theoretical 'age' the clam reaches its maximum shell length (*Lmax*). Since *Lmax* is

dependently related to a maximum point in time (years), it is assumed that *Lmax*

implicates *tmax* (*Lmax* $\Rightarrow$ *tmax*) (Bonel and Lorda 2015). Thus, the theoretical

maximum lifespan of a clam cohort can be approximately expressed as:

$$Lifespan = tmax - t0$$

Time since colonization and key environmental variables

For the same 17 populations and sites studied, we calculated the time elapsed since

population foundation (hereafter time since colonization) as the difference between the sampling dates in the first ecological survey and the first report of introduction. The accuracy of the introduction date is generally very difficult to know, however, many of the studies used report range expansions of a conspicuous invader across well studied regions, where the appearing of the species might have been anticipated and readily detected by ongoing monitoring actions in the initial stages of invasion. Therefore, we consider that the possible error introduced should be moderate to low and the value obtained can be viewed as a good approximation. Moreover, we obtained water temperature and conductivity values for each site averaging monthly mean values for the same period used to estimate growth rates in each study. Other biotic and abiotic variables are known to interact with temperature to drive growth rates of invasive bivalves (e.g. food availability, dissolved oxygen, heavy metal concentration; e.g. (Vohmann et al. 2010; Bonel and Lorda 2015). However, we only considered water temperature and conductivity in the analyses because they have been reported to play a key role on the *Corbicula*'s biology and ecology (Magnussen 2007; Franco et al. 2012; Crespo et al. 2017). Indeed, temperature has emerged as the most important variable explaining the current distribution and, also, predicting that climate change will favor the expansion of *Corbicula*'s into colder areas at higher latitudes (Gama et al. 2017). Likewise, salinity has been considered a key factor influencing the success and velocity of the invasion of new estuarine ecosystems (e.g. Sousa et al., 2006). As *Corbicula* populations considered herein belonged to a broad range of freshwater and estuarine habitats, we therefore included conductivity, which serve as a *proxy* of salinity levels. In order to keep values seasonally unbiased, only entire years were used in subsequent calculations. Whenever these parameters were not reported in the original paper, we obtained values for the same sites and years from the literature and reliable online databases (e. g. https://waterdata.usgs.gov/nwis/qw). When data was only graphically presented, we extracted it following the procedure mentioned above.

Statistical analyses

We ran five separate multivariate weighted linear regressions in order to test ecological processes and characteristics at three different (but non-mutually exclusive) levels: 1) population level, 2) population and individual level, and 3) individual level. In this sense, a response variable at the population level can be considered as an explanatory variable when analyzing a response at the individual level. In all cases, we computed the

variance of the effect size from different populations, which were then used as weights in linear regressions. The inverse-variance weighting allows combining the results from independent measurements giving more importance to the least noisy measurements and *vice versa* (Borenstein et al. 2009).

We were interested in testing individual effects of predictors, which are related to life-history traits of *Corbicula* (hereafter reduced models). However, we added annual mean water temperature and conductivity as quantitative variables in each model (hereafter full models). Preliminary analyses showed that latitudinal effects are strongly correlated with water temperature ($r = -0.84$). Hence, it was excluded from the models to avoid multicollinearity. In some cases, we also included interaction terms between explanatory variables. We used the partial *F*-test to compare models—that is, full models with and without the interaction term, and full *versus* reduced models. As we had strong *a priori* hypotheses as to the direction of effects, we used one-tailed tests. Variables were Ln-transformed to fulfill the models' assumption of normality and homoscedasticity, which were tested using the *gvlma* package (Peña and Slate 2006). Prior to analyses, we standardized explanatory variables to zero mean and a unit of variance in order to get directly comparable effects. All analyses were carried out in R version 3.6.1 (R Development Core Team 2019). Values are given as means ± SE unless otherwise stated.

RESULTS

Literature search

Populations analyzed in this study belonged to Europe, North and South America (Fig. 1; Table 1). These included eight populations in the United States, three in France, three in Argentina, and three in Portugal (see Table 1 for detailed location, bibliographic source, and code number used herein to refer to the populations, for each of the 17 *Corbicula* populations selected, and Table S1 for additional information).

Density, growth, time since colonization, and key environmental variables

Overall, mean density of *Corbicula* from 17 locations worldwide spanned 32 to *ca.* 9,500 ind. m^{-2}, population growth rate (*r*) varied from 0.37 to 189, individual growth rate (*GPI*) ranged 2.53-3.49, age at first reproduction 3.1-22.7 months, and lifespan 2.3-8.0 years (Table 1). On the other hand, time since colonization ranged 2 to 28 years in

the populations studied (Table 1) and yearly water temperature ranged from 13.2 to 20.7

ºC and conductivity from 68 to 8,457 µS/cm (see Table S1 of the Supporting

Information for details). In all cases, we found no evidence that adding an interaction

term with variables of interest nor temperature and conductivity in each model

significantly increased its predictive power (for details see '*Model comparison*' in

Supporting Information). We therefore report estimates and statistics from the reduced

models in Table 2.

At the population level, we found that younger *Corbicula* populations (i.e. newly

established) were characterized by low density and slow population growth rate with

respect to older, long established ones (Fig. 2a, b). We found that density had no

significant effect on population increase (Fig. 2c). When analyzing responses at the

interaction between population and individual processes and characteristics, we found

that clams tended to grow faster (marginally non-significant) in low-density and slow-

growing populations (Fig. 2d, e). In contrast, we observed a strong positive response of

the minimum age at sexual maturity as a function of population growth rate, clearly

indicating that clams reproduce earlier in slow-growing populations (Fig. 2f). This is

directly linked to processes occurring at the individual level, where clams with higher

growth rates reach the minimum reproductive size earlier (10 mm) and have shorter life

span than clams that grow more slowly (Fig. 2g, h).

DISCUSSION

This is the first study that evaluates how population and individual parameters, involved

in the process of range expansion, varied in response to different selective pressures

occurring within the introduced range of the worldwide invasive Asian clam *Corbicula*.

Our main results showed that (i) recently introduced *Corbicula* populations are

characterized by low density and reduced population growth rate, (ii) clams reproduce

earlier in slow-growing populations, and (iii) population growth rate was unfettered by

density (summarized in Fig. 3).

Density of *Corbicula* populations increased with time since colonization, which is

consistent with the standpoint that recently established populations typically occur at

low conspecific density, below the carrying capacity (Cole 1954; Sakai et al. 2001).

Likewise, we found that population growth rate and time since colonization were directly related, suggesting that, in newly colonized areas, populations could have experienced stronger *r*-type selection favoring fitness-related traits effectively increasing the rate of population growth, as it was observed in long established populations (MacArthur and Wilson, 1967; Phillips, 2009; Burton et al., 2010). Population density did not predict changes in the rate of population increase, as variation in population growth remained relatively low even at high-density values. By definition, long established and denser populations tend to be density regulated as they approach to carrying capacity (logistic population growth; e. g. Phillips et al. 2010). Considering the so-called *r*/K selection continuum, this implies that such populations are predicted to shift from the *r*- towards the K-endpoint (MacArthur and Wilson 1967)—that is, selection shifts a population from density-independent to density-dependent regulation. However, *Corbicula*'s population growth rate did not decrease with increasing density despite the fact that some of the populations included in this study exhibit extremely high densities exceeding, in some cases, 100,000 individuals per square meter (e. g. Eng 1979; Table 1).

Why was population growth unfettered by density regulation? A large number of studies have analyzed density dynamics of *Corbicula* and commonly assumed that, following exponential growth, the clam population achieves densities approaching the carrying capacity of the environment (e.g. Sousa et al., 2006). However, to our knowledge, no studies have ever conducted (field or laboratory) experiments to test whether population growth is limited by increasing conspecific density, which, in turn, is expected to reduce local biotic resources. In other words, there is a complete lack of knowledge on the true effect of density on population growth rate. On the contrary, different studies have shown that *Corbicula*'s dynamics are chiefly influenced by environmental variables (e. g. water temperature, conductivity, pH, and/or dissolved oxygen) and seem to be less influenced by population's density (e. g. Crespo et al., 2015; Gama et al., 2016). Indeed, periodic mass mortality due to extreme hydro-meteorological events (e.g. extremely high or low temperatures, drought) could play an important role with infrequent, but catastrophic population declines (>99% in some cases; Mouthon and Parghentanian 2004; Ilarri et al. 2011; McDowell et al. 2017). These events have been described (though not quantified) in several other systems as well (e.g. Vaughn et al. 2015; McDowell and Sousa 2019). Despite that a given

Corbicula population might approach or even grow beyond the carrying capacity of the

ecosystem, evidence from previous studies (and those cited herein) suggests that

population size significantly varies with time. This means that the population is at non-

equilibrium and that such fluctuations (or boom-bust dynamics; Strayer et al., 2017) are

likely driven by density-independent factors, as the above-mentioned hydro-

meteorological events. Thus, it does seem reasonable to infer that *Corbicula*'s

population growth rate is non-density-regulated and will not level off at carrying

capacity, as expected from a highly invasive organism as the Asian clam (McMahon

2002). Our results suggest that this bivalve seems to be far from approaching carrying

capacity, even at extremely high densities, supporting the idea that individuals from

long established denser populations do not face a *K*-selective environment but a less

strongly *r*-selective pressure.

Individuals tended to grow faster in low-density populations experiencing low rates of

growth. Likewise, *Corbicula* clams reproduce earlier in slow-growing populations. In

recently introduced populations, periods of negligible population increase can be driven

by non-mutually exclusive ecological and evolutionary adjustments such as Allee effect,

changing abiotic conditions, and adaptation and selection of new genotypes imposed by

novel environments conditions at the newly habitat (Sakai et al. 2001; Bossdorf et al.

2005; Davis 2005; Ricciardi 2013). Such conditions can lead to a strong selection on

life-history traits that optimize population increase and survival to maturity of offspring

(Stearns 1976, 1977). As predicted by life-history theory, individuals from recently

established populations (typically occurring at the invasion front) face stronger *r*-

selection relative to conspecifics from older populations (MacArthur and Wilson 1967),

favoring traits that accelerate the range expansion. For instance, some empirical

observations have revealed that individuals from recently colonized areas grow faster,

and thus, reproduce earlier, than do conspecifics from older, long-established

populations (e.g. tallow trees: Siemann and Rogers 2001; butterflies: Hanski et al. 2006;

Australian cane toads: Phillips 2009). Accordingly, our results for *Corbicula* clams

showing fast growth and, thus, early reproduction support the idea that such populations

faced strong *r*-selection, which favored traits associated with high reproductive rates in

recently introduced, slow-growing populations.

The increase in individual growth is associated with a decrease in the clams' life span

likely due to trade-offs between different life-history traits. Such trade-offs can be

interpreted as an effective mechanism to bring down generation times, promoting a

rapid build-up of population numbers, which increases the population growth in newly

colonized areas (Cole 1954; Lewontin 1965; Roff 1993) and accelerates primary

invasion and secondary spread of invasive species (Skellam 1951; Phillips et al. 2006;

Lockwood et al. 2007). Collectively, our results suggest that reduced population

increase of the current *Corbicula* population triggers the shortening of the time that the

clams need to reach maturity by favoring clams that grow the fastest. Thus, as early-

maturing clams reproduce earlier, this has the effect of fostering an increase in size of

the future population. In this way, incipient *Corbicula* populations could increase their

chances of overcoming the difficulties associated with low densities and survive to the

population establishment stage, which can clearly subsidize the bivalve's invasive

abilities in newly colonized areas.

All invasive *Corbicula* populations analyzed in this study are characterized by an

extremely low genetic diversity and are mostly fixed for one genotype (see Pigneur et

al. 2012, 2014 for further details). The shift in fitness related traits provides a reasonable

ground for advocating that plasticity facilitated the expression of phenotypic variants in

populations experiencing different environmental circumstances. The question that

arises is whether there is scope for adaptive evolution in genetically depleted

populations of *Corbicula*. One fundamental principle of evolutionary biology is that the

rate of change in response to natural selection is proportional to the amount of additive

genetic variation present (Fisher 1930). Decrease in genetic diversity results in reduced

evolutionary potential required to rapidly adapt to the novel habitat conditions of the

invaded range, among other issues related to critical invasion phases (e.g. initial

colonization, leading edges of range expansion; Schrieber and Lachmuth 2017). It is

still puzzling how *Corbicula*'s genetically depleted populations succeed in their invasion

process, which represents a case of genetic paradox (Pigneur et al. 2014). One possible

explanation is that invasive *Corbicula* might exhibit a certain degree of genetic

polymorphism linked to androgenesis (Pigneur et al. 2012). Indeed, androgenesis in

Corbicula clams combines features of clonal reproduction and the ability of rare genetic

material exchange through 'egg parasitism'. Among other processes, egg parasitism

enables a mixing of different nuclear genomes when the maternal nuclear genome is

incompletely extruded creating triploid individuals (invasive freshwater *Corbicula*

clams produce unreduced spermatozoa; i.e. sperm cells are diploids; Pigneur et al.

2014).

The invasive lineage (Form A/R) considered in our study includes both diploid and

triploid populations, with triploidy being predominant (further details in Pigneur et al.,

2014 and references therein; Etoundi et al., 2019). As polyploidy can buffer deleterious

mutations and can lead to hybrid vigor (heterosis) with increased fitness (e.g. Otto 2007;

Selmecki et al. 2015), polyploid *Corbicula* clams with genotypes associated with broad

tolerance to environmental stressful conditions (usually found at the invasion front) may

be favored during an invasion process, and in the long term a general-purpose genotype

may evolve (Pigneur et al. 2012). Therefore, evolved adaptation to local stressful

conditions in newly colonized areas might be possible. Further studies should focus on

analyzing the level of ploidy within the most common invasive lineage (Form A/R) and

determine whether front/newly established populations show a higher frequency of

triploid individuals, and thus, higher fitness, relative to those from core/long-established

populations.

In conclusion, in this study we provide evidence supporting that different life-history

traits of individuals can respond to different selective pressures occurring on varying

population contexts (e.g. time elapsed since the population foundation, low vs. high

population density, and low vs. high rates of population growth). Our results show that

fitness-related traits, such as individual growth rate and minimum age at sexual

maturity, are maximized in recently established, slow-growing populations, which

experience a stronger *r*-selective environment with respect to long established ones.

While these plastic responses could be interpreted as an effective mechanism to

decrease the risk of stochastic extinction associated with low densities at the invasion

front, population growth rate seems to be independent of population density in a wide

range of environmental conditions within *Corbicula*'s invasive range. Collectively, our

results strongly suggest that plasticity might have favored the expression of traits that

maximizes fitness in populations facing novel environmental conditions. Whether these

are adaptive phenotypic responses remains as an open question that could be addressed

by performing laboratory experiments under common-garden conditions. Although

other relevant variables could not be considered in our models, this study represents a

first step in the attempt to understand in a wide geographical perspective how different

selection pressures affected fitness-related traits of the Asian clam during range shifting.

REFERENCES

Aldridge DW, McMahon RF (1978) Growth, fecundity, and bioenergetics in a natural

population of the Asiatic freshwater clam, Corbicula manilensis Philippi, from

North Central Texas. J Molluscan Stud 44:49–70

Araujo R, Moreno D, Ramos MA (1993) The Asiatic clam *Corbicula fluminea* (Müller,

1774) (Bivalvia: Corbiculidae) in Europe. Am Malacol Bull 10:39–49

Bonel N, Lorda J (2015) Growth and body weight variability of the invasive mussel

Limnoperna fortunei (Mytilidae) across habitat and season. Malacologia 58:129–

145. https://doi.org/10.4002/040.058.0202

Borenstein M, Hedges L V, Higgins JPT, Rothstein HR (2009) Introduction to meta-

analysis, First edit. John Wiley & Sons, Ltd, Chichester, West Sussex

Bossdorf O, Auge H, Lafuma L, et al (2005) Phenotypic and genetic differentiation

between native and introduced plant populations. Oecologia 144:1–11.

https://doi.org/10.1007/s00442-005-0070-z

Brook BW, Bradshaw CJA (2006) Strength of evidence for density dependence in

abundance time series of 1198 species. Ecology 87:1445–1451.

https://doi.org/10.1890/0012-9658(2006)87[1445:SOEFDD]2.0.CO;2

Burton OJ, Phillips BL, Travis JMJ (2010) Trade-offs and the evolution of life-histories

during range expansion. Ecol Lett 13:1210–1220. https://doi.org/10.1111/j.1461-

0248.2010.01505.x

Cao L, Damborenea C, Penchaszadeh PE, Darrigran G (2017) Gonadal cycle of

Corbicula fluminea (Bivalvia: Corbiculidae) in Pampean streams (Southern

Neotropical Region). PLoS One 12:1–16.

https://doi.org/10.1371/journal.pone.0186850

Cataldo D, Boltovskoy D (1999) Population dynamics of *Corbicula fluminea* (Bivalvia)

in the Parana River Delta (Argentina). Hydrobiologia 380:153–163.

https://doi.org/10.1023/A:1003428728693

Cole LC (1954) The population consequences of life history phenomena. Q Rev Biol

29:103–137. https://doi.org/10.1086/400074

Crespo D, Dolbeth M, Leston S, et al (2015) Distribution of *Corbicula fluminea*

(Müller, 1774) in the invaded range: a geographic approach with notes on species

traits variability. Biol Invasions 17:2087–2101. https://doi.org/10.1007/s10530-015-0862-y

Crespo D, Leston S, Martinho F, et al (2017) Survival of *Corbicula fluminea* (Müller, 1774) in a natural salinity and temperature gradient: a field experiment in a temperate estuary. Hydrobiologia 784:337–347. https://doi.org/10.1007/s10750-016-2887-3

Davis HG (2005) r-Selected traits in an invasive population. Evol Ecol 19:255–274. https://doi.org/10.1007/s10682-005-0912-5

Denton ME, Chandra S, Wittmann ME, et al (2012) Reproduction and population structure of *Corbicula fluminea* in an oligotrophic subalpine lake. J Shellfish Res 31:145–152. https://doi.org/10.2983/035.031.0118

Eng LL (1979) Population dynamics of the Asiatic clam, *Corbicula fluminea* (Müller), in the concrete-lined Delta-Mendota Canal of Central California. In: Britton JC (ed) Proceedings, First International Corbicula Symposium. Texan Christian University, Fort Worth, Texas, pp 39–67

Etoundi E, Marescaux J, Vastrade M, et al (2019) Distinct biogeographic origins of androgenetic *Corbicula* lineages followed by genetic captures. bioRxiv 590836. https://doi.org/http://dx.doi.org/10.1101/590836

Fisher RA (1930) The genetical theory of natural selection. Clarendon Press, Oxford

Franco JN, Ceia FR, Patrício J, et al (2012) Population dynamics of *Corbicula fluminea* (Müller, 1774) in mesohaline and oligohaline habitats: invasion success in a Southern Europe estuary. Estuar Coast Shelf Sci 112:31–39. https://doi.org/10.1016/j.ecss.2011.07.014

French JRP, Schloesser DW (1991) Growth and overwinter survival of the Asiatic clam, *Corbicula fluminea*, in the St . Clair River, Michigan. Hydrobiologia 219:165–170

Gama M, Crespo D, Dolbeth M, Anastácio P (2016) Predicting global habitat suitability for *Corbicula fluminea* using species distribution models: the importance of different environmental datasets. Ecol Modell 319:163–169. https://doi.org/10.1016/j.ecolmodel.2015.06.001

Gama M, Crespo D, Dolbeth M, Anastácio PM (2017) Ensemble forecasting of *Corbicula fluminea* worldwide distribution: projections of the impact of climate change. Aquat Conserv Mar Freshw Ecosyst 27:675–684. https://doi.org/10.1002/aqc.2767

Gardner J., Woodall WR, Staats AA, Napoli JF (1976) The invasion of the Asiatic clam

(*Corbicula manilensis* Philippi) in the Altamaha river, Georgia. Nautilus (Philadelphia) 90:117–125. https://doi.org/10.1038/scientificamerican01221846-76b

Gaston KJ (2009) Geographic range limits: achieving synthesis. Proc R Soc B Biol Sci 276:1395–1406. https://doi.org/10.1098/rspb.2008.1480

Gayanilo FC, Sparre P, Pauly D (2002) FAO-ICLARM Stock Assessment Tools (FiSAT). Software version 1.2.0. FAO, Roma

Graney Jr RL, Cherry DS, Cairns Jr J (1980) The influence of substrate, pH, diet and temperature upon cadmium accumulation in the Asiatic clam (*Corbicula fluminea*) in laboratory artificial streams. Nautilus (Philadelphia) 18:833–842

Heinsohn GE (1958) Life history and ecology of the freshwater clam, *Corbicula fluminea*. University of California, Berkeley

Hill JK, Griffiths HM, Thomas CD (2011) Climate change and evolutionary adaptations at species' range margins. Annu Rev Entomol 56:143–159. https://doi.org/10.1146/annurev-ento-120709-144746

Hoegh-Guldberg O, Bruno JF (2010) The impact of climate change on the world's marine ecosystems. Science (80-) 328:1523–1528. https://doi.org/10.1126/science.1189930

Hoenig JM, Hanumara RC (1982) A statistical study of a seasonal growth model for fishes. Univ Rhode Island Dep Comput Sci Tech Report, Narragansett

Hoffmann AA, Sgró CM (2011) Climate change and evolutionary adaptation. Nature 470:479–485. https://doi.org/10.1038/nature09670

Hornbach DJ (1992) Life history traits of a riverine population of the Asian clam *Corbicula fluminea*. Am Modl Nat 127:248–257

Ilarri MI, Antunes C, Guilhermino L, Sousa R (2011) Massive mortality of the Asian clam *Corbicula fluminea* in a highly invaded area. Biol Invasions 13:277–280. https://doi.org/10.1007/s10530-010-9833-5

IPCC (2014) Climate change 2014: impacts, adaptation, and vulnerability. Cambridge University Press, United Kingdom and New York

Ituarte CF (1985) Growth dynamics in a natural population of *Corbicula fluminea* (Bivalvia Sphaeriacea) at Punta Atalaya, Rio de La Plata, Argentina. Stud Neotrop Fauna Environ 20:217–225

Kennedy VS., Van Huekelem L (1985) Gametogenesis and larval production in a population of the introduced Asiatic clam, *Corbicula* sp. (Bivalvia: Corbiculidae),

in Maryland Published by : Marine Biological Laboratory Stable URL : Biol Bull
168:50–60

Komaru A, Kawagishi T, Konishi K (1998) Cytological evidence of spontaneous
androgenesis in the freshwater clam *Corbicula leana* Prime. Dev Genes Evol
208:46–50. https://doi.org/10.1007/s004270050152

Kraemer LR (1979) *Corbicula* (Bivalvia: Sphaeriacea) vs. indigenous mussels
(Bivalvia: Unionacea) in U.S. rivers: a hard case for interspecific competition? Am
Zool 19:1085–1096

Kraemer LR, Lott S (1977) Microscopic anatomy of the visceral mass of *Corbicula*
(Bivalvia: Sphaeriacea). Am Malacol Bull 48–56

Kraemer LR, Swanson C, Galloway M, Kraemer R (1986) Biological basis of behavior
in *Corbicula fluminea*, II. Functional morphology of reproduction and
development and review of evidence for self-fertilization. Am Malacol Bull Spetial
Ed:193–201

Lewontin RC (1965) Selection for colonizing ability. In: Baker H, Stebbins G (eds) The
genetics of colonizing species. Academic Press, New York, pp 79–94

Lockwood JL, Hoopes MF, Marchetti MP (2007) Invasion Ecology. Blackwell
Publishing, Malden, USA; Oxford, UK; Victoria, Australia

Lonhart SI, Jeppesen R, Beas-Luna R, et al (2019) Shifts in the distribution and
abundance of coastal marine species along the eastern Pacific Ocean during marine
heatwaves from 2013 to 2018. Mar Biodivers Rec 12:13.
https://doi.org/10.1186/s41200-019-0171-8

MacArthur RH, Wilson E 0. (1967) The theory of island biogeography. Princeton
University Press

Magnussen E (2007) Interpopulation comparison of growth patterns of 14 fish species
on Faroe Bank : are all fishes on the bank fast-growing? J Fish Bioilogy 71:453–
475. https://doi.org/10.1111/j.1095-8649.2007.01502.x

McDowell WG, Mcdowell WH, Byers JE (2016) Mass mortality of a dominant invasive
species in response to an extreme climate event: implications for ecosystem
function. Limnol Oceanogr. https://doi.org/10.1002/lno.10384

McDowell WG, Sousa R (2019) Mass mortality events of invasive freshwater bivalves:
current understanding and potential directions for future research. Front Ecol Evol
7:. https://doi.org/10.3389/fevo.2019.00331

McMahon RF (2002) Evolutionary and physiological adaptations of aquatic invasive

animals: r selection versus resistance. Can J Fish Aquat Sci 59:1235–1244.

https://doi.org/10.1139/F02-105

McMahon RF, Williams CJ (1986) Growth, life cycle, upper thermal limit and

downstream colonization rates in a natural population of the freshwater bivalve

mollusc, *Corbicula fluminea* (Müller) receiving thermal effluents. Am Malacol

Bull Special Ed:231–239.

Mouthon J (2001) Life cycle and population dynamics of the Asian clam *Corbicula*

fluminea (Bivalvia : Corbiculidae) in the Rhone River at Creys-Malville (France).

Arch für Hydrobiol 151:571–589

Mouthon J, Parghentanian T (2004) Comparison of the life cycle and population

dynamics of two *Corbicula* species, *C. fluminea* and *C. fluminalis* (Bivalvia:

Corbiculidae) in two French canals. Arch für Hydrobiol 161:267–287.

https://doi.org/10.1127/0003-9136/2004/0161-0267

Munro JL, Pauly D (1983) A simple method for comparing the growth of fishes and

invertebrates. Fishbyte 1:5–6

Otto SP (2007) The evolutionary consequences of polyploidy. Cell 131:452–462.

https://doi.org/10.1016/j.cell.2007.10.022

Pauly D, Caddy JF (1985) A modification of Bhattacharya's method for the analysis of

mixtures of normal distributions. Food Agric Organ United Nations 781:

Pauly D, Gaschutz G (1979) A simple method for fitting oscillating length growth data,

with a program for pocket calculators. ICES C 6:24

Pauly D, Munro JL (1984) Once more on the comparison of growth in fish and

invertebrates. Fishbyte 2:

Peña EA, Slate EH (2006) Global validation of linear model assumptions. ournal Am

Stat Assoc 101:341. https://doi.org/10.1038/jid.2014.371

Pfennig DW, Wund MA, Snell-Rood EC, et al (2010) Phenotypic plasticity's impacts

on diversification and speciation. Trends Ecol Evol 25:459–467.

https://doi.org/10.1016/j.tree.2010.05.006

Phillips BL (2009) The evolution of growth rates on an expanding range edge. Biol Lett

5:802–804. https://doi.org/10.1098/rsbl.2009.0367

Phillips BL, Brown GP, Shine R (2010) Life-history evolution in range-shifting

populations. Ecology 91:1617–1627

Phillips BL, Brown GP, Webb JK, Shine R (2006) Invasion and the evolution of speed

in toads. Nature 439:803. https://doi.org/10.1038/439803a

Pigliucci M, Murren CJ, Schlichting CD (2006) Phenotypic plasticity and evolution by

genetic assimilation. J Exp Biol 209:2362–2367. https://doi.org/10.1242/jeb.02070

Pigneur L-M, Etoundi E, Aldridge DC, et al (2014) Genetic uniformity and long-

distance clonal dispersal in the invasive androgenetic *Corbicula* clams. Mol Ecol

23:5102–5116. https://doi.org/10.1111/mec.12912

Pigneur L-M, Hedtke SM, Etoundi E, Doninck K Van (2012) Androgenesis : a review

through the study of the selfish shellfish *Corbicula* spp . Heredity (Edinb)

108:581–591. https://doi.org/10.1038/hdy.2012.3

Rajagopal S, Velde G, Vaate A (2000) Reproductive biology of the Asiatic clams

Corbicula fluminalis and *Corbicula fluminea*. Arch für Hydrobiol 149:403–420

Ramón M, Fernández M, Galimany E (2007) Development of mussel (*Mytilus*

galloprovincialis) seed from two different origins in a semi-enclosed

Mediterranean Bay (N.E. Spain). Aquaculture 264:148–159.

https://doi.org/10.1016/j.aquaculture.2006.11.014

Ricciardi A (2013) Invasive Species. In: Meyers RA (ed) Encyclopedia of Sustainability

Science and Technology. Springer Science+Business Media, New York, pp 5547–

5560

Roff DA (1993) Evolution of life histories: theory and analysis. Springer US

Rosa IC, Pereira JL, Costa R, et al (2014) Dispersal of *Corbicula fluminea* : Factors

influencing the invasive clam's drifting behavior. Ann Limnol - Int J Limnol

50:199–199. https://doi.org/10.1051/limn/2014011

Sakai AK, Allendorf FW, Holt JS, et al (2001) The population biology of invasive

species. Annu Rev Ecol Syst 32:305–332

Schmidlin S, Baur B (2007) Distribution and substrate preference of the invasive clam

Corbicula fluminea in the river Rhine in the region of Basel (Switzerland,

Germany, France). Aquat Sci 69:153–161. https://doi.org/10.1007/s00027-006-

0865-y

Schrieber K, Lachmuth S (2017) The genetic paradox of invasions revisited: the

potential role of inbreeding× environment interactions in invasion success. Biol

Rev 92:939–952. https://doi.org/10.1111/brv.12263

Selmecki A, Maruvka YE, Richmond PA, et al (2015) Polyploidy can drive rapid

adaptation in yeast. Nature 519:349–352. https://doi.org/10.1038/nature14187

Shine R, Brown GP, Phillips BL (2011) An evolutionary process that assembles

phenotypes through space rather than through time. Proc Natl Acad Sci 108:5708–

5711. https://doi.org/10.1073/pnas.1018989108

Siemann E, Rogers WE (2001) Genetic differences in growth of an invasive tree

species. Ecol Lett 4:514–518. https://doi.org/10.1046/j.1461-0248.2001.00274.x

Skellam JG (1951) Random dispersal in theoretical populations. Biometrika 38:196–

218

Somers IF (1988) On A Seasonally Oscillating Growth Function. Fishbyte 6:8–11

Sousa R, Antunes C, Guilhermino L (2008a) Ecology of the invasive Asian clam

Corbicula fluminea (Müller, 1774) in aquatic ecosystems: an overview. Ann

Limnol - Int J Limnol 44:85–94. https://doi.org/10.1051/limn:2008017

Sousa R, Antunes C, Guilhermino L (2006) Factors influencing the occurrence and

distribution of *Corbicula fluminea* (Müller, 1774) in the River Lima estuary. Ann

Limnol - Int J Limnol 42:165–171. https://doi.org/10.1051/limn/2006017

Sousa R, Nogueira AJA, Gaspar MB, et al (2008b) Growth and extremely high

production of the non-indigenous invasive species *Corbicula fluminea* (Müller,

1774): possible implications for ecosystem functioning. Estuar Coast Shelf Sci

80:289–295. https://doi.org/10.1016/j.ecss.2008.08.006

Stearns SC (1976) Life-History tactics: A review of the ideas. Q Rev Biol 51:3–47

Stearns SC (1977) The evolution of life history traits: a critique of the theory and a

review of the data. Annu Rev Ecol Syst 8:145–171.

https://doi.org/10.1146/annurev.es.08.110177.001045

Stites DL, Benke AC, Gillespie DM (1995) Population dynamics, growth, and

production of the Asiatic clam *Corbicula fluminea*, in a blackwater river. Can J

Fish Aquat Sci 52:425–437

Strayer DL, D'Antonio CM, Essl F, et al (2017) Boom-bust dynamics in biological

invasions: towards an improved application of the concept. Ecol Lett 20:1337–

1350. https://doi.org/10.1111/ele.12822

Taylor CC (1958) Cod growth and temperature. J du Cons 23:366–370

Travis J (1994) Evaluating the adaptive role of morphological plasticity. In: Wainwright

PC, Reilly SM (eds) Ecological Morphology: Integrative Organismal Biology.

University of Chicago Press, Chicago, pp 99–122

Vakily JM (1992) Determination and comparison of bivalve growth, with emphasis on

Thailand and other tropical areas

Vaughn CC, Atkinson CL, Julian JP (2015) Drought-induced changes in flow regimes

lead to long-term losses in mussel-provided ecosystem services. Ecol Evol 5:1291–

1305. https://doi.org/10.1002/ece3.1442

Vohmann A, Borcherding J, Kureck A, et al (2010) Strong body mass decrease of the

invasive clam *Corbicula fluminea* during summer. Biol Invasions 12:53–64.

https://doi.org/10.1007/s10530-009-9429-0

Walther G-R, Post E, Convey P, et al (2002) Ecological responses to recent climate

change. Nature 416:389–395. https://doi.org/10.1007/978-3-319-10157-6_4

West-Eberhard MJ (2003) Developmental plasticity and evolution. Oxford University

Press, New York

TABLES

Table 1. Summary of ecological and growth variables (mean ± SD) of worldwide invasive populations of the Asian clam *Corbicula* reviewed from the literature. *Density* is the estimated abundance of clams at the study site (ind./m^2), *Pop.growth* is the population growth rate, *Time* is the number of years elapsed since population colonization at the time of the original study (years), *Ind.growth* is the individual growth rate (*GPI*, a *proxy* for individual growth rate), *mASM* is the minimum age at sexual maturity (months), and lifespan (years), *ID#* is a number assigned arbitrarily for identification purposes. The country, specific study location, and reference to the source paper of the original study used to build the information are shown for each population. See the main text for details on how data was collected and variables were built from the literature. For extended information on each population, see Table S1 of the Supporting Information.

Country	Study location	*Time*	Population parameters		Individual parameters			*ID#*	*Reference*
			Density	*Pop.growth*	*Ind.growth*	*mASM*	*Lifespan*		
Argentina	Punta Atalaya, Río de la Plata	3.5	339±405	2.20±0.01	2.87±0.07	4.30±0.13	3.83±0.57	1	Ituarte (1985)
	Paraná de las Palmas River	25.5	1,053±768	14.44±0.31	2.73±0.01	9.23±1.17	6.82±0.05	2	Cataldo and Boltovskoy (1999)
	Río Negro estuary	15.5	94±8.3	7.85±0.10	2.94±0.23	5.32±1.66	3.55±0.41	3	Hünicken et al. (2019)
France	Saone River	12.5	302±186	14.65±0.01	2.54±0.13	12.73±2.42	7.55±2.17	4	Mouthon (2001)
	Loire Lateral Canal	17.0	992±1209	41.17±0.12	2.80±0.24	8.93±5.95	5.17±3.00	5	Mouthon and Parghentanian (2004)
	Canal of Roanne	17.0	626±897	25.50±0.01	2.77±0.16	8.25±3.43	4.27±1.58	6	Mouthon and Parghentanian (2004)
Portugal	River Minho estuary	16.5	840±387	11.89±0.06	3.32±0.05	3.06±1.20	2.28±0.30	7	Sousa et al. (2008b)
	Mondego estuary	6.9	5,017±5,489	7.67±0.04	–	–	–	8	Franco *et al.*, 2012
	Casal de São Tomé	10.3	3,255±1,115	4.06±0.04	3.14±0.12	3.83±1.69	2.75±0.51	9	Rosa et al. (2014)
USA	Altamaha River	4.5	712±1,613	0.37±0.00	–	–	–	10	Gardner et al. (1976)
	Lake Arlington	2.0	32±11	11.94±0.03	3.49±0.05	5.80±0.60	2.59±0.20	11	Aldridge and McMahon (1978)
	Delta-Mendota canal	28.0	9,470±7067	4.53±0.03	2.77±0.17	9.69±4.07	8.03±2.75	12	Eng (1979)
	New River (Station 1)	2	316±654	9.40±0.02	–	–	–	13	Graney et al. (1980)
	Clear Fork, Trinity River	7.0	4,474±3,899	19.34±0.04	3.00±0.16	6.76±3.62	7.07±2.83	14	McMahon and Williams (1986)
	Mechums River	10.5	669±636	188.52±0.62	2.56±0.15	9.20±4.41	3.31±1.21	15	Hornbach (1992)
	Ogeechee River	10.0	1,398±1,096	6.23±0.00	2.52±0.12	22.72±9.17	6.85±1.63	16	Stites et al. (1995)
	Lake Tahoe	8.0	1,809±723	16.64±0.04	2.80±0.12	6.90±1.90	2.70±1.32	17	Denton et al. (2012)

812 **Table 2.** Summary of results from reduced models of weighted linear regression analyses of invasion and growth variables of global invasive
813 populations of *Corbicula*. Parameter values are means ± SE. We report one-tailed *P* values indicating significant results in bold characters. Note
814 natural logarithmic transformations of variables. For details on descriptive statistics of full models and results on partial *F*-tests see the
815 Supporting Information. All variables were Ln-transformed and standardized to zero mean and a unit of variance (see *Statistical Analyses* for
816 details).

Response variable	Predictor	Estimate ± SE	Intercept ± SE	R^2	*F*-statistics	*num.df, den.df*	*one-tailed* P-value
Density	*Time*	0.454 ± 0.196	-0.906 ± 0.241	0.263	5.34	1, 15	**0.018**
Population growth rate	*Time*	0.936 ± 0.323	-0.518 ± 0.216	0.376	4.03	2, 14	**0.006**
	Density	-0.522 ± 0.407					0.220†
Individual growth rate	*Density*	-0.422 ± 0.265	0.098 ± 0.264	0.283	2.17	2, 11	0.070
	Population growth rate	-0.473 ± 0.344					0.099
minimum Age at sexual maturity	*Individual growth rate*	-0.505± 0.179	-0.192 ± 0.171	0.601	8.28	2, 11	**0.008**
	Population growth rate	0.460 ± 0.208					**0.024**
Lifespan	*Individual growth rate*	-0.718± 0.184	-0.056 ± 0.192	0.559	15.20	1, 12	**0.001**

819 † two-tailed *P*-value.

 FIGURES

 Figure 1. Global distribution of invasive *Corbicula* clams indicating the native range of the genus (dark grey), the distribution of
 of *Corbicula* inside (black/grey shading and black dots) and outside of the genus' native range (color dots), which are mostly represented by
 forms A/R (formerly designed as *Corbicula fluminea sensu lato*; see main text for detailed explanation). Numbered stars are invasive worldwide
 populations reviewed in the present study. Years indicate estimated date of first introduction. See Table 1 and main text for references and details
 on the invasive populations reviewed here. See the Supporting Information for detailed criteria and data used to build the genus native and
 invasive distributions.

Figure 2. Relationships between processes and characteristics at different levels of
organization of 17 global invasive populations of the Asian clam *Corbicula*. At the
population level: a) the effect of time since colonization (*Time,* years) on population
Density (ind./m^2*)* and b, c) the simultaneous effect of *Density* and *Time* on *Population*
growth rate. At the interaction between population and individual level: d, e) the
simultaneous effect of density and population growth rate on individual growth rate, and
f) the effect of *Population growth rate* on the minimum *Age at sexual maturity*
(months). At the individual level: g) the effect of *Individual growth rate* on the
minimum *Age at sexual maturity*, and h) on the *Life span* (years). When plotting the fit
for one variable after testing for simultaneous effects, we removed the effect of the
variable not considered (b, c, d, e). Dashed lines show fits when marginally non-
significant 0.05> α <0.10. Solid lines show linear regression fits when statistically
significant at α < 0.05. The size of circles correspond to the inverse of the variance of
mean value of each observation of the response variable, used to weigh effect sizes in
the regression analyses. All variables were Ln-transformed and standardized to zero
mean and a unit of variance (see *Statistical Analyses* for details).

Figura 3. Summary of the relationships between processes and characteristics at
different levels of organization of 17 global invasive populations of the Asian clam
Corbicula, showing positive (+), negative (–), and no relationships (0) after fitting
weighted linear regressions.Thick solid lines (➡) show linear regression fits when
statistically significant at α < 0.01. Thin solid lines (→) show linear regression fits
when statistically significant at α < 0.05. Dashed lines (– –➤) show fits when
marginally non significant 0.05> α <0.10. Dotted line (······➤) show no significant
relationship α > 0.10. Relationships are shown at three different levels: A) at the
population level, B) the interaction between processes and characteristics at the
population and individual level, and C) at the individual level.

854