Fitness-related traits are maximized in recently introduced, slow-growing populations of a global invasive clam

Leandro A. Hünicken[1,2]*, Francisco Sylvester[2,3], Nicolás Bonel[2,4,5]

[1] Museo Argentino de Ciencias Naturales 'Bernardino Rivadavia', Av. Angel Gallardo 470 (C1405DJR), Buenos Aires, Argentina. Present Address: Centro de Investigación Aplicada y Transferencia Tecnológica en Recursos Marinos ''Almirante Storni'', Güemes 1030, San Antonio Oeste,

R8520CXV Río Negro, Argentina

[2] Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

[3] Instituto para el estudio de la Biodiversidad de Invertebrados, Facultad de Ciencias Naturales, Universidad Nacional de Salta, Av. Bolivia 5150 (4400), Salta, Argentina.

[4] Genética y Ecología Evolutiva, Centro de Recursos Naturales Renovables de la Zona Semiárida (CERZOS– CCT–CONICET/UNS), Camino La Carrindanga Km 7, (B8000CPB) Bahía Blanca, Argentina.

[5] Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175, CNRS—Université de Montpellier, Université Paul- Valéry Montpellier—Ecole Pratique des Hautes Etudes—IRD, 34293 Montpellier cedex 05, France.

* Corresponding author. E-mail: leandrohunicken@gmail.com

Key words: *Corbicula fluminea*, Invasive species, Individual growth rates, Age at first reproduction, Front-edge populations, *r*-selection, Adaptive plasticity.

Author Contributions: LAH, FS, and NB conceived the study. LAH, FS, and NB conducted the literature search and extracted the data. NB designed the methodological approach. LAH and NB performed the data analysis. NB led the writing.

ABSTRACT

Many species are shifting their ranges being forced to rapidly respond to novel stressful environmental conditions. Colonizing individuals experience strong selective forces that favor the expression of life history traits notably affecting dispersal and reproductive rates in newly invaded habitats. Limited information is currently available on trait variation within the invasive range despite being critical for understanding ecological and evolutionary factors that drive the process of range expansion of invasive species. Here we evaluated life history shifts of the widely introduced Asian clam *Corbicula* within its invaded range. Through an exhaustive literature search, we obtained data for 17 invasive *Corbicula* populations from different ecosystems worldwide to test the relationship between population and individual parameters relevant to the process of range expansion. Our main results are that (i) recently introduced *Corbicula* populations are characterized by low conspecific density and low rate of population increase, (ii) clams reproduce earlier in slow-growing populations, and (iii) density had no effect on exponential population increase. All invasive *Corbicula* populations analyzed in this study (Form A/R) are mostly fixed for one genotype. Our results therefore suggest that adaptive phenotypic plasticity favored the expression of traits that maximize fitness in recently established populations facing stronger *r*-selective forces, thereby increasing their chances to overcome difficulties associated with low densities and low population increase in newly invaded areas.

1. INTRODUCTION

The ongoing increase in frequency of extreme climatic events, caused by anthropogenic climate change, and global transport of people and goods is altering ecosystems by triggering a wide array of irreversible consequences (Bossdorf et al., 2005; Hoegh-Guldberg & Bruno, 2010; IPCC, 2014; Walther et al., 2002). Changes in local conditions, in particular those driven by human-induced climate change, are predicted to affect ecological processes across several levels of organization, likely over the coming decades. Such changes are altering the performance of individual organisms, the dynamics of populations and the distribution of species, ultimately modifying ecosystem properties and function (Hoffmann & Sgró, 2011; Lonhart et al., 2019).

Many species are forced to rapidly respond to novel stressful environmental conditions when shifting their distribution and colonizing new areas. If non-native species have difficulties in adapting to such conditions and fail to track abrupt environmental changes, populations become vulnerable to decline and extinction (Hill et al., 2011; Hoffmann & Sgró, 2011). Phenotypic plasticity is the ability of a single genotype to express different phenotypes when exposed to different environmental conditions (Travis, 1994; West-Eberhard, 2003). In a new environment, plasticity rapidly favors the appearance of a novel phenotypic variant, promoting successful colonization and increasing the likelihood for the long-term persistence of populations facing new environmental stress brought about by range shifting (Hill et al., 2011; Pfennig et al., 2010; Pigliucci et al., 2006). Adaptive plasticity that places populations close enough to the new favored phenotypic optimum for directional selection to act provides the first step in adaptation to new environments on ecological time scales (see review in Ghalambor et al. 2007). Individuals at the expanding edge of the species distribution experience strong selective forces (r-selection) that can have profound impacts species life

histories. These forces can, for instance, favor the expression of traits that optimize population

growth and survival from offspring to maturity (e. g. reduced age of maturity and

reproduction, increased growth rate, increased energy allocated to reproduction; Stearns,

1976, 1977), thus affecting dispersal and reproductive rates (Phillips, 2009; Shine et al.,

2011).

Colonization events through either continuous range expansion or jump dispersal are usually

driven by a small number of individuals that are the first to arrive at a new point of the

distribution, where they benefit from increased space open for colonization and low

conspecific density, below carrying capacity (Cole, 1954; Phillips et al., 2010; Sakai et al.,

2001). Hence, the density gradient occurring at the expanding range (i. e. grading from zero to

carrying capacity) creates different selective environments where individuals from a recently

established population face stronger r-selection compared to those from long established ones,

which are exposed to selective K-type forces (e. g. Phillips et al., 2010). According to theory,

low densities favor the expression of life-history traits related with r-type selections (Burton

et al., 2010; MacArthur & Wilson, 1967; Phillips, 2009), leading to exponential (non-density

regulated) population growth in recently established populations while denser, long-

established ones follow a density-regulated population growth (Brook & Bradshaw, 2006).

Recently established populations are characterized by low conspecific density and low rate of

increase, often leading to lag-times between initial colonization and the onset of rapid

population growth (Sakai et al., 2001). The duration of the lag phase is mediated, among

others, by a number of non-mutually exclusive ecological and evolutionary mechanisms (e. g.

low encounter rate for reproduction, density-dependent effects, adaptation, and selection of

new genotypes; Bossdorf et al., 2005; Davis, 2005; Ricciardi, 2013). An effective mechanism

that increases the rate of population growth is early reproduction, which results from fast

growing individuals (Cole, 1954; Lewontin, 1965; Roff, 1993). Individuals that grow faster

will attain reproductive sizes earlier and bring down generation times, promoting a rapid

build-up of population numbers (Cole, 1954; Lewontin, 1965; Phillips, 2009; Roff, 1993) and

accelerating primary invasion and secondary spread of invasive species (Lockwood et al.,

2007; Phillips et al., 2006; Skellam, 1951). Several studies have compared life-history traits

between native and invaded ranges ('home-and-away' comparisons). Such comparisons could

lead to erroneous interpretations of the patterns observed because it would be difficult to

disentangle what were the processes (e.g. propagule bias and/or evolution in populations at

spatial (dis)equilibrium) causing the shifts in life-history traits (Phillips et al., 2010 and

references therein). Interestingly, only a few studies have looked for such trait changes in

introduced species (Gaston, 2009), despite knowledge on life-history trait variation within the

invasive range being critical for better understanding the process of species range expansion.

Bivalves of the genus *Corbicula* (Megerle Von Mühlfeld 1811) are native to Southeast Asia,

the Middle East, Australia and Africa (Araujo et al., 1993) but they have colonized a large

part of the Americas and Europe (Crespo et al., 2015; Mouthon, 2001; Schmidlin & Baur,

2007). The successful invasion of *Corbicula* clams has been mainly attributed to their rapid

growth and maturation, high fecundity, and high dispersal making this bivalve genus one of

the most relevant faunal non-indigenous groups in aquatic ecosystems (reviewed in Sousa et

al. 2008a). Invasive freshwater *Corbicula* populations are constituted by a small number of

lineages composed of hermaphroditic individuals reproducing through androgenesis, for

which the most widespread and abundant genotype in the invaded area is the lineage so-called

Form A/R, widely known as *C. fluminea* (Komaru et al., 1998; Pigneur et al., 2012, 2014).

The current taxonomic status of the invasive *Corbicula* lineages in the Americas and Europe

is still largely unresolved despite several morphological and genetic studies (reviewed in

Pigneur et al. 2014). In the present paper, we deal with this form and refer to it as *Corbicula*

to avoid entering unsettled taxonomic debates.

Here we investigated how different life-history traits of the Asian clam *Corbicula* responded

to different selective pressures occurring at varying population contexts within the introduced

range: (i) newly- vs. long-established, (ii) low vs. high density, and (iii) low vs. high rates of

population increase; all of them being linked to the geographic range shifting of species'

distribution (i. e. front/core populations). To do so, we searched for peer-reviewed articles

providing enough information to estimate time since population introduction as well as

different population and individual processes and characteristics including population density,

individual growth rate, minimum age at sexual maturity, and lifespan of global populations of

this worldwide invader. Different studies suggest that *Corbicula*'s population dynamics are

chiefly regulated by environmental conditions (e. g. Crespo et al., 2015, 2017; Gama et al.,

2017), whereby population declines are typically driven by periodic mass mortality due to

extreme hydro-meteorological events (Ilarri et al., 2011; McDowell et al., 2017; McDowell &

Sousa, 2019). This strongly suggests that invasive *Corbicula* populations would primarily

follow a non-density regulated population increase and, hence, they might not level off at

carrying capacity as happens with many other species (e.g. Brook and Bradshaw, 2006). We

focused our predictions on life history strategies exhibited by individuals from recently

established populations, which typically occur at the invasion front, and likely subjected to

strong *r*-selective pressures with respect to those from older, long-established ones. We tested

the hypotheses that population density and population increase are directly related with time

since introduction. *Corbicula* is widely known to exhibit rapid population growth, being able

to reach extremely high densities (reviewed in McMahon, 2002). We expected density to have

no significant effect on population increase even within a wide range of densities in nature,

though high densities are considered likely to have a negative effect on individual growth

rates. We also asked whether clams that grow faster and reproduce earlier are favored in slow-

growing populations at the expanding front. A related prediction is that clams that grow

rapidly would have a shorter lifespan with respect to individuals exhibiting slow growth.

2. METHODS

2.1 Literature search

We exhaustively searched the literature for peer-reviewed articles on *Corbicula* (*Corbicula*

fluminea sensu lato) providing enough information to estimate the individual growth rate,

population density, rate of exponential population increase, time since colonization, water

temperature, and conductivity of global populations of this clam. To do this, we first

identified relevant studies from ISI Web of Knowledge (Web of Science Core Collection,

1900 through 2017) by conducting a first search using the field tag "topic (TS)" in the

"Advanced search" option and the search string: TS="Corbicula fluminea" OR TS="asia*

clam" (search #1). We conducted a second search including topics TS="population

dynamic*" OR TS="population size*" OR TS="population densit*" OR TS="population

structure" (search #2). We ran a third search using TS="growth" OR TS="individual

growth" OR TS="population growth" (search #3). Finally, we combined searches #2 and #3

into search #4 (#4= #3 OR #2), and restricted the latter by search #1 to obtain a final set of

candidate studies (search #5= #4 AND #1).

This literature search was finished on 27 September 2018 and this protocol identified 215

candidate studies (database search: 215 candidates; gray literature and unpublished data: 0

candidates). To identify papers potentially missing in this list, we subsequently scanned the

references in these papers and conducted a Google Scholar search using keywords mentioned

above and added 8 studies to this candidate pool (Fig. S1). A total of 67 studies that belonged

to research areas unrelated to the topic of the current study (*e. g.*, applied chemistry,

toxicology, paleontology) were excluded from this candidate pool by refining the set based in

'Web of Science Categories'. The resulting studies (n = 156) were finally inspected

individually in detail to assess their eligibility for our analyses according to the following

inclusion criteria: i) Studies that assessed invasive populations of *Corbicula* clams (i. e.

studies exclusively providing estimates for native populations were excluded); ii) Studies that

reported or provided data enough to extract the population density and age/size distribution

for >6-months (see below); iii) Studies that assessed populations in reasonably "average"

natural environments, such that populations from heavily impacted sites (e. g. thermal

plumes) were excluded. We finally retained a sample of 16 studies, covering 17 invasive

populations of *Corbicula* from different ecosystems worldwide (n = 140 out of 156 were

excluded; see Fig. S1 for further details on screening and eligibility criteria). Whenever

population data was only graphically presented, we contacted directly the authors. However,

if data were no longer available or authors did not respond, we used the WebPlotDigitizer

software (http://arohatgi.info/WebPlotDigitizer/) to extract data and accurately estimate

population density and growth rates. This step-by-step criterion for the literature search and

data collection attempted to reduce any potential selection bias in the long-term populations

that might occur because old high-density populations (with higher growth rates) could be

more studied than young low-density populations (with low growth rates), which would be

more likely to be extirpated if population characteristics were not changing with time.

2.2 Population density and exponential population increase

For each population selected from our literature search, we obtained the mean population

density (ind m^2) for the entire study period or extracted temporal values to estimate it

whenever this value was missing in the text. *Corbicula* clams generally exhibit a bivoltine

juvenile release pattern (i. e. producing two generations a year McMahon, 2002; Sousa,

Antunes, et al., 2008), leading to one or more high-density peaks related to recruitment

events yearly. However, density in the field fluctuates almost constantly generating several

peaks and valleys that do not necessarily indicate significant recruitment events. *Corbicula*

populations exhibit rapid growth (Franco et al., 2012; McMahon, 2002; Sousa et al., 2006;

Sousa, Antunes, et al., 2008). Hence, the rate of exponential population increase of

Corbicula can be expressed by deriving the classical exponential growth equation:

208
$$r = \frac{\ln\left(\frac{Nt}{N0}\right)}{t}$$

One of the goals of this study was to compare population growth rate during periods of

substantial growth across populations, and it is such capacity that is of interest to better

understand *Corbicula* invasiveness. Hence, to estimate the rate of exponential population

increase (*r*, hereafter population growth rate), we set a criterion to determine relevant

recruitment peaks in each study by using the average density throughout the study period as

a threshold. By doing so, we excluded time periods where population growth was extremely

low likely due to mass mortality due to extreme hydro-meteorological events (e.g. a period

of dramatic demographic decline), usually characterized by a lag phase (where population

growth can be negligible) followed by an exponential increase (Ricciardi, 2013). Likewise,

this approach allows for exclusion of large time periods resulting from *Corbicula*'s inability

to reproduce year-round as they generally exhibit a uni- or bivoltine juvenile release. We

therefore considered the lowest density value (N_0), the highest peak above the average

density (N_t), and the time period elapsed between the valley and the peak (*t*). Intermediate

density values comprised in each period (i. e. defined by N_0 and N_t) were used to fit the

growth equation and obtain *r*. Population rates of increase were estimated using GraphPad

Prism version 7.00 for Windows, GraphPad Software, La Jolla California USA,

www.graphpad.com. Note that periods defined by non-relevant peaks (i. e. below the

average density threshold) were discarded, as were periods of progressive density decline

between a relevant peak and the next N_0 value (see Fig. S2 for further details on the step-by-

step procedure used to estimate population increase). When populations exhibited more than

one relevant density peak throughout the study period, the population increase rate for such

populations was calculated as the average of single *r* estimations. Then, we standardized

estimates on an annual basis to make them comparable. Note that the calculated *r* values

provide a relative growth rate, that is, one that can be used to compare different populations

to one another, rather than a population growth rate that can be used to forecast future

population sizes.

2.3 Individual growth rates and parameters

For studies that only presented data on shell length-frequency distributions (shell length

[mm] being the greatest distance between the anterior and posterior margins of the shell

valve), we identified age cohorts using that information and applied the Bhattacharya's

method available in FISAT II software (Version 1.2.0, FAO-ICLARM Fish Assessment

Tools; (Gayanilo et al., 2002). To confirm each component of normal distributions from the

modal progression analysis, we used the NORMSEP method also available in the FISAT II

software (Pauly & Caddy, 1985). Once we had cohorts identified for all the studies, we

calculated individual growth parameters using the same equations for all the studies

considered in order to make sure that results were fully comparable across studies. For each

cohort present in each of the populations considered in this study, we fitted the von

Bertalanffy-growth function with seasonal oscillations (hereafter SVBGF) proposed by

Pauly and Gaschutz (1979), Hoenig and Hanumara (1982), and Somers (1988):

$$L_t = L_\infty \, (1 - exp^{\,-K\,[(t-t_0) + T_1 - T_2]}),$$

$$T_1 = C \sin (2\pi \, (t - t_s)) \, / \, 2\pi,$$

and

$$T_2 = C \sin (2\pi \, (t_0 - t_s)) \, / \, 2\pi,$$

where L_t is the predicted shell length at age t, L_∞ is the asymptotic shell length, K is the

growth constant of dimension time (year^{-1} in most seasonally oscillating growth curves)

expressing the rate at which L_∞ is approached, t_0 is the theoretical 'age' the clam has at

shell length zero, C expresses the relative amplitude of the seasonal oscillation and varies

between 0 and 1 (0 indicating lack of summer-winter differences in growth, it was

constrained to be less than or equal to 1 during model fitting), and t_s is the starting point of

the oscillation.

The parameters of the function were estimated by the modeling method available in JMP

statistical software (v9.0 SAS Institute). When preliminary results of the SVBGF failed to

converge to estimate the asymptotic shell length (L_∞), we used the maximum shell length

(*Lmax*) observed or reported at/for each study site to calculate the asymptotic shell length

following the equation suggested by Taylor (1958):

$$Lmax \, / \, 0.95 = L_\infty$$

These values were fixed when we performed a second fit.

Because of the negative correlation between growth parameters (K and L_∞) prevents making

comparisons based on individual parameters (Ramón et al., 2007; Vakily, 1992), we used the

growth-performance index (*GPI*), which reflects the growth rate of a given organism of unit

length. In other words, *GPI* can be viewed as the (theoretical) value of *K* that would occur in

organisms with an L_∞ value of 1 unit of length (Munro & Pauly, 1983), and it was defined by

Pauly and Munro (1984) as:

$$GPI = 2 \log_{10} L_\infty + \log_{10} K$$

We calculated the *GPI* values for cohorts from each study site based on growth parameters

(L_∞ and *K*) obtained after fitting the SVBGF using the *Growth Performance Indices*

application of the FISAT II software (Gayanilo et al., 2002).

2.3.1 Minimum age at sexual maturity

The minimum age at sexual maturity was estimated as the time (months) elapsed until clams

reach the minimum size (shell length) at which invasive *Corbicula* clams become sexually

mature, which, on average, is 10.0 mm ± 0.8 SE (n = 13 reported sizes). This value was

obtained from shell sizes reported in the literature (Aldridge & McMahon, 1978; Cao et al.,

2017; Eng, 1979; French & Schloesser, 1991; Gardner et al., 1976; Heinsohn, 1958; Ituarte,

1985; Kennedy & Van Huekelem, 1985; Kraemer, 1979; Kraemer & Lott, 1977; McMahon,

2002; Rajagopal et al., 2000). We acknowledge that in some conditions and populations,

clams might mature at a different pace and thus introduce a degree of error in such cases.

While this error is unavoidable until more detailed data is available, we are confident that the

10-mm estimate applies to many real-life situations and thus our results based on it are robust.

2.3.2 Lifespan

As *t0* is the theoretical 'age' the clam has at shell length zero, *tmax* can be defined as the

theoretical 'age' the a clam reaches its maximum shell length (*Lmax*). Since *Lmax* is

dependently related to a maximum point in time (years), it is assumed that *Lmax* implicates

tmax (*Lmax* ⇒ *tmax*) (Bonel & Lorda, 2015). Thus, the theoretical maximum lifespan of a

clam cohort can be approximately expressed as:

300   *Lifespan = tmax – t0*

2.4 Time since colonization and key environmental variables

For the same 17 populations and sites studied, we calculated the time elapsed since

population foundation (hereafter time since colonization) as the difference between the

sampling dates in the first ecological survey and the first report of introduction. The

accuracy of the introduction date is generally very difficult to know, however, many of the

studies used report range expansions of a conspicuous invader across well studied regions,

where the appearance of the species might have been anticipated and readily detected by

ongoing monitoring actions in the initial stages of invasion. Therefore, we consider that the

possible error introduced should be moderate to low and the value obtained can be viewed as

a good approximation. Moreover, we obtained water temperature and conductivity values

for each site averaging monthly mean values for the same period used to estimate growth

rates in each study. Other biotic and abiotic variables are known to interact with temperature

to drive growth rates of invasive bivalves including food availability, dissolved oxygen, and

heavy metal concentration (Bonel & Lorda, 2015; Vohmann et al., 2010). However, we only

considered water temperature (ºC) and conductivity (µS/cm) in the analyses because they

have been reported to play a key role on the *Corbicula*'s biology and ecology (Crespo et al.,

2017; Franco et al., 2012; Magnussen, 2007). Indeed, temperature has emerged as the most

important variable explaining the current distribution and, also, predicting that climate

change will favor the expansion of *Corbicula*'s into colder areas at higher latitudes (Gama et

al., 2017). Likewise, salinity has been considered a key factor influencing the success and

velocity of the invasion of new estuarine ecosystems (e.g. Sousa et al., 2006). As *Corbicula*

populations considered herein belonged to a broad range of freshwater and estuarine

habitats, we therefore included conductivity to serve as a *proxy* for salinity levels. In order to

keep values seasonally unbiased, only entire years were used in subsequent calculations.

Whenever these parameters were not reported in the original paper, we obtained values for

the same sites and years from the literature and reliable online databases (e. g.

https://waterdata.usgs.gov/nwis/qw). When data was only graphically presented, we

extracted it following the procedure mentioned above.

2.5 Statistical analyses

We ran five separate multivariate weighted linear regressions in order to test ecological

processes and characteristics at three different (but non-mutually exclusive) levels: 1)

population level, 2) population and individual level, and 3) individual level. In this sense, a

response variable at the population level can be considered as an explanatory variable when

analyzing a response at the individual level. In all cases, we computed the variance of the

effect size from different populations, which were then used as weights in linear regressions.

The inverse-variance weighting allows combining the results from independent measurements

giving more importance to the least noisy measurements and *vice versa* (Borenstein et al.,

2009).

We were interested in testing individual effects of predictors, which are related to life-history

traits of *Corbicula* (hereafter reduced models). However, we added annual mean water

temperature and conductivity as quantitative variables in each model (hereafter full models).

Preliminary analyses showed that latitudinal effects are strongly correlated with water

temperature ($r = -0.84$). Hence, it was excluded from the models to avoid multicollinearity. In

some cases, we also included interaction terms between explanatory variables. We used the

partial *F*-test to compare models—that is, full models with and without the interaction term,

and full *versus* reduced models. As we had strong *a priori* hypotheses as to the direction of

effects, we used one-tailed tests. Variables were Ln-transformed to fulfill the models'

assumption of normality and homoscedasticity, which were tested using the *gvlma* package

(Peña & Slate, 2006). Prior to analyses, we standardized explanatory variables to zero mean

and a unit of variance in order to get directly comparable effects. All analyses were carried

out in R version 3.6.1 (R Development Core Team 2019). Values are given as means ± SE

unless otherwise stated.

357   3. RESULTS

358   3.1 Literature search

359   Populations analyzed in this study belonged to Europe, North and South America (Fig. 1;

360   Table 1). These included eight populations in the United States, three in France, three in

361   Argentina, and three in Portugal (see Table 1 for detailed location, bibliographic source, and

362   code number used herein to refer to the populations, for each of the 17 *Corbicula* populations

363   selected, and Table S1 for additional information).

365   3.2 Density, growth, time since colonization, and key environmental variables

Overall, mean density of *Corbicula* from 17 locations worldwide spanned 32 to *ca*. 9,500 ind.

m^{-2}, population growth rate (*r*) varied from 0.37 to 189, individual growth rate (*GPI*) ranged

2.53-3.49, age at first reproduction 3.1-22.7 months, and lifespan 2.3-8.0 years (Table 1). On

the other hand, time since colonization ranged 2 to 28 years in the populations studied (Table

1) and yearly water temperature ranged from 13.2 to 20.7 ºC and conductivity from 68 to

8,457 µS/cm (see Table S1 of the Supporting Information for details). In all cases, we found

no evidence that adding an interaction term with variables of interest nor temperature and

conductivity in each model significantly increased its predictive power (for details see '*Model*

comparison' in Supporting Information). We therefore report estimates and statistics from the

reduced models in Table 2.

At the population level, we found that younger *Corbicula* populations (i.e. newly established)

were characterized by low density and slow population growth rate with respect to older, long

established ones (Fig. 2a, b). We found that density had no significant effect on population

increase (Fig. 2c). When analyzing responses at the interaction between population and

individual processes and characteristics, we found that clams tended to grow faster

(marginally non-significant) in low-density and slow-growing populations (Fig. 2d, e). In

contrast, we observed a strong positive response of the minimum age at sexual maturity as a

function of population growth rate, clearly indicating that clams reproduce earlier in slow-

growing populations (Fig. 2f). This is directly linked to processes occurring at the individual

level, where clams with higher growth rates reach the minimum reproductive size earlier (10

mm) and have shorter life span than clams that grow more slowly (Fig. 2g, h).

4. DISCUSSION

This is the first study that evaluates how population and individual parameters, involved in

the process of range expansion, varied in response to different selective pressures occurring

within the introduced range of the worldwide invasive Asian clam *Corbicula*. Our main

results showed that (i) recently introduced *Corbicula* populations are characterized by low

density and reduced population growth rate, (ii) clams reproduce earlier in slow-growing

populations, and (iii) population growth rate was unfettered by density (summarized in Fig.

3).

Density of *Corbicula* populations increased with time since colonization, which is consistent

with the standpoint that recently established populations typically occur at low conspecific

density, below the carrying capacity (Cole, 1954; Sakai et al., 2001). Likewise, we found that

population growth rate and time since colonization were directly related, suggesting that, in

newly colonized areas, populations could have experienced stronger *r*-type selection favoring

fitness-related traits effectively increasing the rate of population growth, as it was observed in

long established populations (Burton et al., 2010; MacArthur & Wilson, 1967; Phillips, 2009).

Population density did not predict changes in the rate of population increase, as variation in

population growth remained relatively low even at high-density values. By definition, long

established and denser populations tend to be density regulated as they approach to carrying

capacity (logistic population growth; e. g. Phillips et al., 2010). Considering the so-called *r*/K

selection continuum, this implies that such populations are predicted to shift from the *r*-

towards the K-endpoint (MacArthur & Wilson, 1967)—that is, selection shifts a population

from density-independent to density-dependent regulation. However, *Corbicula*'s population

growth rate did not decrease with increasing density despite the fact that some of the

populations included in this study exhibit extremely high densities exceeding, in some cases,

100,000 individuals per square meter (e. g. Eng 1979; Table 1).

Why was population growth unfettered by density regulation? A large number of studies have

analyzed density dynamics of *Corbicula* and commonly assumed that, following exponential

growth, the clam population achieves densities approaching the carrying capacity of the

environment (e.g. Sousa et al., 2006). However, to our knowledge, no studies have ever

conducted (field or laboratory) experiments to test whether population growth is limited by

increasing conspecific density, which, in turn, is expected to reduce local biotic resources. In

other words, there is a complete lack of knowledge on the true effect of density on population

growth rate. On the contrary, different studies have shown that *Corbicula*'s dynamics are

chiefly influenced by environmental variables (e. g. water temperature, conductivity, pH,

and/or dissolved oxygen) and seem to be less influenced by population's density (e. g. Crespo

et al., 2015; Gama et al., 2016). Indeed, periodic mass mortality due to extreme hydro-

meteorological events (e.g. extremely high or low temperatures, drought) could play an

important role with infrequent, but catastrophic population declines (>99% in some cases;

Ilarri et al., 2011; McDowell et al., 2017; Mouthon & Parghentanian, 2004). These events

have been described (though not quantified) in several other systems as well (e. g. McDowell

& Sousa, 2019; Vaughn et al., 2015). Despite that a given *Corbicula* population might

approach or even grow beyond the carrying capacity of the ecosystem, evidence from

previous studies (and those cited herein) suggests that population size significantly varies with

time. This means that the population is at non-equilibrium and that such fluctuations (or

boom-bust dynamics; Strayer et al., 2017) are likely driven by density-independent factors, as

the above-mentioned hydro-meteorological events. Thus, it does seem reasonable to infer that

Corbicula's population growth rate is non-density-regulated and will not level off at carrying

capacity, as expected from a highly invasive organism as the Asian clam (McMahon 2002).

Our results suggest that this bivalve seems to be far from approaching carrying capacity, even

at extremely high densities, supporting the idea that individuals from long established denser

populations do not face a *K*-selective environment but a less strongly *r*-selective pressure.

Individuals tended to grow faster in low-density populations experiencing low rates of

growth. Likewise, *Corbicula* clams reproduced earlier in slow-growing populations. In

recently introduced populations, periods of negligible population increase can be driven by

non-mutually exclusive ecological and evolutionary adjustments such as Allee effect,

changing abiotic conditions, and adaptation and selection of new genotypes imposed by novel

environments conditions in a new habitat (Bossdorf et al., 2005; Davis, 2005; Ricciardi, 2013;

Sakai et al., 2001). Such conditions can lead to a strong selection on life-history traits that

optimize population increase and survival to maturity of offspring (Stearns, 1976, 1977). As

predicted by life-history theory, individuals from recently established populations (typically

occurring at the invasion front) face stronger r-selection relative to conspecifics from older

populations (MacArthur & Wilson, 1967), favoring traits that accelerate the range expansion.

For instance, some empirical observations have revealed that individuals from recently

colonized areas grow faster, and thus, reproduce earlier, than do conspecifics from older,

long-established populations (e. g. butterflies: Hanski et al., 2006; Australian cane toads:

Phillips, 2009; tallow trees: Siemann & Rogers, 2001). Accordingly, our results for *Corbicula*

clams showing fast growth and, thus, early reproduction support the idea that such

populations faced strong r-selection, which favored traits associated with high reproductive

rates in recently introduced, slow-growing populations.

The increase in individual growth was associated with a decrease in the clams' life span likely

due to trade-offs between different life-history traits. Such trade-offs can be interpreted as an

effective mechanism to bring down generation times, promoting a rapid build-up of

population numbers, increasing population growth in newly colonized areas (Cole, 1954;

Lewontin, 1965; Roff, 1993) and accelerates primary invasion and secondary spread of

invasive species (Lockwood et al., 2007; Phillips et al., 2006; Skellam, 1951). Collectively,

our results suggest that reduced population increase of the current *Corbicula* population

triggers the shortening of the time that the clams need to reach maturity by favoring clams

that grow the fastest. Thus, as early-maturing clams reproduce earlier, this has the effect of

fostering an increase in size of the future population. In this way, incipient *Corbicula*

populations could increase their chances of overcoming the difficulties associated with low

densities and survive to the population establishment stage, which can clearly subsidize the

bivalve's invasive abilities in newly colonized areas.

In conclusion, in this study we have provided evidence supporting the idea that life-history

traits of individuals can respond to selective pressures occurring under varying population

contexts (e.g. varying time since population foundation, low vs. high population density, and

low vs. high rates of population growth). All invasive *Corbicula* populations analyzed in this

study are mostly fixed for one genotype (Form A/R). Our results, therefore, provide

reasonable ground for advocating that clams from recently established, slow-growing

populations experienced a stronger *r*-selective environment than those from long-established

ones; and that adaptive plasticity can maximize the expression of fitness-related traits, such as

individual growth rate and the minimum age at sexual maturity, in these populations. While

these responses could be interpreted as an effective mechanism to decrease the risk of

stochastic extinction associated with low densities at the invasion front, population growth

rate seems to be independent of population density in a wide range of environmental

conditions within *Corbicula*'s invasive range. Although other relevant variables could not be

considered in our models, this study represents a first step in the attempt to understand in a

wide geographical perspective how different selection pressures affect fitness-related traits of

the Asian clam during range shifting.

REFERENCES

Aldridge, D. W., & McMahon, R. F. (1978). Growth, fecundity, and bioenergetics in a natural

population of the Asiatic freshwater clam, *Corbicula manilensis* Philippi, from North

Central Texas. *Journal of Molluscan Studies*, *44*, 49–70.

Araujo, R., Moreno, D., & Ramos, M. A. (1993). The Asiatic clam *Corbicula fluminea*

(Müller, 1774) (Bivalvia: Corbiculidae) in Europe. *American Malacological Bulletin*,

10(1), 39–49.

Bonel, N., & Lorda, J. (2015). Growth and body weight variability of the invasive mussel

Limnoperna fortunei (Mytilidae) across habitat and season. *Malacologia*, *58*(1–2), 129–

145. https://doi.org/10.4002/040.058.0202

Borenstein, M., Hedges, L. V, Higgins, J. P. T., & Rothstein, H. R. (2009). *Introduction to*

meta-analysis (First edit). John Wiley & Sons, Ltd.

Bossdorf, O., Auge, H., Lafuma, L., Rogers, W. E., Siemann, E., & Prati, D. (2005).

Phenotypic and genetic differentiation between native and introduced plant populations.

Oecologia, *144*(1), 1–11. https://doi.org/10.1007/s00442-005-0070-z

Brook, B. W., & Bradshaw, C. J. A. (2006). Strength of evidence for density dependence in

abundance time series of 1198 species. *Ecology*, *87*(6), 1445–1451.

https://doi.org/10.1890/0012-9658(2006)87[1445:SOEFDD]2.0.CO;2

Burton, O. J., Phillips, B. L., & Travis, J. M. J. (2010). Trade-offs and the evolution of life-

histories during range expansion. *Ecology Letters*, *13*, 1210–1220.

https://doi.org/10.1111/j.1461-0248.2010.01505.x

Cao, L., Damborenea, C., Penchaszadeh, P. E., & Darrigran, G. (2017). Gonadal cycle of

Corbicula fluminea (Bivalvia: Corbiculidae) in Pampean streams (Southern Neotropical

Region). *PLoS ONE*, *12*(10), 1–16. https://doi.org/10.1371/journal.pone.0186850

Cataldo, D., & Boltovskoy, D. (1999). Population dynamics of *Corbicula fluminea* (Bivalvia)

in the Parana River Delta (Argentina). *Hydrobiologia*, *380*, 153–163.

https://doi.org/10.1023/A:1003428728693

Cole, L. C. (1954). The population consequences of life history phenomena. *The Quarterly*

Review of Biology, *29*(2), 103–137. https://doi.org/10.1086/400074

Crespo, D., Dolbeth, M., Leston, S., Sousa, R., & Pardal, M. A. (2015). Distribution of

Corbicula fluminea (Müller, 1774) in the invaded range: a geographic approach with

notes on species traits variability. *Biological Invasions*, *17*(7), 2087–2101.

https://doi.org/10.1007/s10530-015-0862-y

Crespo, D., Leston, S., Martinho, F., Pardal, M. A., & Dolbeth, M. (2017). Survival of

Corbicula fluminea (Müller, 1774) in a natural salinity and temperature gradient: a field

experiment in a temperate estuary. *Hydrobiologia*, *784*(1), 337–347.

https://doi.org/10.1007/s10750-016-2887-3

Davis, H. G. (2005). r-Selected traits in an invasive population. *Evolutionary Ecology*, *19*(3),

255–274. https://doi.org/10.1007/s10682-005-0912-5

Denton, M. E., Chandra, S., Wittmann, M. E., Reuter, J., & Baguley, J. G. (2012).

Reproduction and population structure of *Corbicula fluminea* in an oligotrophic

subalpine lake. *Journal of Shellfish Research*, *31*(1), 145–152.

https://doi.org/10.2983/035.031.0118

Eng, L. L. (1979). Population dynamics of the Asiatic clam, *Corbicula fluminea* (Müller), in

the concrete-lined Delta-Mendota Canal of Central California. In J. C. Britton (Ed.),

Proceedings, First International Corbicula Symposium (pp. 39–67). Texan Christian

University.

Franco, J. N., Ceia, F. R., Patrício, J., Modesto, V., Thompson, J., Marques, J. C., & Neto, J.

M. (2012). Population dynamics of *Corbicula fluminea* (Müller, 1774) in mesohaline and

oligohaline habitats: invasion success in a Southern Europe estuary. *Estuarine, Coastal*

and Shelf Science, *112*, 31–39. https://doi.org/10.1016/j.ecss.2011.07.014

French, J. R. P., & Schloesser, D. W. (1991). Growth and overwinter survival of the Asiatic

clam, *Corbicula fluminea*, in the St . Clair River, Michigan. *Hydrobiologia*, *219*, 165–

170.

Gama, M., Crespo, D., Dolbeth, M., & Anastácio, P. (2016). Predicting global habitat

suitability for *Corbicula fluminea* using species distribution models: the importance of

different environmental datasets. *Ecological Modelling*, *319*, 163–169.

https://doi.org/10.1016/j.ecolmodel.2015.06.001

Gama, M., Crespo, D., Dolbeth, M., & Anastácio, P. M. (2017). Ensemble forecasting of

Corbicula fluminea worldwide distribution: projections of the impact of climate change.

Aquatic Conservation: Marine and Freshwater Ecosystems, *27*(3), 675–684.

https://doi.org/10.1002/aqc.2767

Gardner, J. ., Woodall, W. R., Staats, A. A., & Napoli, J. F. (1976). The invasion of the

Asiatic clam (*Corbicula manilensis* Philippi) in the Altamaha river, Georgia. *The*

Nautilus, *90*(3), 117–125. https://doi.org/10.1038/scientificamerican01221846-76b

Gaston, K. J. (2009). Geographic range limits: achieving synthesis. *Proceedings of the Royal*

Society B: Biological Sciences, *276*(1661), 1395–1406.

https://doi.org/10.1098/rspb.2008.1480

Gayanilo, F. C., Sparre, P., & Pauly, D. (2002). FAO-ICLARM Stock Assessment Tools

(FiSAT). Software version 1.2.0. *FAO, Roma*.

Ghalambor, C. ., McKay, J. ., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus non-

adaptive phenotypic plasticity and the potential for contemporary adaptation in new

evironments. *Functional Ecology*, *21*(3), 394–407. https://doi.org/10.1111/j.1365-

2435.2007.01283.x

Graney Jr, R. L., Cherry, D. S., & Cairns Jr, J. (1980). The influence of substrate, pH, diet and

temperature upon cadmium accumulation in the Asiatic clam (*Corbicula fluminea*) in

laboratory artificial streams. *The Nautilus*, *18*(7), 833–842.

Hanski, I., Saastamoinen, M., & Ovaskainen, O. (2006). Dispersal-related life-history trade-

offs in a butterfly metapopulation. *Journal of Animal Ecology*, *75*, 91–100.

https://doi.org/10.1111/j.1365-2656.2005.01024.x

Heinsohn, G. E. (1958). *Life history and ecology of the freshwater clam,* Corbicula fluminea.

University of California, Berkeley.

Hill, J. K., Griffiths, H. M., & Thomas, C. D. (2011). Climate change and evolutionary

adaptations at species' range margins. *Annual Review of Entomology*, *56*, 143–159.

https://doi.org/10.1146/annurev-ento-120709-144746

Hoegh-Guldberg, O., & Bruno, J. F. (2010). The impact of climate change on the world's

marine ecosystems. *Science*, *328*(5985), 1523–1528.

https://doi.org/10.1126/science.1189930

Hoenig, J. M., & Hanumara, R. C. (1982). A statistical study of a seasonal growth model for

fishes. *University of Rhode Island. Department of Computer Science, Technical Report,*

Narragansett.

Hoffmann, A. A., & Sgró, C. M. (2011). Climate change and evolutionary adaptation. *Nature*,

470(7335), 479–485. https://doi.org/10.1038/nature09670

Hornbach, D. J. (1992). Life history traits of a riverine population of the Asian clam

Corbicula fluminea. American Modland Naturalist, *127*(2), 248–257.

Ilarri, M. I., Antunes, C., Guilhermino, L., & Sousa, R. (2011). Massive mortality of the

Asian clam *Corbicula fluminea* in a highly invaded area. *Biological Invasions*, *13*(2),

277–280. https://doi.org/10.1007/s10530-010-9833-5

IPCC. (2014). Climate change 2014: impacts, adaptation, and vulnerability. In C. B. Field &

V. R. Barros (Eds.), *Part A: global and sectoral aspects. Contribution of working group*

II to the fifth assessment report of the Intergovernmental Panel on Climate Change.

Cambridge University Press. https://www.ipcc.ch/pdf/assessment-

report/ar5/wg2/WGIIAR5-FrontMatterA_FINAL.pdf

Ituarte, C. F. (1985). Growth dynamics in a natural population of *Corbicula fluminea*

(Bivalvia Sphaeriacea) at Punta Atalaya, Rio de La Plata, Argentina. *Studies on*

Neotropical Fauna and Environment, *20*(4), 217–225.

Kennedy, V. S. ., & Van Huekelem, L. (1985). Gametogenesis and larval production in a population of the introduced Asiatic clam, *Corbicula* sp. (Bivalvia: Corbiculidae), in Maryland Published by : Marine Biological Laboratory Stable URL : *Biological Bulletin1*, *168*(1), 50–60.

Komaru, A., Kawagishi, T., & Konishi, K. (1998). Cytological evidence of spontaneous androgenesis in the freshwater clam *Corbicula leana* Prime. *Development Genes and Evolution*, *208*(1), 46–50. https://doi.org/10.1007/s004270050152

Kraemer, L. R. (1979). *Corbicula* (Bivalvia: Sphaeriacea) vs. indigenous mussels (Bivalvia: Unionacea) in U.S. rivers: a hard case for interspecific competition? *American Zoologist*, *19*(4), 1085–1096.

Kraemer, L. R., & Lott, S. (1977). Microscopic anatomy of the visceral mass of *Corbicula* (Bivalvia: Sphaeriacea). *American Malacological Bulletin*, 48–56.

Lewontin, R. C. (1965). Selection for colonizing ability. In H. Baker & G. Stebbins (Eds.), *The genetics of colonizing species* (pp. 79–94). Academic Press.

Lockwood, J. L., Hoopes, M. F., & Marchetti, M. P. (2007). *Invasion Ecology*. Blackwell Publishing. http://onlinelibrary.wiley.com/doi/10.1002/cbdv.200490137/abstract

Lonhart, S. I., Jeppesen, R., Beas-Luna, R., Crooks, J. A., & Lorda, J. (2019). Shifts in the distribution and abundance of coastal marine species along the eastern Pacific Ocean during marine heatwaves from 2013 to 2018. *Marine Biodiversity Records*, *12*(1), 13. https://doi.org/10.1186/s41200-019-0171-8

MacArthur, R. H., & Wilson, E. O. (1967). *The theory of island biogeography*. Princeton, NJ.

Magnussen, E. (2007). Interpopulation comparison of growth patterns of 14 fish species on Faroe Bank : are all fishes on the bank fast-growing? *Journal of Fish Bioilogy*, *71*, 453–475. https://doi.org/10.1111/j.1095-8649.2007.01502.x

McDowell, W. G., McDowell, W. H., & Byers, J. E. (2017). Mass mortality of a dominant

invasive species in response to an extreme climate event: Implications for ecosystem

function. *Limnology and Oceanography*, *62*(1), 177–188.

https://doi.org/10.1002/lno.10384

McDowell, W. G., & Sousa, R. (2019). Mass mortality events of invasive freshwater bivalves:

current understanding and potential directions for future research. *Frontiers in Ecology*

and Evolution, *7*(331). https://doi.org/10.3389/fevo.2019.00331

McMahon, R. F. (2002). Evolutionary and physiological adaptations of aquatic invasive

animals: r selection versus resistance. *Canadian Journal of Fisheries and Aquatic*

Sciences, *59*(7), 1235–1244. https://doi.org/10.1139/F02-105

McMahon, R. F., & Williams, C. J. (1986). Growth, life cycle, upper thermal limit and

downstream colonization rates in a natural population of the freshwater bivalve mollusc,

Corbicula fluminea (Müller) receiving thermal effluents. *American Malacological*

Bulletin, *Special Ed*(2), 231–239.

Mouthon, J. (2001). Life cycle and population dynamics of the Asian clam *Corbicula*

fluminea (Bivalvia : Corbiculidae) in the Rhone River at Creys-Malville (France). *Archiv*

Für Hydrobiologie, *151*(4), 571–589.

Mouthon, J., & Parghentanian, T. (2004). Comparison of the life cycle and population

dynamics of two Corbicula species, *C . fluminea* and *C . fluminalis* (Bivalvia:

Corbiculidae) in two French canals. *Archiv Für Hydrobiologie*, *161*(2), 267–287.

https://doi.org/10.1127/0003-9136/2004/0161-0267

Munro, J. L., & Pauly, D. (1983). A simple method for comparing the growth of fishes and

invertebrates. *Fishbyte*, *1*(1), 5–6.

Pauly, D., & Caddy, J. F. (1985). A modification of Bhattacharya's method for the analysis of

mixtures of normal distributions. *Food and Agricultural Organization of the United*

Nations, *781*.

Pauly, D., & Gaschutz, G. (1979). A simple method for fitting oscillating length growth data,

with a program for pocket calculators. *ICES CM*, *6*, 24.

Pauly, D., & Munro, J. L. (1984). Once more on the comparison of growth in fish and

invertebrates. *Fishbyte*, *2*.

Peña, E. A., & Slate, E. H. (2006). Global validation of linear model assumptions. *Ournal of*

the American Statistical Association, *101*(473), 341. https://doi.org/10.1038/jid.2014.371

Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D., &

Moczek, A. P. (2010). Phenotypic plasticity's impacts on diversification and speciation.

Trends in Ecology and Evolution, *25*(8), 459–467.

https://doi.org/10.1016/j.tree.2010.05.006

Phillips, B. L. (2009). The evolution of growth rates on an expanding range edge. *Biology*

Letters, *5*, 802–804. https://doi.org/10.1098/rsbl.2009.0367

Phillips, B. L., Brown, G. P., & Shine, R. (2010). Life-history evolution in range-shifting

populations. *Ecology*, *91*(6), 1617–1627.

Phillips, B. L., Brown, G. P., Webb, J. K., & Shine, R. (2006). Invasion and the evolution of

speed in toads. *Nature*, *439*(7078), 803. https://doi.org/10.1038/439803a

Pigliucci, M., Murren, C. J., & Schlichting, C. D. (2006). Phenotypic plasticity and evolution

by genetic assimilation. *Journal of Experimental Biology*, *209*(12), 2362–2367.

https://doi.org/10.1242/jeb.02070

Pigneur, L.-M., Etoundi, E., Aldridge, D. C., Marescaux, J., Yasuda, N., & Van Doninck, K.

(2014). Genetic uniformity and long-distance clonal dispersal in the invasive

androgenetic *Corbicula* clams. *Molecular Ecology*, *23*(20), 5102–5116.

https://doi.org/10.1111/mec.12912

Pigneur, L.-M., Hedtke, S. M., Etoundi, E., & Doninck, K. Van. (2012). Androgenesis : a

review through the study of the selfish shellfish *Corbicula* spp . *Heredity*, *108*(6), 581–

591. https://doi.org/10.1038/hdy.2012.3

Rajagopal, S., Velde, G., & Vaate, A. (2000). Reproductive biology of the Asiatic clams

Corbicula fluminalis and *Corbicula fluminea*. *Archiv Für Hydrobiologie*, *149*(3), 403–

420.

Ramón, M., Fernández, M., & Galimany, E. (2007). Development of mussel (*Mytilus*

galloprovincialis) seed from two different origins in a semi-enclosed Mediterranean Bay

(N.E. Spain). *Aquaculture*, *264*(1–4), 148–159.

https://doi.org/10.1016/j.aquaculture.2006.11.014

Ricciardi, A. (2013). Invasive Species. In R. A. Meyers (Ed.), *Encyclopedia of Sustainability*

Science and Technology (pp. 5547–5560). Springer Science+Business Media.

https://doi.org/10.1007/978-1-4419-0851-3

Roff, D. A. (1993). *Evolution of life histories: theory and analysis* (D. A. Roff (ed.)). Springer

US.

Rosa, I. C., Pereira, J. L., Costa, R., Gomes, J., de Lourdes Pereira, M., & Gonçalves, F.

(2014). Dispersal of *Corbicula fluminea* : Factors influencing the invasive clam's drifting

behavior. *Annales de Limnologie - International Journal of Limnology*, *50*(3), 199–199.

https://doi.org/10.1051/limn/2014011

Sakai, A. K., Allendorf, F. W., Holt, J. S., Lodge, M., Molofsky, J., With, K. A., Baughman,

S., Cabin, R. J., Cohen, J. E., Ellstrand, N. C., McCauley, D. E., O'Neil, P., Parker, I. M.,

Thompson, J. N., & Weller, S. G. (2001). The population biology of invasive species.

Annual Review of Ecology and Systematics, *32*(1), 305–332.

Schmidlin, S., & Baur, B. (2007). Distribution and substrate preference of the invasive clam

Corbicula fluminea in the river Rhine in the region of Basel (Switzerland, Germany,

France). *Aquatic Sciences*, *69*(1), 153–161. https://doi.org/10.1007/s00027-006-0865-y

Shine, R., Brown, G. P., & Phillips, B. L. (2011). An evolutionary process that assembles

phenotypes through space rather than through time. *Proceedings of the National*

Academy of Sciences, *108*(14), 5708–5711. https://doi.org/10.1073/pnas.1018989108

Siemann, E., & Rogers, W. E. (2001). Genetic differences in growth of an invasive tree

species. *Ecology Letters*, *4*(6), 514–518. https://doi.org/10.1046/j.1461-

0248.2001.00274.x

Skellam, J. G. (1951). Random dispersal in theoretical populations. *Biometrika*, *38*(1/2), 196–

218.

Somers, I. F. (1988). On A Seasonally Oscillating Growth Function. *Fishbyte*, *6*, 8–11.

Sousa, R., Antunes, C., & Guilhermino, L. (2006). Factors influencing the occurrence and

distribution of *Corbicula fluminea* (Müller, 1774) in the River Lima estuary. *Annales de*

Limnologie - International Journal of Limnology, *42*(3), 165–171.

https://doi.org/10.1051/limn/2006017

Sousa, R., Antunes, C., & Guilhermino, L. (2008). Ecology of the invasive Asian clam

Corbicula fluminea (Müller, 1774) in aquatic ecosystems: an overview. *Annales de*

Limnologie - International Journal of Limnology, *44*(2), 85–94.

https://doi.org/10.1051/limn:2008017

Sousa, R., Nogueira, A. J. A., Gaspar, M. B., Antunes, C., & Guilhermino, L. (2008). Growth

and extremely high production of the non-indigenous invasive species *Corbicula*

fluminea (Müller, 1774): possible implications for ecosystem functioning. *Estuarine,*

Coastal and Shelf Science, *80*(2), 289–295. https://doi.org/10.1016/j.ecss.2008.08.006

Stearns, S. C. (1976). Life-History tactics: A review of the ideas. *The Quarterly Review of*

Biology, *51*(1), 3–47.

Stearns, S. C. (1977). The evolution of life history traits: a critique of the theory and a review

of the data. *Annual Review of Ecology and Systematics*, *8*(1), 145–171.

https://doi.org/10.1146/annurev.es.08.110177.001045

Stites, D. L., Benke, A. C., & Gillespie, D. M. (1995). Population dynamics, growth, and

production of the Asiatic clam *Corbicula fluminea*, in a blackwater river. *Canadian*

Journal of Fisheries and Aquatic Sciences, *52*(215), 425–437.

Strayer, D. L., D'Antonio, C. M., Essl, F., Fowler, M. S., Geist, J., Hilt, S., Jarić, I., Jöhnk,

K., Jones, C. G., Lambin, X., Latzka, A. W., Pergl, J., Pyšek, P., Robertson, P., von

Schmalensee, M., Stefansson, R. A., Wright, J., & Jeschke, J. M. (2017). Boom-bust

dynamics in biological invasions: towards an improved application of the concept.

Ecology Letters, *20*(10), 1337–1350. https://doi.org/10.1111/ele.12822

Taylor, C. C. (1958). Cod growth and temperature. *Journal Du Conseil*, *23*, 366–370.

Travis, J. (1994). Evaluating the adaptive role of morphological plasticity. In P. C.

Wainwright & S. M. Reilly (Eds.), *Ecological Morphology: Integrative Organismal*

Biology (pp. 99–122). University of Chicago Press.

Vakily, J. M. (1992). Determination and comparison of bivalve growth, with emphasis on

Thailand and other tropical areas. In *WorldFish*. https://doi.org/10.1007/s13398-014-

0173-7.2

Vaughn, C. C., Atkinson, C. L., & Julian, J. P. (2015). Drought-induced changes in flow

regimes lead to long-term losses in mussel-provided ecosystem services. *Ecology and*

Evolution, *5*(6), 1291–1305. https://doi.org/10.1002/ece3.1442

Vohmann, A., Borcherding, J., Kureck, A., bij de Vaate, A., Arndt, H., & Weitere, M. (2010).

Strong body mass decrease of the invasive clam *Corbicula fluminea* during summer.

Biological Invasions, *12*(1), 53–64. https://doi.org/10.1007/s10530-009-9429-0

Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin,

J.-M., Hoegh-Guldberg, O., & Bairlein, F. (2002). Ecological responses to recent climate

change. *Nature*, *416*(6879), 389–395. https://doi.org/10.1007/978-3-319-10157-6_4

West-Eberhard, M. J. (2003). *Developmental plasticity and evolution*. Oxford University

Press.

TABLES

Table 1. Summary of ecological and growth variables (mean ± SD) of worldwide invasive populations of the Asian clam *Corbicula* reviewed from the literature. *Density* is the estimated abundance of clams at the study site (ind./m^2), *Pop.growth* is the rate of exponential population derived from the classical form of the exponential growth equation, *Time* is the number of years elapsed since population colonization at the time of the original study (years), *Ind.growth* is the individual growth rate (*GPI*, a *proxy* for individual growth rate), *mASM* is the minimum age at sexual maturity (months), and lifespan (years), *ID#* is a number assigned arbitrarily for identification purposes. The country, specific study location, and reference to the source paper of the original study used to build the information are shown for each population. See the main text for details on how data was collected and variables were built from the literature. For extended information on each population, see Table S1 of the Supporting Information.

Country	Study location	Time	Population parameters		Individual parameters			ID#	Reference
			Density	Pop.growth	Ind.growth	mASM	Lifespan		
Argentina	Punta Atalaya, Río de la Plata	3.5	339±405	2.20±0.01	2.87±0.07	4.30±0.13	3.83±0.57	1	Ituarte (1985)
	Paraná de las Palmas River	25.5	1,053±768	14.44±0.31	2.73±0.01	9.23±1.17	6.82±0.05	2	Cataldo and Boltovskoy (1999)
	Río Negro estuary	15.5	94±8.3	7.85±0.10	2.94±0.23	5.32±1.66	3.55±0.41	3	Hünicken et al. (2019)
France	Saone River	12.5	302±186	14.65±0.01	2.54±0.13	12.73±2.42	7.55±2.17	4	Mouthon (2001)
	Loire Lateral Canal	17.0	992±1209	41.17±0.12	2.80±0.24	8.93±5.95	5.17±3.00	5	Mouthon and Parghentanian (2004)
	Canal of Roanne	17.0	626±897	25.50±0.01	2.77±0.16	8.25±3.43	4.27±1.58	6	Mouthon and Parghentanian (2004)
Portugal	River Minho estuary	16.5	840±387	11.89±0.06	3.32±0.05	3.06±1.20	2.28±0.30	7	Sousa et al. (2008b)
	Mondego estuary	6.9	5,017±5,489	7.67±0.04	–	–	–	8	Franco *et al.*, 2012
	Casal de São Tomé	10.3	3,255±1,115	4.06±0.04	3.14±0.12	3.83±1.69	2.75±0.51	9	Rosa et al. (2014)
USA	Altamaha River	4.5	712±1,613	0.37±0.00	–	–	–	10	Gardner et al. (1976)
	Lake Arlington	2.0	32±11	11.94±0.03	3.49±0.05	5.80±0.60	2.59±0.20	11	Aldridge and McMahon (1978)
	Delta-Mendota canal	28.0	9,470±7067	4.53±0.03	2.77±0.17	9.69±4.07	8.03±2.75	12	Eng (1979)
	New River (Station 1)	2	316±654	9.40±0.02	–	–	–	13	Graney et al. (1980)
	Clear Fork, Trinity River	7.0	4,474±3,899	19.34±0.04	3.00±0.16	6.76±3.62	7.07±2.83	14	McMahon and Williams (1986)
	Mechums River	10.5	669±636	188.52±0.62	2.56±0.15	9.20±4.41	3.31±1.21	15	Hornbach (1992)
	Ogeechee River	10.0	1,398±1,096	6.23±0.00	2.52±0.12	22.72±9.17	6.85±1.63	16	Stites et al. (1995)
	Lake Tahoe	8.0	1,809±723	16.64±0.04	2.80±0.12	6.90±1.90	2.70±1.32	17	Denton et al. (2012)

Table 2. Summary of results from reduced models of weighted linear regression analyses of invasion and growth variables of global invasive populations of *Corbicula*. Parameter values are means ± SE. We report one-tailed *P* values indicating significant results in bold characters. Note natural logarithmic transformations of variables. For details on descriptive statistics of full models and results on partial *F*-tests see the Supporting Information. All variables were Ln-transformed and standardized to zero mean and a unit of variance (see *Statistical Analyses* for details).

Response variable	Predictor	Estimate ± SE	Intercept ± SE	R^2	*F*-statistics	*num.df, den.df*	*one-tailed* P-value
Density	*Time*	0.454 ± 0.196	-0.906 ± 0.241	0.263	5.34	1, 15	**0.018**
Population growth rate	*Time*	0.936 ± 0.323	-0.518 ± 0.216	0.376	4.03	2, 14	**0.006**
	Density	-0.522 ± 0.407					0.220†
Individual growth rate	*Density*	-0.422 ± 0.265	0.098 ± 0.264	0.283	2.17	2, 11	0.070
	Population growth rate	-0.473 ± 0.344					0.099
minimum Age at sexual maturity	*Individual growth rate*	-0.505± 0.179	-0.192 ± 0.171	0.601	8.28	2, 11	**0.008**
	Population growth rate	0.460 ± 0.208					**0.024**
Lifespan	*Individual growth rate*	-0.718± 0.184	-0.056 ± 0.192	0.559	15.20	1, 12	**0.001**

† two-tailed *P*-value.

FIGURES

Figure 1. Global distribution of invasive *Corbicula* clams indicating the native range of the genus (dark grey), the distribution of
Corbicula inside (black/grey shading and black dots) and outside of the genus' native range (color dots), which are mostly represented by forms
A/R (formerly designed as *Corbicula fluminea sensu lato*; see main text for detailed explanation). Numbered stars are invasive worldwide
populations reviewed in the present study. Years indicate estimated date of first introduction. See Table 1 and main text for references and details
on the invasive populations reviewed here. See the Supporting Information for detailed criteria and data used to build the genus native and
invasive distributions.

 Figure 2. Relationships between processes and characteristics at different levels of
organization of 17 global invasive populations of the Asian clam *Corbicula*. At the population
level: a) the effect of time since colonization (*Time,* years) on population *Density* (ind./m^2)
and b, c) the simultaneous effect of *Density* and *Time* on *Population growth rate*. At the
interaction between population and individual level: d, e) the simultaneous effect of density
and population growth rate on individual growth rate, and f) the effect of *Population growth
rate* on the minimum *Age at sexual maturity* (months). At the individual level: g) the effect of
Individual growth rate on the minimum *Age at sexual maturity*, and h) on the *Life span*
(years). When plotting the fit for one variable after testing for simultaneous effects, we
removed the effect of the variable not considered (b, c, d, e). Dashed lines show fits when
marginally non-significant 0.05> α <0.10. Solid lines show linear regression fits when
statistically significant at α < 0.05. The size of circles correspond to the inverse of the
variance of mean value of each observation of the response variable, used to weigh effect
sizes in the regression analyses. All variables were Ln-transformed and standardized to zero
mean and a unit of variance (see *Statistical Analyses* for details).

Figura 3. Summary of the relationships between processes and characteristics at different
levels of organization of 17 global invasive populations of the Asian clam *Corbicula*, showing
positive (+), negative (–), and no relationships (0) after fitting weighted linear regressions.
Thick solid lines (➡) show linear regression fits when statistically significant at α < 0.01.
Thin solid lines (⟶) show linear regression fits when statistically significant at α < 0.05.
Dashed lines (– –➤) show fits when marginally non-significant 0.05> α <0.10. Dotted line
(·····➤) show no significant relationship α > 0.10. Relationships are shown at three different
levels: A) at the population level, B) the interaction between processes and characteristics at
the population and individual level, and C) at the individual level.

804