Fitness-related traits are maximized in recently introduced, slow-growing populations of an invasive clam: is this a response to strong *r*-selection?

Leandro A. Hünicken[1,2,3]*, Francisco Sylvester[2,3], Nicolás Bonel[2,4,5]

[1] Centro de Investigación Aplicada y Transferencia Tecnológica en Recursos Marinos Almirante Storni, Escuela Superior de Ciencias Marinas, Universidad Nacional del Comahue, San Martín 247, (8520), San Antonio Oeste, Río Negro, Argentina

[2] Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

[3] Instituto para el estudio de la Biodiversidad de Invertebrados, Facultad de Ciencias Naturales, Universidad Nacional de Salta, Av. Bolivia 5150 (4400), Salta, Argentina.

[4] Laboratorio de Zoología de Invertebrados I, Departamento de Biología, Bioquímica y Farmacia, Universidad Nacional del Sur (UNS), San Juan 670, (B8000ICN) Bahía Blanca, Argentina.

[5] Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175, CNRS—Université de Montpellier, Université Paul-Valéry Montpellier—Ecole Pratique des Hautes Etudes—IRD, 34293 Montpellier cedex 05, France.

* Corresponding author. E-mail: leandrohunicken@gmail.com

Author contributions statement: LAH, FS, and NB conceived the study. LAH, FS, and NB conducted the literature search and extracted the data. NB designed the methodological approach. LAH and NB performed the data analysis. NB led the writing.

ABSTRACT

Many species are shifting their ranges being forced to rapidly respond to novel stressful environmental conditions. Colonizing individuals experience a complex mixture of selective forces that favor rapid evolution of life history traits notably affecting dispersal and reproductive rates in newly invaded habitats. Limited information is currently available on trait variation within the invasive range despite being critical for understanding ecological and evolutionary factors that drive the process of range expansion of invasive species. Here we evaluated life history shifts of the widely introduced Asian clam *Corbicula* within its invaded range. Through an exhaustive literature search, we obtained data for 17 invasive *Corbicula* populations from different ecosystems worldwide to test the relationship between population and individual parameters relevant to the process of range expansion. Our main results show that (i) recently introduced *Corbicula* populations are characterized by low conspecific density and low population growth rate, (ii) clams reproduce earlier in slow-growing populations, and (iii) density had no effect on population increase. Our findings support the standpoint that individuals from recently established populations of *Corbicula* face strong *r*-selective forces that favor traits associated with high reproductive rates, which could be interpreted as an effective mechanism to overcome difficulties associated with low densities and low population growth in newly colonized areas.

Key words: Asian clam, Invasive species, Individual growth, Population increase, *r*-selection.

INTRODUCTION

The ongoing increase in frequency of extreme climatic events, caused by anthropogenic climate change, and global transport of people and goods is altering ecosystems by triggering a wide array of irreversible consequences (Walther et al. 2002; Bossdorf et al. 2005; Hoegh-Guldberg and Bruno 2010; IPCC 2014). Changes in local conditions, in particular those driven by human-induced climate change, are predicted to affect ecological processes across several levels of organization, likely over the coming decades. Such changes are altering the performance of individual organisms, the dynamics of populations and the distribution of species, ultimately modifying ecosystem properties and function (Hoffmann and Sgró 2011; Lonhart et al. 2019). Hence, many species are forced to rapidly respond to novel stressful environmental conditions when shifting their distribution and colonizing new areas.

If non-native species have difficulties in adapting to such conditions and fail to track projected environmental changes, populations become vulnerable to decline and extinction (Hill et al. 2011; Hoffmann and Sgró 2011). These stressful conditions can have a strong effect on species life histories through the selection of traits that optimize population growth and survival from offspring to maturity (e. g. reduced age of maturity and reproduction, increased growth rate, increased energy allocated to reproduction; Stearns, 1976, 1977). Thus, individuals at the expanding edge of the species distribution experience a complex mixture of selective forces that can have profound impacts on the evolution of a species' life history traits affecting, for instance, dispersal and reproductive rates (Phillips 2009; Shine et al. 2011).

Colonization events through either continuous range expansion or jump dispersal are usually driven by a small number of individuals that are the first to arrive at a new point

of the distribution, where they benefit from increased space open for colonization and

low conspecific density, below carrying capacity (Cole 1954; Sakai et al. 2001; Phillips

et al. 2010). Hence, the density gradient occurring at the expanding range (i. e. grading

from zero to carrying capacity) creates different selective environments where

individuals from a recently established population face stronger r-selection compared to

those from long established ones, which are exposed to selective K-type forces (e. g.

Phillips et al., 2010). According to theory, low densities favor the evolution of life-

history traits related with r-type selections (MacArthur and Wilson 1967; Phillips 2009;

Burton et al. 2010), leading to exponential (non-density regulated) population growth in

recently established populations while denser, long-established ones follow a density-

regulated population growth (Brook and Bradshaw 2006).

Recently established populations are characterized by low conspecific density and low

rate of increase, often leading to lag-times between initial colonization and the onset of

rapid population growth (Sakai et al. 2001). The duration of the lag phase is mediated,

among others, by a number of non-mutually exclusive ecological and evolutionary

mechanisms (e. g. low encounter rate for reproduction, density-dependent effects,

adaptation, and selection of new genotypes; Bossdorf et al., 2005; Davis, 2005;

Ricciardi, 2012). An effective mechanism that increase the rate of population growth is

early reproduction, which results from fast growing individuals (Cole 1954; Lewontin

1965; Roff 1993). Individuals that grow faster will attain reproductive sizes earlier and

bring down generation times, promoting a rapid build-up of population numbers (Cole

1954; Lewontin 1965; Roff 1993; Phillips 2009) and accelerating primary invasion and

secondary spread of invasive species (Skellam 1951; Phillips et al. 2006; Lockwood et

al. 2007). Several studies have compared life-history traits between native and invaded

ranges ('home-and-away' comparisons). Such comparisons could lead to erroneous

interpretations of the patterns observed because it would be difficult to disentangle what

were the processes (e.g. propagule bias and/or evolution in populations at spatial

(dis)equilibrium) causing the shifts in life-history traits (Phillips et al., 2010 and

references therein). Interestingly, only few studies have looked for such trait changes on

introduced species across their global invasive range (Gaston 2009) despite knowledge

on life-history trait variation within the invasive range is critical for better

understanding the process of species range expansion, largely induced by climate

change and increasing global transport.

Bivalves of the genus *Corbicula* (Megerle Von Mühlfeld 1811) are native to Southeast

Asia, the Middle East, Australia and Africa (Araujo et al. 1993) but they have colonized

a large part of the Americas and Europe (Mouthon 2001; Schmidlin and Baur 2007;

Crespo et al. 2015). The successful invasion of *Corbicula* clams has been mainly

attributed to their rapid growth and maturation, high fecundity, and high dispersal

making this bivalve genus one of the most relevant faunal non-indigenous groups in

aquatic ecosystems (reviewed in Sousa et al. 2008a). Invasive freshwater *Corbicula*

populations are constituted by a small number of lineages composed of hermaphroditic

individuals, which reproduce through androgenesis, where the most widespread and

abundant lineage is the so-called Form A/R, widely known as *C. fluminea* (Komaru et

al. 1998; Pigneur et al. 2012, 2014). The current taxonomic status of the invasive

Corbicula lineages in the Americas and Europe is still largely unresolved despite

several morphological and genetic studies (reviewed in Pigneur et al. 2014). In the

present paper, we deal with this form and refer to it as *Corbicula* to avoid entering

unsettled taxonomic debates.

Here we investigated how different life-history traits of the Asian clam *Corbicula*

responded to different selective pressures occurring at varying population contexts

within the introduced range: (i) newly- vs. long-established, (ii) low vs. high density,

and (iii) low vs. high population growth rates; all of them being linked to the geographic

range shifting of species' distribution (i. e. front/core populations). To do so, we

searched for peer-reviewed articles providing enough information to estimate time since

population introduction as well as different population and individual processes and

characteristics including population density, population increase, individual growth rate,

minimum age at sexual maturity, and lifespan of global populations of this worldwide

invader. We focus our predictions on life history strategies exhibited by individuals

from recently established populations, which typically occur at the invasion front, and

likely subjected to strong *r*-selective pressures with respect to those from older, long-

established ones. We tested the hypotheses that population density and population

growth increase with time since introduction. *Corbicula* is widely known to exhibit

rapid population growth, being able to reach extremely high densities (reviewed in

Mcmahon, 2002). Different studies suggest that *Corbicula*'s population dynamics are

chiefly regulated by environmental conditions (e.g. Crespo et al., 2015, 2017; Gama et

al., 2017), whereby population declines are typically driven by periodic mass mortality

due to extreme hydro-meteorological events (Ilarri et al. 2011; McDowell et al. 2016;

McDowell and Sousa 2019). This strongly suggests that invasive *Corbicula*

populations primarily follows a non-density regulated population increase and, hence,

they might not approach the carrying capacity as it happens with many other species

(e.g. Brook and Bradshaw, 2006). In this sense, we expected density to have no

significant effect on population increase even within a wide range of densities in nature,

though high densities are considered likely to have a negative effect on individual

growth rates. We also asked whether clams that grow faster and reproduce earlier are

favored in slow-growing populations at the expanding front. A related prediction is that

clams that grow fast would have a shorter lifespan with respect to individuals exhibiting

slow growth.

METHODS

Literature search

We exhaustively searched the literature for peer-reviewed articles on *Corbicula*

(*Corbicula fluminea sensu lato*) providing enough information to estimate the

individual growth rate, population density, population growth rate, time since

colonization, water temperature, and conductivity of global populations of this clam.

To do this, we first identified relevant studies from ISI Web of Knowledge (Web of

Science Core Collection, 1900 through 2017) by conducting a first search using the

field tag "topic (TS)" in the "Advanced search" option and the search string:

TS="Corbicula fluminea" OR TS="asia* clam" (search #1). We conducted a second

search including topics TS="population dynamic*" OR TS="population size*" OR

TS="population densit*" OR TS="population structure" (search #2). We ran a third

search using TS="growth" OR TS="individual growth" OR TS="population growth"

(search #3). Finally, we combined searches #2 and #3 into search #4 (#4= #3 OR #2),

and restricted the latter by search #1 to obtain a final set of candidate studies (search

#5= #4 AND #1).

This literature search was finished on 27 September 2018 and this protocol identified

215 candidate studies (database search: 215 candidates; gray literature and unpublished

data: 0 candidates). To identify papers potentially missing in this list, we subsequently

scanned the references in these papers and conducted a Google Scholar search using

keywords mentioned above and added 8 studies to this candidate pool (Fig. S1). A total

of 67 studies that belonged to research areas unrelated to the topic of the current study

(*e. g.*, applied chemistry, toxicology, paleontology) were excluded from this candidate

pool by refining the set based in 'Web of Science Categories'. The resulting studies (n =

156) were finally inspected individually in detail to assess their eligibility for our

analyses according to the following inclusion criteria: i) Studies that assessed invasive

populations of *Corbicula* clams (i. e. studies exclusively providing estimates for native

populations were excluded); ii) Studies that reported or provided data enough to extract

the population density and age/size distribution for >6-months (see below); iii) Studies

that assessed populations in reasonably "average" natural environments, such that

populations from heavily impacted sites (e. g. thermal plumes) were excluded. We

finally retained a sample of 16 studies, covering 17 invasive populations of *Corbicula*

from different ecosystems worldwide (n = 140 out of 156 were excluded; see Fig. S1 for

further details on screening and eligibility criteria). Whenever population data was only

graphically presented, we contacted directly the authors. However, if data were no

longer available or authors did not respond, we used the WebPlotDigitizer software

(http://arohatgi.info/WebPlotDigitizer/) to extract data and accurately estimate

population density and growth rates.

Population density and population increase

For each population selected from our literature search, we obtained the mean

population density for the entire study period or extracted temporal values to estimate

it whenever this value was missing in the text. *Corbicula* clams generally exhibit a

bivoltine juvenile release pattern (i.e. producing two generations a year; Mcmahon,

2002; Sousa et al., 2008a), leading to one or more high-density peaks related to

recruitment events yearly. However, density in the field fluctuates almost constantly

generating several peaks and valleys that do not necessarily indicated significant

recruitment events. *Corbicula* populations exhibit rapid growth (Mcmahon 2002;

Sousa et al. 2006, 2008a; Franco et al. 2012). Hence, the rate of population increase of

Corbicula can be expressed by deriving the classical exponential growth equation:

182
$$r = \frac{\ln\left(\frac{N_t}{N_0}\right)}{t}$$

To estimate the rate of increase (r), we set a criterion to determine relevant recruitment

peaks in each study by using the average density throughout the study period as a

threshold. We considered the lowest density value (N_0), the highest peak above it (N_t),

and the time period elapsed between the valley and the peak (t). Intermediate density

values comprised in each period (i. e. defined by N_0 and N_t) were used to fit the growth

equation and obtain r. Population rates of increase were estimated using GraphPad

Prism version 7.00 for Windows, GraphPad Software, La Jolla California USA,

www.graphpad.com. Note that periods defined by non-relevant peaks (i. e. below the

average density threshold) were discarded, as were also periods of progressive density

decline between a relevant peak and the next N_0 value (see Fig. S2 for further details

on the step-by-step procedure to estimate population increase). We decided to do this

because the aim of the present study was to compare population growth during periods

of substantial growth across populations. When populations exhibited more than one

relevant density peak throughout the study period, the population increase rate for such

population was calculated as the average of single r estimations. Then, we

standardized estimates by a year to make them comparable.

Individual growth rates and parameters

For studies that only presented data on length-frequency distributions, we identified

age cohorts using that information and applied the Bhattacharya's method available in

FISAT II software (Version 1.2.0, FAO-ICLARM Fish Assessment Tools; (Gayanilo

et al. 2002). To confirm each component of normal distributions from the modal

progression analysis, we used the NORMSEP method also available in the FISAT II

software (Pauly and Caddy 1985). Once we had cohorts identified for all the studies,

we calculated individual growth parameters using the same equations for all the

studies considered to make sure results were fully comparable across studies. For each

cohort present in each of the populations considered in this study, we fitted the von

Bertalanffy-growth function with seasonal oscillations (hereafter SVBGF) proposed

by Pauly and Gaschutz (1979), Hoenig and Hanumara (1982), and Somers (1988):

$$L_t = L_\infty \left(1 - exp^{-K[(t-t_0) + T_1 - T_2]}\right),$$

$$T_1 = C \sin \left(2\pi \left(t - t_s\right)\right) / 2\pi,$$

and

$$T_2 = C \sin \left(2\pi \left(t_0 - t_s\right)\right) / 2\pi,$$

where L_t is the predicted length at age t, L_∞ is the asymptotic length, K is the growth

constant of dimension time (year^{-1} in most seasonally oscillating growth curves)

expressing the rate at which L_∞ is approached, t_0 is the theoretical 'age' the clam have

at length zero, C expresses the relative amplitude of the seasonal oscillation and varies

between 0 and 1 (0 indicating lack of summer-winter differences in growth, it was

constrained to be less than or equal to 1 during model fitting), and t_s is the starting

point of the oscillation.

The parameters of the function were estimated by the modeling method available in

JMP statistical software (v9.0 SAS Institute). When preliminary results of the SVBGF

failed to converge to estimate the asymptotic shell length (L_∞), we used the maximum

shell length (*Lmax*) observed or reported at/for each study site to calculate the

asymptotic shell length following the equation suggested by Taylor (1958):

$$Lmax / 0.95 = L_\infty$$

These values were fixed when we performed a second fit.

Because of the negative correlation between growth parameters (K and L_∞) prevents

making comparisons based on individual parameters (Vakily 1992; Ramón et al. 2007),

we used the growth-performance index (*GPI*), which reflects the growth rate of a given

organism of unit length. In other words, *GPI* can be viewed as the (theoretical) value of

K that would occur in organisms with a L_∞ value of 1 unit of length (Munro and Pauly

1983), and it was defined by Pauly and Munro (1984) as:

$$GPI = 2 \log_{10} L_\infty + \log_{10} K$$

We calculated the *GPI* values for cohorts from each study site based on growth

parameters (L_∞ and K) obtained after fitting the SVBGF using the *Growth*

Performance Indices application of the FISAT II software (Gayanilo et al. 2002).

Minimum age at sexual maturity

The minimum age at sexual maturity was estimated as the time (months) elapsed until

clams reach the minimum size (shell length) at which invasive *Corbicula* clams become

sexually mature, which, on average, is 10.0 mm ± 0.8 SE (n = 13 reported sizes). This

value was obtained from shell sizes reported in the literature (Heinsohn 1958; Gardner

et al. 1976; Kraemer and Lott 1977; Aldridge and Mcmahon 1978; Eng 1979; Kraemer

1979; Ituarte 1985; Kennedy and Van Huekelem 1985; French and Schloesser 1991;

Rajagopal et al. 2000; Mcmahon 2002; Cao et al. 2017). We acknowledge that in some

conditions and populations, clams might mature at a different pace and thus introduce a

degree of error in such cases. While this error is unavoidable until more detailed data is

available, we are confident that the 10-mm estimate applies to many real-life situations

and thus our results based on it are robust.

Lifespan

As *t0* is the theoretical 'age' the clam have at length zero, *tmax* can be defined as the

theoretical 'age' the clam reaches its maximum length (*Lmax*). Since *Lmax* is

dependently related to a maximum point in time (years), it is assumed that *Lmax*

implicates *tmax* (*Lmax* $\Rightarrow$ *tmax*) (Bonel and Lorda 2015). Thus, the theoretical

maximum lifespan of a clam cohort can be approximately express as:

$$Lifespan = tmax - t0$$

Time since colonization and key environmental variables

For the same 17 populations and sites studied, we calculated the time elapsed since

population foundation (hereafter time since colonization) as the difference between the

sampling dates in the first ecological survey and the first report of introduction. The

accuracy of the introduction date is generally very difficult to know, however, many of

the studies used report range expansions of a conspicuous invader across well studied

regions, where the appearing of the species might have been anticipated and readily

detected by ongoing monitoring actions in the initial stages of invasion. Therefore, we

consider that the possible error introduced should be moderated to low and the value

obtained can be view as a good approximation. Moreover, we obtained water

temperature and conductivity values for each site averaging monthly mean values for

the same period used to estimate growth rates in each study. Other biotic and abiotic

variables are known to interact with temperature to drive growth rates of invasive

bivalves (e.g. food availability, dissolved oxygen, heavy metal concentration; e.g.

(Vohmann et al. 2010; Bonel and Lorda 2015). However, we only considered water

temperature and conductivity in the analyses because they have been reported to play a

key role on the *Corbicula*'s biology and ecology (Magnussen 2007; Franco et al. 2012;

Crespo et al. 2017). Indeed, temperature has emerged as the most important variable

explaining the current distribution and, also, predicting that climate change will favor

the expansion of *Corbicula*'s into colder areas at higher latitudes (Gama et al. 2017).

Likewise, salinity has been considered a key factor influencing the success and

velocity of the invasion of new estuarine ecosystems (e.g. Sousa et al., 2006). As

Corbicula populations considered herein belonged to a broad range of freshwater and

estuarine habitats, we therefore included conductivity, which serve as a *proxy* of

salinity levels. In order to keep values seasonally unbiased, only entire years were used

in subsequent calculations. Whenever these parameters were not reported in the

original paper, we obtained values for the same sites and years from the literature and

reliable online databases (e. g. https://waterdata.usgs.gov/nwis/qw). When data was

only graphically presented, we extracted it following the procedure mentioned above.

Statistical analyses

We ran five separate multivariate weighted linear regressions in order to test ecological

processes and characteristics at three different (but non-mutually exclusive) levels: 1)

population level, 2) population and individual level, and 3) individual level. In this

sense, a response variable at the population level can be considered as an explanatory

variable when analyzing a response at the individual level. In all cases, we computed the

variance of the effect size from different populations, which were then used as weights

in linear regressions. The inverse-variance weighting allows combining the results from

independent measurements giving more importance to the least noisy measurements and

vice versa (Borenstein et al. 2009).

We were interested in testing individual effects of predictors, which are related to life-

history traits of *Corbicula* (hereafter reduced models). However, we added annual mean

water temperature and conductivity as quantitative variables in each model (hereafter

full models). Preliminary analyses showed that latitudinal effects are strongly correlated

with water temperature ($r = -0.84$). Hence, it was excluded from the models to avoid

multicollinearity. In some cases, we also included interaction terms between

explanatory variables. We used the partial F-test to compare models—that is, full

models with and without the interaction term, and full *versus* reduced models. As we

had strong *a priori* hypotheses as to the direction of effects, we used one-tailed tests.

Variables were Ln-transformed to fulfill the models' assumption of normality and

homoscedasticity, which were tested using the *gvlma* package (Peña and Slate 2006).

Prior to analyses, we standardized explanatory variables to zero mean and a unit of

variance in order to get directly comparable effects. All analyses were carried out in R

version 3.6.1 (R Development Core Team 2019). Values are given as means ± SE unless

otherwise stated.

RESULTS

Literature search

Populations analyzed in this study belonged to Europe, North and South America (Fig.

1; Table 1). These included eight populations in the United States, three in France, three

in Argentina, and three in Portugal (see Table 1 for detailed location, bibliographic

source, and code number used herein to refer to the populations, for each of the 17

Corbicula populations selected, and Table S1 for additional information).

Density, growth, time since colonization, and key environmental variables

Overall, mean density of *Corbicula* from 17 locations worldwide spanned 32 to *ca.*

9,500 ind. m^{-2}, population rate of increase (*r*) varied from 0.37 to 189, individual growth

rate (*GPI*) ranged 2.53-3.49, age at first reproduction 3.1-22.7 months, and lifespan 2.3-

8.0 years (Table 1). On the other hand, time since colonization ranged 2 to 28 years in

the populations studied (Table 1) and yearly water temperature ranged from 13.2 to 20.7

°C and conductivity from 68 to 8,457 µS/cm (see Table S1 of the Online Resource for

details). In all cases, we found no evidence that adding an interaction term with

variables of interest nor temperature and conductivity in each model significantly

increase its predictive power (for details see '*Model comparison*' in Online Resource).

We therefore report estimates and statistics from the reduced models in Table 2.

At the population level, we found that younger *Corbicula* populations (i.e. newly

established) were characterized by low density and slow population growth rate with

respect to older, long established ones (Fig. 2a, b). We found that density had no

significant effect on population increase (Fig. 2c). When analyzing responses at the

interaction between population and individual processes and characteristics, we found

that clams tended to grow faster (marginally non-significant) in low-dense and slow-

growing populations (Fig. 2d, e). In contrast, we observed a strong positive response of

the minimum age at sexual maturity as a function of population increase, clearly

indicating that clams reproduce earlier in slow-growing populations (Fig. 2f). This is

directly linked to processes occurring at the individual level, where clams with higher

growth rates reach the minimum reproductive size earlier (10 mm) and live less than

clams that grow more slowly (Fig. 2g, h).

DISCUSSION

This is the first study that evaluates how population and individual parameters, involved

in the process of range expansion, varied in response to different selective pressures

occurring within the introduced range of the worldwide invasive Asian clam *Corbicula*.

Our main results showed that (i) recently introduced *Corbicula* populations are

characterized by low density and reduced rate of increase, (ii) clams reproduce earlier in

slow-growing populations, and (iii) population growth was unfettered by density

(summarized in Fig. 3).

Density of *Corbicula* populations increased with time since colonization, which is

consistent with the standpoint that recently established populations typically occur at

low conspecific density, below the carrying capacity (Cole 1954; Sakai et al. 2001).

Likewise, we found that population growth increased with time since colonization,

suggesting that, in newly colonized areas, populations could have experienced stronger

r-type selection favoring fitness-related traits effectively increasing the rate of

population growth, as it was observed in long established populations (MacArthur and

Wilson, 1967; Phillips, 2009; Burton et al., 2010). Population density did not predict

changes in the rate of population increase, as variation in population growth remained

relatively low even at high-density values. By definition, long established and denser

populations tend to be density regulated as they approach to carrying capacity (logistic

population growth; e. g. Phillips et al. 2010). Considering the so-called *r*/K selection

continuum, this implies that such population is predicted to shift from the *r*- towards the

K-endpoint (MacArthur and Wilson 1967)—that is, selection shifts a population from

density-independent to density-dependent regulation. However, *Corbicula*'s population

growth rate did not decrease with increasing density despite the fact that some of the

populations included in this study exhibit extremely high densities exceeding, in some

cases, 100,000 individuals per square meter (e. g. Eng 1979; Table 1).

Why population growth was unfettered by density regulation? A large number of

studies have analyzed density dynamics of *Corbicula* and commonly assumed that,

following exponential growth, the clam population achieves densities approaching the

carrying capacity of the environment (e.g. Sousa et al., 2006). However, to our

knowledge, no study have ever conducted (field or laboratory) experiments to test

whether population growth is limited by increasing conspecific density, which, in turn,

is expected to reduce local biotic resources. In other words, there is a complete lack of

knowledge on the true effect of density on population growth. On the contrary, different

studies have shown that *Corbicula*'s dynamics are chiefly influenced by environmental

variables (e. g. water temperature, conductivity, pH, and/or dissolved oxygen) and seem

to be less influenced by population's density (e. g. Crespo et al., 2015; Gama et al.,

2016). Indeed, periodic mass mortality due to extreme hydro-meteorological events (e.g.

extremely high or low temperatures, drought) could play an important role with

infrequent, but catastrophic population declines (>99% in some cases; Mouthon and

Parghentanian 2004; Ilarri et al. 2011; McDowell et al. 2017). These events have been

described (though not quantified) in several other systems as well (e.g. Vaughn et al.

2015; McDowell and Sousa 2019; Fey et al. 2015). Despite a given *Corbicula*

population might approach or even grow beyond the carrying capacity of the ecosystem,

evidence from previous studies (and those cited herein) suggests that population size

significantly varies with time. This means that the population is at non-equilibrium and

that such fluctuations (or boom-bust dynamics; Strayer et al., 2017) are likely driven by

density-independent factors, as the above-mentioned hydro-meteorological events.

Thus, it does seem reasonable to infer that *Corbicula*'s population growth rate is non-

density-regulated and do not level off at carrying capacity, as expected from a highly

invasive organism as the Asian clam (McMahon 2002). Our results suggest that this

bivalve seems to be far from approaching carrying capacity, even at extremely high

densities, supporting the idea that individuals from long established denser populations

do not face a *K*-selective environment but a less strongly *r*-selective pressure.

Individuals tended to grow faster in low-density populations experiencing low rates of

increase. Likewise, *Corbicula* clams reproduce earlier in slow-growing populations. In

recently introduced populations, periods of negligible population increase can be driven

by non-mutually exclusive ecological and evolutionary adjustments such as Allee effect,

changing abiotic conditions, and adaptation and selection of new genotypes imposed by

novel environments conditions at the newly habitat (Sakai et al. 2001; Bossdorf et al.

2005; Davis 2005; Ricciardi 2012). Such conditions can lead to a strong selection on

life-history traits that optimize population increase and survival to maturity of offspring

(Stearns 1976, 1977). As predicted by life-history theory, individuals from recently

established populations (typically occurring at the invasion front) face stronger *r*-

selection relative to conspecifics from older populations (MacArthur and Wilson 1967),

favoring traits that accelerate the range expansion. For instance, some empirical

observations have revealed that individuals from recently colonized areas grow faster,

and thus, reproduce earlier, than do conspecifics from older, long-established

populations (e. g. tallow trees: Siemann and Rogers 2001; butterflies: Hanski et al.

2006; Australian cane toads: Phillips 2009). Accordingly, our results for *Corbicula*

clams showing fast growth and, thus, early reproduction support the standpoint that

such populations faced strong *r*-selection, which favored traits associated with high

reproductive rates in recently introduced, slow-growing populations.

The increase in individual growth is associated with a decrease of the clams' life span,

supporting the idea of compensatory responses (i. e. trade-off) between different life-

history traits. Such trade-offs can be interpreted as an effective mechanism to bring

down generation times, promoting a rapid build-up of population numbers, which

increases the population growth in newly colonized areas (Cole 1954; Lewontin 1965;

Roff 1993) and accelerate primary invasion and secondary spread of invasive species

(Skellam 1951; Phillips et al. 2006; Lockwood et al. 2007). Collectively, our results

suggest that reduced population increase of the current *Corbicula* population triggers

the shortening of the time that the clams need to reach maturity by favoring clams that

grow the fastest. Hence, as early-maturing clams reproduce earlier, this has the effect of

fostering the increase in size of the future population. In this way, incipient *Corbicula*

populations can increase their chances of overcoming the difficulties associated with

low densities and survive to the population establishment stage, which can clearly

subsidize the bivalve's invasive abilities in newly colonized areas.

One can argue that the issues associated to low densities (e.g. Allee effect) would not

apply to *Corbicula* populations as it has been largely assumed that this invasive bivalve

reproduces mainly by self-fertilization, meaning that they do not need to find a partner

to mate. However, cross-fertilization seems to be the norm in *Corbicula* clams whereas

self-fertilization takes place on a less regular basis (see pp. 200, Kraemer et al., 1986),

suggesting that this bivalve is rather a facultative selfer. In line with their findings,

Hünicken et al. (2019) reported extremely low population densities at the southernmost

invasion front in Patagonia where *Corbicula* has been established in the area for more

than 15 years. This suggests that this clam would be predominately outcrosser, and not

selfer because, otherwise, they would have shown an increase in density with time since

colonization, which was not the case. Notably, despite the evidence found by Kraemer

et al. (1986) and their conclusions, somewhat the idea of self-fertilization as the norm

(and not as the exception), has been spread in the literature as an effective mechanism

by which a single individual would be capable of founding new populations (reviewed

in Mcmahon, 2002). Thus, experimental assays are needed to clearly determine under

which conditions invasive *Corbicula* would reproduce by selfing.

Is evolutionary adaptation to local conditions possible in *Corbicula*? All invasive

Corbicula populations analyzed in this study are constituted by androgenetic individuals

belonging to the same lineage (Form A/R), which are distributed all across Europe and

the Americas and characterized by an extremely low genetic diversity (see Pigneur et al.

2012, 2014 for further details). The question that arises is whether there is scope for

evolution in genetically depleted populations of *Corbicula*. One fundamental principle

of evolutionary biology is that the rate of change in response to natural selection is

proportional to the amount of additive genetic variation present (Fisher 1930). Decrease

in genetic diversity results in reduced evolutionary potential required to rapidly adapt to

the novel habitat conditions of the invaded range, among other issues related to critical

invasion phases (e.g. initial colonization, leading edges of range expansion; Schrieber

and Lachmuth 2017). It is still puzzling how *Corbicula*'s genetically depleted

populations succeed in their invasion process, which represents a case of genetic

paradox (Pigneur et al. 2014). One possible explanation is that invasive *Corbicula*

might exhibit a certain degree of genetic polymorphism linked to androgenesis (Pigneur

et al. 2012). Indeed, androgenesis in *Corbicula* clams combines features of clonal

reproduction and the ability of rare genetic material exchange through 'egg parasitism'.

Among other processes, egg parasitism enables a mixing of different nuclear genomes

when the maternal nuclear genome is incompletely extruded creating triploid

individuals (invasive freshwater *Corbicula* clams produce unreduced spermatozoa; i.e.

sperm cells are diploids; Pigneur et al. 2014).

The invasive lineage (Form A/R) considered in our study includes both diploid and

triploid populations, with triploidy being predominant (further details in Pigneur et al.,

2014 and references therein; Etoundi et al., 2019). As polyploidy can buffer deleterious

mutations and can lead to hybrid vigor (heterosis) with increased fitness (e.g. Otto 2007;

Selmecki et al. 2015), polyploid *Corbicula* clams with genotypes associated with broad

tolerance to environmental stressful conditions (usually found at the invasion front) may

be favored during an invasion process, and in the long term a general-purpose genotype

may evolve (Pigneur et al. 2012). Therefore, evolved adaptation to local stressful

conditions in newly colonized areas might be possible, though evolved responses might

be limited by low levels of quantitative genetic variation (e.g. Dybdahl and Drown,

2011). Hence, further studies should focus on analyzing the level of ploidy within the

most common invasive lineage (From A/R) and determine whether front/newly

established populations show a higher frequency of triploids individuals, and thus,

higher fitness, relative to those from core/long-established populations.

In conclusion, in this study we provide evidence supporting that different life history

traits of individuals can respond to different selective pressures occurring at varying

population contexts (e.g. time elapsed since the population foundation, low vs. high

population density, and low vs. high rates of increase). Our results show that fitness-

related traits, such as individual growth rate and minimum age at sexual maturity, are

maximized in recently established, slow-growing populations, which experience a

stronger *r*-selective environment with respect to long established ones. While these

responses might be interpreted as an effective mechanism to decrease the risk of

stochastic extinction associated with low densities at the invasion front, population

increase seems to be independent of population density in a wide range of

environmental conditions within *Corbicula*'s invasive range. Whether trait shifts

resulted from plasticity or genetically based evolutionary change remain as an open

question that could be addressed by conducting laboratory experiments under common-

garden conditions. Although other relevant variables could not be considered in our

models, this study represents a first step in the attempt to understand in a wide

geographical perspective whether evolvability of fitness-related traits increases

invasiveness of *Corbicula* during processes of range shift. Knowledge of rapid

evolutionary adaptation is of high concern as a steadily increasing number species are

forced to rapidly respond to novel stressful conditions as consequence of increasing

global transport and human-induced climate change.

ACKNOWLEDGEMENTS

We would like to thank Tim Janicke, Maxime Dubart, and Ronaldo Sousa for valuable

comments on previous versions of this manuscript. This project was partially financed

by PICT 2016-0631 from the Argentine Agencia Nacional de Promoción Científica y

Tecnológica to FS. NB was partially supported by the "Programa de Financiamiento

Parcial de Estadías en el Exterior para Investigadores Asistentes", National Scientific

and Technical Research Council CONICET (Res. Nº 1236/08; 4118/16)".

REFERENCES

Aldridge DW, Mcmahon RF (1978) Growth, fecundity, and bioenergetics in a natural

population of the Asiatic freshwater clam, *Corbicula manilensis* Philippi, from

North Central Texas. J Molluscan Stud 44:49–70

Araujo R, Moreno D, Ramos MA (1993) The Asiatic clam *Corbicula fluminea* (Müller,

1774) (Bivalvia: Corbiculidae) in Europe. Am Malacol Bull 10:39–49

Bonel N, Lorda J (2015) Growth and body weight variability of the invasive mussel

Limnoperna fortunei (Mytilidae) across habitat and season. Malacologia 58:129–

145. https://doi.org/10.4002/040.058.0202

Borenstein M, Hedges L V, Higgins JPT, Rothstein HR (2009) Introduction to meta-

analysis, First edit. John Wiley & Sons, Ltd, Chichester, West Sussex

Bossdorf O, Auge H, Lafuma L, Rogers WE, Siemann E, Prati D (2005) Phenotypic and

genetic differentiation between native and introduced plant populations. Oecologia

144:1–11. https://doi.org/10.1007/s00442-005-0070-z

Brook BW, Bradshaw CJA (2006) Strength of evidence for density dependence in

abundance time series of 1198 species. Ecology 87:1445–1451.

https://doi.org/10.1890/0012-9658(2006)87[1445:SOEFDD]2.0.CO;2

Burton OJ, Phillips BL, Travis JMJ (2010) Trade-offs and the evolution of life-histories

during range expansion. Ecol Lett 13:1210–1220. https://doi.org/10.1111/j.1461-

0248.2010.01505.x

Cao L, Damborenea C, Penchaszadeh PE, Darrigran G (2017) Gonadal cycle of

Corbicula fluminea (Bivalvia: Corbiculidae) in Pampean streams (Southern

Neotropical Region). PLoS One 12:1–16.

https://doi.org/10.1371/journal.pone.0186850

Cataldo D, Boltovskoy D (1999) Population dynamics of *Corbicula fluminea* (Bivalvia)

in the Parana River Delta (Argentina). Hydrobiologia 380:153–163.

https://doi.org/10.1023/A:1003428728693

Cole LC (1954) The population consequences of life history phenomena. Q Rev Biol

29:103–137. https://doi.org/10.1086/400074

Crespo D, Dolbeth M, Leston S, Sousa R, Pardal MA (2015) Distribution of *Corbicula*

fluminea (Müller, 1774) in the invaded range: a geographic approach with notes on

species traits variability. Biol Invasions 17:2087–2101.

https://doi.org/10.1007/s10530-015-0862-y

Crespo D, Leston S, Martinho F, Pardal MA, Dolbeth M (2017) Survival of *Corbicula*

fluminea (Müller, 1774) in a natural salinity and temperature gradient: a field

experiment in a temperate estuary. Hydrobiologia 784:337–347.

https://doi.org/10.1007/s10750-016-2887-3

Davis HG (2005) r-Selected traits in an invasive population. Evol Ecol 19:255–274.

https://doi.org/10.1007/s10682-005-0912-5

Denton ME, Chandra S, Wittmann ME, Reuter J, Baguley JG (2012) Reproduction and

population structure of *Corbicula fluminea* in an oligotrophic subalpine lake. J

Shellfish Res 31:145–152. https://doi.org/10.2983/035.031.0118

Dybdahl MF, Drown DM (2011) The absence of genotypic diversity in a successful

parthenogenetic invader. Biol Invasions 13:1663–1672.

https://doi.org/10.1007/s10530-010-9923-4

Eng LL (1979) Population dynamics of the Asiatic clam, *Corbicula fluminea* (Müller),

in the concrete-lined Delta-Mendota Canal of Central California. In: Britton JC

(ed) Proceedings, First International Corbicula Symposium. Texan Christian

University, Fort Worth, Texas, pp 39–67

Etoundi E, Marescaux J, Vastrade M, Debortoli N, Hedtke SM, Pigneur L-M, Vurgo J,

Flot J-M, Van Doninck K (2019) Distinct biogeographic origins of androgenetic

Corbicula lineages followed by genetic captures. bioRxiv 590836.

https://doi.org/http://dx.doi.org/10.1101/590836

Fey SB, Siepielski AM, Nusslé S, Cervantes-Yoshida K, Hwan JL, Huber ER, Fey MJ,

Catenazzi A, Carlson SM (2015) Recent shifts in the occurrence, cause, and

magnitude of animal mass mortality events. Proc Natl Acad Sci U S A 112:1083–

1088. https://doi.org/10.1073/pnas.1414894112

Fisher RA (1930) The genetical theory of natural selection. Clarendon Press, Oxford

Franco JN, Ceia FR, Patrício J, Modesto V, Thompson J, Marques JC, Neto JM (2012)

Population dynamics of *Corbicula fluminea* (Müller, 1774) in mesohaline and

oligohaline habitats: invasion success in a Southern Europe estuary. Estuar Coast

Shelf Sci 112:31–39. https://doi.org/10.1016/j.ecss.2011.07.014

French JRP, Schloesser DW (1991) Growth and overwinter survival of the Asiatic clam,

Corbicula fluminea, in the St . Clair River, Michigan. Hydrobiologia 219:165–170

Gama M, Crespo D, Dolbeth M, Anastácio P (2016) Predicting global habitat suitability

for *Corbicula fluminea* using species distribution models: the importance of

different environmental datasets. Ecol Modell 319:163–169.

https://doi.org/10.1016/j.ecolmodel.2015.06.001

Gama M, Crespo D, Dolbeth M, Anastácio PM (2017) Ensemble forecasting of

Corbicula fluminea worldwide distribution: projections of the impact of climate

change. Aquat Conserv Mar Freshw Ecosyst 27:675–684.

https://doi.org/10.1002/aqc.2767

Gardner J., Woodall WR, Staats AA, Napoli JF (1976) The invasion of the Asiatic clam

(*Corbicula manilensis* Philippi) in the Altamaha river, Georgia. Nautilus

(Philadelphia) 90:117–125. https://doi.org/10.1038/scientificamerican01221846-

76b

Gaston KJ (2009) Geographic range limits: achieving synthesis. Proc R Soc B Biol Sci

276:1395–1406. https://doi.org/10.1098/rspb.2008.1480

Gayanilo FC, Sparre P, Pauly D (2002) FAO-ICLARM Stock Assessment Tools

(FiSAT). Software version 1.2.0. FAO, Roma

Graney Jr RL, Cherry DS, Cairns Jr J (1980) The influence of substrate, pH, diet and

temperature upon cadmium accumulation in the Asiatic clam (*Corbicula fluminea*)

in laboratory artificial streams. Nautilus (Philadelphia) 18:833–842

Heinsohn GE (1958) Life history and ecology of the freshwater clam, *Corbicula*

fluminea. University of California, Berkeley

Hill JK, Griffiths HM, Thomas CD (2011) Climate change and evolutionary adaptations

at species' range margins. Annu Rev Entomol 56:143–159.

https://doi.org/10.1146/annurev-ento-120709-144746

Hoegh-Guldberg O, Bruno JF (2010) The impact of climate change on the world's

marine ecosystems. Science (80-) 328:1523–1528.

https://doi.org/10.1126/science.1189930

Hoenig JM, Hanumara RC (1982) A statistical study of a seasonal growth model for

fishes. Univ Rhode Island Dep Comput Sci Tech Report, Narragansett

Hoffmann AA, Sgró CM (2011) Climate change and evolutionary adaptation. Nature

470:479–485. https://doi.org/10.1038/nature09670

Hornbach DJ (1992) Life history traits of a riverine population of the Asian clam

Corbicula fluminea. Am Modl Nat 127:248–257

Hünicken LA, Abrameto M, Bonel N (2019) *Corbicula* at its southernmost invasion

front in Patagonia: unusual low density and asymmetric trait responses to varying

environmental conditions. J Molluscan Stud 84:1–11.

https://doi.org/10.1016/0038-1098(79)91043-3

Ilarri MI, Antunes C, Guilhermino L, Sousa R (2011) Massive mortality of the Asian

clam *Corbicula fluminea* in a highly invaded area. Biol Invasions 13:277–280.

https://doi.org/10.1007/s10530-010-9833-5

IPCC (2014) Climate change 2014: impacts, adaptation, and vulnerability. Cambridge

University Press, United Kingdom and New York

Ituarte CF (1985) Growth dynamics in a natural population of *Corbicula fluminea*

(Bivalvia Sphaeriacea) at Punta Atalaya, Rio de La Plata, Argentina. Stud Neotrop

Fauna Environ 20:217–225

Kennedy VS., Van Huekelem L (1985) Gametogenesis and larval production in a

population of the introduced Asiatic clam, *Corbicula* sp. (Bivalvia: Corbiculidae),

in Maryland Published by : Marine Biological Laboratory Stable URL : Biol Bull

168:50–60

Komaru A, Kawagishi T, Konishi K (1998) Cytological evidence of spontaneous

androgenesis in the freshwater clam *Corbicula leana* Prime. Dev Genes Evol

208:46–50. https://doi.org/10.1007/s004270050152

Kraemer LR (1979) *Corbicula* (Bivalvia: Sphaeriacea) vs. indigenous mussels

(Bivalvia: Unionacea) in U.S. rivers: a hard case for interspecific competition? Am

Zool 19:1085–1096

Kraemer LR, Lott S (1977) Microscopic anatomy of the visceral mass of *Corbicula*

(Bivalvia: Sphaeriacea). Am Malacol Bull 48–56

Kraemer LR, Swanson C, Galloway M, Kraemer R (1986) Biological basis of behavior

in *Corbicula fluminea*, II. Functional morphology of reproduction and

development and review of evidence for self-fertilization. Am Malacol Bull Spetial

Ed:193–201

Lewontin RC (1965) Selection for colonizing ability. In: Baker H, Stebbins G (eds) The

genetics of colonizing species. Academic Press, New York, pp 79–94

Lockwood JL, Hoopes MF, Marchetti MP (2007) Invasion Ecology. Blackwell

Publishing, Malden, USA; Oxford, UK; Victoria, Australia

Lonhart SI, Jeppesen R, Beas-Luna R, Crooks JA, Lorda J (2019) Shifts in the

distribution and abundance of coastal marine species along the eastern Pacific

Ocean during marine heatwaves from 2013 to 2018. Mar Biodivers Rec 12:13.

https://doi.org/10.1186/s41200-019-0171-8

MacArthur RH, Wilson E 0. (1967) The theory of island biogeography. Princeton

University Press

Magnussen E (2007) Interpopulation comparison of growth patterns of 14 fish species

on Faroe Bank : are all fishes on the bank fast-growing? J Fish Bioilogy 71:453–

475. https://doi.org/10.1111/j.1095-8649.2007.01502.x

McDowell WG, Mcdowell WH, Byers JE (2016) Mass mortality of a dominant invasive

species in response to an extreme climate event: implications for ecosystem

function. Limnol Oceanogr. https://doi.org/10.1002/lno.10384

McDowell WG, McDowell WH, Byers JE (2017) Mass mortality of a dominant

invasive species in response to an extreme climate event: Implications for

ecosystem function. Limnol Oceanogr 62:177–188.

https://doi.org/10.1002/lno.10384

McDowell WG, Sousa R (2019) Mass mortality events of invasive freshwater bivalves:

current understanding and potential directions for future research. Front Ecol Evol

7:. https://doi.org/10.3389/fevo.2019.00331

Mcmahon RF (2002) Evolutionary and physiological adaptations of aquatic invasive

animals: r selection versus resistance. Can J Fish Aquat Sci 59:1235–1244.

https://doi.org/10.1139/F02-105

McMahon RF, Williams CJ (1986) Growth, life cycle, upper thermal limit and

downstream colonization rates in a natural population of the freshwater bivalve

mollusc, *Corbicula fluminea* (Müller) receiving thermal effluents. Am Malacol

Bull Special Ed:231–239.

Mouthon J (2001) Life cycle and population dynamics of the Asian clam *Corbicula*

fluminea (Bivalvia : Corbiculidae) in the Rhone River at Creys-Malville (France).

Arch für Hydrobiol 151:571–589

Mouthon J, Parghentanian T (2004) Comparison of the life cycle and population

dynamics of two *Corbicula* species, *C. fluminea* and *C. fluminalis* (Bivalvia:

Corbiculidae) in two French canals. Arch für Hydrobiol 161:267–287.

https://doi.org/10.1127/0003-9136/2004/0161-0267

Munro JL, Pauly D (1983) A simple method for comparing the growth of fishes and

invertebrates. Fishbyte 1:5–6

Otto SP (2007) The evolutionary consequences of polyploidy. Cell 131:452–462.

https://doi.org/10.1016/j.cell.2007.10.022

Pauly D, Caddy JF (1985) A modification of Bhattacharya's method for the analysis of

mixtures of normal distributions. Food Agric Organ United Nations 781:

Pauly D, Gaschutz G (1979) A simple method for fitting oscillating length growth data,

with a program for pocket calculators. ICES C 6:24

Pauly D, Munro JL (1984) Once more on the comparison of growth in fish and

invertebrates. Fishbyte 2:21

Peña EA, Slate EH (2006) Global validation of linear model assumptions. J Am Stat

Assoc 101:341. https://doi.org/10.1038/jid.2014.371

Phillips BL (2009) The evolution of growth rates on an expanding range edge. Biol Lett

5:802–804. https://doi.org/10.1098/rsbl.2009.0367

Phillips BL, Brown GP, Shine R (2010) Life-history evolution in range-shifting

populations. Ecology 91:1617–1627

Phillips BL, Brown GP, Webb JK, Shine R (2006) Invasion and the evolution of speed

in toads. Nature 439:803. https://doi.org/10.1038/439803a

Pigneur L-M, Etoundi E, Aldridge DC, Marescaux J, Yasuda N, Van Doninck K (2014)

Genetic uniformity and long-distance clonal dispersal in the invasive androgenetic

Corbicula clams. Mol Ecol 23:5102–5116. https://doi.org/10.1111/mec.12912

Pigneur L-M, Hedtke SM, Etoundi E, Doninck K Van (2012) Androgenesis : a review

through the study of the selfish shellfish *Corbicula* spp . Heredity (Edinb)

108:581–591. https://doi.org/10.1038/hdy.2012.3

Rajagopal S, Velde G, Vaate A (2000) Reproductive biology of the Asiatic clams

Corbicula fluminalis and *Corbicula fluminea*. Arch für Hydrobiol 149:403–420

Ramón M, Fernández M, Galimany E (2007) Development of mussel (*Mytilus*

galloprovincialis) seed from two different origins in a semi-enclosed

Mediterranean Bay (N.E. Spain). Aquaculture 264:148–159.

https://doi.org/10.1016/j.aquaculture.2006.11.014

Ricciardi A (2012) Invasive Species. In: Meyers RA (ed) Encyclopedia of Sustainability

Science and Technology. Springer Science+Business Media, New York, pp 5547–

5560

Roff DA (1993) Evolution of life histories: theory and analysis. Springer US

Rosa IC, Pereira JL, Costa R, Gomes J, de Lourdes Pereira M, Gonçalves F (2014)

Dispersal of *Corbicula fluminea* : Factors influencing the invasive clam's drifting

behavior. Ann Limnol - Int J Limnol 50:199–199.

https://doi.org/10.1051/limn/2014011

Sakai AK, Allendorf FW, Holt JS, Lodge M, Molofsky J, With KA, Baughman S,

Cabin RJ, Cohen JE, Ellstrand NC, McCauley DE, O'Neil P, Parker IM,

Thompson JN, Weller SG (2001) The population biology of invasive species.

734   Annu Rev Ecol Syst 32:305–332

735 Schmidlin S, Baur B (2007) Distribution and substrate preference of the invasive clam

736   *Corbicula fluminea* in the river Rhine in the region of Basel (Switzerland,

737   Germany, France). Aquat Sci 69:153–161. https://doi.org/10.1007/s00027-006-

738   0865-y

739 Schrieber K, Lachmuth S (2017) The genetic paradox of invasions revisited: the

740   potential role of inbreeding× environment interactions in invasion success. Biol

741   Rev 92:939–952. https://doi.org/10.1111/brv.12263

742 Selmecki A, Maruvka YE, Richmond PA, Guillet M, Shoresh N, De S, Kishony R,

743   Michor F, Dowell R, Pellman D (2015) Polyploidy can drive rapid adaptation in

744   yeast. Nature 519:349–352. https://doi.org/10.1038/nature14187

745 Shine R, Brown GP, Phillips BL (2011) An evolutionary process that assembles

746   phenotypes through space rather than through time. Proc Natl Acad Sci 108:5708–

747   5711. https://doi.org/10.1073/pnas.1018989108

748 Siemann E, Rogers WE (2001) Genetic differences in growth of an invasive tree

749   species. Ecol Lett 4:514–518. https://doi.org/10.1046/j.1461-0248.2001.00274.x

750 Skellam JG (1951) Random dispersal in theoretical populations. Biometrika 38:196–

751   218

752 Somers IF (1988) On A Seasonally Oscillating Growth Function. Fishbyte 6:8–11

753 Sousa R, Antunes C, Guilhermino L (2008a) Ecology of the invasive Asian clam

754   *Corbicula fluminea* (Müller, 1774) in aquatic ecosystems: an overview. Ann

755   Limnol - Int J Limnol 44:85–94. https://doi.org/10.1051/limn:2008017

756 Sousa R, Antunes C, Guilhermino L (2006) Factors influencing the occurrence and

757   distribution of *Corbicula fluminea* (Müller, 1774) in the River Lima estuary. Ann

758   Limnol - Int J Limnol 42:165–171. https://doi.org/10.1051/limn/2006017

Sousa R, Nogueira AJA, Gaspar MB, Antunes C, Guilhermino L (2008b) Growth and

extremely high production of the non-indigenous invasive species *Corbicula*

fluminea (Müller, 1774): possible implications for ecosystem functioning. Estuar

Coast Shelf Sci 80:289–295. https://doi.org/10.1016/j.ecss.2008.08.006

Stearns SC (1976) Life-History tactics: a review of the ideas. Q Rev Biol 51:3–47.

https://doi.org/10.1086/409052

Stearns SC (1977) The evolution of life history traits: a critique of the theory and a

review of the data. Annu Rev Ecol Syst 8:145–171.

https://doi.org/10.1146/annurev.es.08.110177.001045

Stites DL, Benke AC, Gillespie DM (1995) Population dynamics, growth, and

production of the Asiatic clam *Corbicula fluminea*, in a blackwater river. Can J

Fish Aquat Sci 52:425–437

Strayer DL, D'Antonio CM, Essl F, Fowler MS, Geist J, Hilt S, Jarić I, Jöhnk K, Jones

CG, Lambin X, Latzka AW, Pergl J, Pyšek P, Robertson P, von Schmalensee M,

Stefansson RA, Wright J, Jeschke JM (2017) Boom-bust dynamics in biological

invasions: towards an improved application of the concept. Ecol Lett 20:1337–

1350. https://doi.org/10.1111/ele.12822

Taylor CC (1958) Cod growth and temperature. J du Cons 23:366–370

Vakily JM (1992) Determination and comparison of bivalve growth, with emphasis on

Thailand and other tropical areas

Vaughn CC, Atkinson CL, Julian JP (2015) Drought-induced changes in flow regimes

lead to long-term losses in mussel-provided ecosystem services. Ecol Evol 5:1291–

1305. https://doi.org/10.1002/ece3.1442

Vohmann A, Borcherding J, Kureck A, bij de Vaate A, Arndt H, Weitere M (2010)

Strong body mass decrease of the invasive clam *Corbicula fluminea* during

summer. Biol Invasions 12:53–64. https://doi.org/10.1007/s10530-009-9429-0

Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin J-M,

Hoegh-Guldberg O, Bairlein F (2002) Ecological responses to recent climate

change. Nature 416:389–395. https://doi.org/10.1007/978-3-319-10157-6_4

TABLES

Table 1. Summary of ecological and growth variables (mean ± SD) of worldwide invasive populations of the Asian clam *Corbicula* reviewed from the literature. *Density* is the estimated abundance of clams at the study site (ind./m^2), *Pop.growth* is the rate of population increase, *Time* is the number of years elapsed since population colonization at the time of the original study (years), *Ind.growth* is the individual growth rate (*GPI*, a *proxy* for individual growth rate), *mASM* is the minimum age at sexual maturity (months), and lifespan (years), *ID#* is a number assigned arbitrarily for identification purposes. The country, specific study location, and reference to the source paper of the original study used to build the information are shown for each population. See the main text for details on how data was collected and variables were built from the literature. For extended information on each population, see Table S1 of the Online Resource.

Country	Study location	Time	Population parameters		Individual parameters			ID#	Reference
			Density	Pop.growth	Ind.growth	mASM	Lifespan		
Argentina	Punta Atalaya, Río de la Plata	3.5	339±405	2.20±0.01	2.87±0.07	4.30±0.13	3.83±0.57	1	Ituarte (1985)
	Paraná de las Palmas River	25.5	1,053±768	14.44±0.31	2.73±0.01	9.23±1.17	6.82±0.05	2	Cataldo and Boltovskoy (1999)
	Río Negro estuary	15.5	94±8.3	7.85±0.10	2.94±0.23	5.32±1.66	3.55±0.41	3	Hünicken et al. (2019)
France	Saone River	12.5	302±186	14.65±0.01	2.54±0.13	12.73±2.42	7.55±2.17	4	Mouthon (2001)
	Loire Lateral Canal	17.0	992±1209	41.17±0.12	2.80±0.24	8.93±5.95	5.17±3.00	5	Mouthon and Parghentanian (2004)
	Canal of Roanne	17.0	626±897	25.50±0.01	2.77±0.16	8.25±3.43	4.27±1.58	6	Mouthon and Parghentanian (2004)
Portugal	River Minho estuary	16.5	840±387	11.89±0.06	3.32±0.05	3.06±1.20	2.28±0.30	7	Sousa et al. (2008b)
	Mondego estuary	6.9	5,017±5,489	7.67±0.04	–	–	–	8	Franco *et al.*, 2012
	Casal de São Tomé	10.3	3,255±1,115	4.06±0.04	3.14±0.12	3.83±1.69	2.75±0.51	9	Rosa et al. (2014)
USA	Altamaha River	4.5	712±1,613	0.37±0.00	–	–	–	10	Gardner et al. (1976)
	Lake Arlington	2.0	32±11	11.94±0.03	3.49±0.05	5.80±0.60	2.59±0.20	11	Aldridge and McMahon (1978)
	Delta-Mendota canal	28.0	9,470±7067	4.53±0.03	2.77±0.17	9.69±4.07	8.03±2.75	12	Eng (1979)
	New River (Station 1)	2	316±654	9.40±0.02	–	–	–	13	Graney et al. (1980)
	Clear Fork, Trinity River	7.0	4,474±3,899	19.34±0.04	3.00±0.16	6.76±3.62	7.07±2.83	14	McMahon and Williams (1986)
	Mechums River	10.5	669±636	188.52±0.62	2.56±0.15	9.20±4.41	3.31±1.21	15	Hornbach (1992)
	Ogeechee River	10.0	1,398±1,096	6.23±0.00	2.52±0.12	22.72±9.17	6.85±1.63	16	Stites et al. (1995)
	Lake Tahoe	8.0	1,809±723	16.64±0.04	2.80±0.12	6.90±1.90	2.70±1.32	17	Denton et al. (2012)

800 **Table 2.** Summary of results from reduced models of weighted linear regression analyses of invasion and growth variables of global invasive
801 populations of *Corbicula*. Parameter values are means ± SE. We report one-tailed *P* values indicating significant results in bold characters. Note
802 natural logarithmic transformations of variables. Variable names as per Table 1. For details on descriptive statistics of full models and results on
803 partial *F*-tests see the Online Resource.

Response variable	Predictor	Estimate ± SE	Intercept ± SE	R^2	*F*-statistics	*num.df, den.df*	*one-tailed* P-value
Ln(*Density*)	Ln(*Time*)	0.454 ± 0.196	-0.906 ± 0.241	0.263	5.34	1, 15	**0.018**
Ln(*Pop.growth*)	Ln(*Time*)	0.936 ± 0.323	-0.518 ± 0.216	0.376	4.03	2, 14	**0.006**
	Ln(*Density*)	-0.522 ± 0.407					0.220†
Ln(*Ind.growth*)	Ln(*Density*)	-0.422 ± 0.265	0.098 ± 0.264	0.283	2.17	2, 11	0.070
	Ln(*Pop.growth*)	-0.473 ± 0.344					0.099
Ln(*mASM*)	Ln(*Ind.growth*)	-0.505± 0.179	-0.192 ± 0.171	0.601	8.28	2, 11	**0.008**
	Ln(*Pop.growth*)	0.460 ± 0.208					**0.024**
Ln(*Lifespan*)	Ln(*Ind.growth*)	-0.718± 0.184	-0.056 ± 0.192	0.559	15.20	1, 12	**0.001**

806 † two-tailed *P*-value

FIGURE LEGENDS

Figure 1. Global distribution of invasive *Corbicula* clams indicating the native range of the genus (dark grey), the distribution of of *Corbicula* inside (black/grey shading and black dots) and outside of the genus' native range (color dots), which are mostly represented by forms A/R (formerly designed as *Corbicula fluminea sensu lato*; see main text for detailed explanation). Numbered stars are invasive worldwide populations reviewed in the present study. Years indicate estimated date of first introduction. See Table 1 and main text for references and details on the invasive populations reviewed here. See the Online Resource for detailed criteria and data used to build the genus native and invasive distributions.

Figure 2. Relationships between processes and characteristics at different levels of organization of 17 global invasive populations of the Asian clam *Corbicula*. At the population level: a) the effect of time since colonization (*Time,* years) on population density *(Density*, ind./m^2*)* and b, c) the simultaneous effect of *Density* and *Time* on population increase (*Pop.growth*¸% rate of increase). At the interaction between population and individual level: d, e) the simultaneous effect of density and population increase on individual growth (*Ind.growth*, GPI), and f) the effect of population increase on the minimum age at sexual maturity (*mASM*, months). At the individual level: g) the effect of individual growth on the minimum age at sexual maturity, and h) on the *Life span* (years). When plotting the fit for one variable after testing for simultaneous effects, we removed the effect of the variable not considered (b, c, d, e). Dashed lines show fits when marginally significant $0.05 > \alpha < 0.10$. Solid lines show linear regression fits when statistically significant at $\alpha < 0.05$. The size of circles correspond to the inverse of the variance of mean value of each observation of the response variable, used to weigh effect sizes in the regression analyses (see main text for details). Note natural logarithmic transformation of variables

Figura 3. Summary of the relationships between processes and characteristics at different levels of organization of 17 global invasive populations of the Asian clam *Corbicula*, showing positive (+), negative (–), and no relationships (0) after fitting weighted linear regressions.Thick solid lines (➡) show linear regression fits when statistically significant at $\alpha < 0.01$. Thin solid lines (→) show linear regression fits when statistically significant at $\alpha < 0.05$. Dashed lines (– ➤) show fits when marginally significant $0.05 > \alpha < 0.10$. Dotted line (······➤) show no significant relationship $\alpha > 0.10$. Relationships are shown at three different levels: A) at the population level, B) the interaction between processes and characteristics at the population and individual level, and C) at the individual level.

.

FIGURES

Figure 1

Figure 2

855 **Figure 3**