The gut microbiota influences how circulating immune cells in humans change from one day to the next.

Jonas Schluter[1], Jonathan U. Peled[2,3], Bradford P. Taylor[1], Melody Smith[2,3], Kate A. Markey[2,3], Ying Taur[4], Rene Niehus[5], Anna Staffas[6], Anqi Dai[2], Emily Fontana[4], Luigi A. Amoretti[4], Roberta J. Wright[4], Sejal Morjaria[4], Maly Fenelus[7], Melissa S. Pessin[7], Nelson J. Chao[8], Meagan Lew[8], Lauren Bohannon[8], Amy Bush[8], Anthony D. Sung[8], Tobias M. Hohl[4], Miguel-Angel Perales[2,3], Marcel R.M. van den Brink[2,3], Joao B. Xavier[1]

1) Computational and Systems Biology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA

2) Adult Bone Marrow Transplantation Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

3) Weill Cornell Medical College, New York, NY, USA

4) Infectious Disease Service, Department of Medicine, and Immunology Program, Sloan Kettering Institute, New York, NY, USA.

5) Harvard University, T.H. Chan School of Public Health, 677 Huntington Avenue, Boston, MA 02115

6) Sahlgrenska Cancer Center, Department of Microbiology and Immunology, Institute of Biomedicine, University of Gothenburg, SE-405 30, Gothenburg, Sweden

7) Department of Laboratory Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

8) Division of Hematologic Malignancies and Cellular Therapy, Duke University School of Medicine

Correspondence: jonas.schluter@gmail.com, xavierj@mskcc.org

ABSTRACT

The gut microbiota influences the development and homeostasis of the mammalian immune system[1–3], can alter immune cell compositions in mice[4–7], and is associated with responses to immunotherapy that rely on the activity of peripheral immune cells[8–12]. Still, our understanding of how the microbiota modulates circulatory immune cells remains limited, particularly in humans where a lack of manipulative experiments makes inference challenging. Here we overcome this challenge by studying hundreds of hospitalized—and closely monitored—bone marrow transplantation patients as they recover from chemotherapy-induced immune ablation. This aggressive treatment causes large shifts in both circulatory immune cell and microbiota populations, allowing the relationships between the two to be studied simultaneously over time with unprecedented resolution. We analyzed daily changes in white blood cell counts from 2,235 patients, and 10,680 longitudinal fecal microbiota samples to identify bacterial genera consistently associated with those changes. Bayesian inference and validation across different patient cohorts revealed consistent associations between intestinal bacteria and peripheral immune cell dynamics in the context of immunomodulatory medications, clinical metadata and homeostatic feedbacks between peripheral immune cells. The quantification of validated microbiota associations enabled us to contrast the potency of fermentatively active, obligate anaerobic bacteria with that of medications with known immunomodulatory mechanism, and this way estimate the microbiota potential to alter peripheral immune cell dynamics directly in patients. Our analysis establishes and quantifies the link between the intestinal microbiota and immune cell dynamics in humans, with implications for microbiota-driven modulation of immunity and immunotherapies that rely on circulatory immune cells.

MAIN TEXT

Experiments in mice provide mounting evidence that the mammalian intestinal microbiome influences the development[1–3] and homeostasis of its host's immune system[4–7,13–15]. In humans, inflammatory bowel diseases correlate with functional dysbiosis in the gut microbiota[16,17]. Furthermore, children born preterm and at term have different gut microbiome compositions and differ in the development of immune cell populations in their blood[18]. The composition of the gut microbiota may also influence the success of immunotherapies[8–11]. Immune checkpoint inhibitor therapy relies on activation of circulating T-cells and its success has, independently, been associated with abundances of intestinal anaerobic genera such as *Akkermansia*[9] and *Faecalibacterium*[10]. It is therefore an intriguing prospect to augment treatments such as cancer immunotherapy[19], including the burgeoning field of chimeric antigen receptor (CAR) T-cell therapy[20], by leveraging microbiome-driven immune system modulation. However, our understanding of how the microbiota influences the dynamic changes in immune cell populations in humans, and how this compares to deliberate immunomodulatory interventions remains limited.

To overcome this limitation, we here quantified the associations between the microbiome and day-by-day changes in peripheral immune cell counts. We investigated the dynamics of immune reconstitution after allogeneic hematopoietic cell therapy (HCT) within the universe of all 2,926 patients who underwent HCT at Memorial Sloan Kettering for various hematological malignancies, including leukemia, between 2003 and 2019 (**Figure 1A, Table S1**). The conditioning regimen of radiation and chemotherapy administered to HCT patients is the most severe perturbation to the immune system deliberately performed in humans and thus offers a unique opportunity to investigate dynamic links between gut microbiota and immune system directly in humans. Conditioning depletes circulating white blood cell counts (**Figure 1A**). This can lead to prolonged periods of neutropenia (<500 neutrophils per µl blood) followed by reconstitution after transplanted stem cells have matured sufficiently to release granulocytes from the bone marrow (neutrophil engraftment is defined as 3 consecutive days with >500 neutrophils per µl blood, **Figure 1A-C**). The blood of each patient is carefully monitored throughout this recovery, and medications are administered to modulate the

70   immune cell dynamics, including granulocyte-colony stimulating factor (G-CSF) to increase

71   neutrophil counts, and immunosuppressants such as mycophenolate mofetil or tacrolimus to prevent

Figure 1: Immune reconstitution and microbiome dynamics after allogeneic hematopoietic cell transplantation (HCT). A) Three major periods of HCT—immunoablation during chemotherapeutic conditioning before HCT, defined as day 0, post-HCT neutropenia, and reconstitution following neutrophil engraftment—lead to recovery trajectories with large variability between patients. Shown are the average counts (shaded: ±1 standard deviation, σ) of neutrophils, lymphocytes and monocytes per day relative to HCT from patients transplanted between 2003 and 2019 (A), contrasted with two individual patients (B,C) representative of the recovery trajectories for different stem cell graft source; patient 1 who received a peripheral blood stem cell graft, PBSC (shaded region: mean±1 standard deviation of all PBSC patients), and patient 2 who received a graft of umbilical cord blood (shaded region: mean±1 standard deviation of all cord patients). Fecal samples collected during HCT reveal the loss of microbial diversity reported previously[22,23] (D-F) and commensal families (G-I), often replaced by Enterococcaceae domination. J,K). Administration of immunomodulatory medications for the two example patients.

complications such as graft-vs-host disease (**Figure 1 J,K**). To investigate if the composition of the

gut microbiota affects the dynamics of circulating white blood cells in our patients, we analyzed

detailed blood and clinical metadata of our patients between 3 days before HCT and until 100 days

post neutrophil engraftment (excluding pediatric patients, and other exclusion criteria: N=2,235,

supplementary methods). During this period patients are monitored carefully, and our analysis

included over 140,000 host phenotype measurements in the form of complete blood counts which

quantify the most abundant white blood cells—neutrophils, lymphocytes, monocytes, eosinophils—as

well as platelet counts (**Figure 1, S1**). We started collecting patients' fecal microbiota data in 2009[21],

from which we obtained 10,680 high frequency, longitudinal microbiota compositions from 1,294 of

our HCT patients.

We have shown previously that HCT patients lose gut microbiota biodiversity and commensal

microbial families during their treatment (**Figure 1D-I**)[22,23], and, furthermore, recently associated

impaired white blood cell recovery after bone-marrow transplantation in mice with a depleted

intestinal flora[24] In our patients, microbial diversity usually recovers slowly during white blood cell

reconstitution (**Figure 1D**); however, microbiota recovery as well as immune reconstitution can vary

strongly between patients and treatment types (**Figure 1B,C,J,K**). This variation is illustrated by the

distinct trajectories of patient 1 who received a graft of peripheral blood stem cells (PBSC), retained

high microbiota diversity and engrafted earlier (**Figure 1B,E,H**), and patient 2 who received a graft of

umbilical cord blood (cord), lost microbiota diversity and engrafted later (**Figure 1C,F,I**). Low

microbiota diversity at the time of neutrophil engraftment has been associated with 5-fold increased

transplant-related mortality[25], suggesting that that the joint recovery of the microbiota and white blood

cells in circulation is critical for clinical outcomes.

To detect a causal link between the microbiota and white blood cell recovery dynamics, we

first used data from a recent prospective randomized trial of autologous fecal microbiota

transplantation (auto-FMT), which represents a deliberate microbiota manipulation in our patients[22]

(supplementary methods). Twenty-four patients (**Figure 2A, Table S2)** underwent randomization,

resulting in 10 untreated control and 14 treated patients, including patient 3 in **Figure S2**. To

investigate if auto-FMT affected white blood cell reconstitution, we compared the 24 patients'

neutrophil, lymphocyte and monocyte counts post-engraftment (i.e. when the transplanted

hematopoietic cells begin producing new white blood cells, **Figure 2A**). FMT procedures were

conducted at variable time points relative to neutrophil engraftment, but overall, we observed higher

counts of each white blood cell type in patients who received an auto-FMT during the first 100 days

post neutrophil engraftment ($p < 0.01$, **Figure 2B,S3-5**).

Figure 2: FMT-treated patients showed higher counts of white blood cells. A) Absolute counts of neutrophils (blue), lymphocytes (green) and monocytes (orange) in 24 patients enrolled in a randomized controlled trial to receive an autologous fecal microbiota transplant post-neutrophil engraftment (FMT treatment: red vertical line, control: black vertical line). B) Results from a linear mixed effects model with random effects per patient and per day relative to neutrophil engraftment show elevated neutrophil, lymphocyte and monocyte counts in patients receiving auto-FMT after their treatment as compared to control patients, bars and confidence intervals for the averages of observed white blood cell counts without FMT and post-FMT (***: $p < 0.001$, **: $p < 0.01$).

The increased white blood cell counts in patients receiving auto-FMT indicated that this deliberate microbiota manipulation affected circulatory immune cell dynamics, supporting a causal role of the microbiota. The increase could be due to the reconstitution of a complex microbiota that

we saw in these patients[22] and the associated metabolic capabilities[6,7,24], or it could be a systemic

response to a severe therapy which introduced billions of intestinal organisms at once via an enema

(no enema was administered to control patients[22]). Moreover, while our analysis used a mixed-effects

model which accounted for patient-specific HCT treatments in this randomized patient cohort, chance

differences in extrinsic factors such as different immunomodulatory drug exposures may have

affected this result due to the small cohort size. Therefore, we next investigated the link between the

gut microbiota composition and the dynamics of white blood cell recovery in our large observational

cohort of HCT patients.

126   Homeostasis of circulatory white blood cell counts is a complex, dynamic process:

neutrophils, lymphocytes and monocytes are formed and released into the blood *de novo* by

differentiation of hematopoietic progenitor cells from the bone marrow, and they can be mobilized

from thymus and lymph nodes (lymphocytes), spleen, liver and lungs (neutrophils); they can also

migrate from the blood to other tissues when needed[26]. These processes are dynamic sources and

sinks of circulatory white blood cells, and they can be modulated by drugs administered to patients

receiving HCT. To identify factors associated with these dynamic source- and sink-processes—

including the microbiota—we developed a two-stage approach analyzing the changes of white blood

cell counts between two days (i.e. the rates of cell count increases and decreases). Stage 1 served as a

feature selection stage where we used data of 1,096 patients (after filtering for qualifying samples and

applying exclusion criteria, see supplementary methods) *without* available microbiome information to

identify associations between clinical metadata, including immunomodulatory medications (methods),

and changes of white blood cell counts from one day to the next (**Figure 3A**). Stage 2 was performed

on data from an independent cohort of 841 different patients at MSK from whom concurrent

microbiome samples were available. Stage 2, our main analysis, sought to reveal associations between

the abundance of microbial taxa and the daily changes in blood cell counts in context of

immunomodulatory medications, additional clinical metadata and the current state of the blood itself.

143   In stage 1 we calculated the changes in neutrophil, lymphocyte and monocyte counts during

patients' recovery from >20,000 pairs of post-engraftment blood samples separated by a single day

(**Figure 3B, S6-8,** supplementary methods). Using a cross-validated feature selection approach, we

detected medications and HCT treatment parameters that were associated with different rates of

change in neutrophil, lymphocyte and monocyte counts, including, as expected, G-CSF and the graft

stem cell sources (**Figure S6-S8, Table S1**).

During stage 2 we sought to identify associations between bacterial taxa of the gut microbiota

and the dynamics of immune cells in circulation. For this, we performed Bayesian inferences using

data from different sets of patients *with* available microbiome samples. Stage 1 had identified—as

expected—that stem cell graft sources are associated with immune reconstitution kinetics (e.g. cord

on average slower compared with PBSC[27]), and we therefore stratified our patients by graft source in

stage 2. The model of stage 2 now included microbial genera as additional predictors of observed

changes in white blood cell counts to the medications selected in stage 1, clinical features

(conditioning intensity, age, sex), and the current state of the blood in the form of counts of

neutrophils, lymphocytes, monocytes, eosinophils, and platelets. The data comprised 841 patients, but

approximately 60% of the stool samples paired with a daily change in white blood cell counts were

taken before neutrophil engraftment (**Figure 3B, Table S1,** supplementary methods), i.e. when blood

cells counts were zero. In total, the analyzed 2,615 post-engraftment observations of changes in

neutrophil counts during immune reconstitution (lymphocytes: 2,006, monocytes: 2,534) with paired

stool samples which provided a large sample of observed white blood cell dynamics (**Figure 3B,**

Table S3,S4). We first focused on the data from the largest (**Figure 3C**) cohort—patients who

received a PBSC graft—and withheld the other cohorts (bone marrow, BM; T-cell depleted graft

(*ex-vivo*) by CD34+selection, TCD; cord blood, cord) to use as independent validation cohorts. For

this validation, we analyzed TCD, BM, and cord patients' data in the same way as PBSC patients'

data and compared the resulting posterior coefficient distributions (methods). We assigned

coefficients obtained from the PBSC cohort a validation score (v-score) between 0 and 3, representing

the number of times that the focal coefficient was validated in the other cohorts; but, conservatively,

the score was always set to zero if we observed counter-evidence among any of the other data sets, i.e.

evidence that coefficients had the opposite sign. Finally, we analyzed data from another patient cohort

consisting of 493 bone marrow transplantation recipients treated at Duke University including 9,603

blood samples and a total of 629 microbiota samples from 218 patients, albeit with lower sampling

density, and we used the results from this analysis for further validation.

Notably, as a verification of our approach, we detected associations between the

administration of immunomodulators and increased or decreased rates of immune cell count changes

consistent with the known biological mechanism of these medications (**Figure 3C,S9A**). The

strongest association we found across all predictors is consistent with the well-known neutrophil-

increasing effect of G-CSF[28]; G-CSF administration—used to accelerate recovery from

chemotherapy-induced neutropenia[28]—was associated with a 140% increase in the rate of neutrophil

changes from one day to the next ([114%, 170%], 95 percent probability density interval [HPDI95]).

This finding was observed in all MSK validation data sets (v-score=3, **Figure 3D)**, as well as among

Duke University patients (**Figure S10**). We furthermore found a G-CSF-associated increase of 43%

([30%, 58%]HPDI95, v-score=3) in monocyte rates, and, although smaller, in lymphocyte rates (16%,

[5%, 27%]HPDI95, v-score=2). Both neutrophil and lymphocyte rates decreased following the

exposure to immunosuppressive medications (cetirizine -18%, [-35%,+5%]HPDI95, mycophenolate

mofetil -8% [15%,+1%]HPDI95, respectively). Finally, less intensive chemotherapeutic conditioning

regimens (reduced intensity and non-ablative conditioning) were associated with higher average

lymphocyte count increases during immune reconstitution (**Figure S9C)**

Beyond associating medications in agreement with their known biological mechanism, our

analysis detected associations between the current count of white blood cells and their rate of change:

a negative association among lymphocytes, negative associations between counts of neutrophils and

lymphocytes with the rates of monocytes, and a negative association between the counts of platelets

and the rates of neutrophils (**Figure 3E**). Conversely, we found positive associations between

monocytes and the rates of each of the investigated white blood cell subsets. These associations

suggest a feedback network (**Figure 3F**) of circulatory white blood cell counts, reflecting potentially

complex homeostatic regulation mechanisms of immune cell compositions in the blood[26]. More

importantly, the associations quantified for medications and potential homeostatic feedbacks provided

a benchmark against which we could compare associations from gut microbial taxa.

Figure 3: Bayesian inference conducted in context of the host state (white blood cell and platelet counts)
and clinical variables including immunomodulatory medications reveals the microbiota potential to affect
daily changes in peripheral white blood cell counts. A) Cartoon of the model: observed changes in white
blood cell counts between two consecutive days are associated with the current state of the host in the form of
blood cell counts in circulation, the administration of immunomodulatory medications, patient and clinical
metadata, and the state of the microbiome. B) Visualization of the dynamic white blood cell data; scatter plot of
the principal components (PC) of observed daily changes of neutrophils, lymphocytes and monocytes without
(grey, see **Figures S6-8**) and with (orange) concurrent longitudinal microbiome data (bold: post engraftment
sample counts). C) PBSC patients (N=312) provided the most blood samples with simultaneous microbiome
data (n=995) relative to TCD, BM and cord patients, who were used as validation data sets. D-G) PBSC patient
data inference results; bars show the posterior parameter estimate distributions (thin: 95% highest posterior
density intervals, HPDI95, thick: HPDI50) for the jointly inferred associations between treatments (D), white
blood cells counts (E, revealing a potential homeostatic regulation network, F), and log-fecal microbiota genus
abundances (G) with the observed daily changes in neutrophils, lymphocytes and monocytes. v-score: number
of validation cohort confirming associations, *always* set to zero if invalidated in any of the TCD, BM, or cord
cohorts.

We identified microbial genera that consistently associated with increases or decreases in

white blood cell counts by first using data from the PBSC patients and then validating the associations

in the other cohorts (**Figure 3G**). Higher abundances of *Faecalibacterium* (+8%, [+1%,

+14%]HPDI95 per $\log_{10}$), *Ruminococcus 2* (+5%, [-0%, +10%]HPDI95) and *Akkermansia* (+4%,

[1%, 7%]HPDI95) were associated with greater neutrophil increases, whereas increased *Rothia* (-3%,

[-7%, +0%]HPDI95), and *Clostridium sensu stricto 1* (-3%, [-6%, 0%]HPDI95) abundances

associated with reduced neutrophil rates. These results were validated in univariate analyses

conducted in the Duke University cohort (**Figure S10**). *Staphylococcus* was positively associated with

lymphocyte rates (+4%, [+1%, +6%]HPDI95) and, again, *Ruminococcus 2* was also associated with

faster lymphocyte increases (+5%, [+1%, +9%]HPDI95). Both *Faecalibacterium* as well as

Ruminococcus 2 also associated with increases in monocytes, and while this association was validated

in other cohorts (v-score 3 and 1, respectively), there was higher uncertainty of the association

estimate (HPDI50>0). Again, *Clostridium sensu strictu 1* (-3% [-5%,-1%]HPDI95) associated

consistently with decreased rates of monocytes. The associations we identified—and validated in

other cohorts—between microbial taxa in the gut and daily changes in white blood cell counts support

the idea that hematopoiesis and mobilization respond to the composition of the gut microbiome,

influencing systemic immunity[29].

Most of the taxa that strongly associated with white blood cell dynamics were obligate

anaerobes. *Rothia*, was a notable exception: this aerobic genus is typically found in the oral cavity[30]

but can become an opportunistic pathogen in immunosuppressed patients and is not known to provide

metabolic functions to the host[31]. Obligate anaerobes, on the other hand, produce metabolic

byproducts such as short-chain fatty acids[32,33] and bacterial cell-wall molecules[1,34,35] that modulate

immune responses and granulopoiesis[6]. Nutritional support from the intestinal microbiota improved

hematopoietic reconstitution in a mouse model[24]. To identify a similar association in our patients, we

estimated a microbiota potency by multiplying the $\log_{10}$-abundances of microbial genera in a sample

with their corresponding posterior coefficients. We analyzed shotgun metagenomics sequences from

124 of the samples and observed that samples with positive microbiota potency were associated with

enrichment in cholate degradation and vitamin-B1 synthesis related pathways, as well as butanoate

formation (**Figure S11**). Our findings are in line with evolutionary theory[36] that essential but broadly

available microbial traits such as the production of B vitamins, secondary bile acid metabolism, and

fermentation to short-chain fatty acids[37] could be co-opted by the host's immune system as part of the

homeostatic interplay between immune system and a complex microbiota[38,39]. For example,

Ruminococcus 2 is a genus that contains *R. bromii*, a keystone species necessary for the efficient

release of energy from complex starch in the normal diet[40]. Reassuringly, the genera

Faecalibacterium, *Ruminococcus 2* and *Akkermansia* that we associated with faster rates of white

blood cells were among those best reconstituted by auto-FMT[22], potentially explaining why we found

higher counts of neutrophils, monocytes and lymphocytes in patients who received the auto-FMT

(**Figure 2B**).

The associations we reveal are interpretable as potential effectors on sources and sinks of

white blood cell counts in circulation. Intestinal bacteria may affect white blood cell counts in

circulation by influencing either their sources in the bone marrow or their cytokine profiles[41] and

proliferation rates in the blood, their sinks in different organs, or both. Our analysis cannot distinguish

the effects of microbial communities on *de novo* hematopoiesis, for example, in isolation from their

impact on other sources and sinks. Unlike the plausible role of obligate anaerobe fermenters in

augmenting hematopoiesis via nutritional support[24], the positive association detected between

Staphylococcus and lymphocyte dynamics could instead result from reduced migration of T cells out

of circulation into the gut epithelium[42], especially since high abundances of *Staphylococcus* are

associated with low gut microbiota diversity (p<0.001, **Figure S12**), which indicates a depleted

microbiota.

Nevertheless, our approach allows us to leverage the chronology of events and assess

"mathematical causality"[43]. Of course, due to the observational nature of these data there are risks of

confounding that could explain some of the associations found, but the close temporal

correspondence[43] between microbiota and blood cell dynamics, and the validation across cohorts

reduces the number of plausible confounders. Our results, therefore, quite naturally suggest candidate

microbial taxa to manipulate if we seek to steer complex hematopoietic dynamics and utilize the

microbiota as an immunomodulatory component of the human body. *Faecalibacterium*,

Ruminococcus 2, and *Akkermansia* most strongly associated with increases in white blood cell counts

from one day to the next. Intriguingly, members of *Faecalibacterium* and *Ruminococcus* in one

study[10], and *Akkermansia* in another[9], were identified as enriched in patients with better responses to

anti–PD-1 immunotherapy, which suggests a disagreement between the two studies[44]. Our results

consolidate these findings, and, furthermore, allow us to quantitatively relate the potency of

manipulating these intestinal commensals to that of immunomodulatory drugs. While these genera are

common in the gut microbiota of healthy people[17], the abundance of each genus can drop below

detection in our patients during the intestinal damage related to HCT[23]. Therefore, realistic ranges of

3-5 orders of magnitude in bacterial abundances (**Figure S13**) can yield effect sizes similar to that of

homeostatic feedbacks between white blood cells and several immunomodulatory medications (e.g. a

change in *Ruminococcus 2* from below detection to 1% relative abundance associated with a +67%

and +63% increases in neutrophil and lymphocyte rates, respectively). The relatively small effect

sizes of intestinal microbes, in concert and over time, could therefore have significant impact on

systemic immunity even in individuals with less severely injured microbiomes.

In sum, our work links the gut microbiota to the dynamics of the human immune system via

peripheral white blood cells populations. We were able to infer and quantify this link because we

aggregated an unprecedented resource: vast high-frequency longitudinal data of blood cell counts in

conjunction with microbiota samples and detailed clinical metadata. The links between microbiota

components and immune cell dynamics could be used in the future to improve immunotherapy and

help identify microbiota treatments to cure inflammatory diseases[9,10,45–49].

296
Bibliography

1.   Mazmanian, S. K., Liu, C. H., Tzianabos, A. O. & Kasper, D. L. An immunomodulatory
molecule of symbiotic bacteria directs maturation of the host immune system. *Cell* **122,**
107–118 (2005).
2.   Gomez de Agüero, M. *et al.* The maternal microbiota drives early postnatal innate
immune development. *Science* **351,** 1296–1302 (2016).
3.   Sonnenberg, G. F. & Artis, D. Novel connections and precision approaches. *Nat Rev*
Immunol **19,** 75–76 (2019).
4.   Geva-Zatorsky, N. *et al.* Mining the human gut microbiota for immunomodulatory
organisms. *Cell* **168,** 928–943.e11 (2017).
5.   Tan, T. G. *et al.* Identifying species of symbiont bacteria from the human gut that, alone,
can induce intestinal Th17 cells in mice. *Proc Natl Acad Sci U S A* **113,** E8141–E8150
(2016).
6.   Balmer, M. L. *et al.* Microbiota-derived compounds drive steady-state granulopoiesis via
MyD88/TICAM signaling. *J Immunol* **193,** 5273–5283 (2014).
7.   Deshmukh, H. S. *et al.* The microbiota regulates neutrophil homeostasis and host
resistance to Escherichia coli K1 sepsis in neonatal mice. *Nat Med* **20,** 524–530 (2014).
8.   Tanoue, T. *et al.* A defined commensal consortium elicits CD8 T cells and anti-cancer
immunity. *Nature* **565,** 600–605 (2019).
9.   Routy, B. *et al.* Gut microbiome influences efficacy of PD-1-based immunotherapy
against epithelial tumors. *Science* **359,** 91–97 (2018).
10. Gopalakrishnan, V. *et al.* Gut microbiome modulates response to anti-PD-1
immunotherapy in melanoma patients. *Science* **359,** 97–103 (2018).
11. Vétizou, M. *et al.* Anticancer immunotherapy by CTLA-4 blockade relies on the gut
microbiota. *Science* **350,** 1079–1084 (2015).
12. Matson, V. *et al.* The commensal microbiome is associated with anti-PD-1 efficacy in
metastatic melanoma patients. *Science* **359,** 104–108 (2018).
13. Atarashi, K. *et al.* ATP drives lamina propria T(H)17 cell differentiation. *Nature* **455,**
808–812 (2008).
14. Ivanov, I. I. *et al.* Specific microbiota direct the differentiation of IL-17-producing T-
helper cells in the mucosa of the small intestine. *Cell Host Microbe* **4,** 337–349 (2008).
15. Cahenzli, J., Köller, Y., Wyss, M., Geuking, M. B. & McCoy, K. D. Intestinal microbial
diversity during early-life colonization shapes long-term IgE levels. *Cell Host Microbe*
14, 559–570 (2013).
16. Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory bowel
diseases. *Nature* **569,** 655–662 (2019).
17. Integrative HMP (iHMP) Research Network Consortium. The integrative human
microbiome project. *Nature* **569,** 641–648 (2019).
18. Olin, A. *et al.* Stereotypic immune system development in newborn children. *Cell* **174,**
1277–1292.e14 (2018).
19. Brandi, G. & Frega, G. Microbiota: Overview and Implication in Immunotherapy-Based
Cancer Treatments. *Int J Mol Sci* **20,** (2019).
20. Xin Yu, J., Hubbard-Lucey, V. M. & Tang, J. The global pipeline of cell therapies for
cancer. *Nat Rev Drug Discov* **18,** 821–822 (2019).
21. Taur, Y. *et al.* Intestinal domination and the risk of bacteremia in patients undergoing

allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis* **55,** 905–914 (2012).

22. Taur, Y. *et al.* Reconstitution of the gut microbiota of antibiotic-treated patients by
autologous fecal microbiota transplant. *Sci Transl Med* **10,** (2018).

23. Morjaria, S. *et al.* Antibiotic-Induced Shifts in Fecal Microbiota Density and
Composition during Hematopoietic Stem Cell Transplantation. *Infect Immun* **87,** (2019).

24. Staffas, A. *et al.* Nutritional Support from the Intestinal Microbiota Improves
Hematopoietic Reconstitution after Bone Marrow Transplantation in Mice. *Cell Host*
Microbe **23,** 447–457.e4 (2018).

25. Taur, Y. *et al.* The effects of intestinal tract bacterial diversity on mortality following
allogeneic hematopoietic stem cell transplantation. *Blood* **124,** 1174–1182 (2014).

26. Scheiermann, C., Frenette, P. S. & Hidalgo, A. Regulation of leucocyte homeostasis in
the circulation. *Cardiovasc Res* **107,** 340–351 (2015).

27. Thompson, P. A. *et al.* Umbilical cord blood graft engineering: challenges and
opportunities. *Bone Marrow Transplant* **50 Suppl 2,** S55-62 (2015).

28. Gabrilove, J. L. *et al.* Effect of granulocyte colony-stimulating factor on neutropenia and
associated morbidity due to chemotherapy for transitional-cell carcinoma of the
urothelium. *N Engl J Med* **318,** 1414–1422 (1988).

29. Belkaid, Y. & Hand, T. W. Role of the microbiota in immunity and inflammation. *Cell*
157, 121–141 (2014).

30. Hall, M. W. *et al.* Inter-personal diversity and temporal dynamics of dental, tongue, and
salivary microbiota in the healthy oral cavity. *npj Biofilms and Microbiomes* **3,** 2 (2017).

31. Ramanan, P., Barreto, J. N., Osmon, D. R. & Tosh, P. K. Rothia bacteremia: a 10-year
experience at Mayo Clinic, Rochester, Minnesota. *J Clin Microbiol* **52,** 3184–3189
(2014).

32. Ulven, T. Short-chain free fatty acid receptors FFA2/GPR43 and FFA3/GPR41 as new
potential therapeutic targets. *Front Endocrinol (Lausanne)* **3,** 111 (2012).

33. Smith, P. M. *et al.* The microbial metabolites, short-chain fatty acids, regulate colonic
Treg cell homeostasis. *Science* **341,** 569–573 (2013).

34. Hergott, C. B. *et al.* Peptidoglycan from the gut microbiota governs the lifespan of
circulating phagocytes at homeostasis. *Blood* **127,** 2460–2471 (2016).

35. Ladinsky, M. S. *et al.* Endocytosis of commensal antigens by intestinal epithelial cells
regulates mucosal T cell homeostasis. *Science* **363,** (2019).

36. Foster, K. R., Schluter, J., Coyte, K. Z. & Rakoff-Nahoum, S. The evolution of the host
microbiome as an ecosystem on a leash. *Nature* **548,** 43–51 (2017).

37. Yoshii, K., Hosomi, K., Sawane, K. & Kunisawa, J. Metabolism of dietary and microbial
vitamin B family in the regulation of host immunity. *Front. Nutr.* **6,** 48 (2019).

38. Zaza, G. *et al.* Impact of maintenance immunosuppressive therapy on the fecal
microbiome of renal transplant recipients: Comparison between an everolimus- and a
standard tacrolimus-based regimen. *PLoS ONE* **12,** e0178228 (2017).

39. McLoughlin, K., Schluter, J., Rakoff-Nahoum, S., Smith, A. L. & Foster, K. R. Host
selection of microbiota via differential adhesion. *Cell Host Microbe* **19,** 550–559 (2016).

40. Ze, X., Duncan, S. H., Louis, P. & Flint, H. J. Ruminococcus bromii is a keystone
species for the degradation of resistant starch in the human colon. *ISME J* **6,** 1535–1543
(2012).

41. Schirmer, M. *et al.* Linking the human gut microbiome to inflammatory cytokine
production capacity. *Cell* **167,** 1125–1136.e8 (2016).

42. Fu, Y.-Y. *et al.* T Cell Recruitment to the Intestinal Stem Cell Compartment Drives
Immune-Mediated Intestinal Damage after Allogeneic Transplantation. *Immunity* **51,**
90–103.e3 (2019).

43. Gerber, G. K. The dynamic microbiome. *FEBS Lett* **588,** 4131–4139 (2014).

44. Jobin, C. Precision medicine using microbiota. *Science* **359,** 32–34 (2018).
45. Zitvogel, L., Ma, Y., Raoult, D., Kroemer, G. & Gajewski, T. F. The microbiome in
cancer immunotherapy: Diagnostic tools and therapeutic strategies. *Science* **359,** 1366–
1370 (2018).
46. Hsiao, E. Y. *et al.* Microbiota modulate behavioral and physiological abnormalities
associated with neurodevelopmental disorders. *Cell* **155,** 1451–1463 (2013).
47. Lakritz, J. R. *et al.* Gut bacteria require neutrophils to promote mammary tumorigenesis.
Oncotarget **6,** 9387–9396 (2015).
48. Lakritz, J. R. *et al.* Beneficial bacteria stimulate host immune cells to counteract dietary
and genetic predisposition to mammary cancer in mice. *Int J Cancer* **135,** 529–540
(2014).
49. Chen, H. *et al.* A forward chemical genetic screen reveals gut microbiota metabolites
that modulate host physiology. *Cell* **177,** 1217–1231.e18 (2019).
50. Caporaso, J. G. *et al.* Ultra-high-throughput microbial community analysis on the
Illumina HiSeq and MiSeq platforms. *ISME J* **6,** 1621–1624 (2012).
51. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon
data. *Nat Methods* **13,** 581–583 (2016).
52. Murali, A., Bhargava, A. & Wright, E. S. IDTAXA: a novel approach for accurate
taxonomic classification of microbiome sequences. *Microbiome* **6,** 140 (2018).
53. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res* **41,** D590-6 (2013).
54. Pinheiro, J. C., Bates, D. M., DebRoy, S. S. & Sarkar, D. Nlme: Linear and Nonlinear
Mixed Effects Models. (2013).
55. Tibshirani, R. Regression shrinkage and selection via the lasso. *Journal of the Royal
Statistical Society: Series B (Methodological)* **58,** 267–288 (1996).
56. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine
Learning Research* (2011).
57. Salvatier, J., Wiecki, T. V. & Fonnesbeck, C. Probabilistic programming in Python using
PyMC3. *PeerJ Computer Science* **2,** e55 (2016).
58. Hoffman, M. D. & Gelman, A. The No-U-Turn Sampler: Adaptively Setting Path
Lengths in Hamiltonian Monte Carlo. *Journal of Machine Learning Research* (2014).
59. Franzosa, E. A. *et al.* Species-level functional profiling of metagenomes and
metatranscriptomes. *Nat Methods* **15,** 962–968 (2018).
426
427

Supplementary Information

Appended and available online.

Acknowledgments and Conflicts of Interest

We thank Marc Lipsitch, Kevin R. Foster, Sandra B. Andersen, Jonathan Kevin Sia, Eric G. Pamer, Sibylle Mitschka and the members of the Xavier lab for helpful discussion and comments on the manuscript. This work was supported by the National Institutes of Health (NIH) grant U01 AI124275 and grant R01 AI137269 to JBX, by the MSKCC Cancer Center Core Grant P30 CA008748, the Parker Institute for Cancer Immunotherapy at Memorial Sloan Kettering Cancer Center, the Sawiris Foundation, the Society of Memorial Sloan Kettering Cancer Center, MSKCC Cancer Systems Immunology Pilot Grant and Empire Clinical Research Investigator Program. MS received funding from the Burroughs Wellcome Fund Postdoctoral Enrichment Program, the Damon Runyon Physician-Scientist Award, and the Robert Wood Johnson Foundation. MRMvdB and JUP received financial support from Seres Therapeutics. TMH is investigator in the Pathogenesis of Infectious Diseases from the Burroughs Wellcome Fund, and funded via an award from Geoffrey Beene Foundation, and NIH RO1 AI093808. M-AP has received honoraria from AbbVie, Bellicum, Bristol-Myers Squibb, Incyte, Merck, Novartis, Nektar Therapeutics, and Takeda; has received research support for clinical trials from Incyte, Kite (Gilead) and Miltenyi Biotec; and serves on data and safety monitoring boards for Servier and Medigene and scientific advisory boards for MolMed and NexImmune. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

J.S. and J.B.X. wrote the manuscript. J.S. and J.B.X. designed the analyses with expert help from R.N. J.U.P. and Y.T. contributed to the clinical data preparation, B.P.T. provided the 16S data processing pipelines, A.D. provided the shotgun processing pipelines. All authors contributed to the writing and interpretation of the results.

SUPPLEMENTARY INFORMATION

Methods

Complete blood count collection and characterization

Absolute white blood cells count data were obtained from routine complete blood counts ordered by

clinicians during normal clinical practice, used to obtain informative diagnostic and monitoring

information. Blood samples received in the clinical hematology laboratory were analyzed using

Sysmex XN automated hematology analyzers (Sysmex, Lincolnshire, IL) and, when needed based on

specific flags and parameters as per MSKCC standard operating procedures, were validated manually

using the Sysmex DI-60 Slide Processing System or CellaVision DM9600 Automated Digital

Morphology System (Sysmex, Lincolnshire, IL).

16S rRNA gene amplification and multiparallel sequencing

For each sample, duplicate 50-µl PCRs were performed, each containing 50 ng of purified DNA, 0.2 mM

deoxynucleotide triphosphates, 1.5 mM MgCl2, 2.5 U Platinum Taq DNA polymerase, 2.5 µl of 10× PCR

buffer, and 0.5 µM of each primer designed to amplify the V4-V5: 563F (5′-nnnnnnnn-NNNNNNNNNNNN-

AYTGGGYDTAAAGNG-3′) and 926R (5′-nnnnnnnn-NNNNNNNNNNNN-CCGTCAATTYHTTTRAGT-3′).

A unique 12-base Golay barcode (Ns) precedes the primers for sample identification[50], and one to eight

additional nucleotides were placed in front of the barcode to offset the sequencing of the primers. Cycling

conditions were 94°C for 3 min, followed by 27 cycles of 94°C for 50 s, 51°C for 30 s, and 72°C for 1 min. For

the final elongation step, 72°C for 5 min was used. Replicate PCRs were pooled, and amplicons were purified

using the QIAquick PCR Purification Kit (Qiagen). PCR products were quantified and pooled at equimolar

amounts before Illumina barcodes and adaptors were ligated, using the Illumina TruSeq Sample Preparation

protocol. The completed library was sequenced on an Illumina MiSeq platform following the Illumina

recommended procedures with a paired-end 250 × 250 bp kit

Sequence analysis

The 16S (V4-V5) paired-end reads were merged and demultiplexed. Amplicon sequence variants

(ASVs) were identified using the Divisive Amplicon Denoising Algorithm (DADA2) pipeline

including filtering and trimming of the reads[51]. Reads were trimmed to the first 180 bp or the first

484   point with a quality score Q<2. Reads were removed if they contained ambiguous nucleotides (N) or

485   if two or more errors were expected based on the quality of the trimmed read. We assigned taxonomy

486   to ASVs using a 8-mer based classifier trained by IDTaxa[52] using the SILVA database[53].

Diversity calculations

489   Microbiome alpha-diversity was measured by the inverse Simpson (IS) index of a sample. It was

490   calculated by $IS_i = \frac{1}{\sum_{j=1}^{N} p_{ij}^2}$, where p is the relative abundance of the jth ASV out of N total ASVs in

491   sample i.

Linear mixed-effects model of white blood cell counts

To study the effect of auto-FMT on white blood cells, we investigated the white blood cell counts of

24 enrolled patients of this trial from the day of neutrophil engraftment until 100 days after. FMT

occurred on different days relative to neutrophil engraftment. Thus, we performed an analogous

analysis to that conducted in the original publication that demonstrated how FMT re-established a

diverse microbiome in the post-FMT period[22]. To answer if white blood cell counts differed post-

FMT, we used a linear mixed effects model of white blood cell counts, y, modeled as a function of the

FMT treatment as well as patient and timepoint specific random effects. We included random

intercept terms for each day i and each patient j, and a fixed effects term for the post-FMT period with

associated coefficient "armpost", using the indicator variable "FMT", that is 1 when a patient was

from the FMT treated arm of the trial and day was greater than or equal to the day of the FMT

procedure. We conducted independent analyses for neutrophil, lymphocyte and monocyte counts.

This resulted in the following model of a cell count, y, for patient j on day i:

$$y_{ij} = \beta_0 + armpost * FMT_{ij} + day_i + patient_j + \varepsilon_{ij}, \qquad i = 0, \ldots, D, j = 0, \ldots, P$$

with prior distributions $day_i \sim \mathcal{N}(0, \sigma_{day}^2)$, and $patient_j \sim \mathcal{N}(0, \sigma_{patient}^2)$, independent error

$\varepsilon_{ij} \sim \mathcal{N}(0, \sigma^2)$ and fixed intercept β_0, for the D days post neutrophils engraftment and P patients,

(D=100, P=24). For convenience of those interested in reanalyzing our data, the part of our data

concerning the auto-FMT analysis is available in tidy format (data SD1).[54]

Dynamic systems analyses

We analyzed factors associated with the observed changes of absolute counts of neutrophils, lymphocytes and monocytes between two days. In the following we describe how chronology of events and biological samples were encoded, and the models used to infer a role of medications, clinical parameters and the microbiome on dynamics of white blood cells.

To reveal factors that associate with day-to-day changes in white blood cell counts, we started from a first-order differential equation of white blood cell (W) dynamics:

$$\frac{d(W)}{dt} = W\left(gr + \sum_{j=1}^{P} \beta_j X_j\right)$$

Where gr represents the intercept, i.e. the base line rate of change during immune reconstitution, and β_j are the to-be-estimated coefficients of the P predictors $X_j, j \in P$, of the white blood cell dynamics. This equation was then linearized to

$$\frac{d\, ln(W)}{dt} = gr + \sum_{j=1}^{P} \beta_j X_j$$

And we parameterized the corresponding discrete difference equation:

$$\frac{\Delta \ln(W)}{\Delta t} = gr + \sum_{j=1}^{P} \beta_j X_j$$

where $\Delta\ln(W)$ is the log-difference between single days of neutrophils, lymphocytes or monocytes counts, and $\Delta t=1$ for all intervals. Predictors include the counts of neutrophils, lymphocytes, monocytes, eosinophils and platelets during an interval (homeostatic feedbacks), immunomodulatory medication and clinical observations such as a blood stream infection and the onset of graft versus host disease, HCT parameters such as graft types and conditioning regimens, and, additionally, the microbiota composition in "stage 2" of our analysis (supplementary methods for data exclusion and additional details on interval definitions). Importantly, by parameterizing a dynamic equation and analyzing *rates of change*, our coefficient estimates have an immediate causal interpretation within our modeling framework (i.e. a $\beta_j > 0$ implies that higher levels of the corresponding X_j *increases* the

respective white blood cell type, *W*). To differentiate such results from other associations, analyses of

this type have been termed "mathematical causality"[43].

Stage 1 analysis: Feature selection. Identifying medications and clinical observations associated with

white blood cell dynamics from patients without microbiome data

Stage 1 uses data of patients without any available microbiome samples and the following model of

white blood cell changes, y:

$$y(x|\beta, gr) = gr + \sum_{j=1}^{p} \beta_j X_j + \varepsilon,$$

with intercept, *gr*, and residual error $\varepsilon \sim \mathcal{N}(0, \sigma^2)$. The predictors, *X*, include dummy variables for the

HCT graft type, patients' age on the date of HCT, sex, 13 most frequently observed positive blood

cultures with remaining other blood stream infections grouped into a separate category "other

infections", an indicator for the onset of graft versus host disease, administrations of 55 different,

most common immunomodulatory medications and platelet transfusion events, and HCT conditioning

intensity regimens as well as the log-transformed geometric mean counts of neutrophils, lymphocytes,

monocytes, eosinophils and platelets during the respective interval. We used elastic net regression[55]

for feature selection using the sklearn package for the Python programming language[56]. For elastic net

regression with 50% L1-penalty, predictors were scaled between zero and 1, and we used 10-fold

cross validation (i.e. leaving out 10% of patients at each cross-validation step) to choose the

regularization strength, λ, solving for

$$argmin_{gr,\beta} \left\{ \frac{1}{2N} \sum_{i=1}^{N} (y_i - gr - \sum_{j=1}^{p} x_{ij}\beta_j)^2 + \frac{1}{2}\lambda \sum_{j=1}^{p} |\beta_j| + \frac{1}{2}\lambda \sum_{j=1}^{p} \beta_j^2 \right\}$$

554
Stage 1 yielded a sparse coefficient matrix of predictors used to design the model in stage 2.

Expanded analysis on patients with microbiome data – stage 2

To identify associations between microbiota and white blood cell dynamics, we conducted an

analogous, Bayesian regression using the package PyMC3 for the Python programming language[57].

Stage 1 identified important difference between transplant types, and we therefore stratified our data

into 4 cohorts according to their stem cell graft source. Using data independently from each cohort,

we applied "no U-turn" sampling[58] to produce 10,000 posterior samples from 5 independent MCMC

chains that parameterized the model:

$$y \sim \mathcal{N}(\mu, \sigma^2)$$

$$\mu = gr + \sum_{j=1}^{\hat{P}} x_j \beta_j$$

with uninformative prior distributions

$$gr \sim \mathcal{N}\ (mean = 0,\ standard\ deviation = \ 100)$$

$$\beta_j \sim \mathcal{N}\ (mean = 0,\ standard\ deviation = \ 100)$$

$$\sigma \sim Half\,Cauchy\ (\ beta = 2)$$

where y is the observed daily change of a focal white blood cell type as in stage 1 with normal

distributed mean, μ, and σ, the model uncertainty. μ was a function of the baseline growth rate, gr,

and predictors, $\hat{P}$: medications with non-zero coefficients in stage 1, the white blood cell counts,

patient age and sex, and HCT conditioning intensities; additionally, $\hat{P}$ now included the log-

abundances of microbial genera as measured by 16S sequencing from DNA in the stool collected on

the second day of a daily interval (see supplementary methods for details). We considered taxa that

were among the 100 most abundant, or had reached maximum relative abundances of at least 10%,

and selected those who were non-zero in more than 75% of our samples. White blood cell counts and

microbiota data present during a daily interval were log-transformed, and zeros were filled with half

of the minimum observed non zero counts (i.e. 0.5e3 and 2e-6, respectively). We focused on the

largest cohort (PBSC) and used the independent inference results from TCD, BM, and cord cohorts

for validation.

Validation score

Coefficients learnt from the PBSC patient cohort were assigned a "validation score" based on the

results obtained from the other three MSK patient cohorts. Our requirements for validation were

conservative; we required evidence from our validation data sets as well as absence of counter

evidence. For regression results from each of the validation graft type cohorts, i.e. TCD, BM, and

cord, we checked if a coefficient had more than 75% probability (50%HPDI) to have the same sign as

the mean of the PBSC coefficient posterior for a given predictor. If so, this was considered evidence

of validation, and we summed the evidence over the three validation sets (i.e. maximum score of 3, 1

from each of TCD, BM, and cord cohorts). Conversely, if we found more than 75% probability

among any of the validation data sets that a given predictor had the *opposite* sign as the posterior

mean calculated from PBSC data, this was considered counter evidence and the validation score was

always set to zero.

Validation on data from Duke University

We analyzed 9,603 blood samples with 25,581 associated administrations of immunomodulatory

medications, and 741 microbiota samples from Duke as an orthogonal data set to validate our

findings. The temporal resolution of this data was much lower, and after filtering for samples from the

relevant post neutrophil engraftment period, and by requiring daily intervals, 83 valid, complete data

points were available. Using these data, we correlated daily blood cell changes individually with those

predictors that achieved more than 95% probability density in the positive or negative domain in the

PBSC data regression. For each of these predictors, we present the sign of slopes and Bonferroni

corrected *p*-values from individual linear regressions.

Shotgun sequencing

Sequencing of 124 post-neutrophil engraftment was conducted on the Illumina HiSeq platform. For

details and the processing of the FASTQ files, see supplementary methods. We used the HUMAnN2

pipeline[59] with default settings for functional profiling of our samples, with the UniRef90 data base

and ChocoPhlAn for alignment, and we renormalized our samples by library depth to copies per

million. We used MetaCyc to obtain stratified and unstratified pathway abundances.

Statistical analysis of shotgun data

We calculated the predicted microbiota potency score for each sample and separately for neutrophils,

lymphocytes and monocytes, by multiplying the abundances of taxa in each of the 124 samples with

the corresponding posterior coefficients obtained from the PBSC inference. To distinguish the sets of

metabolic functions that separate samples with positive and negative predicted potencies, we

converted the pathway abundances into presence and absences profiles. We performed a linear

discriminant analysis between positive and negative potency samples with a least squares solver and

automatic shrinkage using the Ledoit-Wolf lemma using the sklearn package for the Python

programming language[56]. To assess differences in the presence or absence of pathways between

samples with positive and negative potency, we used Fisher's exact test for each pathway.

623
624
625
626

628
629

630
631
632 Figure S1: Eosinophil counts and platelet counts per day relative to HCT; line: mean, shaded: ± standard
633 deviation)
634

635
636
637 Figure S2: HCT patient who received an autologous fecal microbiota transplant (auto-FMT, dashed red line)
638 that restored commensal microbial families and ecological diversity in the gut microbiota, with concurrent cell
639 counts of peripheral neutrophils, lymphocytes and monocytes and immunomodulatory drug administrations.

640
641

642
643
644 Figure S3: Neutrophil counts in 24. FMT trial patients. Thin lines: raw data (blue: post-FMT); thick black: mean
645 per day, thick blue: mean+post-FMT coefficient. Means and confidence intervals (shaded region) from linear
646 mixed effects model (methods).

647
648
649 Figure S4: Lymphocyte counts in 24. FMT trial patients. Thin lines: raw data (blue: post-FMT); thick black:
650 mean per day, thick blue: mean+post-FMT coefficient. Means and confidence intervals (shaded region) from
651 linear mixed effects model (methods).
652

653
654
655 Figure S5: Monocyte counts in 24. FMT trial patients. Thin lines: raw data (blue: post-FMT); thick black: mean

per day, thick blue: mean+post-FMT coefficient. Means and confidence intervals (shaded region) from linear mixed effects model (methods).

Figure S6: "Stage 1" regression on neutrophil dynamics on patients without microbiome data. Coefficients from 10-fold cross-validated elastic net regression daily changes in neutrophils. gr: intercept; TCD: T-cell depleted graft (ex-vivo) by CD34+selection; PBSC: peripheral blood stem cells; BM: bone marrow; cord: umbilical cord blood; REDUCE: reduced-intensity conditioning regimen; F: female; N: patients, n: samples (daily changes in neutrophils).

Figure S7: "Stage 1" regression on lymphocytes dynamics on patients without microbiome data. Coefficients from 10-fold cross-validated elastic net regression daily changes in lymphocytes. gr: intercept; TCD: T-cell depleted graft (ex-vivo) by CD34+selection; PBSC: peripheral blood stem cells; BM: bone marrow; cord: umbilical cord blood REDUCE: reduced-intensity conditioning regimen; F: female. N: patients, n: samples (daily changes in lymphocytes).

R^2: 0.084
N: 1,095
n:20,019

gr	4.74e-01
G-CSF	2.73e-01
monocytes	1.83e-01
eosinophils	2.76e-02
F	6.00e-03
age	3.74e-03
REDUCE	1.29e-03
hydrocortisone	-1.59e-03
prednisone	-5.80e-03
cetirizine	-7.02e-03
tacrolimus	-1.39e-02
TCD	-1.45e-02
diphenhydramine	-1.50e-02
octreotide	-1.66e-02
platelet_transfusion	-1.73e-02
cytomegalovirus immune globulin	-2.24e-02
BM	-2.50e-02
mycophenolate mofetil	-3.82e-02
dexamethasone	-3.94e-02
cyclosporine	-4.08e-02
cord	-4.60e-02
hydroxyzine	-5.40e-02
immune globulin	-5.57e-02
platelets	-1.57e-01
lymphocytes	-2.75e-01
neutrophils	-4.93e-01

Effect on
Δ monocytes

Figure S8: "Stage 1" regression on lymphocytes dynamics on patients without microbiome data. Coefficients from 10-fold cross-validated elastic net regression daily changes in lymphocytes. gr: intercept; TCD: T-cell depleted graft (ex-vivo) by CD34+selection; PBSC: peripheral blood stem cells; BM: bone marrow; cord: umbilical cord blood REDUCE: reduced-intensity conditioning regimen; F: female. N: patients, n: samples (daily changes in lymphocytes).

681
682
683   Figure S9: Additional coefficient estimates from the Bayesian regression, see also Figure 3. REDUCE: reduced-
684   intensity conditioning regimen; NONABL: non-myeloablative conditioning regimen. F: female

Figure S10: Validation analysis of predictors on white blood cell dynamics using data from patients treated at DukeHealth. Individual univariate regressions of microbiome and clinical predictors identified in stage 2 of our analysis on daily changes in neutrophils, lymphocytes and monocyte. Bonferroni corrected p-values: ***<0.001, **<0.01, *<0.05; p>0.05: n.s. Sign of coefficients from MSK PBSC patients for comparison.

Figure S11: Functional analysis of microbiota samples. To distinguish samples predicted to increase rates of white blood cells, a microbiota potency score was calculated from posterior coefficients (Figure 3, methods) and the relative abundance of taxa in samples. Bars show linear discriminant analysis (LDA) scores of MetaCyc pathway profiles from 124 shotgun sequenced samples that distinguished positive and negative potency samples the most (LDA-score magnitude in the 95th percentile). Highlighted pathways are discussed in the main text. For each pathway, we tested differences between positive and negative potency samples using Fisher's exact test; p-value <0.001: ***, <0.01:**, <0.05:*.

Figure S12: The relative non-zero abundance of Staphylococcus is inversely related to microbiome alpha diversity, shaded: 95% confidence intervals.

723
724
725 Figure S13: Abundance profiles of the two genera, *Faecalibacterium* and *Ruminococcus 2*, most strongly
726 associated with white blood cell increase; number of times detected (left) and log_{10} abundance distribution when
727 above detection (right).

735 Table S1: Data set summary and patient characteristics. HCT-graft types: TCD: T-cell depleted graft (*ex-vivo*)
736 by CD34+selection; PBSC: peripheral blood stem cells; BM: bone marrow; cord: umbilical cord blood;
737 Conditioning intensity: Bacigalupo classification, graded categories from most to least intense (ABLATIVE,
738 REDUCE, NONABL).
739

patients		2,926
HCT therapies*		3,060
blood samples	total	450,635
	between HCT-day -21 and HCT-day 183	193,396
Disease	Leukemia	1,635
	Non-Hodgkin's Lymphoma	415
	Multiple Myeloma	170
	Hodgkin's disease	88
	other	752
HCT graft type	TCD	1,106
	PBSC unmodified	959
	BM unmodified	617
	cord	378
Conditioning intensity	ABLATIVE	65%
	REDUCE	21%
	NONABL	13%
Gender	M	58%
	F	42%
Age of adults (years)	25%-tile	39
	mean	50
	75%-tile	62
Microbiome samples	total	12,633
	- from patients with blood data	10,680
	- of those, post engraftment	4,179
	- of those with daily change in WBC	2,615
	patients with microbiome sample	1,290

740 *) some patient received several HCTs
741
742
743 Table S2: Patient and HCT characteristics of 24 patients enrolled in the randomized controlled FMT trial.

	control	FMT treated
N patients	10	14
ABLATIVE	6	7
REDUCE	4	7
BM unmodified	1	3
PBSC unmodified	3	4
TCD	5	3
cord	1	4

752 Table S3: Patient and HCT characteristics of the subset of patients who donated microbiota samples.

patients		1,294
Disease distribution	Leukemia	51%
	Non-Hodgkin's Lymphoma	15%
	Multiple Myeloma	8%
	Hodgkin's disease	3%
	other	23%
HCT graft type	TCD	37%
	PBSC unmodified	38%
	BM unmodified	9%
	cord	16%
Conditioning intensity	ABLATIVE	55%
	REDUCE	34%
	NONABL	11%
Gender	M	59%
	F	41%
Age (years)	25%-tile	46
	mean	54
	75%-tile	65

753
754
755 Table S4: Patient and HCT characteristics of the subset of patients who did not donate microbiota samples.

patients		1,010
Disease distribution	Leukemia	53%
	Non-Hodgkin's Lymphoma	17%
	Multiple Myeloma	5%
	Hodgkin's disease	5%
	other	20%
HCT graft type	TCD	40%%
	PBSC unmodified	31%
	BM unmodified	17%
	cord	12%
Conditioning intensity	ABLATIVE	65%
	REDUCE	14%
	NONABL	21%
Gender	M	58%
	F	42%
Age (years)	25%-tile	36
	mean	47
	75%-tile	59

756
757
758 Table S5: Patient and HCT characteristics of the Duke University patient cohort.
759

patients		493
Disease distribution	Lymphoma	11%
	Leukemia	50%
	Non-Hodgkin's Lymphoma	4%
	Multiple Myeloma	8%

		Hodgkin's disease	4%
		other	24%
HCT graft type		TCD	0%
		PBSC unmodified	72%
		BM unmodified	11%
		cord	16%
Conditioning intensity		ABLATIVE	92%
		NONABL	7%
Gender		M	65%
		F	35%
Age (years)		25%-tile	41
		mean	49
		75%-tile	57

Supplementary Data

The data used in our study is organized in supplementary tables (data-tables.zip), with corresponding filenames (italic):

1. *cGENUS.csv*: relative taxon abundances in fecal microbiota samples from 12,633 stool samples
2. *cHCTMETA.csv*: HCT characteristics
3. *cINFECTIONS.csv*: positive blood culture results
4. *cIVS.csv*: diversity (inverse Simpson index) for each fecal microbiota sample
5. *cMED.csv*: medication data
6. *cPIDMETA.csv*: anonymized patient demographics
7. *cWBC.csv*: absolute counts of neutrophils, lymphocytes, monocytes, eosinophils, and platelets with indication if included in analyses
8. *cDUKE__GENUS.csv*: relative taxon abundances in fecal microbiota samples from 12,633 stool samples
9. *cDUKE__WBC.csv*: absolute counts of neutrophils, lymphocytes, monocytes, eosinophils, and platelets with indication if included in analyses
10. *cDUKE__MED.csv*: medication data
11. FMT_analysis.csv: convenience table for Figure 2