The gut microbiota influences circulatory immune cell dynamics in humans

Jonas Schluter[1], Jonathan U. Peled[2,3], Bradford P. Taylor[1], Melody Smith[2,3], Ying Taur[4], Sejal

Morjaria[4], Maly Fenelus[5], Melissa S. Pessin[5], Tobias M. Hohl[4], Miguel-Angel Perales[2,3], Marcel R.M.

van den Brink[2,3], Joao B. Xavier[1]

1) Computational and Systems Biology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer

Center, New York, NY, USA

2) Adult Bone Marrow Transplantation Service, Department of Medicine, Memorial Sloan Kettering Cancer

Center, New York, NY, USA

3) Weill Cornell Medical College, New York, NY, USA

4) Infectious Disease Service, Department of Medicine, and Immunology Program, Sloan Kettering Institute,

New York, NY, USA.

5) Department of Laboratory Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

Correspondence: schluetj@mskcc.org, xavierj@mskcc.org

ABSTRACT

The gut microbiota influences the development and homeostasis of the mammalian immune system[1–3]

and can alter immune cell compositions in mice[4–7]. However, our understanding of how the

microbiota modulates immunity remains limited, particularly in humans where a lack of manipulative

experiments makes inference challenging. Here we overcome this challenge by studying hundreds of

hospitalized—and closely monitored—bone marrow transplantation patients as they recover from

chemotherapy-induced immune ablation. This aggressive treatment causes large shifts in both white

blood cell and microbiota populations allowing the relationships between the two to be studied

simultaneously over time, with unprecedented resolution. Our analysis shows that the ecological

diversity of the gut microbiota had an immunosuppressive effect on circulating lymphocyte counts

similar in magnitude to that of anti-inflammatory and immunosuppressive drugs administered to

cancer patients. Moreover, by controlling for drug treatments and clinical metadata, we identified

several microbiota components strongly associated with white blood cell dynamics: Streptococcaceae

associated with lymphocyte increase and Actinomycetaceae with lymphocyte suppression,

Prevotellaceae with monocyte increase and Enterobacteriaceae with monocyte suppression,

Ruminococcaceae with neutrophil increase and Lachnospiraceae and Bacteroidaceae with neutrophil

suppression. Our analysis establishes a direct link between the intestinal microbiota and systemic

immunity in humans, with implications on clinical outcomes and microbiota-based therapies.

MAIN TEXT

Experiments in mice provide mounting evidence that the mammalian intestinal microbiome influences the development[1–3] and homeostasis of its host's immune system[4–10]. Further evidence comes from a recent study showing that white blood cell compositions differ between children born preterm and at term along with their microbiome composition[11]. However, our understanding of the influence of the microbiota on the adult immune system remains limited. To overcome this, we here investigated the dynamics of immune reconstitution after allogeneic hematopoietic cell therapy (HCT) in hundreds of patients at Memorial Sloan Kettering, treated for various hematological malignancies, including leukemia (**Figure 1A, Table S1**).

The conditioning regimen of radiation and chemotherapy administered to HCT patients is the most severe perturbation to the immune system deliberately performed in humans and presents a unique opportunity to investigate the dynamic links between gut microbiota and immune system directly in humans. Immune ablation during conditioning (**Figure 1A**) depletes circulating white blood cell counts. This is followed by characteristic white blood cell trajectories with prolonged periods of neutropenia (<500 neutrophils per μl blood) and reconstitution of white blood cell counts once the transplanted cells engraft (**Figure 1A-C**). The blood of individual patients is monitored daily throughout this recovery, and medications are administered to ensure successful immune reconstitution and prevent graft-vs-host disease (**Figure 1 G,H**).

To investigate whether the composition of the gut microbiota could affect the dynamics of circulating white blood cells in our patients, we combined high frequency, longitudinal microbiota compositions measured for hundreds of patients[12] with the detailed blood and metadata from thousands of patients. We compiled over 75,000 administrations of a range of medications, clinical events such as blood stream infections, and over 100,000 host physiology measurements in the form of the most abundant white blood cells—neutrophils, lymphocytes, monocytes, eosinophils—as well as platelet counts (**Figure 1, S1**).

Figure 1: Immune reconstitution after hematopoietic cell transplantation (HCT). A) Three major periods of HCT: immunoablation during conditioning with chemotherapy and radiation before HCT, defined as day 0, post-HCT neutropenia, and reconstitution following neutrophil engraftment lead to repeatedly observed recovery trajectories. Shown are the average counts (shaded: ± std) of neutrophils, lymphocytes and monocytes per day relative to HCT from 2,047 patients transplanted between 2003 and 2018, with count ranges from healthy adults indicated (A), and two individual patients (B,C); patient 1 who received a peripheral blood stem cell graft, and patient 2 who received a graft of umbilical cord blood. For 460 patients, fecal samples were collected during HCT, with varying coverage, revealing loss of microbial diversity (D-F) and commensal families (G-I), often replaced by Enterococcaceae domination (shaded: ± std). J,K) Administration of immunomodulatory medications for the two example patients.

We have shown previously that HCT patients lose gut microbiota biodiversity and commensal microbial families during their treatment (**Figure 1D,G-I**)[12]. A recent study in mice demonstrated that such antibiotic-induced gut decontamination can negatively affect white blood cell recovery after HCT[13]. In our patients, microbial diversity usually recovers slowly during white blood cell reconstitution (**Figure 1D**); however, microbiota recovery as well as immune reconstitution can vary strongly between patients and treatment types. This variation is illustrated by the distinct trajectories

of two patients: patient 1 who received a graft of peripheral blood stem cells, retained high microbiota

diversity and engrafted earlier (**Figure 1B,E,H**), and patient 2 who received a graft of umbilical cord

blood, lost microbiota diversity and engrafted later (**Figure 1C,F,I**). Low microbiota diversity at the

time of neutrophil engraftment has independently been associated with 5-fold increased transplant-

related mortality[14], suggesting that that the joint recovery of microbiota diversity and white blood

cells in circulation is critical for clinical outcomes.

To identify a causal link between the microbiota and white blood cell recovery dynamics, we first used data from a recent prospective clinical trial of autologous fecal microbiota transplantation (auto-FMT) in our patients[12]. We performed this deliberate microbiota manipulation in 24 patients, randomized to 10 untreated control and 14 treated patients, including patient 3 in **Figure 2A**. To ask if auto-FMT affected white blood cell reconstitution, we compared the 24 patients' neutrophil, lymphocyte and monocyte counts post-engraftment (i.e. when the transplanted hematopoietic cells begin producing new white blood cells). We observed higher counts of each white blood cell type in the period after randomization into control and treatment groups in patients who received an auto-FMT (p<0.01, **Figure 2B**).

Figure 2: HCT patient who received an autologous fecal microbiota transplant (auto-FMT, dashed red line)
that restored commensal microbial families and ecological diversity in the gut microbiota (A), with concurrent
cell counts of peripheral neutrophils, lymphocytes and monocytes and immunomodulatory drug
administration. B) Results from a linear mixed effects model with random effects per patient and per day
relative to neutrophil engraftment show elevated neutrophil, lymphocyte and monocyte counts in patients
receiving auto-FMT after their treatment as compared to control patients (***: p<0.01).
108

The increased white blood cell counts in patients receiving auto-FMT pointed towards a

causal role of this severe microbiota manipulation on circulatory immune cell dynamics. This increase

could be due to the re-establishment of a complex microbiota and the associated metabolic

capabilities[6,7,13], or it could be a systemic response of the body to the sudden influx of billions of

intestinal organisms. Moreover, while our mixed-effects analysis accounted for patient-specific HCT

treatments such as their graft source, chance differences in extrinsic factors such as

immunomodulatory drug administration could have affected this result due to the small cohort size.

Therefore, we next investigated the link between the gut microbiota composition and the dynamics of

white blood cell recovery using our large observational cohort of HCT patients.

Homeostasis of mammalian blood cell counts is complex: neutrophils, lymphocytes and

monocytes are formed and released into the blood *de novo* by differentiation of hematopoietic

progenitor cells from the bone marrow, and they can be mobilized from thymus and lymph nodes

(lymphocytes), spleen, liver and lungs (neutrophils); they can also migrate from the blood to other

tissues when needed[15]. These processes are dynamic sources and sinks of circulatory white blood

cells, and they can be modulated by drugs administered to patients receiving HCT. To identify drivers

of these dynamics—including the microbiota—we developed a two-stage approach. We first used a

dataset of patients *without* microbiome information to identify medications, clinical observations and

other metadata that drive changes in white blood cell counts from one day to the next. This prior

information fed into and informed our main analysis which we performed on set of different patients

from whom concurrent microbiome samples were available.

We first used the longitudinal data of over 1,500 patients from whom no microbiome samples

were collected (**Figure 3A**). We analyzed the counts of neutrophils, lymphocytes and monocytes

during patients' recovery (i.e. from 5 days before neutrophil engraftment until day 100), observing re-

establishment of homeostasis 'as it happened'. We used 3-fold cross-validated (lasso) regression to

identify factors consistently associated with increases and decreases in neutrophil, lymphocyte and

monocyte counts among these patients, including their stem cell graft source, medications as well as

various clinical and demographic data (**Figure S2,S3, Table S1**). We then used the identified effects

as prior information for a Bayesian inference to contextualize potential contributions of the gut

microbiota. The set of patients with microbiota data was smaller (446 patients, **Figure 3A**), but they

provided an unbiased sampling of the white blood cell dynamics observed across patients lacking

microbiota data (**Figure 3B**).

Notably, as a benchmark, this approach detected the expected strong effects of

immunomodulatory medications (**Figure 3C,S4**). Most obviously, we identified the administration of

granulocyte-stimulating factor (GCSF)—administered to accelerate recovery from chemotherapy-

induced neutropenia[16]—as the strongest driver of neutrophil and monocyte increases from one day to

the next, yielding an estimated increase of 97% ([76%, 121%], 90 percent probability density interval

[HDI90]) of neutrophils and an increase of 41% ([28%, 54%] HDI90) of monocytes. Lymphocytes,

too, increased during GCSF administration, although less so (9%, [-1%, 19%] HDI90). The strongest

associations with lymphocyte dynamics resulted from administration anti-inflammatory steroids that

are known to induce lymphocytopenia by removing circulatory lymphocytes (prednisone[17], -16%

([-27%, -3%], HDI90).

Beyond correctly recognizing the causal roles of medications with known biological

mechanisms, our analysis detected feedbacks between white blood cell types. Specifically, we

identified a negative feedback loop between neutrophils and lymphocytes, a suppression of

monocytes by neutrophils and platelets (**Figure 3D**), and an additional suppression from platelet

transfusion events (**Figure 3C**). Monocytes positively fed-back on each of the investigated white

blood cell counts, including a positive feedback of monocytes onto themselves. Interestingly, this

suggests a homeostatic feedback network of circulatory white blood cell counts.

Figure 3: Results from Bayesian inference of coefficients in a model of peripheral white blood cell dynamics. A) Cartoon of the two-stage model inferred from patients' data: observed changes in white blood cell counts between two days are driven by the current state of the host, including the current counts of white blood cells in circulation, the effects of immunomodulatory medications administered, patient metadata and the state of the microbiome. Host state and medications were used in a prior analysis on patients without microbiota samples (brown outline, results in Figure S3), and informed the main analysis presented in panels C-F on patients with concurrent microbiome information. B) Visualization of the dynamic white blood cell data used; scatter plot of the principal components (PC) of observed daily changes of neutrophils, lymphocytes and monocytes without (grey) and with (green) concurrent longitudinal microbiome data. C-F) Regression results; bars show the posterior parameter estimate distributions (highest density intervals, HDI, color intensity indicate credibility of results) for the jointly inferred effects of immunomodulatory medications and platelet transfusion (C), homeostatic regulation network of white blood cells counts and associated posteriors (D), microbiota alpha diversity (inverse Simpson index, E) and log-microbial family abundances (F) on the observed daily changes in neutrophils, lymphocytes and monocytes.

To investigate the immunomodulatory role of the gut microbiota ecosystem, we included the inverse Simpson diversity index—an ecological index calculated at the OTU level. Additionally, we also included the relative abundances of OTUs combined at the family level where many important physiological traits are shared across member species[18]. We found that high diversity suppressed circulatory lymphocyte counts (**Figure 3E**). A raise in microbiota diversity of 5 inverse Simpson

index units corresponded to a -6% ([-11%, 1%] HDI90) suppression of lymphocytes. This is similar in

magnitude to the effect of immunosuppressive drug administrations to prevent graft-vs-host disease

(mycophenolate mofetil, -10% [-20%, 1%] HDI90) and that of anti-inflammatory steroids (**Figure**

3C). Whether this apparent immunosuppressive effect of a complex microbiota is an indicator for

good recovery and re-acquisition of homeostatic regulation among our patients, or indeed, may be

problematic in the special situation of an immunosuppressed population, requires further

investigation. Nonetheless, high microbiota diversity has a strong positive association with better

survival of HCT patients[14].

Our analysis revealed several commensal microbial families that modulated white blood cell

dynamics (**Figure 3F**). Bacteroidaceae increased lymphocytes, Prevotellaceae increased monocytes,

and Ruminococcaceae increased neutrophils. The effect sizes of individual families were relatively

small, suggesting that potential immunomodulatory microbiota treatments would need to be

prolonged or simultaneously target multiple microbial families. A notable exception would be a

reduction in Streptococcaceae: Streptococcaceae dominations reached maximum relative abundances

of over 90% in our patient cohort while their median abundance was <4%. Our results suggest that an

increase in Streptococcaceae from this median level to >90% domination results in daily lymphocyte

increases of ~6% or, conversely, that resolving a Streptococcaceae domination could have an

immunosuppressive effect. This response of circulatory lymphocyte counts to Streptococcaceae

domination could reflect reduced lymphocyte migration to the gut epithelium or indicate a gut

dysbiosis-induced acute immune response that may influence the risk of gut domination-born

infections[7,19].

Microbiota homeostasis is characterized by high abundances of obligate anaerobes. Our

analysis revealed that two major families of commensal obligate anaerobes, Bacteroidaceae and

Lachnospiraceae, suppressed neutrophils (**Figure 3F**). Neutrophil-associated immunity in mice

responds to gut bacteria, which could affect tumor progression in distal organs with reduced overall

tumor burden upon systemic neutrophil depletion[20]. Intriguingly, our finding that common commensal

families can modulate circulatory immune cells in human patients supports the idea established in

mice that microbiota modulation could influence systemic immunity to fight tumor progression[20,21].

The most immediate interface between mammalian hosts and their microbiota are the various

epithelia of the body. In mice, individual microbial species trigger specific immune responses at the

gut epithelium[9,22,23]. Also, various microbial traits, including the production of metabolic byproducts

such as short-chain fatty acids[24,25] and bacterial cell wall molecules[1,26,27] modulate immune responses,

including granulopoiesis[6]. These traits are shared by many different microbial species, and therefore

plausible candidates for the human body to sense gut homeostasis or dysbiosis. High-resolution,

longitudinal microbiome data[12] coupled with a large observational dataset of immune reconstitution in

cancer patients enabled us to detect immunomodulatory effects of the microbiota ecosystem. Our

approach accounted for patient differences such as the source of transplanted stem cells and yielded

features that consistently associated with increases or decreases of white blood cells from one day to

the next. Intriguingly, systemic effects of microbiota ecosystem features support predictions from

evolutionary theory that hosts evolved to sense, respond to and control broad ecological features such

as diversity and family abundances, rather than specific strains[28,29].

In sum, our work shows evidence of a link between the gut microbiota and the human

immune system via peripheral white blood cell dynamics. We were able to infer this link because we

leveraged an unprecedented resource: a vast dataset of high-resolution, longitudinal human phenotype

data in the form of blood cell counts, compiled with detailed metadata. The effects inferred between

the microbiota and the temporal changes in circulatory white blood counts indicate a systemic,

homeostatic link that could help understand the modulating role of a 'healthy' microbiota during

immunotherapy[30–33], and holds potential for microbiota therapies[20,21,34].

257 Bibliography

1. Mazmanian, S. K., Liu, C. H., Tzianabos, A. O. & Kasper, D. L. An immunomodulatory molecule of
symbiotic bacteria directs maturation of the host immune system. *Cell* **122,** 107–118 (2005).
2. Gomez de Agüero, M. *et al.* The maternal microbiota drives early postnatal innate immune
development. *Science* **351,** 1296–1302 (2016).
3. Sonnenberg, G. F. & Artis, D. Novel connections and precision approaches. *Nat Rev Immunol* **19,**
75–76 (2019).
4. Geva-Zatorsky, N. *et al.* Mining the human gut microbiota for immunomodulatory organisms.
Cell **168,** 928–943.e11 (2017).
5. Tan, T. G. *et al.* Identifying species of symbiont bacteria from the human gut that, alone, can
induce intestinal Th17 cells in mice. *Proc Natl Acad Sci U S A* **113,** E8141–E8150 (2016).
6. Balmer, M. L. *et al.* Microbiota-derived compounds drive steady-state granulopoiesis via
MyD88/TICAM signaling. *J Immunol* **193,** 5273–5283 (2014).
7. Deshmukh, H. S. *et al.* The microbiota regulates neutrophil homeostasis and host resistance to
Escherichia coli K1 sepsis in neonatal mice. *Nat Med* **20,** 524–530 (2014).
8. Atarashi, K. *et al.* ATP drives lamina propria T(H)17 cell differentiation. *Nature* **455,** 808–812
(2008).
9. Ivanov, I. I. *et al.* Specific microbiota direct the differentiation of IL-17-producing T-helper cells
in the mucosa of the small intestine. *Cell Host Microbe* **4,** 337–349 (2008).
10. Cahenzli, J., Köller, Y., Wyss, M., Geuking, M. B. & McCoy, K. D. Intestinal microbial diversity
during early-life colonization shapes long-term IgE levels. *Cell Host Microbe* **14,** 559–570 (2013).
11. Olin, A. *et al.* Stereotypic immune system development in newborn children. *Cell* **174,** 1277–
1292.e14 (2018).
12. Taur, Y. *et al.* Reconstitution of the gut microbiota of antibiotic-treated patients by autologous
fecal microbiota transplant. *Sci Transl Med* **10,** (2018).
13. Staffas, A. *et al.* Nutritional Support from the Intestinal Microbiota Improves Hematopoietic
Reconstitution after Bone Marrow Transplantation in Mice. *Cell Host Microbe* **23,** 447–457.e4
(2018).
14. Taur, Y. *et al.* The effects of intestinal tract bacterial diversity on mortality following allogeneic
hematopoietic stem cell transplantation. *Blood* **124,** 1174–1182 (2014).
15. Scheiermann, C., Frenette, P. S. & Hidalgo, A. Regulation of leucocyte homeostasis in the
circulation. *Cardiovasc Res* **107,** 340–351 (2015).
16. Gabrilove, J. L. *et al.* Effect of granulocyte colony-stimulating factor on neutropenia and
associated morbidity due to chemotherapy for transitional-cell carcinoma of the urothelium. *N
Engl J Med* **318,** 1414–1422 (1988).
17. Fauci, A. S. & Dale, D. C. Alternate-day prednisone therapy and human lymphocyte
subpopulations. *J Clin Invest* **55,** 22–32 (1975).
18. Goldford, J. E. *et al.* Emergent simplicity in microbial community assembly. *Science* **361,** 469–
474 (2018).
19. Taur, Y. *et al.* Intestinal domination and the risk of bacteremia in patients undergoing allogeneic
hematopoietic stem cell transplantation. *Clin Infect Dis* **55,** 905–914 (2012).
20. Lakritz, J. R. *et al.* Gut bacteria require neutrophils to promote mammary tumorigenesis.
Oncotarget **6,** 9387–9396 (2015).
21. Lakritz, J. R. *et al.* Beneficial bacteria stimulate host immune cells to counteract dietary and
genetic predisposition to mammary cancer in mice. *Int J Cancer* **135,** 529–540 (2014).
22. Ivanov, I. I. *et al.* Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* **139,**
485–498 (2009).
23. Tanoue, T. *et al.* A defined commensal consortium elicits CD8 T cells and anti-cancer immunity.
Nature **565,** 600–605 (2019).

24.   Ulven, T. Short-chain free fatty acid receptors FFA2/GPR43 and FFA3/GPR41 as new potential
therapeutic targets. *Front Endocrinol* **3,** 111 (2012).
25.   Smith, P. M. *et al.* The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell
homeostasis. *Science* **341,** 569–573 (2013).
26.   Hergott, C. B. *et al.* Peptidoglycan from the gut microbiota governs the lifespan of circulating
phagocytes at homeostasis. *Blood* **127,** 2460–2471 (2016).
27.   Ladinsky, M. S. *et al.* Endocytosis of commensal antigens by intestinal epithelial cells regulates
mucosal T cell homeostasis. *Science* **363,** (2019).
28.   Schluter, J. & Foster, K. R. The evolution of mutualism in gut microbiota via host epithelial
selection. *PLoS Biol* **10,** e1001424 (2012).
29.   Foster, K. R., Schluter, J., Coyte, K. Z. & Rakoff-Nahoum, S. The evolution of the host
microbiome as an ecosystem on a leash. *Nature* **548,** 43–51 (2017).
30.   Zitvogel, L., Ma, Y., Raoult, D., Kroemer, G. & Gajewski, T. F. The microbiome in cancer
immunotherapy: Diagnostic tools and therapeutic strategies. *Science* **359,** 1366–1370 (2018).
31.   Routy, B. *et al.* Gut microbiome influences efficacy of PD-1-based immunotherapy against
epithelial tumors. *Science* **359,** 91–97 (2018).
32.   Gopalakrishnan, V. *et al.* Gut microbiome modulates response to anti-PD-1 immunotherapy in
melanoma patients. *Science* **359,** 97–103 (2018).
33.   Hsiao, E. Y. *et al.* Microbiota modulate behavioral and physiological abnormalities associated
with neurodevelopmental disorders. *Cell* **155,** 1451–1463 (2013).
34.   Chen, H. *et al.* A forward chemical genetic screen reveals gut microbiota metabolites that
modulate host physiology. *Cell* (2019). doi:10.1016/j.cell.2019.03.036
35.   Wheeler, D. L. *et al.* Database resources of the National Center for Biotechnology Information.
Nucleic Acids Res **35,** D5-12 (2007).
36.   Anasetti, C. *et al.* Peripheral-blood stem cells versus bone marrow from unrelated donors. *N*
Engl J Med **367,** 1487–1496 (2012).
37.   Davies, S. M. *et al.* Engraftment and survival after unrelated-donor bone marrow
transplantation: a report from the national marrow donor program. *Blood* **96,** 4096–4102
(2000).
38.   Tibshirani, R. Regression shrinkage and selection via the lasso. *J. Royal Stat. Soc.*
(Methodological) **58,** 267–288 (1996).
39.   Salvatier, J., Wiecki, T. V. & Fonnesbeck, C. Probabilistic programming in Python using PyMC3.
PeerJ Computer Science **2,** e55 (2016).
340
341

Supplementary Information

Appended and available online.

Acknowledgments and Conflicts of Interest

We thank Kevin R. Foster, Sandra B. Andersen, Rene Niehus, Katharine Coyte, Jonathan Kevin Sia, Eric G. Pamer, and the Xavier lab for helpful discussion and comments on the manuscript. This work was supported by the National Institutes of Health (NIH) grant U01 AI124275 and grant R01 AI137269 to JBX. This research was further supported by the MSKCC Cancer Center Core Grant P30 CA008748, the Parker Institute for Cancer Immunotherapy at Memorial Sloan Kettering Cancer Center, the Sawiris Foundation, the Society of Memorial Sloan Kettering Cancer Center, MSKCC Cancer Systems Immunology Pilot Grant, Empire Clinical Research Investigator Program. MS received funding from the Burroughs Wellcome Fund Postdoctoral Enrichment Program, the Damon Runyon Physician-Scientist Award, and the Robert Wood Johnson Foundation. MRMvdB and JUP received financial support from Seres Therapeutics. TMH is investigator in the Pathogenesis of Infectious Diseases from the Burroughs Wellcome Fund, and funded via an award from Geoffrey Beene Foundation, and NIH RO1 AI093808. M-AP has received honoraria from AbbVie, Bellicum, Bristol-Myers Squibb, Incyte, Merck, Novartis, Nektar Therapeutics, and Takeda; has received research support for clinical trials from Incyte, Kite (Gilead) and Miltenyi Biotec; and serves on data and safety monitoring boards for Servier and Medigene and scientific advisory boards for MolMed and NexImmune. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

J.S., B.P.T. and J.B.X. wrote the manuscript. J.S. and J.B.X. designed the analyses. J.U.P. and Y.T. contributed to the data preparation. All authors contributed to the writing and interpretation of the results.

SUPPLEMENTARY INFORMATION

Methods

Complete blood count collection and characterization

Absolute white blood cells count data were obtained from routine complete blood counts ordered by

clinicians during normal clinical practice, used to obtain informative diagnostic and monitoring

information. Blood samples received in the clinical hematology laboratory were analyzed using

Sysmex XN automated hematology analyzers (Sysmex, Lincolnshire, IL) and, when needed based on

specific flags and parameters as per MSKCC standard operating procedures, were validated manually

using the Sysmex DI-60 Slide Processing System or CellaVision DM9600 Automated Digital

Morphology System (Sysmex, Lincolnshire, IL).

Microbial sample collection, data availability and ecological diversity calculation

The microbiota data analyzed here was obtained from a previous publication [12]. Briefly, stool samples

were collected from patients and sequenced using primers for the V4 and V5 region of the 16S gene.

OTU clustering was performed using UPARSE and classification was conducted using NCBI BLAST

on the refseqRNA database[35]. Diversity as measured by the inverse Simpson index of a sample was

calculated by $IS = \frac{1}{\sum_{j=1}^{N} p_{ij}^2}$, where p is the relative abundance of the jth OTU out of N total OTUs in

sample i.

Linear mixed-effects model of white blood cell counts

We used a linear mixed effects model to analyze white blood cell counts before and after auto-FMT

with the following model equation: *value ~ armpost + (1|day) +(1|patientid)*, where *value* is the

count of a given white blood cell on a given *day* relative to neutrophil engraftment, and *armpost* is a

binary variable that is 1 when a patient was from the FMT treated arm of the trial and *day* was greater

than or equal to the day of FMT procedure.

Dynamic systems analyses

We analyzed factors associated with the observed changes of absolute counts of neutrophils,

lymphocytes and monocytes between two days.

Interval data calculation

For a given daily interval, a medication was considered present for at least part of the interval when it

was administered on either endpoint. For the microbiota and blood stream infection data, data from

the end day were considered for the interval. Homeostatic feedback calculations used the geometric

mean of the white blood cell counts between the two endpoints. We included only those covariates

that were present during at least 10 intervals. The stem cell graft source is a major determinant of

engraftment times and can affect recovery dynamics[36,37], and we therefore included intercept terms for

unmodified peripheral blood stem cell grafts (PBSC), bone marrow (BM), T-cell depleted graft

(*ex-vivo*) by CD34+selection (TCD) and cord blood (cord).

Identification of effectors on white blood cell dynamics from patients without microbiome data

To reveal factors that affect day-to-day white blood cell count changes, we used L1-regularized

(lasso) regression[38]. Stronger regularization leads to fewer predictors in the final model; we here used

3-fold cross validation leaving out 33% of patients at each cross-validation step to choose the

regularization strength, yielding a sparse coefficient matrix of white blood cell changes. We used this

to parameterize a linearized differential equation of daily blood cell count changes,

$$\frac{\Delta \ln(W_i)}{\Delta t} = gr + \sum_{j=0}^{WBCs} s_{ij} W_j + \sum_{i=0}^{drugs, clinical\ obs, HCT\ parameters, patient\ demographics} \beta_i X_i$$

where $\Delta\ln(W_i)$ is the log difference between single days of neutrophils, lymphocytes or monocytes

counts, and $\Delta t=1$ for all intervals. gr is the intercept and represents the base line rate of change, and *s*

is the coefficient for homeostatic feedback of the current blood state, i.e. the counts of blood cells W_j

(neutrophils, lymphocytes, monocytes, eosinophils, platelets) on the rate of change of W_i. X_i are

indicator variables for clinical observations (e.g. bacteremia), patient gender, the drug administrations

and the type of cells that were transplanted (BM, PBSC, TCD, cord). Continuous variables were zero

centered and standardized to unit variance. After cross-validated regression, the resulting coefficient

matrix was used to inform priors of a hierarchical Bayesian model.

Expanded analysis on patients with microbiome data

We inferred effects of medications, clinical parameters and patients' demographics onto white blood

cell rates of change in an analogous Bayesian regression using the package PyMC3 for the Python

programming language[39]. We incorporated the above results as prior information, now additionally

including a linear effect from log-transformed family relative abundances (F_i). We combined the

relative OTU abundances at the family level, and only included families that at least once reached a

relative abundance of >1% and had relative abundances >0.1% in more than 10% of all samples. We

also included a linear effect of the zero-centered inverse Simpson diversity index (*IVS*) calculated at

the OTU level and standardized to unit variance. This data set was smaller, and we therefore used a

hierarchical model set up with random intercepts and partial pooling per transplanted cell type (*HCT*)

and patient gender (*SEX*), leading to following model (other parameters as before)

438
$$\frac{\Delta \ln(W_i)}{\Delta t} = gr + (1|HCT) + (1|SEX) + \sum_{j=0}^{WBCs} s_{ij}W_j + \sum_{i=0}^{drugs,patient\ age} \beta_i X_i + \sum_{k=0}^{families} \beta_i F_i + \beta_i IVS$$

The prior parameter distributions of the resulting computational graph were informed with the

regression results from the cross-validated analysis of patients without microbiota data (no reuse of

data). When a coefficient was non-zero, it's prior distribution was a permissive, wide Normal

distribution centered on the coefficient (e.g. for the activation of neutrophils from monocytes, we used

Normal(μ=6.4e-02, std=100^2) as prior. A coefficient that yielded a coefficient of zero after the cross-

validated regression of non-microbiota patients would be assigned a regularizing prior centered

around zero (e.g. Normal(μ=0, std=0.01)). Hierarchy was included by assigning regularized random

intercepts for transplanted cell types and patient gender. For example, the reduced baseline rate of

monocytes among cord patients identified in the non-microbiota regression would lead to a prior of

Normal(μ=-3.5e-02, std=1) with partially pooled information across all transplanted cell types via a

shared variance with prior Exponential(lambda=10). The resulting graph of the model is visualized in

Figure S2.

Supplementary Figures

452
453

454
455
456   Figure S1: Eosinophil counts and platelet counts per day relative to HCT; line: mean, shaded: ± std)
457
458
459

460
461   Figure S2: Computational graph of the hierarchical linear model of daily changes in white blood cell counts
462   (*dW*) modeled as a function of graft source (HCT), patient characteristics (sex, age), medications (*Medications,*
463   *platelet transfusions*), homeostatic feedbacks (*Wj*), log-relative abundances of microbiota families (*M*), and
464   ecological diversity (*IVS*).
465
466
467

Day-to-day fold change

N: patients, n: observations

Figure S3: Coefficients from independent, L1-penalized (lasso), 3-fold cross-validated regressions of daily changes in neutrophils, lymphocytes and monocytes. gr: intercept; TCD: T-cell depleted graft (ex-vivo) by CD34+selection; PBSC: peripheral blood stem cells; BM: bone marrow; cord: umbilical cord blood.

473
474 Figure S4: Additional coefficient estimates from the Bayesian regression, see also Figure 3.
475
476
477 Table S1: Patient characteristics. TCD: T-cell depleted graft (*ex-vivo*) by CD34+selection; PBSC: peripheral blood
478 stem cells; BM: bone marrow; cord: umbilical cord blood.
479

Graft sources	%-age
PBSC	39
TCD	38
cord	14
BM	8

Age	years
mean	51
std	14
min	18
max	79

Sex	%-age
Female	59
Male	41

Conditioning regimen	%-age
ABLATIVE	56
REDUCE	29
NONABL	15

480
Supplementary Data

482 The data used in our study is organized in supplementary tables (data-tables.zip), with corresponding
483 filenames (italic):

1. *alphadiversity.csv*: inverse Simpson diversity index per sample
2. *hctmeta.csv*: HCT metadata
3. *infections.csv*: positive blood culture results
4. *microbiome.csv*: relative abundances of major microbial families
5. *sample_id_link.csv*: lookup table to raw microbiota sequencing data[12]
6. *pidmeta.csv*: anonymized patient demographics
7. *treatments.csv*: medication administration/treatment events
8. *wbc.csv*: absolute counts of neutrophils, lymphocytes, monocytes, eosinophils, and platelets