1 **O-glycosylation regulates plant developmental transitions downstream of miR156**

2 Krishna Vasant Mutanwad[1], Alexandra Baekelandt[2,3], Nicole Neumayer[1], Claudia Freitag[1],
3 Isabella Zangl[1], Dirk Inzé[2,3] and Doris Lucyshyn[1*]
4
5 [1]Department of Applied Genetics and Cell Biology, University of Natural Resources and Life
6 Sciences, Vienna, Muthgasse 18 1190 Vienna, Austria.
7 [2]Ghent University, Department of Plant Biotechnology and Bioinformatics, Technologiepark
8 71, 9052 Ghent, Belgium.
9 [3]VIB Center for Plant Systems Biology, Technologiepark 71, 9052 Ghent, Belgium.
10
11 [*]Correspondence: doris.lucyshyn@boku.ac.at

12
13 Keywords: *Arabidopsis thaliana*, O-glycosylation, plant development, phase change, flowering
14 time, SPL
15

SUMMARY

The timing of plant developmental transitions is decisive for reproductive success and thus
tightly regulated. The transition from juvenile to adult vegetative and later to the reproductive
phase is controlled by an endogenous pathway regulated by miR156, targeting the
SQUAMOSA PROMOTER BINDING PROTEIN (SBP/SPL) family of transcription factors.
SPLs regulate a number of developmental processes, such as trichome formation, leaf shape
and floral transition. Such complex regulatory pathways often involve post-translational
modifications (PTMs), integrating a range of internal and external signals. One of these PTMs
is O-glycosylation, the attachment of a single monosaccharide to serine or threonine of nuclear
and cytoplasmic proteins, which is found on a number of very diverse proteins. O-
GlcNAcylation is the most common type of cytosolic O-glycosylation, but in plants also O-
fucose modification occurs. Here we show that mutants defective in the O-fucosyltransferase
SPINDLY (SPY) show accelerated developmental transitions. Genetic analysis shows that
this effect is independent of miR156 levels, but partly dependent on functional SPLs. In a
phenotyping analysis, we found that SPY and SPLs also control leaf growth, as loss of function
mutants showed defects in cell expansion, while SPL9 also regulates cell division in rosette
leaves. Moreover, SPLs interact directly with SPY and are O-glycosylated. Our results show
that O-glycosylation is involved at several steps in the regulation of developmental transitions
and organ growth in *Arabidopsis thaliana*.

INTRODUCTION
During their life cycle, plants undergo developmental transitions by changing from the juvenile
to adult vegetative and later the reproductive phase [1-3]. Each of these irreversible transitions
is marked by specific morphological and developmental changes, and while they are governed
by an internal developmental program, the timing is flexible and can be influenced by the
environment. Both of these phase transitions are controlled by small RNAs, the families of
miR156 and miR172. miR156 is encoded by eight genes, and there is at least partial
redundancy with the closely related miR157, which is encoded by four genes [4]. miR156
directly targets a group of transcription factors within the family of SQUAMOSA PROMOTER
BINDING PROTEINs (SBP/SPL), which then control both transitions by inducing a number of
transcription factors regulating adult leaf traits, floral transition and floral meristem identity, as
well as the expression of miR172 [5-14]. The family of SPLs contains 16 members, but only

well as the expression of miR172 [5-14]. The family of SPLs contains 16 members, but only
10 of them are targeted by miR156. Interestingly, regulation occurs both by transcript cleavage
as well as translation repression, reflecting the high level of complexity in the regulatory
pathways of developmental transitions [4, 15]. Additionally, the decrease in miR156
expression throughout development is strongly influenced by metabolism: the accumulation
of sugars during plant growth leads to a decrease in miR156 and thus an increase in miR172,
which then additionally feeds back negatively on the expression of miR156 [16, 17]. Thereby
as plants grow and build up biomass, the increasing availability of sugars leads to a release
of the repression of SPL transcription factors and thus the developmental switch from the
juvenile to adult and later reproductive phase. At the same time, organ growth is promoted by
a transition from the proliferative phase characterized by cell divisions, to a phase where cell
expansion is driving growth. This is regulated by GROWTH-REGULATING FACTORs (GRFs),
a group of transcription factors regulated by miR396 [18-20]

The timing of transitions is extremely important for plants to ensure reproductive
success. Thus, the transition from the vegetative to the reproductive phase is controlled not
only by the internal developmental program described above, but also by seasonal cues and
environmental conditions, such as day length, ambient temperature or light quality, which are
integrated at several levels. Environmental factors, most importantly day length, are perceived
in the leaves, where they induce the expression of the floral integrator *FLOWERING LOCUS*
T (*FT*). FT is a small, mobile protein, moving through the vasculature to the shoot apical
meristem, where it interacts with the transcription factor FLOWERING LOCUS D (FD) to
induce flowering [21-23]. In the shoot apical meristem, the floral integrators SOC1
(SUPRESSOR OF OVEREXPRESSION OF CONSTANS 1), LFY (LEAFY) and FT regulate
the transcriptional network that underlies flowering time control, as reviewed in [24]. A role for
gibberellins (GAs) in the induction of flowering time has already been suggested in the 1950s
[25, 26], and research on the long-day plant *Arabidopsis thaliana* has shown that GA is
necessary for floral induction under short photoperiods [27]; In long days, its effect is mostly
masked by the much earlier responding photoperiod pathway [28]. Gibberellins have since
been placed at several points within the flowering time regulatory network [11-13, 29-33], with
SOC1, *LFY* and/or *SPLs* being direct targets. The effect of GA on flowering time is mediated
by DELLA proteins, a small family of five transcriptional repressors that are negatively
regulated by GA [34]. In the absence of GA, DELLA proteins bind to a number of transcription
factors, thereby preventing their ability to bind their target genes [35-37]. The miR156-
regulated aging pathway is independent of photoperiod, but is affected by gibberellins, as
SPL15 is directly inhibited by the interaction with the DELLA protein RGA in the absence of
GA [12].

O-glycosylation of nucleocytoplasmic proteins is an abundant post-translational
modification, with diverse targets. In plants, two O-glycosyltransferases are described in this
context, the O-GlcNAc Transferase (OGT) SECRET AGENT (SEC) and the Protein O-
Fucosyltransferase (POFUT) SPINDLY (SPY) [38, 39]. These enzymes use UDP-GlcNAc (N-
acetylglucosamine) and GDP-fucose, respectively, to transfer the single sugar moiety to
serine or threonine residues on their target proteins. Among these targets are the GA
signaling-repressing DELLA proteins as well as a number of other transcriptional regulators
[38-40]. So far, only few examples of regulation by O-glycosylation have been characterized
in plants. While *sec*-mutants show only very subtle phenotypes, *spy*-mutants show a range of
developmental defects, and most of them have been explained by enhanced GA signaling
[38, 39, 41-43]. One of the most prominent features of *spy*-mutants is early flowering. In long
photoperiods, SPY acts together with GIGANTEA (GI) to repress expression of *CONSTANS*

(*CO*) and *FT* [44]. SPY also strongly represses flowering in short photoperiods, which has
been explained by its role in regulating GA signaling [41, 42].
Here, we present a further characterization of the role of SPY in the control of floral
transitions and show an additional function of O-glycosylation in the regulation of the transition
from juvenile to adult vegetative, and then to the reproductive phase. A combination of genetic
and phenotypic analysis suggests a direct regulation of SPL-transcription factors downstream
of miR156 by O-glycosylation. We also observed that leaf growth is strongly affected by SPY
and SPLs at the level of cell expansion. While there was also an effect of SPL9 on cell division,
we did not see an indication for control of cell division by glycosylation. We describe effects of
protein O-glycosylation that are potentially independent of gibberellins, with redundancy
between O-GlcNAc and O-fucose modification, but a much stronger effect of O-fucosylation.
107
RESULTS
The O-fucosyltransferase SPINDLY delays flowering in long and short photoperiods
In order to characterize the effect of O-glycosylation on flowering time regulation, we
studied flowering time in O-glycosylation mutants. As strong *spy* alleles show severely
reduced fertility [39, 41], we used the T-DNA-insertion line *spy-22* (SALK_090582) for all our
experiments. This line showed strongly reduced expression of SPY (Fig.S1A-B) and was
flowering early at 8.6 ±1.4 total rosette leaves (TRL), compared to 13.1 ± 1.1 in the wild-type
Col-0 (Figure 1A-B, Table 1). The early flowering of *spy-22* was complemented by a
SPY::SPY:Flag (*SPY:Flag*) construct (Fig.S1C-D). Recently, modest early flowering of *sec-
5* was reported [45], which we also observed, with a very small difference to the wild-type
(11.8 ± 1.6 TRL in *sec-5* compared to 13.1 ± 1.1 in Col-0; Figure 1A-B, Table 1). Accordingly,
expression of the major floral integrator gene *FT* was up-regulated in *spy-22* but not *sec-5*,
(Figure 1C). When grown in short photoperiods (8 h light / 16 h dark), where *FT* is not
expressed, *spy-22* also displayed strong early flowering with 22.7 ± 0.5 TRL compared to 66.0
± 7.9 in Col-0, which we did not observe in *sec-5*, flowering at 61 ± 5.0 TRL (Figure 1D-E,
Table 1). Early flowering of *spy-22* was only partially suppressed by *ft-10*, resulting in a
phenotype of *ft-10 spy-22* (17.4 ± 1.4 TRL) intermediate between *ft-10* (47.7 ± 2.9 TRL) and
the wild-type Col-0 (14.6 ± 1.3 TRL), while *ft-10 sec-5* (46.2 ± 6.4 TRL) was comparable to *ft-
10* (Figure 1F, Table 1). This confirms previously shown results with mutants in the L*er*-0
background [44], indicating that SPY regulates flowering by regulating *FT* and additionally in
an *FT*-independent manner. This suggests that additional factors independent of the
photoperiod pathway may be affected by SPY, while SEC appears to play a minor role.
Therefore, we also generated crosses with the late flowering Col-0 FRI. This line carries an
active FRIGIDA (FRI) allele introgressed from the ecotype Sf-2, and consequently expresses
high levels of the floral repressor *FLOWERING LOCUS C* (*FLC*) leading to very low levels of
FT expression and requirement of vernalization for floral induction [46, 47]. Late flowering of
Col-0 FRI (74.2 ± 7.8 TRL) was strongly accelerated in *spy-22* FRI (26.0 ± 1.4 TRL, Figure
1G), but it was not affected by *sec-5* (data not shown). Levels of *FLC* expression were
maintained high in that line, and *FT* was repressed (Figure 1H), suggesting that SPY
represses flowering in Col-0 FRI independently and in addition of the vernalization pathway
and FT. Thus, our data indicate that SPY represses floral transition only partly via FT and the
photoperiod pathway. Additionally, SPY is necessary to maintain the vernalization requirement
of Col-0 FRI.
141
142

Mutants in SPY show accelerated transition from juvenile to adult phase

Bypassing of the vernalization requirement of Col-0 FRI has previously been shown in lines expressing an miRNA-resistant stabilized version of SQUAMOSA PROMOTOR-BINDING PROTEIN-LIKE 3 (rSPL3) that lacks an miRNA-binding site [9]. SPL-transcription factors regulate the juvenile to adult vegetative phase transition as well as flowering time; thus, we analyzed also the juvenile to adult vegetative phase change in O-glycosylation mutants. This transition is morphologically marked by the formation of trichomes on the abaxial side of leaves and changes in leaf morphology, an increase in the ratio of length to width of leaf blades, and the onset of serration [48] [49]. Based upon appearance of trichomes on the abaxial side of rosette leaves, we observed that *spy-22* plants consistently transitioned early from the juvenile to the adult stage at 3.5 ± 0.6 juvenile leaves (JL) compared to 5.8 ± 0.6 JL in Col-0, while *sec-5* (5.9 ± 0.7 JL) did not show significant differences compared to the wild-type (Figure 2A, Table 2). In order to dissect the juvenile to adult phase transition from floral transition, we included *ft-10-spy-22* in this analysis. We found that *ft-10* had a delayed juvenile-to-adult transition, with 9.2 ± 1.0 JL, which had been shown before [50]. On the other hand, *ft-10 spy-22* was similar to *spy-22* with 3.5 ± 0.6 JL, even though *ft-10 spy-22* (17.4 $\pm$ 1.3 TRL) showed an extended adult vegetative phase and was flowering considerably later than *spy-22* (7.1 $\pm$ 0.6 TRL) (Figure 2B, Table 2). This suggests that phase transition is regulated independently of daylength-dependent flowering time in *spy-22*. Juvenile-to-adult phase transition is regulated by an internal developmental program, orchestrated by a balance of counteracting miR156 and miR172 [7, 51]. Thus, we analyzed transcript levels of primary miRNA156a, miRNA156b and miRNA156c in 5-day old seedlings, but did not find differences between the wild-type and *spy-22* (Figure 2C).

Levels of miR156 are, among other pathways, regulated by the accumulation of sugars in the leaves of growing plants [16, 17]. Cellular nutrient availability, especially increased sugar levels, enhance global levels of O-glycosylation in animals [52-55]. To test if O-glycosylation might be involved in regulation of miR156 abundance in response to sugar levels in plants, we treated 5-day old Col-0, *spy-22* and *sec-5* seedlings with 50 mM sucrose by transferring them to liquid media with and without added sucrose for 24 h. While we could reproduce the results of a strong decrease in miR156a, miR156b and miR156c in response to sucrose treatment as shown before [16, 17], we did not see any significant differences between *spy-22* and the wild-type after the treatment (Figure 2D).

We then compared expression of mature miR156 and miR172 in juvenile and adult leaves of soil-grown plants. Leaves were harvested for all lines at the timepoint when the wild-type plants started to bolt. We observed balanced levels of miR156 and miR172 between juvenile and adult leaves in Col-0. *spy-22*, which was already flowering, showed a strong shift in the balance towards miR172, with a more than 4-fold increase in adult leaves compared to the juvenile leaves (Figure 2E). *sec-5* showed the same tendency, but at a lower level, while *ft-10* and *ft-10 spy-22*, which were not flowering yet when the samples were taken, showed lower levels of miR172 in adult compared to juvenile leaves (Figure 2E). This observation correlates with the flowering time of these lines (Figure 1D and 2B, Table 1) and their developmental stage at the timepoint of sampling, with miR172 levels being increased in the lines that were already in the reproductive phase and decreased in the lines remaining in the vegetative stage. We did not observe a change in the ratio of miR156 in juvenile leaves compared to adult leaves in *spy-22* or *sec-5* in comparison to the wild-type (Figure 2E). We repeatedly observed lower levels of miR156 in adult leaves of *ft-10*, but not in *ft-10 spy-22*,

again correlating with the fact that *ft-10* is the line with the longest vegetative phase in this
dataset and was far from transition at the timepoint of sampling.

Taken together, we did not see a change in levels of miR156 in *spy-22* or *sec-5*, but we
observed an increase in miR172 levels in *spy-22* and sec-5 as a consequence of accelerated
floral transition. Our results suggest that there is no involvement of O-fucosylation in regulating
the levels or balance of miR156 and miR172 during developmental phase transitions.

SPY regulates phase transitions via SPL transcription factors independently of miR156
To further test for a potential interaction between miR156 and O-glycosylation in the
regulation of developmental transitions we generated *35S::MIRNA156a* lines. Overexpression
of *miRNA156a* in Col-0 led to a strong delay of the juvenile-to-adult phase transition and
extremely delayed flowering, as shown before [7, 9] (Figure 3A-D). We then crossed
35S::MIRNA156a Col-0-with *spy-22* and *sec-5*, and the offspring of these crosses displayed
a strong repression of the miR156 overexpression phenotype. While *35S::MIRNA156a* Col-0
formed 51.8 ± 6.8 leaves, *35S::MIRNA156a spy-22* was flowering at 9.2 ± 1.2 TRL and
35S::MIRNA156a sec-5 at 11.3 ±1.2 TRL (Figure 3 A - C). A comparison of expression levels
of miR156 in juvenile leaves *35S::MIRNA156a spy-22* and *35S::MIRNA156a sec-5* with the
respective parent lines *spy-22, sec-5, 35S::MIRNA156a* Col-0 and the wildtype Col-0 showed
very high levels of mature miR156 in the crossed lines, with miR156 levels at the same level
in *35S::MIRNA156a* Col-0 and *35S::MIRNA156a spy-22*, and weaker overexpression in
35S::MIRNA156a sec-5 (Figure 3D). These data show that the effect of overexpressing
miR156 is strongly suppressed in both *spy-22* and *sec-5* and that the early flowering of these
mutants is independent of miR156 levels.

miR156 targets the family of SPL-transcription factors, which induce juvenile and adult
phase transitions and floral transition [4, 7, 8, 51]. We therefore generated crosses of the O-
glycosylation mutants with *spl9-4 spl15-1*, a line displaying delayed phase change, to test if
SPLs are necessary for early developmental transitions of *spy-22*. We chose this line, as it
has been characterized to show the strongest effect in delaying phase transitions among
combinations of *SPL*-double mutants previously described [8], and both SPL9 as well as
SPL15 directly interact with the DELLA protein RGA [11, 12], which is itself a target of SPY
and SEC [38, 39]. We included *SPL9::rSPL9:GFP* [9] in our analysis, a line carrying an miR
resistant version of SPL9 that consequently displays very early phase transitions. The early
juvenile-to-adult transition of *spy-22* (3.5 ± 0.6 JL) was partly suppressed in the *spy spl9/15*
triple cross (5.4 ± 0.7 JL), and flowering time was also affected (7.8 ± 1.1 TRL in *spy-22,* and
10.2 ± 1.5 TRL in *spy spl9/15*, Figure 4A-B, Table 4). In *sec spl9/15*, the number of juvenile
leaves was slightly increased compared to *sec-5* (5.9 ± 0.7 JL in *sec-5* and 6.8 ± 0.7 JL in *sec*
spl9/15), and *spl9/15* also delayed flowering of *sec-5* in the triple mutant (11.4 ± 1.03 TRL in
sec-5 and 16.3 ± 1.5 TRL in *sec spl9/15*) indicating that SPY and SEC might have common
functions in this pathway (Figure 4A-B, Table 4). We did not see a full suppression of the early
flowering of *spy-22* in *spy spl9/15* to the level of *spl9-4/15-1*. SPLs are encoded by a gene
family with partially overlapping functions [8, 15], and potentially higher order *SPL*-mutants
could lead to a stronger effect. Also, the effect of SPY on the photoperiod pathway and *FT*
expression function independently and in parallel to the aging pathway, accelerating flowering
time in parallel to SPLs.

We also observed additive effects in shoot architecture in the triple crosses (Figure 4B).
While *spy spl9/15* showed a strong loss of apical dominance compared to the wildtype, we
observed the opposite in *sec spl9/15*, with a strong increase in apical dominance. While we

did not follow up or quantify these observations yet, this might be an indication that SPY and
SEC function redundantly as well as independently in different pathways.
We also observed the plastochron length by quantifying the number of leaves formed in
regular intervals in parallel to determining the absolute leaf number (Figure 4C-D).
SPL9::rSPL9:GFP was previously described as having a long plastochron, producing fewer
leaves per day than wildtype, while *spl9/15* has a short plastochron [9], which we also
observed. In this analysis, both *spy-22* and *sec-5* show a slightly longer plastochron compared
to the wildtype, which was suppressed in both triple crosses with *spl9/15* (Figure 4C-D).
Taken together, our results suggest that SPY and SEC both negatively regulate phase
change and flowering independently of miR156, potentially in interaction with SPLs.
246
247 SPY and SPL9/15 are involved leaf size determination
While quantifying flowering time of the distinct mutants, we noticed additional shoot growth
phenotypes that were not observed in the parent lines. In order to analyze these in more detail
and dissect putative additive effects, we determined individual leaf areas as well as rosette
areas and rosette growth rates (Figure 5). At 25 days after sowing (DAS), *spy-22* showed
strongly decreased leaf areas, while *sec-5* had only slightly smaller leaves (Figure 5A, B). The
spl9/15 mutant had produced more leaves than the wildtype at this timepoint, correlating to
the shorter plastochron (Figure 4C and 4D). Additionally, *spl9/15* showed a strong decrease
in the areas of the first 7 leaves, while the leaf areas from leaf 9 onwards were increased
compared to the wild-type Col-0, suggesting a developmental shift. This shift was also
observed in the triple crosses with *spy-22* and *sec-5*. Additionally, we observed a clear additive
effect for the decrease in leaf size, suggesting that O-glycosylation and SPL9 and SPL15
regulate leaf size independently from each other, but the developmental shift seen in *spl9/15*
is epistatic in the triple crosses (Figure 5B). *SPL9::rSPL9:GFP* showed the highest increase
in leaf size (with leaf 1 and 2 showing an increase in leaf area of 121% compared to Col-0)
and a developmental shift compared to the wild-type in the opposite direction of *spl9/15*. Also
in contrary to *spl9/15*, the number of leaves produced in *SPL9::rSPL9:GFP* at 25 DAS was
strongly reduced to 3.9 ± 0.3 leaves, compared to 13.0 ± 1.0 leaves in the wildtype, similar to
the increased plastochron observed in (Figure 4C-D), and [9]. Accordingly, the rosette area
calculated as the sum of the individual leaf areas (mm^2) was drastically reduced in
SPL9::rSPL9:GFP (Figure 5A,C). At 25 DAS, *spy-22* and *spy-22 spl9/15* also showed a strong
reduction in rosette area, while rosettes of *sec-5* were only slightly reduced (Figure 5A-B). The
rosette areas of *spl9/15* and *sec spl9/15* were not significantly affected (Figure 5 A-C).
In a time course analysis, rosette growth correlated to the plastochron length in the
analyzed lines (Figure 4C and D), with *SPL9::rSPL9:GFP* showing the slowest growth (Figure
5C-D). This is mainly resulting from the low number of leaves, as the individual leaf size for
the first developing leaves was increased strongly in this line compared to Col-0 (Figure 5B).
In *spl9/15,* the decrease in leaf size (Figure 5B) was partly compensated by the higher number
of leaves due to shorter plastochron (Figure 4C and D). A positive correlation of plastochron
and leaf size has also previously been shown in [56]. *spy-22* and *spy spl9/15* showed a
considerably slower rosette growth compared to the wildtype, due to decreased leaf size. *sec-*
5 and *sec spl9/15* showed rosette growth rates comparable to the wildtype (Figure 5D).
To determine the cellular effect of the differences in leaf areas observed in O-
glycosylation and *spl* mutants, we determined epidermal cell size and number in the first leaf
pair (L1/2) of all lines. Leaves were harvested at 25 DAS, a timepoint when wild-type Col-0
leaves are considered mature. The leaves were cleared and microscopic drawings of the cells
were used to determine pavement-, guard- and total cell number and individual pavement cell

area (Figure 6A). Total cell number (guard and pavement cells) was strongly increased in
SPL9::rSPL9:GFP (Figure 6B, Fig. S2A-B), but there was no change in pavement cell area
(Figure 6C), suggesting that the increase in leaf area in *rSPL9* mutants is due to increased
cell division. In all other lines, guard- and pavement cell numbers were not significantly
changed compared to the wildtype (Figure 6B, Fig. 6SA-B). When measuring individual
pavement cell areas (PCA), we found a decrease in *spy-22* (-34%) and *spy spl9/15* (-47%)
compared to the wild-type, with a clear additive effect. PCA was also decreased in *spl9/15* (-
30%) and *sec spl9/16* (-23%) mutants, though not statistically significant. The measurements
go in line with the decrease in leaf and rosette area described above, suggesting that the leaf
area decrease in *spy-22*, *spl9/15* and *spy spl9/15* results from additive defects in cell
expansion (Figure 5B). However, we cannot exclude that the observed effects are not due to
slower leaf growth rates on cellular level in these lines. Analysis of mature leaves in *spy-22*
and *spy spl9/15*, taking samples at a later timepoint, could further address this question.
Overall, however, our data suggest that cell division is not affected in these mutants. Given
the increased cell division in *SPL9::rSPL9:GFP* (Figure 6A-C), we cannot exclude that
redundancy among SPLs masks potential effects on cell division in *spl9/15*. Analysis of higher
order mutants would address this question.

The distribution of cell sizes indicates fewer cells in the category of the biggest cells in
spy-22, and especially in *spy spl9/15* (Figure 6D), correlating with the results of the decreases
in leaf area (Figure 5B). The number of cells in the smallest category is increased in these
lines, suggesting a developmental shift at cellular level. Again, however, we cannot exclude
that this is potentially due to cells not yet having reached maturity in these lines.

We then also determined stomatal index (SI) and stomata density (SD), corresponding
to the proportion of stomata per pavement cells and the density of the stomata, respectively.
For SI, we only observed a slight increase in *SPL9::rSPL9:GFP*, suggesting that more stomata
are initiated, while all the other lines did not show significant differences compared with
wildtype (Figure 6F). On the other hand, SD was significantly increased in *spy-22* and *spy
spl9/15* (Figure 6G), likely to result from the decrease in average pavement cell area in these
lines at the timepoint used for phenotypic analysis (Figure 6C).

Overall, our data suggest that SPL9 positively regulates cell division, while we did not
see an effect on cell division in the other analyzed lines. We also observed a reduction in
pavement cell area in *spy-22* and *spy spl9/15,* either due to defects in cell expansion or
delayed growth rates.

SPY O-fucosylates SPL transcription factors
Several SPLs were recently shown to be O-GlcNAc modified, among them also SPL8,
which does not carry a miR156 recognition site [40]. We therefore tested if SPL15, which is
important for the control of flowering time, directly interacts with SPY and can be O-
fucosylated. *35S::FLAG:SPY* was co-infiltrated with *35S::SPL15:HA* and *35S::SPL8:HA* as a
positive control, respectively, in *N. benthamiana* leaves for co-immunoprecipitation. When
analyzing protein expression after infiltration of *N. benthamiana* leaves, we could clearly see
a shift of SPL8 and SPL15 when co-infiltrated with SPY compared to the independent
infiltration, suggesting a modification of SPL8 and SPL15 by SPY (Figure 7A). Interestingly
we also consistently observed a stabilization of SPL8 and SPL15 in the samples co-infiltrated
with SPY compared to single infiltrations (Figure 7A, a representative example of several
biological repeats is shown.) When precipitating Flag:SPY with an anti-Flag antibody, we could
pull down SPL15, indicating protein-protein interaction (Figure 7B). These results indicate that

SPY directly interacts with, and glycosylates SPL15. Together with our data from genetic
analysis, this suggests that SPY inhibits SPL-activity by glycosylation.
333

DISCUSSION
A multitude of different post-translational modifications (PTMs) confer complexity to
the regulation of protein function and stability. They are essential components of signalling
pathways in the course of development, often integrating environmental changes or fine-
tuning crosstalk between different regulatory pathways. O-glycosylation of cytosolic and
nuclear proteins is the modification of a number of very diverse proteins with a single
monosaccharide on serine or threonine residues – which is in contrast to N-glycosylation
events that involve the formation of branched carbohydrate chains of varying composition in
the secretory pathway [57]. Most organisms carry only one type of cytosolic O-glycosylation,
with O-GlcNAcylation being the most common and best described example that is very well
conserved among all kingdoms. Yeast is lacking an O-GlcNAc transferase, but uses instead
O-mannose modification for the same molecular function. Many of the target proteins are
however conserved between yeast and animal cells [58, 59], suggesting that the molecular
function is more conserved than the type of monosaccharide involved. Plants are exceptional
in that they use two different sugars attached to the same protein targets [39]. Currently the
only other organism described to use fucosylation as well as GlcNAc modification is
Toxoplasma gondii, but probably with distinct targets for the two different modifications [59].
In Arabidopsis, only few O-glycosylated targets have been characterized so far. O-
glycosylation of the transcription factors TEOSINTE BRANCHED, CYCLOIDEA AND PCF 14
(TCP14) and TCP15 by SPY increases their stability during cytokinin signalling [60, 61]. SPY
was also implicated in the integration of reactive oxygen species signalling during root
development [62]. Further protein interactions between SPY and GIGANTEA [44], and SPY
and SWI3C [63] have been identified. However, by far the best characterised O-glycosylated
protein in plants is the DELLA protein RGA. The current model suggests that O-GlcNAc and
O-fucose modification have opposite effects on DELLAs during GA signalling, with O-
GlcNAcylation leading to a closed conformation of DELLAs, rendering them less active. On
the other hand, fucosylation leads to an open conformation of RGA, facilitating the interaction
with target transcription factors, thus increasing DELLA activity [38, 39]. However, a number
of open questions remain, even in the context of GA signalling, such as the observation that
the double knockout of SPY and SEC is embryonic lethal even using weaker alleles, while the
single mutants don't show drastic developmental phenotypes [64]. This suggests that there
might be targets where both modifications have the same effect, or even both modifications
are necessary at the same time, though not identified so far. A proteomic study using lectin
weak affinity chromatography (LWAC) with glucosamine-binding wheat germ agglutinin
(WGA) revealed O-GlcNAc modification of many different proteins, among them a number of
transcriptional regulators, including SPLs [40]. SPL9 and SPL15 directly interact with RGA
[11, 12, 32], and this interaction is likely to be affected by glycosylation of RGA as it was shown
for other transcription factors such as BRASSINAZOLE-RESISTANT 1(BZR1), JASMONATE-
ZIM-DOMAIN PROTEIN 1 (JAZ1), PHYTOCHROME INTERACTING FACTOR 3 and 4 (PIF3,
PIF4) [38, 39]. Here, we show that SPL15 additionally directly interacts with and is modified
by SPY, suggesting that there is an additional, potentially independent effect of glycosylation
on SPL15 function. Moreover, we see an additive effect in *spy spl9/15* on leaf size as well as
pavement cell size, indicating that SPLs might additionally be regulated by other factors in the
determination of leaf size.

Our data indicate that the activity of SPLs is stabilized in both O-glycosylation mutants
despite overexpression of *miR156a*, as the late flowering phenotype of *35S::miR156a* is
almost completely suppressed in *spy-22 35S::miR156a* and *sec-5 35S::miR156a*. This is
supported by the genetic analysis using *spl9/15* crosses with *spy-22* and *sec-5*, where the
early flowering of *spy-22* and *sec-5* is partly suppressed, suggesting that functional SPLs are
necessary for the accelerated flowering time in O-glycosylation mutants. It has previously
been shown that miR156 overexpression lines are less sensitive to GA treatment in terms of
flowering time, keeping their extreme late flowering phenotype even when sprayed with GA
[9, 11]. *spy*-mutants are often described as showing constitutive gibberellin signalling, but in
contrast to gibberellin treatment of miR156 overexpression lines, we see a strong suppression
of late flowering in *spy-22 35S::MIR156a* mutants. Moreover, albeit SPY and SEC have
opposite effects in gibberellin signalling [38, 39], *sec-5* as well as *spy-22* suppress the late
flowering of *35S::MIR156a*, and to a lesser extent developmental transitions, suggesting that
the effect of glycosylation on SPLs might be independent of GA signalling and DELLAs, as
well as miR156, with a degree of redundancy between O-GlcNAc and O-fucose.
Phenotyping of leaf and rosette growth as well as cellular analysis of our lines suggest
a role for both SPY and SPL9/SPL15 in leaf and cell size, with a clear additive effect. Again,
we observed a potentially redundant function for SPY and SEC in this respect, although the
phenotypes of *sec-5* were very subtle, and not statistically significant. While we did not see an
effect of SPY on cell division, we observed a strong positive effect on cell division in
SPL9::rSPL9:GFP. On the other hand, total cell number was not significantly decreased in
spl9/15. This may be due to redundancy and potentially higher order *spl* mutants might show
a defect in cell division. However, the results of the leaf and rosette growth analysis might
reflect different plastochron length or rates of growth and/or development rather than organ
size regulation in the analyzed lines. Sampling of leaves at a later timepoint, when the leaves
of all lines are fully developed and have reached maturity, might reveal additional cellular
effects in these mutants and lead to additional conclusions. It will be interesting to see in
further studies, if (1) SPY indeed regulates cell expansion, but not cell division, via SPLs, or if
(2) the differences we see in leaf size are a result of different developmental timing and leaf
growth rates in *spy-22* and *spl9/15* mutants, or (3) if the effect of SPY on cell expansion is
completely independent of SPLs, for instance by interaction with other targets.
Overall, we suggest a model, where O-glycosylation regulates developmental
transitions on multiple levels (Figure 7C). In long photoperiods, SPY suppresses expression
of *FT*, via interaction with GI [44]. Additionally, glycosylation of RGA regulates its interaction
with transcription factors regulating flowering time, such as PIFs [35, 39, 65] and potentially
also SPLs [11, 12, 32]. Our data add an additional level of regulation by direct glycosylation
of SPLs, controlling them independently of miR156, and potentially also independently of
gibberellins.
416
417
418 **MATERIAL AND METHODS**
419
Plant Material and growth conditions
Arabidopsis thaliana ecotype Col-0 of was used as wild type. Lines *spy-22* (SALK_090582),
sec-5 (SALK_034290), *spl9-4 spl15-1* (N67865, [14]), *ft-10* (GK-290E08, [66]), FRI SF-2
(N6209) and pSPL9::GFP-rSPL9 (N9954, [9]), all in Col-0 background, were obtained from
the European Arabidopsis Stock Centre (NASC) (Scholl et al., 2000). Seeds were surface
sterilized with 70% ethanol and transferred ½ Murashige and Skoog medium (2.15 g/L MS

Salts, 0.25 g/L MES, pH 5.7, 1% agar). Seeds were stratified in the dark at 4°C for 48 hours.
Based upon the experiment the seedlings were germinated and grown in either long day (LD,
16 hours light / 8 hours dark) or short day (SD, 8 hours light / 16 hours dark) conditions at
22°C. For studying phase change and flowering time, seedlings were transferred to soil at 5
days after germination. For all phenotyping experiments, transgenic lines and their wild-type
controls were sown in pots filled with soil. After sowing, the seeds were stratified for 3 days at
4°C and grown at 21°C under a 16 h day/8 h night regime for 25 days. Experiments were
performed with transgenic and wild type seeds harvested from plants that were grown in
parallel.
435
Flowering time and phase transition quantification
Flowering time was quantified by counting the total number of rosette leaves (TRL) produced.
Phase transitions were studied by observing the appearance of abaxial trichomes at the lower
side of rosette leaves. Rosette leaves without any abaxial trichomes were grouped as juvenile
leaves (JL) and rosette leaves with abaxial trichomes were considered adult. Arabidopsis
rosettes were harvested and individual leaves were taped to white paper and scanned for
representations. Scanned pictures were edited to black and white using Paint 3D.
443
RNA extraction and qPCR
For expression analysis, seedlings were grown on ½ MS plates. For studying the primary
miRNA expression in response to sucrose treatment, sterilized seeds were germinated in 50
ml ½ MS liquid media with shaking at 140 rpm. At 5 DAG, seedlings were transferred to ½ MS
media supplemented with and without 50 mM sucrose, and seedlings were harvested after 24
h. 200 mg of total plant material was used for total RNA extraction using SV Total Isolation
System (Promega). 1µg of RNA was used for cDNA synthesis with the iScript™ cDNA
Synthesis kit (Bio-Rad). GoTaq® qPCR Master Mix (Promega) was used for quantitative real-
time PCR, primers are listed in Table 5 and data was analyzed with Bio-Rad CFX Manager
and Microsoft Excel for relative quantification using the 2(-ΔΔC(T))-method [67]. For technical
repeats, every sample was done in triplicates, representative results from one of at least two
biological replicates are shown (as given in the figure legends). For studying mature miRNA
expression, rosette leaves were harvested at the point when Col-0 plants were flowering.
Harvested rosette leaves were separated into juvenile and adult leaves, ground in liquid
nitrogen and total RNA from approximately 150 mg of ground leaf material was extracted with
a ReliaPrep miRNA Cell and Tissue Miniprep System kit (Promega). A stem loop pulsed RT-
PCR protocol [8, 68] was used to study the expression of mature miRNA using a Thermo
Scientific RevertAid First Strand cDNA synthesis kit and stem-loop primers for miR156 and
miR172 listed in Table 5, small nuclear RNA snoR101 was used as internal reference. Equal
amounts of each RNA and 1 µM stem loop primer were incubated at 65°C for 5 minutes with
10mM dNTPs. RiboLock RNase Inhibitor and a 5X reaction buffer were added and
amplification was done for 60 cycles with 30 s of 30°C, 30 s of 42°C and 50°C for 1 s
respectively. PCR products were diluted 4x in nuclease free water, qPCR was done with the
GoTaq® qPCR Master Mix (Promega). For technical repeats, every sample was done in
triplicates. Following qPCR, the data was analysed with Bio-Rad CFX Manager and Microsoft
Excel. Representative results from one of three biological replicates are shown.
470
Plasmid construction and generation of transgenic lines
miR156a overexpressing and *SPY::SPY:FLAG* constructs were generated by amplifying the
genomic DNA from Col-0, and for transient overexpression, *SPY*, *SPL8* and *SPL15* were

amplified from cDNA to generate the respective tagged constructs using Q5 high fidelity DNA
polymerase (NEB). The primers (listed in Table 5) contained 5'-overhangs binding to the
linearized, NcoI / XhoI-digested backbone of the cloning vector pENTR™ 4. PCR products
were excised and purified from agarose gel using GeneJET Gel Extraction Kit (Thermo Fisher)
and cloned into Gateway™ pENTR™ 4 by mixing the linearized vector backbone and PCR
product in a 1:1 ratio using Gibson assembly (NEB), before transformation into DH10B electro-
competent *E. coli* cells. Plasmids containing the gene of interest were extracted using
GeneJET Plasmid Miniprep Kit (Thermo Fisher) and confirmed by sequencing.
Plant expression vectors were generated using the above created entry clones and destination
vectors pK7WG2D (Karimi et al., 2002) for *35S::MIR156a*, pEarleyGate202 for
35S::FLAG:SPY, pEarleyGate201 for *35S::HA:SPL8* or *35S::HA:SPL15* and pEarleyGate302
for *SPY::SPY:FLAG* (Earley et al., 2006). Recombination of the entry clones with the
destination vectors was done using Gateway LR Clonase ll enzyme mix. Positive colonies with
the plasmid of interest were selected for spectinomycin (150µg/mL) resistance for miRNA
overexpressing pK7WG2D constructs, and kanamycin (50µg/mL) resistance for
pEarleyGate201, pEarleyGate202 and pEarleyGate302 constructs, respectively on LB
medium. Plasmids carrying the gene of interest were extracted from overnight bacterial culture
using GeneJET Plasmid Miniprep Kit (Thermo Fisher) and confirmed by sequencing. Correct
plasmids were transformed to *Agrobacterium tumefaciens* strain GV3101 (pMP90) before
transformation of plants by floral dipping [69]. *35S:MIR156a* constructs were transformed to
Col-0, then crossed to *spy-22* and *sec-5* backgrounds, and *SPY::SPY:FLAG* was introduced
into *spy-22* background.
496
<u>Rosette growth and leaf series analysis</u>
For the projected rosette measurements over time, pots containing the plants were positioned
in a randomized way for 25 DAS on a platform, named MIRGIS, allowing fully automatic
phenotyping. Pictures were acquired on a daily interval and image processing and analysis
were performed using the ImageJ software (http://rsb.info.nih.gov/ij/). To construct the
projected rosette area and rosette growth curves, the area measurements of similar transgenic
lines plants were compiled. For the leaf series analysis, individual leaves (cotyledons and
rosette leaves) of 25 day-old plants were dissected at the base of the petiole, oriented from
old (cotyledons) to young on a thin layer of 1% w/v agar covering the bottom of a square plate
(22.5 x 22.5 cm) and, if necessary, cut to flatten them. The leaf series were imaged and the
individual leaf areas were measured with the ImageJ software (http://rsb.info.nih.gov/ij/). The
real rosette area was defined as the sum of the individual plant leaves. The growth experiment
was performed in three independent biological repeats containing 8 plants per experiment.
510
<u>Cellular analysis</u>
Following the leaf series analysis at 25 DAS, the first leaf pairs (L1/2; 16 leaves) were cleared
with 100% ethanol, stored in lactic acid and mounted on microscopic slides. The leaves were
photographed with a dark-field binocular microscope and the leaf blade areas were
determined with the ImageJ software. For each line, three leaves with an area closest to the
median and the average were selected for detailed cellular analysis. Abaxial (lower) epidermal
cells (~150 cells) at the tip of the leaves were drawn with a binocular microscope (Leica,
DMLB) fitted with a drawing tube and a differential interference contrast objective. The
microscopic drawings were scanned for digitalization and guard, pavement and total cell
numbers, as well as pavement cell area, stomatal density, stomatal index were calculated with

the ImageJ software, as described in [19]. The cellular analysis experiment was performed in
three independent biological repeats.
523

Transient protein expression in *N. benthamiana*
For transient expression in *N. benthamiana*, *Agrobacterium tumefaciens* strain, GV3101
(pMP90) expressing *35S::FLAG:SPY* and *35S::HA:SPL8* or *-SPL15* constructs were
cultivated overnight using rifampicin (50μg/mL), gentamycin (50μg/mL) and kanamycin
(50μg/mL) selection. Bacterial cells were harvested, washed and resuspended in infiltration
medium (500 mM MES, 20 mM Na_3PO_4, 100 μM acetosyringone, 50 mg/L D-glucose) to
OD600 of 0.5. For co-infiltration, equal amounts of the respective cultures were mixed. After
infiltration on the abaxial side of a leaf of a five-week old *N. benthamiana* plant, the plants
were maintained in the growth chamber for 3 days before harvesting.
533

Protein extraction and co-immunoprecipitation
For protein extraction, 1 g of agro-infiltrated leaves were frozen in liquid nitrogen, ground and
taken up in 50 mM Tris-HCl pH 7.6, 150 mM NaCl, 1% Triton X-100, 1x plant protease inhibitor
cocktail (Sigma) and 20 μM PUGNAc in the ratio 2:1. 100μL extract was stored for further
analysis as input. 50 μL of anti-FLAG beads (Miltenyi Biotec) were added to the remaining
plant extract and the samples were incubated on a rotor at 4°C for 30 minutes. A μMACS
column was placed in the magnetic field of a μMACS separator and washed with above
mentioned 200 μL extraction buffer. Plant extract containing the anti-FLAG beads was added
to the column and 100 μL of the flow-through were collected and saved for further analysis.
The column was rinsed 4 times with 200 μL TBS (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1x
plant protease inhibitor cocktail (Sigma), 20 μM PuGNAc, 1.5 mM DTE) and once with 100 μL
wash buffer (20 mM Tris-HCl pH 7.5). For elution, 20μL 0.1 M glycine, pH 2.3 was loaded onto
to the column and incubated for 3-5 minutes. After adding 60μL 0.1 M glycine the first eluate
was collected and neutralized with 20 μL 0.5 M Tris, pH 8.0. In the next step 20 μL of pre-
heated 95°C hot Laemmli buffer (10% glycerol, 60 mM Tris-HCl pH 6.8, 2% SDS, 0.1%
bromophenol blue, 5% β-mercaptoethanol) was loaded to the column and incubated for 5
minutes. Subsequently, 80 μL of pre-heated 95°C hot Laemmli buffer were added and the
eluate was collected and used for SDS-PAGE analysis and Western blotting.
552

Western Blotting and antibody dilutions
Input samples and eluates from the co-immunoprecipitation from both individual and co-
infiltrated samples were subjected to SDS PAGE, and transferred to a PVDF membrane (Roth
Roti®-PVDF, pore size 0.45 μm) using the wet transfer method in the Mini-Protean Tetra-
System (Bio-Rad). The membranes were washed with PBST (137 mM NaCl, 2.7 mM KCl, 10
mM Na_2HPO_4, 1.8 mM KH_2PO_4, 0.1% (v/v) Tween 80) for 10 minutes, blocked with 3% milk
for 1 hour at room temperature and probed with anti-FLAG M2 monoclonal antibody (mouse,
Sigma F1804, 1:2000 in 3% milk/PBST), or blocked with 5% BSA for 1 hour at room
temperature and probed with anti-HA antibody (rabbit, Cell Signaling Technology 3742S,
1:1000 in 5% BSA/PBST) respectively. After probing at 4°C overnight, the membranes were
washed with PBST before incubation with the respective secondary antibodies, goat anti-
mouse HRP (1:10000) (Dianova 115-035-164), and goat anti-rabbit HRP (1:20000, Agrisera
A S09 602) for 1 hour at room temperature. After washing, Bio-Rad Clarity Western ECL
substrate was used for chemiluminescence detection on a Fusion Solo S (Vilber).
567

568 Data and statistical analysis

We used Excel for analysis of gene expression, and GraphPad Prism 5 and R (*ggplot*2, R Core Team, https://www.r-project.org/) for statistical analysis and generating graphs for quantification of rosette leaves. In the leaf quantification graphs, the mean is shown and error bars represent standard error, n is given in the respective tables. For statistical analysis, one-way ANOVA followed by Tukey's multiple comparison test were done. For the leaf series analysis, statistical tests were performed with the mixed models and plm procedure using SAS Enterprise Guide 7 software as described previously in [70]. An F-test ($p<0.05$) was used to assess equality of variances. For each leaf, pairwise comparisons amongst the transgenic line or between the transgenic lines and the wild type control were performed using a Tukey-Kramer post-hoc analysis ($p < 0.05$). For the statistical analysis of the leaf blade area of L1/2 and the cellular data, ANOVA tests were performed in R (version 3.3.2) (https://www.r-project.org) followed by Tukey's multiple comparison test ($p < 0.05$). For all statistical tests, the transgenic line was included as the fixed factor in the model. All experiments were performed in three biological replicates and the repeat factor was included as a random factor.

ACKNOWLEDGEMENTS
We gratefully acknowledge Marie-Theres Hauser for sharing equipment, Lukas Grinninger for technical support and Julia König for supplying *N. benthamiana* plants. We thank Barbara Korbei, Vinod Kumar and Jürgen Kleine-Vehn for valuable input. This work was supported by the Austrian Academy of Sciences ÖAW (DOC-fellowship to K.V.M. and APART fellowship to D.L.) and the Austrian Science Fund FWF (P-29051). A.B. received funding from the Research Foundation Flanders (FWO) under the project number 3G038719.

AUTHOR CONTRIBUTIONS:
D.L. conceived the project, D.L., K.V.M designed experiments and wrote the manuscript. A.B. designed and performed phenotyping and cellular analysis, K.V.M. performed all other experiments with the help of N.N., C.F. and I.Z.

DECLARATION OF INTERESTS
The authors declare no competing interests.

REFERENCES

1.   Poethig, R.S. (2003). Phase change and the regulation of developmental timing in plants. Science *301*, 334-336.
2.   Baurle, I., and Dean, C. (2006). The timing of developmental transitions in plants. Cell *125*, 655-664.
3.   Huijser, P., and Schmid, M. (2011). The control of developmental phase transitions in plants. Development *138*, 4117-4129.
4.   He, J., Xu, M., Willmann, M.R., McCormick, K., Hu, T., Yang, L., Starker, C.G., Voytas, D.F., Meyers, B.C., and Poethig, R.S. (2018). Threshold-dependent repression of SPL gene expression by miR156/miR157 controls vegetative phase change in Arabidopsis thaliana. PLoS genetics *14*, e1007337.
5.   Fouracre, J.P., and Poethig, R.S. (2016). The role of small RNAs in vegetative shoot development. Current opinion in plant biology *29*, 64-72.
6.   Poethig, R.S. (2009). Small RNAs and developmental timing in plants. Curr Opin Genet Dev *19*, 374-378.

7. Wu, G., Park, M.Y., Conway, S.R., Wang, J.W., Weigel, D., and Poethig, R.S. (2009).
The sequential action of miR156 and miR172 regulates developmental timing in
Arabidopsis. Cell *138*, 750-759.
8. Xu, M., Hu, T., Zhao, J., Park, M.Y., Earley, K.W., Wu, G., Yang, L., and Poethig, R.S.
(2016). Developmental Functions of miR156-Regulated SQUAMOSA PROMOTER
BINDING PROTEIN-LIKE (SPL) Genes in Arabidopsis thaliana. PLoS genetics *12*,
e1006263.
9. Wang, J.W., Czech, B., and Weigel, D. (2009). miR156-regulated SPL transcription
factors define an endogenous flowering pathway in Arabidopsis thaliana. Cell *138*,
738-749.
10. Yang, L., Conway, S.R., and Poethig, R.S. (2011). Vegetative phase change is mediated
by a leaf-derived signal that represses the transcription of miR156. Development
138, 245-249.
11. Yu, S., Galvao, V.C., Zhang, Y.C., Horrer, D., Zhang, T.Q., Hao, Y.H., Feng, Y.Q., Wang,
S., Schmid, M., and Wang, J.W. (2012). Gibberellin regulates the Arabidopsis floral
transition through miR156-targeted SQUAMOSA promoter binding-like transcription
factors. The Plant cell *24*, 3320-3332.
12. Hyun, Y., Richter, R., Vincent, C., Martinez-Gallegos, R., Porri, A., and Coupland, G.
(2016). Multi-layered Regulation of SPL15 and Cooperation with SOC1 Integrate
Endogenous Flowering Pathways at the Arabidopsis Shoot Meristem. Dev Cell *37*,
254-266.
13. Jung, J.H., Ju, Y., Seo, P.J., Lee, J.H., and Park, C.M. (2012). The SOC1-SPL module
integrates photoperiod and gibberellic acid signals to control flowering time in
Arabidopsis. The Plant journal : for cell and molecular biology *69*, 577-588.
14. Schwarz, S., Grande, A.V., Bujdoso, N., Saedler, H., and Huijser, P. (2008). The
microRNA regulated SBP-box genes SPL9 and SPL15 control shoot maturation in
Arabidopsis. Plant Mol Biol *67*, 183-195.
15. Xing, S., Salinas, M., Höhmann, S., Berndtgen, R., and Huijser, P. (2010). miR156-
Targeted and Nontargeted SBP-Box Transcription Factors Act in Concert to Secure
Male Fertility in Arabidopsis. The Plant Cell *22*, 3935-3950.
16. Yang, L., Xu, M., Koo, Y., He, J., and Poethig, R.S. (2013). Sugar promotes vegetative
phase change in Arabidopsis thaliana by repressing the expression of MIR156A and
MIR156C. Elife *2*, e00260.
17. Yu, S., Cao, L., Zhou, C.M., Zhang, T.Q., Lian, H., Sun, Y., Wu, J., Huang, J., Wang, G.,
and Wang, J.W. (2013). Sugar is an endogenous cue for juvenile-to-adult phase
transition in plants. eLife *2*, e00269.
18. Czesnick, H., and Lenhard, M. (2015). Size control in plants--lessons from leaves and
flowers. Cold Spring Harb Perspect Biol *7*, a019190.
19. Andriankaja, M., Dhondt, S., De Bodt, S., Vanhaeren, H., Coppens, F., De Milde, L.,
Muhlenbock, P., Skirycz, A., Gonzalez, N., Beemster, G.T., et al. (2012). Exit from
proliferation during leaf development in Arabidopsis thaliana: a not-so-gradual
process. Dev Cell *22*, 64-78.
20. Debernardi, J.M., Mecchia, M.A., Vercruyssen, L., Smaczniak, C., Kaufmann, K., Inze,
D., Rodriguez, R.E., and Palatnik, J.F. (2014). Post-transcriptional control of GRF
transcription factors by microRNA miR396 and GIF co-activator affects leaf size and
longevity. The Plant journal : for cell and molecular biology *79*, 413-426.

21. Jaeger, K.E., and Wigge, P.A. (2007). FT protein acts as a long-range signal in Arabidopsis. Current biology : CB *17*, 1050-1054.

22. Wigge, P.A., Kim, M.C., Jaeger, K.E., Busch, W., Schmid, M., Lohmann, J.U., and Weigel, D. (2005). Integration of spatial and temporal information during floral induction in Arabidopsis. Science *309*, 1056-1059.

23. Mathieu, J., Warthmann, N., Kuttner, F., and Schmid, M. (2007). Export of FT protein from phloem companion cells is sufficient for floral induction in Arabidopsis. Current biology : CB *17*, 1055-1060.

24. Andres, F., and Coupland, G. (2012). The genetic basis of flowering responses to seasonal cues. Nat Rev Genet *13*, 627-639.

25. Wittwer, S.H., and Bukovac, M.J. (1957). Gibberellin Effects on Temperature and Photoperiodic Requirements for Flowering of Some Plants. Science *126*, 30-31.

26. Brian, P.W. (1958). Role of gibberellin-like hormones in regulation of plant growth & flowering. Nature *181*, 1122-1123.

27. Wilson, R.N., Heckman, J.W., and Somerville, C.R. (1992). Gibberellin Is Required for Flowering in Arabidopsis thaliana under Short Days. Plant Physiol *100*, 403-408.

28. Reeves, P.H., and Coupland, G. (2001). Analysis of flowering time control in Arabidopsis by comparison of double and triple mutants. Plant Physiol *126*, 1085-1091.

29. Moon, J., Suh, S.S., Lee, H., Choi, K.R., Hong, C.B., Paek, N.C., Kim, S.G., and Lee, I. (2003). The SOC1 MADS-box gene integrates vernalization and gibberellin signals for flowering in Arabidopsis. The Plant journal : for cell and molecular biology *35*, 613-623.

30. Achard, P., Herr, A., Baulcombe, D.C., and Harberd, N.P. (2004). Modulation of floral development by a gibberellin-regulated microRNA. Development *131*, 3357-3365.

31. Porri, A., Torti, S., Romera-Branchat, M., and Coupland, G. (2012). Spatially distinct regulatory roles for gibberellins in the promotion of flowering of Arabidopsis under long photoperiods. Development *139*, 2198-2209.

32. Yamaguchi, N., Winter, C.M., Wu, M.F., Kanno, Y., Yamaguchi, A., Seo, M., and Wagner, D. (2014). Gibberellin acts positively then negatively to control onset of flower formation in Arabidopsis. Science *344*, 638-641.

33. Galvao, V.C., Collani, S., Horrer, D., and Schmid, M. (2015). Gibberellic acid signaling is required for ambient temperature-mediated induction of flowering in Arabidopsis thaliana. The Plant journal : for cell and molecular biology *84*, 949-962.

34. Harberd, N.P., Belfield, E., and Yasumura, Y. (2009). The angiosperm gibberellin-GID1-DELLA growth regulatory mechanism: how an "inhibitor of an inhibitor" enables flexible response to fluctuating environments. The Plant cell *21*, 1328-1339.

35. de Lucas, M., Daviere, J.M., Rodriguez-Falcon, M., Pontin, M., Iglesias-Pedraz, J.M., Lorrain, S., Fankhauser, C., Blazquez, M.A., Titarenko, E., and Prat, S. (2008). A molecular framework for light and gibberellin control of cell elongation. Nature *451*, 480-484.

36. Feng, S., Martinez, C., Gusmaroli, G., Wang, Y., Zhou, J., Wang, F., Chen, L., Yu, L., Iglesias-Pedraz, J.M., Kircher, S., et al. (2008). Coordinated regulation of Arabidopsis thaliana development by light and gibberellins. Nature *451*, 475-479.

37. Li, Q.F., Wang, C., Jiang, L., Li, S., Sun, S.S., and He, J.X. (2012). An interaction between BZR1 and DELLAs mediates direct signaling crosstalk between brassinosteroids and gibberellins in Arabidopsis. Science signaling *5*, ra72.

38. Zentella, R., Hu, J., Hsieh, W.P., Matsumoto, P.A., Dawdy, A., Barnhill, B., Oldenhof,
H., Hartweck, L.M., Maitra, S., Thomas, S.G., et al. (2016). O-GlcNAcylation of master
growth repressor DELLA by SECRET AGENT modulates multiple signaling pathways in
Arabidopsis. Genes Dev *30*, 164-176.
39. Zentella, R., Sui, N., Barnhill, B., Hsieh, W.P., Hu, J., Shabanowitz, J., Boyce, M.,
Olszewski, N.E., Zhou, P., Hunt, D.F., et al. (2017). The Arabidopsis O-
fucosyltransferase SPINDLY activates nuclear growth repressor DELLA. Nature
chemical biology.
40. Xu, S.L., Chalkley, R.J., Maynard, J.C., Wang, W., Ni, W., Jiang, X., Shin, K., Cheng, L.,
Savage, D., Huhmer, A.F., et al. (2017). Proteomic analysis reveals O-GlcNAc
modification on proteins with key regulatory functions in Arabidopsis. Proceedings of
the National Academy of Sciences of the United States of America *114*, E1536-E1543.
41. Silverstone, A.L., Tseng, T.-S., Swain, S.M., Dill, A., Jeong, S.Y., Olszewski, N.E., and
Sun, T.-p. (2007). Functional Analysis of SPINDLY in Gibberellin Signaling in
Arabidopsis. Plant Physiology *143*, 987-1000.
42. Swain, S.M., Tseng, T.-s., and Olszewski, N.E. (2001). Altered Expression of SPINDLY
Affects Gibberellin Response and Plant Development. Plant Physiology *126*, 1174-
1185.
43. Tseng, T.-S., Swain, S.M., and Olszewski, N.E. (2001). Ectopic Expression of the
Tetratricopeptide Repeat Domain of SPINDLY Causes Defects in Gibberellin
Response. Plant Physiology *126*, 1250-1258.
44. Tseng, T.-S., Salomé, P.A., McClung, C.R., and Olszewski, N.E. (2004). SPINDLY and
GIGANTEA Interact and Act in Arabidopsis thaliana Pathways Involved in Light
Responses, Flowering, and Rhythms in Cotyledon Movements. The Plant Cell Online
16, 1550-1563.
45. Xing, L., Liu, Y., Xu, S., Xiao, J., Wang, B., Deng, H., Lu, Z., Xu, Y., and Chong, K. (2018).
Arabidopsis O-GlcNAc transferase SEC activates histone methyltransferase ATX1 to
regulate flowering. EMBO J *37*.
46. Clarke, J.H., and Dean, C. (1994). Mapping FRI, a locus controlling flowering time and
vernalization response in Arabidopsis thaliana. Molecular & general genetics : MGG
242, 81-89.
47. Lee, I., and Amasino, R.M. (1995). Effect of Vernalization, Photoperiod, and Light
Quality on the Flowering Phenotype of Arabidopsis Plants Containing the FRIGIDA
Gene. Plant Physiol *108*, 157-162.
48. Yu, S., Lian, H., and Wang, J.W. (2015). Plant developmental transitions: the role of
microRNAs and sugars. Curr Opin Plant Biol *27*, 1-7.
49. Telfer, A., Bollman, K.M., and Poethig, R.S. (1997). Phase change and the regulation
of trichome distribution in Arabidopsis thaliana. Development *124*, 645-654.
50. Willmann, M.R., and Poethig, R.S. (2011). The effect of the floral repressor FLC on
the timing and progression of vegetative phase change in Arabidopsis. Development
138, 677-685.
51. Wu, G., and Poethig, R.S. (2006). Temporal regulation of shoot development in
Arabidopsis thaliana by miR156 and its target SPL3. Development *133*, 3539-3547.
52. Olivier-Van Stichelen, S., Dehennaut, V., Buzy, A., Zachayus, J.-L., Guinez, C., Mir, A.-
M., El Yazidi-Belkoura, I., Copin, M.-C., Boureme, D., Loyaux, D., et al. (2014). O-
GlcNAcylation stabilizes β-catenin through direct competition with phosphorylation
at threonine 41. The FASEB Journal *28*, 3325-3338.

53.   Walgren, J.L., Vincent, T.S., Schey, K.L., and Buse, M.G. (2003). High glucose and
757   insulin promote O-GlcNAc modification of proteins, including alpha-tubulin.
758   American journal of physiology. Endocrinology and metabolism *284*, E424-434.
759   54.   Peng, C., Zhu, Y., Zhang, W., Liao, Q., Chen, Y., Zhao, X., Guo, Q., Shen, P., Zhen, B.,
760   Qian, X., et al. (2017). Regulation of the Hippo-YAP Pathway by Glucose Sensor O-
761   GlcNAcylation. Mol Cell *68*, 591-604 e595.
762   55.   Hart, G.W. (2014). Nutrient Regulation of Cellular Metabolism & Physiology by O-
763   GlcNAcylation. Journal of Biological Chemistry.
764   56.   Wang, J.W., Schwab, R., Czech, B., Mica, E., and Weigel, D. (2008). Dual effects of
765   miR156-targeted SPL genes and CYP78A5/KLUH on plastochron length and organ size
766   in Arabidopsis thaliana. Plant Cell *20*, 1231-1243.
767   57.   Strasser, R. (2016). Plant protein glycosylation. Glycobiology *26*, 926-939.
768   58.   Halim, A., Larsen, I.S., Neubert, P., Joshi, H.J., Petersen, B.L., Vakhrushev, S.Y., Strahl,
769   S., and Clausen, H. (2015). Discovery of a nucleocytoplasmic O-mannose
770   glycoproteome in yeast. Proceedings of the National Academy of Sciences of the
771   United States of America *112*, 15648-15653.
772   59.   Bandini, G., Haserick, J.R., Motari, E., Ouologuem, D.T., Lourido, S., Roos, D.S.,
773   Costello, C.E., Robbins, P.W., and Samuelson, J. (2016). O-fucosylated glycoproteins
774   form assemblies in close proximity to the nuclear pore complexes of Toxoplasma
775   gondii. Proceedings of the National Academy of Sciences of the United States of
776   America *113*, 11567-11572.
777   60.   Steiner, E., Efroni, I., Gopalraj, M., Saathoff, K., Tseng, T.S., Kieffer, M., Eshed, Y.,
778   Olszewski, N., and Weiss, D. (2012). The Arabidopsis O-linked N-acetylglucosamine
779   transferase SPINDLY interacts with class I TCPs to facilitate cytokinin responses in
780   leaves and flowers. The Plant cell *24*, 96-108.
781   61.   Steiner, E., Livne, S., Kobinson-Katz, T., Tal, L., Pri-Tal, O., Mosquna, A., Tarkowska,
782   D., Mueller, B., Tarkowski, P., and Weiss, D. (2016). The Putative O-Linked N-
783   Acetylglucosamine Transferase SPINDLY Inhibits Class I TCP Proteolysis to Promote
784   Sensitivity to Cytokinin. Plant physiology *171*, 1485-1494.
785   62.   Cui, H., Kong, D., Wei, P., Hao, Y., Torii, K.U., Lee, J.S., and Li, J. (2014). SPINDLY,
786   ERECTA, and Its Ligand STOMAGEN Have a Role in Redox-Mediated Cortex
787   Proliferation in the Arabidopsis Root. Molecular plant.
788   63.   Sarnowska, E.A., Rolicka, A.T., Bucior, E., Cwiek, P., Tohge, T., Fernie, A.R., Jikumaru,
789   Y., Kamiya, Y., Franzen, R., Schmelzer, E., et al. (2013). DELLA-interacting SWI3C core
790   subunit of switch/sucrose nonfermenting chromatin remodeling complex modulates
791   gibberellin responses and hormonal cross talk in Arabidopsis. Plant Physiol *163*, 305-
792   317.
793   64.   Hartweck, L.M., Genger, R.K., Grey, W.M., and Olszewski, N.E. (2006). SECRET AGENT
794   and SPINDLY have overlapping roles in the development of Arabidopsis thaliana L.
795   Heyn. Journal of Experimental Botany *57*, 865-875.
796   65.   Kumar, S.V., Lucyshyn, D., Jaeger, K.E., Alos, E., Alvey, E., Harberd, N.P., and Wigge,
797   P.A. (2012). Transcription factor PIF4 controls the thermosensory activation of
798   flowering. Nature *484*, 242-245.
799   66.   Yoo, S.K., Chung, K.S., Kim, J., Lee, J.H., Hong, S.M., Yoo, S.J., Yoo, S.Y., Lee, J.S., and
800   Ahn, J.H. (2005). CONSTANS activates SUPPRESSOR OF OVEREXPRESSION OF
801   CONSTANS 1 through FLOWERING LOCUS T to promote flowering in Arabidopsis.
802   Plant Physiol *139*, 770-778.

803   67.   Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data
804   using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods (San
805   Diego, Calif.) *25*, 402-408.
806   68.   Varkonyi-Gasic, E., Wu, R., Wood, M., Walton, E.F., and Hellens, R.P. (2007).
807   Protocol: a highly sensitive RT-PCR method for detection and quantification of
808   microRNAs. Plant Methods *3*, 12.
809   69.   Clough, S.J., and Bent, A.F. (1998). Floral dip: a simplified method for Agrobacterium-
810   mediated transformation of Arabidopsis thaliana. The Plant journal : for cell and
811   molecular biology *16*, 735-743.
812   70.   Blomme, J., Inze, D., and Gonzalez, N. (2014). The cell-cycle interactome: a source of
813   growth regulators? J Exp Bot *65*, 2715-2730.
814
815

TABLES

Table 1:Rosette leaf numbers for graphs shown in Figure 1

		Rosette leaves (average ± SD)	n
LD	Col-0	13.1 ± 1.1	51
	spy-22 [a]	8.6 ± 1.4	46
	sec-5 [a]	11.8 ± 1.6	49
SD	Col-0	61.0 ± 6.1	33
	spy-22	20.7 ± 3.1	33
	sec-5	60.5 ± 6.7	30
LD		Rosette leaves (average ± SD)	n
	Col-0	14.6 ± 1.3	13
	ft-10	47.6 ± 2.9	9
	spy-22	7.1 ± 0.6	13
	sec-5	14.9 ± 1.5	11
	ft-10 spy-22	17.4 ± 1.3	11
	ft-10 sec -5	46.2 ± 6.4	8
		Rosette leaves (average ± SD)	n
	Col-0	14.7 ± 1.3	13
	spy-22	7.1 ± 0.7	13
	FRI	74.2 ± 7.8	12
	FRI *spy-22*	26.0 ± 1.4	10

average ± SD is given
One-way ANOVA, Tukeys Multiple Comparison Test:
[a] significantly lower than Col-0 (*** p ≤ 0.001)

Table 2: Rosette leaf numbers for graphs shown in Figure 2

		Juvenile leaves	n	Rosette leaves	n
LD	Col-0	5.8 ± 0.6	42	12.6 ± 1.2	56
	spy-22 [a]	3.5 ± 0.6 ***	33	7.8 ± 1.1 ***	45
	sec-5 [a]	5.9 ± 0.7 ns	31	11.4 ± 1.03 ***	45
		Juvenile leaves	n	Rosette leaves	n
	Col-0	5.0 ± 0.6	22	14.6 ± 1.3	13
	spy-22	3.6 ± 0.5	19	7.1 ± 0.6	13
	ft-10	9.2 ± 1.0	20	47.6 ± 2.9	9
	ft-10 spy-22	3.5 ± 0.6	16	17.4 ± 1.3	11

average ± SD is given
One-way ANOVA, Tukeys Multiple Comparison Test:
[a] significantly lower than Col-0 (*** p ≤ 0.001)
[b] significantly less rosette leaves than Col-0 (* p ≤ 0.05)

Table 3: Rosette leaf numbers for graphs shown in Figure 3

	Rosette leaves (average ± SD)	n
Col-0	10 ± 0.9	12
spy-22	8 ± 0.6	10
sec-5	9.8 ± 0.9	11
35S::MIR156a Col-0	51.8 ± 6.8	11
35S::MIR156a spy-22	9 ± 1.2	24
35S::MIR156a sec-5	11.3 ± 1.2	22

average ± SD is given

843 **Table 4: Rosette leaf numbers for graphs shown in Figure 4**
844

		Juvenile leaves	n	Rosette leaves	n
LD	Col-0	5.8 ± 0.6	42	12.6 ± 1.2	56
	spy-22 [d]	3.5 ± 0.6	33	7.8 ± 1.1 [***]	45
	sec-5 [d]	5.9 ± 0.7 [ns]	31	11.4 ± 1.03 [***]	45
	spl9-4/15-1 [e]	8.8 ± 0.6	37	19.6 ± 2.2	49
	spy-22 spl9-4/15-1 [a b]	5.4 ± 0.7	42	10.2 ± 1.5	54
	sec-5 spl9-4/15-1 [a c]	6.8 ± 0.7	46	16.3 ± 1.5	60
	SPL9::rSPL9:GFP	1.3 ± 0.5	37	5.4 ± 1.1	50

855
average ± SD is given
One-way ANOVA, Tukeys Multiple Comparison Test:
[a] significantly more juvenile and rosette leaves than *spy-22* or *sec-5* (*** $p \leq 0.001$),
[b] significantly less juvenile and rosette leaves than *spl9-4/15-1* (*** $p \leq 0.001$)
[c] significantly less juvenile leaves than *spl9-4/15-1* (** $p \leq 0.01$)
[d] significantly less rosette leaves than Col-0 (*** $p \leq 0.001$)
[e] significantly more rosette leaves than Col-0 (*** $p \leq 0.001$)
863
864
865 **Table 5: Distribution of pavement cell circularity and area**
866

Distribution of pavement cell area							
Cell area (x10³ µm²)	Col-0	*spy-22*	*sec-5*	*spl9/15*	*spy-22 spl9/15*	*sec-5 spl9/15*	*rSPL9*
< 4	26,82	32,73	28,96	28,94	38,77	31,49	23,18
4-8	36,59	45,05	38,03	39,90	48,27	39,94	42,88
8-12	22,74	18,77	21,68	23,97	12,35	20,58	21,36
12-16	10,83	3,15	8,74	4,62	0,62	5,84	6,29
16-20	2,49	0,30	2,43	2,57	0,00	2,15	3,81
20-24	0,53	0,00	0,16	0,00	0,00	0,00	1,66
>24	0,00	0,00	0,00	0,00	0,00	0,00	0,83
Sum (%)	100,00	100,00	100,00	100,00	100,00	100,00	100,00

898 **Table 6: Oligonucleotides used in this study**
899

ID	Sequence	Orientation	Purpose
At5g60390 (EF1α)	TGAGCACGCTCTTCTTGCTTTCA	forward	qPCR, reference
	GGTGGTGGCATCCATCTTGTTACA	reverse	
pri-miR156a	CTTCGTTCTCTATGTCTCAATCTCTC	forward	qPCR [16]
	TGATTAAAGGCTAAAGGTCTCCTC	reverse	
pri-miR156b	GTGATAATGAGTGATGACTGATG	forward	qPCR [16]
	GAAAACGTGACCGGGACCGAATCG	reverse	
pri-miR156c	GTGATAATGAGTGATGACTGATG	forward	qPCR [16]
	GAAAACGTGACCGGGACCGAATCG	reverse	
FT	CATTTTATGATACGAGTAACGAACGGTG	forward	qPCR
	CACTCTCATTTTCCTCCCCCTCTC	reverse	
FLC	TCATGTGGGAGCAGAAGCTG	forward	qPCR
	CCGCCGATTTAAGGTGGCTA	reverse	
miR156	GTCGTATCCAGTGCAGGGTCCGAGGTATTCGCACTGGATACGACGTGCTCA	stem-loop for RT	qPCR/miRNA-quantification [8]
miR172	GTCGTATCCAGTGCAGGGTCCGAGGTATTCGCACTGGATACGACATGCAG	stem-loop for RT	qPCR/miRNA-quantification [8]
snoR101	CTTCACAGGTAAGTTCGCTTG	forward	qPCR/miRNA-quantification [8]
	AGCATCAGCAGACCAGTAGTT	reverse	
miR172	CGGCGGAGAATCTTGATGATGC	forward	qPCR/miRNA-quantification [8]
	GTGCAGGGTCCGAGGT	reverse	
miR156	GCGGCGGTGACAGAAGAGAGT	forward	qPCR/miRNA-quantification [8]
	GTGCAGGGTCCGAGGT	reverse	

ID	Sequence	Orientation	Purpose
spy-22	GTTAAACCCTAAGTATCGGAC	forward	Genotyping
	TTGGCATAAGAAAGTGTATC	reverse	
	ATTTTGCCGATTTCGGAAC	T-DNA	
sec-5	CACGCCTGGCTCTTGCTCATCAG	forward	Genotyping
	GCGGATTGCACGATCAGTGTC	reverse	
	ATTTTGCCGATTTCGGAAC	T-DNA	
spl 15-1	TGTTGGTGTCTGAAGTTGCTG	forward	Genotyping
	TCCACCGAGTCTTCTTCACTC	reverse	
	TGGTTCACGTAGTGGGCCATCG	T-DNA	
spl 9-4	TGGTTCCTCCACTGAGTCATC	forward	Genotyping
	GCTCATTATGACCAGCGAGTC	reverse	
	TAGCATCTGAATTTCATAACCAATCTCGATACAC	T-DNA	
ft-10	TAAGCTCAATGATATTCCCGTACA	forward	Genotyping
	CAGGTTCAAAACAAGCCAAGA	T-DNA	
	CCCATTTGGACGTGAATGTAGACAC	reverse	
Col-0 FRI	ATGAGATTGCCGGTGCTTT	forward	Genotyping
	TGGTCGATGATGTCAACAAAA	reverse	

ID	Sequence	Orientation	Purpose
miRNA156	TACAAAAAAGCAGGCTCCACTCTTTGTCTTCTCCAGTTAAAAC	forward	Cloning 35::miRNA156a
	GCTGGGTCTAGATATCTCGACAAGAGAGACAGAGAAAG	reverse	
SPY	TACAAAAAAGCAGGCTCCACAGACAAGAGGGTTTTATAACTC	forward	Cloning SPY-promotor + Flag
	TGTCGTCATCGTCTTTGTAGTCCATTTTTTTTGTAACTAAAATCTTG	reverse	
	CTACAAAGACGATGACGACAAGGTGGGACTGGAAGATGATAC	forward	Cloning Flag + SPY-ORF
	GCTGGGTCTAGATATCTCGACTAGCTAGTGGAGTCCATTC	reverse	
SPL8	TACAAAAAAGCAGGCTCCACCTCCTCCACCCCTTCCG	forward	Cloning 35S::HA:SPL8
	GCTGGGTCTAGATATCTCGATCCGCTGGAGAAAAACATTG	reverse	
SPL15	TACAAAAAAGCAGGCTCCACCACCATGGAGTTGTTAATGTGTTC	forward	Cloning 35S::HA:SPL8
	GCTGGGTCTAGATATCTCGATCAAAGAGACCAATTGAAATG	reverse	

904 **FIGURES**
905
906 Figure 1
907

908
909
910
Figure 1. SPY suppresses flowering in long and short photoperiods.
(**A**) Representative pictures of wildtype Col-0, *spy-22* and *sec-5* grown in LD conditions, and
(**B**) total rosette leaf numbers of these lines. (**C**) Relative expression levels of *FT* in 10-day old
seedlings of *spy-22* and *sec-5* grown in LD conditions. (**D**) Representative pictures of wildtype
Col-0, *spy-22* and *sec-5* grown in SD conditions, and (**E**) total rosette leaf numbers of these
lines. (**F**) Total rosette leaf numbers of wildtype Col-0, *spy-22, sec-5, ft-10, ft-10 spy-22* and
ft-10 sec-5 grown in LD conditions. (**G**) Total rosette leaf numbers of wildtype Col-0, *spy-22,*
Col-0 FRI and FRI *spy-22* grown in LD conditions, and (**H**) relative expression levels of *FLC*
and *FT* in 10-day old seedlings of Col-0, *spy-22,* Col-0 FRI and FRI *spy-22* grown in LD
conditions.
For expression analysis, an average of three biological repeats ± SEM is shown, n > 20. For
comparison of rosette leaf numbers, one-way ANOVA with Tukey's multiple comparison was
done, n is given in Table 1 (*** p ≤ 0.001).

Figure 2
925

926
927

Figure 2. SPY regulates juvenile to adult phase transitions
(**A**) Leaf scans (left panel) and juvenile and total leaf numbers (right panel) of wildtype Col-0,
spy-22 and *sec-5* grown in LD conditions. One-way ANOVA with Tukey's multiple comparison
was done, n is given in Table 2. (*** p ≤ 0.001, ns: not significant). (**B**) Juvenile and total leaf
numbers of wildtype Col-0, *spy-22*, *ft-10* and *ft-10 spy-22* grown in LD conditions. (**C**) Relative
expression levels of primary *miR156a*, *-b* and *-c* in Col-0 and *spy-22* in 5-day old seedlings,
an average of three biological repeats +/- SEM is shown, n > 20. (**D**) Relative expression levels
of primary *miR156a*, *-b*, and *-c* in Col-0 and *spy-22* in 5-day old seedlings treated with 50 mM
sucrose for 24 h. An average of two biological repeats is shown, n > 20. (**E**) Relative
expression of mature *miR156* and *miR172* in juvenile and adult leaves of Col-0, *spy-22*, *sec-
5*, *ft-10* and *ft-10 spy-22*. Representative results of one out of three biological repeats (average
± SD) are shown.

Figure 3

Figure 3. Overexpression phenotypes of miR156 are suppressed in *spy-22* and *sec-5*
(A, B) Representative pictures of 4-week-old (A) and 7-week-old (B) plants of wildtype Col-0, *spy-22* and *sec-5* crossed with *35S::miRNA156a.* **(C)** Total rosette leaves of *35S::miRNA156a* Col-0*, 35S::miRNA156a spy-22* and *35S::miRNA156a sec-5*, n is given in Table 3. **(D)** Relative expression levels of mature *miR156* in juvenile leaves of wildtype Col-0, *spy-22*, *sec-5, 35S::miRNA156a* Col-0*, 35S::miRNA156a spy-22* and *35S::miRNA156a sec-5*. One biological repeat (average ± SD) is shown.

973   Figure 4
974

975
976
Figure 4: O-glycosylation represses phase transitions partly via SPL9 and SPL15
(**A**) Juvenile and total rosette leaf numbers of plant lines indicated in the graph, grown in LD
conditions. Averages ± SEM are shown. For comparison of rosette leaf numbers, one-way
ANOVA with Tukey's multiple comparison was done, n is given in Table 4 ([a] significantly more
juvenile and rosette leaves than *spy-22* or *sec-5* (*** p ≤ 0.001), [b] significantly fewer juvenile
and rosette leaves than *spl9-4/15-1* (*** p ≤ 0.001) [c] significantly fewer juvenile leaves than
spl9-4/15-1 (** p ≤ 0.01). Full statistical comparison is given in Table 4.
(**B**) Representative pictures of 6-week-old plants with leaf numbers shown in (A). **(C,D)**
Rosette leaf numbers over time of plants quantified in (A), averages ± SD are given. **(E)** Leaf
scans of total rosette leaves of the indicated lines.
987

Figure 5

Figure 5. SPY and SPL9/SPL15 regulate leaf growth
(**A**) Representative pictures of 25-day-old rosettes of plants used for leaf and rosette area measurements. (**B**) Leaf areas of lines indicated in the graph. Averages ± SEM of three biological repeats using 8 plants per line for each repeat are shown. (**C**) Real rosette area, calculated from the leaf area measurements depicted in (B). For statistical analysis, one-way ANOVA with Tukey's multiple comparison was done, significant differences to the wildtype Col-0 are shown (*** $p \leq 0.001$, * $p \leq 0.05$). (**D**) growth rates of rosette areas (mm^2) of the indicated lines.

999 Figure 6
1000

1001
1002

Figure 6. Cellular analysis of *spy-22, sec-5, spl9/15* **triple mutants and**
*SPL9::rSPL9:GFP***.**

(A) Representative drawings of cell margins of the indicated lines, as used for the analysis of
cell numbers, areas, circularity, SI and SD. **(B)** Total epidermal cell number and **(C)** pavement
cell area of lines indicated in the graph. **(D)** Distribution of pavement cell area of the indicated
lines. **(E)** Stomatal Index (SI) and **(F)** stomata density (SD) of the indicated lines. Averages ±
SEM of three biological repeats using 3 plants per line for each repeat are shown (n = 9). For
statistical analysis, one-way ANOVA with Tukey's multiple comparison was done, significant
differences to the wildtype Col-0 are shown (*** $p \leq 0.001$, * $p \leq 0.05$).

1022 Figure 7

1023
1024

Figure 7. SPY interacts with SPLs.
(**A**) Western Blot of protein extracts from transient expression in *N. benthamiana*,
35S::SPY:Flag, *35S::SPL8:HA*, *35S::SPL15:HA* single infiltrations, as well as *35S::SPY:Flag*,
35S::SPL8:HA and *35S::SPY:Flag*, *35S::SPL15:HA* co-infiltration is shown. Protein extracts
were blotted and probed with anti HA-antibody to visualize SPL8 and SPL15. (**B**) Samples
after co-infiltration of transiently expressed *35S::SPY:Flag* and *35S::SPL15:HA* were used for
immunoprecipitation of SPY using an anti-Flag antibody, co-immunoprecipitation of SPL15 is
shown with an anti-HA antibody. (**C**) Current working model suggesting a role for O-
glycosylation at several levels in the regulation of developmental transitions in *Arabidopsis*
thaliana.

1035
1036
1037

1038   Supplemental Figure 1.

1039
Supplemental Figure 1. Description of mutant lines *spy-22* and *sec-5*
(**A**) Structure of *SPY* and *SEC*, 5'- and 3'-UTR are shown in light grey and exons in dark grey
boxes, the lines represent introns. The position of the T-DNA insertion in *spy-22*
(SALK_090582) and *sec-5* (SALK_034290) are given. (**B**) Relative expression levels of *SPY*
and *SEC* in *spy-22* and *sec-5 seedlings,* an average of three biological repeats ± SEM is
presented, n > 20. (**C**) Juvenile and total rosette leaf numbers of wildtype Col-0, *spy-22* and
spy-22 SPY::SPY:Flag (*SPY:Flag*) grown in LD conditions. (**D**) Representative pictures of
wildtype Col-0, *spy-22,* and *spy-22 SPY::SPY:Flag* (*SPY:Flag*) grown in LD conditions.

1067 Supplemental Figure 2.

1068
1069 **Supplemental Figure 2.**

1070 (**A**) Pavement and (**B**) guard cell numbers of lines indicated in the graph. Averages ± SEM of
1071 three biological repeats using 3 plants per line for each repeat are shown (n = 9). For statistical
1072 analysis, one-way ANOVA with Tukey's multiple comparison was done, significant differences
1073 to the wildtype Col-0 are shown (*** $p \leq 0.001$).

1103   Supplemental Figure 3

1104
1105
1106   **Supplemental Figure 3. Full membranes of blots shown in Figure 7.**
1107