An *In-Silico* Investigation of Menthol Metabolism

Taweetham Limpanuparb*, Wanutcha Lorpaiboon and Kridtin Chinsukserm

Science Division, Mahidol University International College, Mahidol University, Nakhon Pathom 73170,

Thailand

* taweetham.lim@mahidol.edu

ABSTRACT

Prevalence of mentholated products for consumption has brought great importance to studies on menthol's metabolic pathways to ensure safety, design more potent derivatives, and identify therapeutic benefits. Proposed pathways of (-)-menthol metabolism based on metabolites found experimentally in previous works by Yamaguchi, Caldwell & Farmer, Madyastha & Srivatsan and Hiki et al. were not in agreement. This *in silico* approach is based on the three *in vivo* studies and aims to resolve the discrepancies. Reactions in the pathways are conjugation with glucuronic acid/sulfate, oxidation to alcohol, aldehyde & carboxylic acid, and formation of a four-membered/five-membered ring. Gas-phase structures, standard Gibbs energies and SMD solvation energies at B3LYP/6-311++G(d,p) level were obtained for 102 compounds in the pathways. This study provides a more complete picture of menthol metabolism by combining information from three experimental studies and filling missing links in previously published pathways.

Introduction

(-)-Menthol or 1*S*,3*R*,4*S*-menthol is a naturally occurring compound found in plants of the *Mentha* genus commonly known as mint. It is the most abundant in nature among the 8 possible stereoisomers, and make up at least 50% of peppermint (*Mentha piperita*) oil and 70-80% of corn mint (*Mentha arvensis*) oil [1]. (-)-Menthol, commonly referred to as menthol, has characteristic minty smell and flavor and exerts a cooling sensation when applied to the skin and mucosal membranes [2]. Other isomers differ slightly in odor and physical characteristics and do not possess the cooling action [3, 4].

Menthol finds a wide range of applications from personal care products, medications, and confectionery to pesticides and cigarettes. The popularity of the compound as a flavoring agent ranks third most important after vanilla and citrus [5], and the annual production of menthol in India alone is in excess of 200 thousand metric tons [6]. Mentholated products can be purchased as prescribed or over-the-counter medications as alleviators of common cold and respiratory conditions [7], inhibitors of growth of foodborne pathogens [8], and analgesics [9].

Considering its wide range of applications, mechanisms of action of menthol were relatively unknown until recently. The cooling sensation is a result of the activation of transient receptor potential melastatin-8 (TRPM8), an ion channel selective to temperature, voltage, and menthol [10]. Experimental evidence also show that (-)-menthol can selectively activate κ-opioid receptors in mice and, as a result, leads to its analgesic properties [9].

In addition, chemical derivatives of menthol with enhanced activity have been successfully synthesized [11].

However, health effects of mentholated cigarettes is of great concern, not only because the improved taste may

facilitate initiation or inhibit quitting but also because metabolism of menthol via this route of administration has not

been well studied [12, 13].

A few studies have been conducted on toxicological effects of menthol which supports the generally accepted

belief that it is safe and nontoxic. No signs of toxicity were observed in rats exposed to continuous doses of up to

800 mg/kg/day for 28 days [5], and chronic exposure to high concentrations of menthol vapor was not reported to

have toxic effects in rats [14]. *In vitro* studies on various animal tissues report deterioration of biological

membranes at concentrations 0.32-0.76 mM [15]. The recommended daily intake for humans of 0-0.2 mg/kg

proposed by the WHO [16] is not supported by any toxicological data but was set to err on the side of safety

knowing that higher doses taken may not have produced adverse side effects.

To the best of our knowledge, three *in vivo* studies by Yamaguchi, Caldwell & Farmer [17], Madyastha & Srivatsan

[18] and Hiki et al. [19] have identified metabolites of menthol in humans and animals. Metabolites were identified

by GC/MS from the urine and bile of rats treated with oral doses of 500 [17] and 800 [18] mg menthol/kg body

weight. Over the course of 48 hours, a majority of the doses were excreted in the urine and feces. A more recent

randomized, double-blind, placebo-controlled study in human by Hiki et al. [19] was conducted by directly spraying

0.8% (-)-menthol solution at escalating doses of 10-40 mL onto the gastric mucosa. Blood and urine of the

participants were sampled over a 24-hour period and analyzed with GC/MS for menthol metabolites. In total, 72

metabolites were identified or proposed in this human study alone, compared to 9 and 18 metabolites in the

previous two experiments. (See S3 File for the full list of metabolites in the first worksheet in the spreadsheet file.)

In vitro investigation of metabolism in human liver microsomes revealed that the same key reactions in the

metabolic pathway in rats occur in the microsomes [20, 21].

FIGURE 1 Metabolic pathway of menthol in rats and in human, an adaptation from Yamaguchi, Caldwell, & Farmer [17], Madyastha & Sirvastan

[18] and Hiki et al. [19]. Red, Green, and Blue numbers indicate that menthol metabolites were found in both rats and human, only in rats, and

only in human respectively. Gray and Black numbers indicate menthol metabolites proposed by previous experiments and by this paper

respectively. Arrows to the right and arrows upward indicate oxidation reactions $+ \frac{1}{2} O_2$ and $- H_2$ respectively. Downward arrows indicate

conjugation with sulfate. Dashed arrows indicate reactions of four-membered ring metabolites. Diagonal arrows toward top left indicate

dehydration reaction. Main pathways are shown on the left and pathways containing glucuronide metabolites with similar possible connections

are shown on the right. Lists of compounds and reactions are provided in Table 1 and Table 2, respectively.

This *in silico* investigation is based on the metabolites identified experimentally by the three *in vivo* studies [17-19].

We aim to resolve discrepancies and missing links found in these three studies by proposing more complete

pathways in Figure 1 where all 73 experimentally identified metabolites, 5 previously proposed intermediates and

24 newly proposed intermediates are included. Possible reactions involved in the pathways are conjugation with

glucuronic acid/sulfate, oxidation to alcohol, aldehyde & carboxylic acid, and formation of a four/five-membered

ring at position 3, 7, 8, 9 and 10 of the parent compound. In this paper, we calculated Gibbs energies of reactions

and associated them with the type, the position and the step of reaction in the pathways.

Computational Details

Gas-phase structures were calculated at the B3LYP/6-311++G(d,p) level of theory and were confirmed to be at

minimum energy on the electronic potential energy surfaces by frequency calculations. The solvation energies in

water of the gas-phase structures were calculated with the SMD model [22] through single-point energy

calculations. The calculation of Gibbs energies in solution phase is the same as in our previous work [23, 24]

where there is a correction for the standard state of liquid water (~ 55 mol/L) [25-28]. All quantum chemical

calculations were performed using the Q-Chem 5.1 program package [29]. (Shell script, spreadsheet templates,

and Mathematica [30] notebook used were modified from our previous work [23, 24]. All output files and other

associated codes to obtain the standard Gibbs energies of the reaction are provided in S1 File and S2 File

respectively.) The abbreviated names for each of the metabolites in this study are as in Table 1. For simplicity,

we based the naming system of menthol metabolites on their five substitutable positions, namely position 3, 7, 8, 9

and 10. A menthol metabolite is referred to as a five-character sequence named according to its substituted

functional groups at these positions with the abbreviation explained below in Table 1.

All DFT calculations were completed with no imaginary frequencies, showing that each of the structures obtained

from gas-phase calculations were minima on the potential energy surfaces. The lowest energy structure of (-)-

menthol is a chair conformer of hexane where all three substituent groups are in equatorial positions as shown in

Table 1. This is consistent with previous computational result at B3LYP/6-31G(d,p) level [31]. Benchmark

calculations were also performed at MP2/6-311++G(d,p) level for metabolites along the most likely pathways in

Figure 2. Reaction energies obtained from MP2 and B3LYP are in good agreement. (Coefficient of determination

r^2=0.9999 and mean absolute error (MAE) of 2.25 kcal/mol. See the third worksheet of the Excel file in S3 File for

details.) These results are in line with earlier studies and confirm that B3LYP/6-311++G** yields acceptable

results at a reasonable computational cost [23, 32].

Results

The present study has combined the different published metabolic pathways of menthol and offers the relative

stabilities of each metabolite based on thermodynamic calculations for each step involved as reported in Figures 1

to 4. Reaction energies were computed with the relevant additional reagents (oxygen, sulfate group, hydronium

ion and glucuronic acid) and product (water and hydrogen) added to the scheme. They may not be the actual

compound in the reactions but they serve as simple reference points for the thermodynamic calculations for

reactions of interest. The full list of compounds and reaction energies is in the first worksheet of the Excel file in

S3 File.

TABLE 1 Abbreviations for the nomenclature of menthol metabolites referred to by the present study and a
list of 102 compounds in this study grouped by molecular formula.
104

Position / Group	3	7	8	9	10	
Original form	o	e	e	e	e	
Alkan**e**	-	e	e	e	e	
Alco**ho**l	o	o	o	<u>o</u>	o	
Aldeh**y**de	y	y	-	y	y	
Carbo**x**ylic acid	-	x	-	x	-	
Dehydration	-4D for four-membered ring formation at positions 3 and 8					
Aldol reaction	-5A for four-membered ring formation at positions 3 and 9					
Glucuronic acid	O	<u>O</u>,X	-	<u>O,X,Y</u>	-	
Sulfate group	s	-	-	-	-	

- An underlined indicates that there are *R* and *S* stereoisomers due to the substitution.
- Substitution at position 9 leads to a new chiral center if it is not the same as 10.
- Substitution at position 10 is forced to have lower or the same oxidation state for the carbon atom when compared to position 9.
- Dashes are where substitution with the functional group at that respective position cannot occur

105
106

formula: $C_{10}H_{20}O$ (molar mass 156.26, total 1 compound)

1 oeeee   (in chair form)

formula: $C_{10}H_{20}O_2$ (molar mass 172.26, total 4 compounds)

2 ooeee   **3** oeoee   **4** oeeoe-*R*   **5** oeeoe-*S*

formula: $C_{10}H_{20}O_3$ (molar mass 188.26, total 6 compounds)

6 oooee   **7** ooeoe-*R*   **8** ooeoe-*S*   **9** oeooe-*R*   **10** oeooe-*S*   **11** oeeoo

formula: $C_{10}H_{18}O_2$ (molar mass 170.25, total 4 compounds)

12 oyeee-*R*   **13** oyeee-*S*   **14** oeeye-*R*   **15** oeeye-*S*

formula: $C_{10}H_{18}O_3$ (molar mass 186.25, total 28 compounds)

16 oxeee-*R*   **17** oxeee-*S*   **18** oeexe-*R*   **19** oeexe-*S*   **20** oyoee-*R*   **21** oyoee-*S*

22 oyeoe-*RR*   **23** oyeoe-*RS*   **24** oyeoe-*SR*   **25** oyeoe-*SS*

26 ooeye-*R*   **27** ooeye-*S*   **28** oeoye-*R*   **29** oeoye-*S*   **30** oeeyo-*R*   **31** oeeyo-*S*

32 ooeye-5A-*RR*   **33** ooeye-5A-*RS*   **34** ooeye-5A-*SR*   **35** ooeye-5A-*SS*

36 oeoye-5A-*RR*   **37** oeoye-5A-*RS*   **38** oeoye-5A-*SR*   **39** oeoye-5A-*SS*

40 oeeyo-5A-*RR*   **41** oeeyo-5A-*RS*   **42** oeeyo-5A-*SR*   **43** oeeyo-5A-*SS*

formula: $C_{10}H_{18}O_4$ (molar mass 202.25, total 2 compounds)

44 oxoee-*R*   **45** oxoee-*S*

formula: $C_{10}H_{16}O_3$ (molar mass 184.23, total 5 compounds)

46 oyeye-*RR*   **47** oyeye-*RS*   **48** oyeye-*SR*   **49** oyeye-*SS*   **50** oeeyy

formula: $C_{10}H_{18}O$ (molar mass 154.25, total 1 compound)

51 oeoee-4D

formula: $C_{10}H_{18}O_2$ (molar mass 170.25, total 1 compound)

52 oooee-4D

formula: $C_{10}H_{16}O_2$ (molar mass 168.23, total 2 compounds)

53 oyoee-4D-*R*   **54** oyoee-4D-*S*

formula: $C_{10}H_{16}O_3$ (molar mass 184.23, total 2 compounds)

55 oxoee-4D-*R*   **56** oxoee-4D-*S*

formula: $C_{10}H_{20}O_4S$ (molar mass 236.33, total 1 compound)

57 seeee

formula: $C_{10}H_{20}O_5S$ (molar mass 252.33, total 3 compounds)

58 soeee   **59** seeoe-*R*   **60** seeoe-*S*

formula: $C_{10}H_{20}O_6S$ (molar mass 268.33, total 2 compounds)

61 soeoe-*R*   **62** soeoe-*S*

formula: $C_{16}H_{28}O_7$ (molar mass 332.39, total 1 compound)

63 Oeeee

formula: $C_{16}H_{28}O_8$ (molar mass 348.39, total 8 compounds)

64 Ooeee   **65** oOeee   **66** Oeoee   **67** oeOee   **68** Oeeoe-*R*   **69** Oeeoe-*S*   **70** oeeOe-*R*   **71** oeeOe-*S*

formula: C$_{16}$H$_{28}$O$_9$ (molar mass 364.39, total 6 compounds)

72 Ooeoe-*R* **73** Ooeoe-*S* **74** oOeoe-*R* **75** oOeoe-*S* **76** ooeOe-*R* **77** ooeOe-*S*

formula: C$_{16}$H$_{26}$O$_8$ (molar mass 346.37, total 4 compounds)

78 Oyeee-*R* **79** Oyeee-*S* **80** Oeeye-*R* **81** Oeeye-*S*

formula: C$_{16}$H$_{26}$O$_9$ (molar mass 362.37, total 16 compounds)

82 Oxeee-*R* **83** Oxeee-*S* **84** Oeexe-*R* **85** Oeexe-*S* **86** oXeee-*R* **87** oXeee-*S* **88** oeeXe-*R* **89** oeeXe-*S*

90 oOeye-5A-*RR* **91** oOeye-5A-*RS* **92** oOeye-5A-*SR* **93** oOeye-5A-*SS* **94** ooeYe-5A-*RR* **95** ooeYe-5A-*RS* **96** ooeYe-5A-*SR* **97** ooeYe-5A-*SS*

formula: C$_{16}$H$_{24}$O$_9$ (molar mass 360.36, total 5 compounds)

98 Oyeye-*RR* **99** Oyeye-*RS* **100** Oyeye-*SR* **101** Oyeye-*SS* **102** Oeeyy

Notes

- Compounds are highlighted in yellow when their *RS* designator changes from their parent compounds.
- G stands for a glucuronyl group:

108 **TABLE 2** Representative of oxidation reactions to alcohol, aldehyde and carboxylic acid, ring formation
109 (dehydration reaction and aldol reaction) and conjugation with glucuronic acid/sulfate group.

Abbreviation/explanation	Example
o1 for oxidation from alkane to alcohol	**1** oeeee + 1/2 O$_2$ → **2** ooeee
o2 for oxidation from aldehyde to carboxylic acid	**12** oyeee-*R* + 1/2 O$_2$ → **16** oxeee-*R*
o3 for oxidation form alcohol to aldehyde	**6** oooee → H$_2$ + **20** oyoee-*R*
4D for dehydration (four-membered ring formation)	**3** oeoee → **51** oeoee-4D + H$_2$O
5A for aldol reaction (five-membered ring formation)	**30** oeeyo-*R* → **40** oeeyo-5A-*RR*
g for conjugation with glucuronic acid	**1** oeeee + GOH → **63** Oeeee + H$_2$O
s for conjugation with sulfate	**1** oeeee + 2H$_3$O$^+$ + SO$_4^{2-}$ → **57** seeee + 3H$_2$O

110
FIGURE 2 Average Gibbs energies of reaction in the gas phase and in solution for various reactions at the B3LYP/6-

311++G(d,p) level of theory: (o1) oxidation from alkane to alcohol, (o2) oxidation from aldehyde to carboxylic acid, (o3) oxidation

from alcohol to aldehyde, (4D) dehydration or four-membered ring formation, (5A) aldol reaction or five-membered ring

formation, (g) conjugation with glucuronic acid and (s) conjugation with sulfate at five different positions of (-)-menthol.

Representative reactions of each type are shown in Table 2.

FIGURE 3 Relative stability of 102 (-)-menthol metabolite compounds at the B3LYP/6-311++G(d,p) level of theory.

FIGURE 4 Lowest Gibbs energy diagram for the metabolism of (-) menthol at the B3LYP/6-311++G(d,p) level of theory.

Discussion
118
119

The standard Gibbs energies for each reaction in both gas and solution phase listed in the first worksheet of

the spreadsheet of S3 File are summarized in Figure 2. With the exception of sulfation (s-3), reaction

energies in gas phase are lower in magnitude but have the same sign as their equivalents in solution phase.

Oxidation reactions (o1 and o2, addition of $\frac{1}{2}O_2$) and conjugation with sulfate are the most exergonic and

should occur easily. This may be explained by the fact that oxidation tends to introduce polar functional

groups whose interactions with water serve to stabilize the compound. The reaction energies of sulfation are

very exothermic in gas phase but not solution phase. This large difference in energy is explained by the

presence of charged species in the reactant side which is greatly stabilized by the solvent – water – when

compared to the product side that receives little stabilization upon solvation. The most endergonic reactions

are four-membered ring formation (4D) and oxidation from alcohol to aldehyde (o3, removal of H_2). The four-

membered ring formation was proposed based on experimental evidence [18] published in 1988 and should

be verified in further experiment. Difference in reaction energies due to position effect can be mostly

explained by steric hindrance (i.e. g-8 has the highest reaction energy.) and inductive effect (i.e. o-8

producing secondary alcohol is the most exergonic.).

Gas phase energies were used to calculate solution phase energies, but as discussed previously they are

not representative of the reactions that occur in biological systems. Since these reactions happen in solution,

only the energies in solution phase were considered for each metabolite. Standard Gibbs energies for each

metabolite are shown relative to the parent compound (**1** oeeee) in Figure 3. The step numbers correspond

to the number of reactions required to generate each compound from **1** oeeee according to Figure 1. The

first step from the parent compound tends to be the most exergonic with an average at -38.6 kcal/mol and

the average reaction energy decreases monotonically to around -3.1 kcal/mol at the fifth step.

Compounds with the lowest energy from each step were identified and are shown in Figure 4 with additional

intermediates for completion. In general, the metabolite with the lowest relative energy in a step was the

starting material for the lowest energy metabolite in the next.

• This energy diagram (Figure 4) is in agreement with major aspects of published metabolic

pathways, in particular the conversion of menthol to *p*-menthane-3,8-diol (**3** oeooe). Partly due to

increased solubility, the compounds **3** oeooe and its glucuronic acid conjugates, **66** Oeooe/**67**

oeOee, were found to be major metabolites excreted in the urine of both rats and humans [17-19].

In contrast, *p*-menthane-3,7-diol (**2** ooeee) and *p*-menthane-3,9-diol (**4** oeeoe-*R*, **5** oeeoe-*S*)

excreted from both rats and humans in small quantities. Figure 2 reports that oxidation from alkane

to aldehyde at either position 7, 8, 9, or 10 is equally exothermic with a slight preference for position

8. Published evidence that **3** oeoee is formed as a product of enzymatic activity [18] and this

observed thermodynamic preference explain the disproportionately large amount of **3** oeoee isolated

experimentally compared to its isomers.

• Oxidation from alcohol to aldehyde is an endothermic reaction, hence metabolites containing

aldehyde groups are either not detected or detected in small quantities and serve as intermediates

to products of intramolecular aldol condensation to form cyclic ethers or further exothermic oxidation

to carboxylic acid. In rats, no metabolites containing aldehyde groups were detected in the plasma,

urine, bile, or feces. [17, 18] The published metabolic pathways show a direct conversion from

alcohol to carboxylic acid. Only the most recent study conducted by Hiki et al. [19] reported

detection of aldehyde menthol glucuronides in human urine at very low levels; the pathway proposed

by Hiki et al shows further conversion to cyclic ethers and carboxylic acid. Since oxidation is a

stepwise process, Figure 4 shows this stepwise conversion from **6** oooee to **44** oxoee-*R*.

Concluding remarks

In this study, gas-phase structures of menthol and its metabolites (a total of 102 compounds and 151

reactions) were obtained by quantum calculations at B3LYP/6-311++G(d,p) level. The standard Gibbs

energies of their respective reactions in solution were calculated with the SMD solvation model and

corrected for standard state conditions. The lowest energy diagram (Figure 4) reported was largely in

agreement with previously published experimental results. Information obtained in this study opens

possibilities for further investigation of the pharmacological effects of menthol and its metabolites. Given that

oxidation metabolites of menthol are energetically favorable, potency and toxicity of these oxidized

derivatives should be further investigated. Different stereoisomer of menthol as well as MD-based

approaches could also be explored in future research.

Supporting information

174 S1 File All Q-CHEM output files (zip)

175 S2 File Wolfram Mathematica notebook, shell script and template (zip)

176 S3 File Microsoft Excel spreadsheet and associated files (zip)

Acknowledgement

178 We appreciate helpful suggestions from Assoc. Prof. Dr. Yuthana Tantirungrotechai.

Author Contributions

[This is to be copied from Editorial Manager after acceptance of the manuscript.]

References

1. Eccles R. Menthol and related cooling compounds. Journal of Pharmacy and Pharmacology. 1994;46(8):618-30.

2. Watson H, Hems R, Rowsell D, Spring D. New compounds with the menthol cooling effect. J Soc Cosmet Chem. 1978;29(185):200.

3. Swandulla D, Carbone E, Schäfer K, Lux H. Effect of menthol on two types of Ca currents in cultured sensory neurons of
vertebrates. Pflügers Archiv. 1987;409(1-2):52-9.

4. Swandulla D, Schäfer K, Lux H. Calcium channel current inactivation is selectively modulated by menthol. Neuroscience letters.
1986;68(1):23-8.

5. Thorup I, Würtzen G, Carstensen J, Olsen P. Short term toxicity study in rats dosed with pulegone and menthol. Toxicology letters.
1983;19(3):207-10.

6. Statistica. Production volume of menthol in India from FY 2013 to FY 2017 (in thousand metric tons) 2019. Available from:
https://www.statista.com/statistics/727911/india-menthol-production-volume.

7. Patel T, Ishiuji Y, Yosipovitch G. Menthol: a refreshing look at this ancient compound. Journal of the American Academy of
Dermatology. 2007;57(5):873-8.

8. Karapinar M, Aktu ŞE. Inhibition of foodborne pathogens by thymol, eugenol, menthol and anethole. International Journal of Food
Microbiology. 1987;4(2):161-6.

9. Galeotti N, Mannelli LDC, Mazzanti G, Bartolini A, Ghelardini C. Menthol: a natural analgesic compound. Neuroscience letters.
2002;322(3):145-8.

10. Brauchi S, Orio P, Latorre R. Clues to understanding cold sensation: thermodynamics and electrophysiological analysis of the cold
receptor TRPM8. Proceedings of the National Academy of Sciences. 2004;101(43):15494-9.

11. Sherkheli MA, Vogt-Eisele AK, Bura D, Márques LRB, Gisselmann G, Hatt H. Characterization of selective TRPM8 ligands and their
structure activity response (SAR) relationship. Journal of Pharmacy & Pharmaceutical Sciences. 2010;13(2):242-53.

12. Giovino GA, Sidney S, Gfroerer JC, O'Malley PM, Allen JA, Richter PA, et al. Epidemiology of menthol cigarette use. Nicotine &
Tobacco Research. 2004;6(Suppl_1):S67-S81.

13. Ahijevych K, Garrett BE. Menthol pharmacology and its potential impact on cigarette smoking behavior. Nicotine & Tobacco
Research. 2004;6(Suppl_1):S17-S28.

14. Rakieten N, Rakieten ML, Boykin M. Effects of menthol vapor on the intact animal with special reference to the upper respiratory
tract. Journal of the American Pharmaceutical Association. 1954;43(7):390-2.

15. Bernson VS, Pettersson B. The toxicity of menthol in short-term bioassays. Chemico-biological interactions. 1983;46(2):233-46.

16. Joint FAO/WHO Expert Committee on Food Additives (World Health Organization & Food and Agriculture Organization of the
United Nations), editor Evaluation of certain food additives: twentieth report of the Joint FAO/WHO Expert Committee on Food
Additives; meeting held in Rome from 21 to 29 April 1976. Geneva2009.

17. Yamaguchi T, Caldwell J, Farmer PB. Metabolic fate of [3H]-l-menthol in the rat. Drug Metabolism and Disposition. 1994;22(4):616-
24.

18. Madyastha KM, Srivatsan V. Studies on the metabolism of l-menthol in rats. Drug Metabolism and Disposition. 1988;16(5):765-72.

19. Hiki N, Kaminishi M, Hasunuma T, Nakamura M, Nomura S, Yahagi N, et al. A phase I study evaluating tolerability,
pharmacokinetics, and preliminary efficacy of L menthol in upper gastrointestinal endoscopy. Clinical Pharmacology & Therapeutics.
2011;90(2):221-8.

20. Miyazawa M, Marumoto S, Takahashi T, Nakahashi H, Haigou R, Nakanishi K. Metabolism of (+)-and (-)-menthols by CYP2A6 in
human liver microsomes. Journal of oleo science. 2011;60(3):127-32.

21. Gelal A, Jacob P, Yu L, Benowitz NL. Disposition kinetics and effects of menthol. Clinical Pharmacology & Therapeutics.
1999;66(2):128-35.

22. Marenich AV, Cramer CJ, Truhlar DG. Universal solvation model based on solute electron density and on a continuum model of the
solvent defined by the bulk dielectric constant and atomic surface tensions. The Journal of Physical Chemistry B. 2009;113(18):6378-
96.

23. Limpanuparb T, Roongruangsree P, Areekul C. A DFT investigation of the blue bottle experiment: E∘ half-cell analysis of
autoxidation catalysed by redox indicators. Royal Society Open Science. 2017;4(11):170708.

24. Limpanuparb T, Noorat R, Tantirungrotechai Y. In Silico Investigation of Mitragynine and 7-Hydroxymitragynine Metabolism. BMC
Research Notes. 2019;12:451.

25. Ho J, Ertem MZ. Calculating free energy changes in continuum solvation models. The Journal of Physical Chemistry B.
2016;120(7):1319-29.

26. Ho J, Coote ML. A universal approach for continuum solvent pKa calculations: are we there yet? Theoretical Chemistry Accounts.
2010;125(1-2):3.

27. Camaioni DM, Schwerdtfeger CA. Comment on "Accurate experimental values for the free energies of hydration of H+, OH-, and
H3O+". The Journal of Physical Chemistry A. 2005;109(47):10795-7.

28. Bryantsev VS, Diallo MS, Goddard Iii WA. Calculation of solvation free energies of charged solutes using mixed cluster/continuum
models. The Journal of Physical Chemistry B. 2008;112(32):9709-19.

29. Shao Y, Gan Z, Epifanovsky E, Gilbert AT, Wormit M, Kussmann J, et al. Advances in molecular quantum chemistry contained in
the Q-Chem 4 program package. Molecular Physics. 2015;113(2):184-215.

30. Wolfram Research Inc. Mathematica. 2019.

31. Härtner J, Reinscheid UM. Conformational analysis of menthol diastereomers by NMR and DFT computation. Journal of Molecular
Structure. 2008;872(2-3):145-9.

32. Wiberg KB. Basis set effects on calculated geometries: 6-311++G** vs. aug-cc-pVDZ. Journal of Computational Chemistry.
2004;25(11):1342-6.

46-49 oycyc
50 oeeyy

16-17 oxccc
18-19 oeexe
20-21 oyoee
22-25 oycoc
26-27 oocyc
28-29 oeoye
30-31 oeeyo

53-54 oyoee-4D

55-56 oxocc-4D

12-13 oyeee
14-15 occyc

32-35 ooeye-5A
36-39 oeoye-5A
40-43 occyo-5A

44-45 oxoee

51 oeoee-4D

52 ooocc-4D

6 oooee
7-8 ooeoe
9-10 ocooc
11 occoo

1 occcc

2 ooccc
3 ococc
4-5 oeeoe

57 sccsc

58 soeee
59-60 sccoc

61-62 sococ

63 Oeeee