Genomic hypervariability of phage Andromeda is unique among known

dsDNA viruses

Damian J. Magill[1†], Leonid A. Kulakov[1], Timofey A. Skvortsov[2]*

[1] Queen's University Belfast, School of Biological Sciences,

19 Chlorine Gardens, Belfast BT9 5DL, Northern Ireland

[2] Queen's University Belfast, School of Pharmacy, Medical Biology Centre,

97 Lisburn Road, Belfast BT9 7BL, Northern Ireland

* Correspondence: Timofey A. Skvortsov, Queen's University Belfast, School of Pharmacy,

Medical Biology Centre, 97 Lisburn Road, Belfast, BT9 7BL, Northern Ireland; email:

t.skvortsov@qub.ac.uk

[†] Present address: DuPont (Danisco), Nutrition & Health, Route de Buxières – CS 10010,

86220 Dangé-Saint-Romain, France

Keywords: bacteriophage; phage; genome hypervariability; quasispecies;

Autographivirinae.

Abstract

A new lytic bacteriophage Andromeda, specific to the economically important plant pathogen *Pseudomonas syringae*, was isolated and characterised. It belongs to the *Podoviridae* family, *Autographivirinae* subfamily and possesses a linear dsDNA genome of 40,008 bp with four localised nicks. Crucially, Andromeda's genome has no less than 80 hypervariable sites (SNPs), which show genome wide distribution resulting in heterogenous populations of this phage reminiscent of those of RNA virus quasispecies. Andromeda has no nucleotide sequence homology to phage phiNFS, a member of *phiKMVviruses*, in which a similar phenomenon was discovered. We show that Andromeda and Andromeda-related phages form a group within the *Autographivirinae,* designated here as the "ExophiKMVviruses". The "ExophiKMVviruses" were revealed to share conservation of gene order with core *phiKMVviruses* despite their sequence-based relationship to SP6-related phages. Our findings suggest that genomic hypervariability might be a feature that occurs among various *Autographivirinae* groups.

Introduction

The *phiKMVviruses* constitute a ubiquitously distributed group of bacteriophages closely related to *Teseptimaviruses* and *Zinderviruses*. These three large groups and a number of smaller ones fall under the *Autographivirinae* subfamily, the major difference between them being the localisation of their single subunit RNA polymerase enzymes. With respect to *phiKMVviruses* in particular, the majority of phages infect *Pseudomonas* strains, with first representatives being described in 1980s [1]. *phiKMVviruses* are of interest for various reasons. Those infecting *Pseudomonas aeruginosa* exhibit a number of interesting features including the presence of four localised single-strand interruptions (nicks) associated with the consensus sequence 5'-CGACNNNNNCCTACTCCGG-3' [1, 2]. The more recent study revealed that at least some phiKMV phages exhibit hypervariability associated with specific genomic regions [3]. These hypervariable loci were found in genes associated with phage adsorption/host specificity and may therefore influence the evolutionary pathway of the phages [3].

This study describes the biological properties of a novel phage, Andromeda, which infects *Pseudomonas syringae*. The presence of a large number of hypervariable sites in the Andromeda's genome and its relationship to *phiKMVviruses* are investigated.

Materials and Methods

Phage Isolation and Purification

Bacteriophage Andromeda was isolated by enrichment of 100 ml of water from River Lagan (Belfast, Northern Ireland) in cultures of *Pseudomonas syringae* pv *syringae* Van Hall followed by plating using the double agar overlay plaque assay [4]. Pure isolates of the phage were obtained by conducting three rounds of phage propagation from individual plaques. To obtain high titre preparations of the phage, cultures of *P. syringae* (OD_{600} = 0.3) were inoculated at a MOI = 0.01. Chloroform was added immediately after culture lysis (~4 hours) to maximise yield. Resistance of the phage particles to chloroform treatment was determined in separate experiments. The phage lysate was concentrated by PEG precipitation, and purified using CsCl density gradient centrifugation as described by Sambrook and Russell [4]. Bands were extracted and dialysis ensued with modified SM buffer (100 mM NaCl, 50 mM Tris, 10 mM, 8 mM $MgSO_4$, 10 mM $CaCl_2$) using Amicon Ultra-15 100kDa filters (Millipore). Phage concentrates were treated with SDS and

proteinase K, after which phenol-chloroform DNA extraction followed by ethanol

precipitation was used to obtain phage DNA [4]. DNA was further purified twice prior to

sequencing using the MoBio DNA Purification Kit according to the manufacturer's

instructions.

Electron Microscopy

Carbon-coated formovar grids were subjected to hydrophilic treatment with poly-L-lysine

for 5 min. Upon drying, CsCl purified and dialysed phage suspensions were deposited on

the grids and successively stained with 2% uranyl acetate (pH 4.5). Samples were

observed using a Phillips CM100 transmission electron microscope at 100 KeV.

Determination of Adsorption Coefficient

Determination of adsorption efficiency of phages was carried out using the method

described in [5]. All adsorption assays were conducted in triplicate with final adsorption

constant calculated from the mean of the replicate constants.

One-step Growth Curves

Determination of bacteriophage one-step growth curves was conducted according to the

established protocol. Bacterial cultures were grown until $OD_{600} = 0.3$ (10^8 CFU/ml) and

inoculated with phage to give MOI $\geq$ 1. Adsorption was allowed for 5 min at 37^{o}C followed

by incubation with aeration at the same temperature. Duplicate 100 µl samples were taken

every 5 minutes for 90 minutes. One of each samples were treated with 150 µl of

chloroform and both were plated on the lawns of *P. syringae* using the double agar overlay

plaque assay. Burst size was calculated by dividing the average of the values following the

growth phase of the curve (the step) by the average of the baseline values before the first

increase was observed in titre. The average of triplicates was then taken as the burst size.

Latent periods were taken as the point in the graph of the untreated phage extraction

whereby a rise in titre was observed, whereas the eclipse period was determined from the

chloroform treated curve.

Visualisation of localised single-strand interruptions

Detection of localised nicks were conducted as described earlier [2]. Phage DNA (3 µg)

underwent overnight ligation at 4^{o}C in a final volume of 20 µl. NaOH was added to

reactions to a final concentration of 50 mM. This was also done for 1.8 µg of unligated

DNA. Both samples were incubated for 5 min on ice and immediately analysed by gel

electrophoresis (0.9% agarose).

Library Preparation and Sequencing

Sequencing libraries were prepared from 50 ng of phage genomic DNA using the Nextera

DNA Sample Preparation Kit (Illumina, USA) at the University of Cambridge Sequencing

Facility. A 1% PhiX v3 library spike-in was used as a quality control for cluster generation

and sequencing. Sequencing of the resulting library was carried out from both ends (2x300

bp) with the 600-cycle MiSeq Reagent Kit v3 on MiSeq (Illumina, USA) and the adapters

trimmed from the resulting reads at the facility.

Sequence Assembly and Bioinformatics

Sequencing reads underwent initial quality checking using FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) followed by stringent

trimming parameters. Reads with a Q-score < 20, containing N's, and with a final length <

40bp were discarded using Trimmomatic v0.32 [6]. Assembly was carried out using

Geneious R8 (Biomatters, New Zealand) and checked for potential misassembly by

comparison with output from SPAdes v3.7.0 [7]. Genome length contigs were input to the

scaffolding tool SSPACE to extend these in order to ensure the resolution of genome ends

[8]. In addition, the sequence obtained was utilised as a reference for read mapping using

bbmap v35.x (https://jgi.doe.gov/data-and-tools/bbtools/) and subsequent manual

inspection in IGV to check for potentially ambiguous regions [9]. ORFs were classified

using the intrinsic Artemis [10] ORF classification and GeneMark.hmm [11] followed by

manual inspection, ORF borders correction and verification of ribosomal binding sites.

Gene annotation was carried out using BLASTp, Delta-BLAST, HMMER, InterProScan

and HHpred [12–15]. tRNAs were classified using tRNA-scanSE and Aragorn [16, 17], and

rho-independent terminators were identified using Arnold [18] with an energy cutoff of -10

employed, following by manual inspection of the output from subsequent Mfold analysis

and adjustment as necessary [19]. To identify putative regulatory sequences, 100 bp

upstream of every ORF was extracted using an in-house script and analysed using both

neural network promoter prediction for prokaryotic sigma 70 promoters and Multiple EM for

Motif Elicitation to identify phage specific promoters [20]. Phage Andromeda assembled

genomic sequence was deposited to NCBI Genbank (Genbank accession number:

KX458241; RefSeq accession number: NC_031014). A reannotated genome of phage

Andromeda in Genbank format can be found in Supplementary File 1.

Phylogenetic analysis

In order to taxonomically annotate the newly isolated phage and determine its phylogenetic relationships with other bacteriopahges, two conserved Andromeda proteins, the single subunit RNA polymerase (RNAP, YP_009279548.1) and the terminase large subunit (TerL, YP_009279565.1), were used to generate phylogenetic trees. BLASTp search with each of these proteins against *Podoviridae* available in RefSeq at the moment of writing was conducted and only phage genomes containing copies of both TerL and RNAP with high enough similarity (e-value ≤ 1e-5; query coverage ≥ 75%; percent identity ≥ 25%) to corresponding Andromeda's proteins were retained for subsequent analyses; 147 complete phage genomes were selected in total (Supplementary Table 1). Alignments of the amino acid sequences of RNAP and TerL were carried out in MAFFT using L-INS-I settings [21], TrimAl was used to for automated alignment trimming [22]. The inference of phylogenies was then carried out using IQ-Tree [23], which was allowed to compute the optimal protein model to be used. Visualisation of the resulting tree file was subsequently performed with the Interactive tree of life (IToL) [24]. The GGgenes package (https://CRAN.R-project.org/package=gggenes) for R programming environment (http://www.R-project.org/) was used to generate a series of genome maps.

Molecular Modelling

Molecular models were constructed from corresponding amino acid sequences using the I-Tasser package [25] and visualised with Chimera [26]. Fidelity of models was assessed using the validation tools of the Whatif server (https://swift.cmbi.umcn.nl/whatif/) and checked for stereochemical clashes within Chimera. Ramachandran analysis was subsequently carried out as an additional point of quality assessment. Superimposition and quantitative similarity of protein structures was provided by the TM-Align algorithm [27].

Results and Discussion

Isolation and Biological Characteristics

Bacteriophage Andromeda was isolated from the River Lagan as described in Materials and Methods. Upon infection of *Pseudomonas syringae,* phage Andromeda yields large, clear plaques of ~3 mm in size. Electron microscopic analysis of Andromeda revealed a virion morphology characteristic of *Podoviridae* (Fig. 1). Specifically, Andromeda was found to possess a capsid of icosahedral symmetry with a length of 61 nm (±1.5 nm) and width of 62 nm (±1 nm) and a detectable tail structure.

Analysis of the growth characteristics revealed an adsorption coefficient of 1.48×10^{-9} ml min^{-1} (Fig. 2a), eclipse period of 20 – 25 min, and latent period of 25 – 30 min (Fig. 2b). Upon completion of its infection cycle, Andromeda yields on average 70 particles per cell.

Genome Structure and Gene Regulation

Phage DNA was sequenced as described in Materials and Methods to a per base average coverage of 2149. The assembly and subsequent analysis were conducted and these revealed a linear dsDNA genome of 40,008 bp (Supplementary File 1).

The GC content of the genome is 58.23% and using cumulative GC skew analysis, the putative origin of replication was found to be at 3001 bp. 46 ORFs were predicted within the Andromeda genome resulting in a total coding potential of 95.1%. Six ORFs utilise TTG, and 1 utilises GTG as start codons with the rest beginning with ATG. All ORFs are encoded off the same strand. Analysis of the Andromeda genome by both the PhageTerm [28] and Li's methods [29] revealed that this phage is terminally redundant and likely utilises a headful mechanism of packaging similar to that of phage P1 (+ve strand analysis (p = 1.17e-4), -ve strand analysis (p = 2.81e-6)). Based on nucleotide homology, the closest relative to Andromeda is the *P. tolaasii* phage Bf7 [30] with which it shares 83% identity across 85% of its genome (Fig. 3). Of the 46 ORFs, 25 were functionally assigned and are arranged within a strict modular architecture. Eight ORFs encode products associated with DNA replication, recombination, and repair. In particular, the presence of a single subunit RNA polymerase of T3/T7 type was detected (Fig. 3; RNAP), which is located downstream of the DNA polymerase (DNAP). This RNA polymerase is conserved within *Autographivirinae* and exhibits similarity to that of phiNFS, particularly in the N- and C- termini, the specificity loop, and the palm and finger domains. Indeed, molecular modelling of both polymerases and their superimposition with TM-Align revealed remarkably similar structures (Supplementary Figure 1) with TM-score of 0.94. This, along

with the characteristic position of the RNA polymerase, initially suggested the inclusion of

Andromeda within the *phiKMVviruses*. Eleven ORFs in Andromeda encode structural

proteins including typical gene products observed in members of the *Autographivirinae*

such as tail tubular proteins. The remaining six functionally assigned ORFs encode the

terminase small and large subunits as well as the components of the lysis machinery, with

the lysozyme belonging to the GH24 family of proteins.

With respect to gene regulation in Andromeda, nine σ^{70} promoters were predicted, of

which three precede the early gene region. This appears to represent a strong signal

permitting the rapid transcription of early genes. Due to the presence of an RNA

polymerase, we attempted to predict whether phage specific promoters were present. No

statistically significant elements were found and of the obtained weak predictions, these

were largely found on the non-coding strand. It is possible that Andromeda utilises σ^{70}

elements in the generation of large polycistronic mRNAs. In regard to transcriptional

termination, only one rho-independent terminator was found, downstream of gp5. No tRNA

genes were discovered in this phage, a feature typical of the *Autographivirinae* group of

phages.

Localised single strand interruptions have been reported in *phiKMVviruses* infecting *P.*

aeruginosa [2] and *P. putida* phage tf [31]. A search of the Andromeda genome failed to

find the consensus associated with nicks in *P. aeruginosa* phages or *P. putida* phage.

Upon subjecting Andromeda genomic DNA to NaOH denaturation, faint but distinguishable

bands were observable on agarose gels which disappear when conducting this reaction

after ligation (Supplementary Figure 2). These results prove that localised nicks containing

adjacent 5'-phosphate and 3'-hydroxyl groups are present in the Andromeda genome. The

low intensity of the bands observed may suggest that only a proportion of phage DNA

molecules is nicked. This was previously demonstrated for T5 phage of *Escherichia coli*

where a number of major (present in over 50% of molecules) and minor (lower presence)

canonical nicks are known [32–34].

Genetic Heterogeneity in the Andromeda Genome

It has previously been discovered that several members of the *phiKMVviruses* display a

high level of genome variability that is localised to specific regions [3]. This is also true in

case of Andromeda, which was found to possess the highest number of such variants. In-

depth analysis of this genomic feature was conducted in order to gain insights into the

nature of the changes and the potential mechanisms at play.

Analysis of the deep sequencing data of the Andromeda genome revealed that there are

at least 80 positions (SNP) where nucleotide substitutions are detected with frequencies
ranging from 1% to 12.48%. Detailed information with respect to each variant is provided
in Supplementary Table 2. Cumulative analysis of these frequencies shows that lower
frequency variants predominate, with a high level of tapering observed above 3% (Fig. 4).
The mutations are distributed genome-wide, which is dissimilar to the pattern reported
previously in other *phiKMVviruses* where a distinct bias of SNPs distribution (e.g. in genes
responsible for phage adsorption) was observed [3]. On a cumulative basis however, it is
clear that regions of high variability are observable, which suggests either the consecutive
introduction of errors in a single event such as polymerase slippage, or that these
represent hotspots for mutation (Fig. 4).
Further analysis of the nature of variants revealed that 73/80 (91.3%) lie within coding
regions which is unsurprising given that the coding potential of the genome is 95.1 % (Fig.
5). With respect to those variants lying within coding regions, 55 of these result in an
amino acid change. Of the 18 silent mutations, 17 were found to exist within the third
codon position. This position is overall the predominant location of variants with the
second codon position having a similar abundance (Fig. 5). Interestingly, these 17 silent
mutations constitute only half of the total variants observed at the third position. The
predominance of missense mutations in general is extremely interesting, particularly in the
third position. Analysis of the proportion of transversion and transition mutations revealed
an unexpected abundance of the former, with 78.75% of all variants being of this type
(including those in non-coding regions). Most transversions are observed in the second
codon position, closely followed by the third. In general, whilst there are more potential
pathways to transversion mutations, they are in reality less common relative to transitions.
The exchange of a single ring base for a double ring and vice-versa is a significant change
and is more likely to result in an alteration at the amino acid level and indeed, one that is
less likely to be conservative. The predominance of transversions here explains the high
level of non-silent mutations observed across all codon positions and is a unique
occurrence not previously reported in other biological systems. The additional abundance
of transversion mutations may also further implicate the DNA polymerase as a responsible
factor in variant generation through some form of biochemical bias. This is additionally
supported by the presence of an insertion mutation in the Andromeda genome within
ORF14 (Supplementary Table 2). This variant lies within an 8 base poly-G tract that may
have given rise to a slippage event during replication and results in a +1 frameshift and
truncated ORF14 product.
The possible involvement of the phage DNA polymerases in generation of hypervariability

in genomes of *phiKMVviruses* was discussed previously [3]. Phage Andromeda's DNA

polymerase is similar to those of *phiKMVviruses* with respect to absence of the

thioredoxin-biding domain; however, the genome-wide distribution of variable sites in its

genome suggest that different mechanisms of diversity generation might be in play in case

of Andromeda.

Phylogenetic relationships of Andromeda

Autographivirinae is a subfamily within *Podoviridae,* which at the time of writing included

several genera, with *Teseptimaviruses, Drulisviruses, Friunaviruses, Przondoviruses,*

Zinderviruses and *phiKMVviruses* being the largest. Certain features of Andromeda, such

as genomic architecture and gene order, presence of nicks and hypervariable sites in its

DNA, and the similarity of its gene products such as the RNA/DNA polymerases, tail

tubular proteins, major capsid protein, terminase subunits, and lysis machinery, implicates

this phage as a member of *Autographivirinae*, the *phiKMVviruses* in particular.

Nevertheless, it has no detectable nucleotide identity to currently recognised members of

phiKMVviruses and has recently been assigned to *Bifseptivirus* genus within *Podoviridae*

alongside with *Pseudomonas* phage Bf7 by the International Committee on Taxonomy of

Viruses (ICTV proposal 2018.063B.R.Bifseptvirus, approved on 24 February 2019). This

highlights the somewhat problematic nature of taxonomic classification of Andromeda and

in order to better understand its phylogenetic standing, we conducted an analysis of

Andromeda using two conserved proteins – RNA polymerase (RNAP) and terminase large

subunit (TerL). BLASTp searches of the NCBI RefSeq database for RNAP and TerL

similar to those of Andromeda (see Materials and Methods) resulted in the retrieval of 147

complete viral genomes, including 46 recognised by the ICTV (Supplementary Table 1).

Almost all the phages used for phylogenetic analyses belong to *Autographivirinae* (except

Pseudomonas phages Andromeda and Bf7 and *Aquamicrobium* phage P14) and infect a

wide range of hosts from *Cyanobacteria* to *Pseudomonas.* Reconstruction of phylogenetic

trees with IQ-Tree using RNAP and TerL as markers resulted in trees of similar

morphology (Fig. 6a and 6b). It can be clearly seen that *Teseptimaviruses* and

Przondoviruses cluster separately from the majority of other phages, while Andromeda

belongs to a loose cluster of *Autographivirinae* phages comprising several smaller genera

with only few members (e.g. *Bifseptivirus* – *Pseudomonas* phages Andromeda and Bf7,

Aqualcavirus – *Aquamicrobium* phage P14) and unclassified *Autographivirinae* (e.g.

Ralstonia phage RSB3). Although we do not intend to propose any novel classification, we

designate this cluster of viruses as "ExophiKMVviruses" for the purposes of discussing this

group and to highlight its distinctive position relative to other *Autographivirnae*.

Despite not currently being classified by ICTV as *Autographivirinae*, phage Andromeda as

well as Bf7 could be included as members of this group based on the presence of a single

subunit RNAP in their genomes and their position within the cluster of

"ExophiKMVviruses". When we analyse the gene order of representatives of each genus

(Fig. 7), we find that the "ExophiKMVviruses", despite being most closely related to

Zinderviruses based on sequence identity percent of marker genes, show the downstream

localisation of the single-subunit RNA polymerase as is typical for "true" *phiKMVviruses.*

This is a feature conserved across the group (Fig. 7). Analysis of genome size distribution

of the putative *Autographivirinae* (Fig. 8) show that "ExophiKMVviruses" possess a

genome size range similar to that of other *phiKMVviruses*. We have shown previously that

phiKMVviruses such as *Pseudomonas* phage phiNFS lack the thioredoxin binding domain

in their DNA polymerases [3] that is responsible for heightened processivity and fidelity of

Teseptimaviruses [35, 36]. Molecular modelling and superimposition of the Andromeda

and T7 DNA polymerases shows the absence of this domain in Andromeda and highlights

another difference between *phiKMVviruses* and "Exophikmvviruses" in comparison to

Teseptimaviruses (Supplementary Figure 3). Taken together, it seems that the

"ExophiKMVviruses" might occupy an interface group between the *phiKMVviruses* and the

Teseptimaviruses , although sequence based classification alone being insufficient to

adequately resolve the relationships of the *Autographivirinae* as a whole*.*

Our analysis of the Andromeda phage has once again highlighted the existence of a novel

phenomenon within *Pseudomonas* phages that results in the production of highly

heterogeneous populations. Whilst error-prone polymerase activity is almost certainly to be

involved in the generation of genomic hypervariability, it unlikely to be the only constituent

at play, with intriguing patterns of codon position distribution and localisation suggesting

the involvement of some post-replication mechanisms. The taxonomic position of

Andromeda suggests that this phenomenon is not unique to *phiKMVviruses* and could

potentially include "Exophikmvviruses" and/or *Autographivirinae* as a whole.

Authors and contributors

Conceptualisation: D.J.M. and L.A.K.; data collection and analysis: D.J.M. and T.A.S.;

writing – original draft preparation: D.J.M., T.A.S. and L.A.K.; writing – review and editing:

T.A.S., D.J.M. and L.A.K

Funding information

This work received no specific grant from any funding agency

Conflicts of interest

The authors declare that there are no conflicts of interest.

Ethical statement

No experimental work with humans or animals was performed in this study.

References

1. Kulakov LA, Ksenzenko VN, Kochetkov VV, Mazepa VN, Boronin AM. DNA homology and adsorption specificity of *Pseudomonas aeruginosa* virulent bacteriophages. *MGG Mol Gen Genet* 1985; **200**: 123–127.

2. Kulakov LA, Ksenzenko VN, Shlyapnikov MG, Kochetkov VV, Del Casale A, Allen CCR, et al. Genomes of "phiKMV-like viruses" of *Pseudomonas aeruginosa* contain localized single-strand interruptions. *Virology* 2009; **391**: 1–4.

3. Magill DJ, Kucher PA, Krylov VN, Pleteneva EA, Quinn JP, Kulakov LA. Localised genetic heterogeneity provides a novel mode of evolution in dsDNA phages. *Sci Rep* 2017; **7**.

4. Sambrook Joseph, Russell DW. Molecular cloning : a laboratory manual. 2001. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

5. Clokie MRJ, Kropinski AM (eds). Bacteriophages: methods and protocols. Volume 1. 2009. Humana Press, New York.

6. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 2014; **30**: 2114–2120.

7. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *J Comput Biol* 2012; **19**: 455–477.

8. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. Scaffolding pre-assembled contigs using SSPACE. *Bioinforma Oxf Engl* 2011; **27**: 578–579.

9. Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* 2013; **14**: 178–192.

10. Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. Artemis: an integrated platform for visualization and analysis of high-throughput sequence-based

experimental data. *Bioinformatics* 2012; **28**: 464–469.

11. Lukashin AV, Borodovsky M. GeneMark.hmm: New solutions for gene finding. *Nucleic*

Acids Res 1998; **26**: 1107–1115.

12. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search

tool. *J Mol Biol* 1990; **215**: 403–410.

13. Boratyn GM, Schäffer AA, Agarwala R, Altschul SF, Lipman DJ, Madden TL. Domain

enhanced lookup time accelerated BLAST. *Biol Direct* 2012; **7**: 12–12.

14. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5:

genome-scale protein function classification. *Bioinforma Oxf Engl* 2014; **30**: 1236–

1240.

15. Zimmermann L, Stephens A, Nam S-Z, Rau D, Kübler J, Lozajic M, et al. A

Completely Reimplemented MPI Bioinformatics Toolkit with a New HHpred Server at

its Core. *J Mol Biol* 2018; **430**: 2237–2243.

16. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA

genes in genomic sequence. *Nucleic Acids Res* 1997; **25**: 955–964.

17. Laslett D, Canback B. ARAGORN, a program to detect tRNA genes and tmRNA

genes in nucleotide sequences. *Nucleic Acids Res* 2004; **32**: 11–16.

18. Naville M, Ghuillot-Gaudeffroy A, Marchais A, Gautheret D. ARNold: A web tool for

the prediction of Rho-independent transcription terminators. *RNA Biol* 2011; **8**: 11–13.

19. Zuker M. Mfold web server for nucleic acid folding and hybridization prediction.

Nucleic Acids Res 2003; **31**: 3406–3415.

20. Bailey TL, Johnson J, Grant CE, Noble WS. The MEME Suite. *Nucleic Acids Res*

2015; **43**: W39–W49.

21. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol Biol Evol* 2013; **30**: 772–780.

22. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 2009; **25**:

1972–1973.

23. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol*

2015; **32**: 268–274.

24. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. *Nucleic Acids Res* 2019; **47**: W256–W259.

25. Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y. The I-TASSER Suite: protein

structure and function prediction. *Nat Methods* 2015; **12**: 7–8.

26. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al.

UCSF Chimera—A visualization system for exploratory research and analysis. *J*

Comput Chem 2004; **25**: 1605–1612.

27. Zhang Y, Skolnick J. TM-align: a protein structure alignment algorithm based on the

TM-score. *Nucleic Acids Res* 2005; **33**: 2302–2309.

28. Garneau JR, Depardieu F, Fortier L-C, Bikard D, Monot M. PhageTerm: a tool for fast

and accurate determination of phage termini and packaging mechanism using next-

generation sequencing data. *Sci Rep* 2017; **7**: 8292.

29. Li S, Fan H, An X, Fan H, Jiang H, Chen Y, et al. Scrutinizing Virus Genome Termini

by High-Throughput Sequencing. *PLOS ONE* 2014; **9**: e85806.

30. Sajben-Nagy E, Maróti G, Kredics L, Horváth B, Párducz Á, Vágvölgyi C, et al.

Isolation of new Pseudomonas tolaasii bacteriophages and genomic investigation of

the lytic phage BF7. *FEMS Microbiol Lett* 2012; **332**: 162–169.

31. Glukhov AS, Krutilina AI, Shlyapnikov MG, Severinov K, Lavysh D, Kochetkov VV, et

al. Genomic Analysis of Pseudomonas putida Phage tf with Localized Single-Strand

DNA Interruptions. *PLOS ONE* 2012; **7**: e51163.

32. Bujard H. LOCATION OF SINGLE-STRAND INTERRUPTIONS IN THE DNA OF

BACTERIOPHAGE T5+. *Proc Natl Acad Sci U S A* 1969; **62**: 1167–1174.

33. Hayward GS, Smith MG. The chromosome of bacteriophage T5: I. Analysis of the

single-stranded DNA fragments by agarose gel electrophoresis. *J Mol Biol* 1972; **63**:

383–395.

34. Johnston JV, Nichols BP, Donelson JE. Distribution of 'minor' nicks in bacteriophage

T5 DNA. *J Virol* 1977; **22**: 510–519.

35. Magill DJ, McGrath JW, O'Flaherty V, Quinn JP, Kulakov LA. Insights into the

structural dynamics of the bacteriophage T7 DNA polymerase and its complexes. *J*

Mol Model 2018; **24**: 144.

36. Tabor S, Huber HE, Richardson CC. Escherichia coli thioredoxin confers processivity

on the DNA polymerase activity of the gene 5 protein of bacteriophage T7. *J Biol*

Chem 1987; **262**: 16212–16223.

Legends to Figures

Fig. 1. Electron microscopic image of bacteriophage Andromeda.

Fig. 2. Growth characteristics of bacteriophage Andromeda. A. Determination of adsorption coefficient. B. One-step growth curves: blue line – infected *P. syringae* cells treated with chloroform (intracellular phage); red line – untreated infected *P. syringae* cells (extracellular phage). All experiments were performed in triplicates, error bars represent one standard deviation.

Fig. 3. Genetic organisation of Andromeda and its homology to Pseudomonas bacteriophage Bf7. Areas of substantial similarity between two genomes are shown as trapezia connecting two genome regions coloured according to average nucleotide identity levels of these regions (as calculated by BLASTn). Arrows on the genome map represent identified genes and demonstrate the direction of their transcription. They are coloured to reflect common functions of the encoded products. Putative genes with unidentified function are shown in grey colour. DNAP – DNA polymerase, RNAP – RNA polymerase, MCP – major capsid protein, TerL – terminase large subunit.

Fig. 4. The presence and distribution of variable sites in phage Andromeda. Each dot indicates an SNP, variant frequency is given by its position along the y-axis. The histogram on the top shows uneven distribution of variable sites in the phage's genome, the histogram on the right demonstrates the distribution of variant frequencies.

Fig. 5. Summary statistics of SNPs in bacteriophage Andromeda with respect to their position in codons and their effect on amino acids.

Fig. 6. Analysis of phylogenetic relationships of phage Andromeda. A. Consensus phylogenetic tree based on multiple sequence alignment of RNAP amino acid sequences of 147 RefSeq phages; LG+F+R6 substitution model. B. Consensus phylogenetic tree based on multiple sequence alignment of TerL amino acid sequences of 147 RefSeq phages; LG+R5 substitution model. Consensus trees were constructed from 1000 bootstrap trees, numbers in parentheses are bootstrap supports (%). Both trees were rooted with a cluster of 11 cyanophages as outgroup. "Exophikmvviruses" are outlined by a red box.

Fig. 7. Comparison of positions of RNA (blue) and DNA (red) polymerases in representatives of different genera. In *Zinderviruses* (SP6), *Teseptimaviruses* (T7) and *Przondoviruses* (K11) DNA polymerase is adjacent to structural genes and is positioned downstream of RNA polymerase, while for the rest of the viruses the opposite is true.

504   **Fig. 8.** Genome size distribution among viruses of different taxonomic groups analysed in this

505   study.

Legends to Supplementary Data

Supplementary Figure 1. Superimposition of three-dimensional structures of RNA polymerases from Pseudomonas phages Andromeda (salmon) and phiNFS (purple).

Supplementary Figure 2. Visualisation of the localised single stranded interruptions in the Andromeda's genome. (A) GeneRuler 1 kb Plus DNA Ladder ladder; (B) Untreated phage DNA; (C) NaOH treated DNA, major DNA hydrolysis products are labelled (1-3); (D) NaOH treated DNA which was preliminarily ligated by T4 DNA ligase.

Supplementary Figure 3. Superimposition of three-dimensional structures of DNA polymerases from Pseudomonas phage Andromeda (salmon) and Enterobacteria phage T7 (blue, model derived from PDB 1SKR).

Supplementary Table 1. NCBI RefSeq phage genomes used in this study.

Supplementary Table 2. Detailed description of SNPs and indels in phage Andromeda genome.

Supplementary File 1. Phage Andromeda annotated genome (Genbank format).

20 nm

Pseudomonas phage Andromeda

DNAP RNAP MCP TerL

Pseudomonas phage Bf7

5000 10000 15000 20000 25000 30000 35000 40000

Nucleotide identity

100% 68%

DNA replication/recombination/repair

Phage morphogenesis

DNA packaging and processing

Lysis

Bacteriophage Andromeda SNP Summary Statistics

Tree scale: 0.1

outgroup

"Exophikmvviruses"

A

Teseptimavirus
Zindervirus
phiKMVvirus
Friunavirus
Drulisvirus
Przondovirus

B