Angiosperm speciation speeds up near the poles

J. IGEA[1][*], A. J. TANENTZAP[1]

[1] Department of Plant Sciences, University of Cambridge, Downing St, Cambridge,

CB2 3EA, UK

[*] Correspondence: ji247@cam.ac.uk

<u>**Abstract**</u>

Recent evidence has questioned whether the Latitudinal Diversity Gradient

(LDG), whereby species richness increases towards the Equator, results from

higher rates of speciation in the tropics. Allowing for time heterogeneity in

speciation rate estimates for over 28,000 angiosperm species, we found that the

LDG does not arise from variation in speciation rates because lineages speciated

faster outside the tropics. These results were consistently retrieved using two

other methods to test the association between occupancy of tropical habitats and

speciation rates. Our speciation rate estimates were robust to the effects of both

undescribed species and missing taxa. Overall, our results show that speciation

rates follow an opposite pattern to global variation in species richness. Greater

ecological opportunity in the temperate zones, stemming from less saturated

communities and greater environmental change, may ultimately explain these

results.

<u>Keywords:</u> latitudinal diversity gradient, speciation, angiosperms, macroevolution,

biogeography, biodiversity

Introduction

Biodiversity on Earth is very unevenly distributed. The Latitudinal Diversity Gradient (LDG), whereby species richness increases towards the Equator, is the most prominent example of this unevenness. This pervasive pattern has fascinated biologists for more than two centuries[1–4], but its underlying causes remain largely unknown.

A major class of explanations proposes that elevated rates of speciation generate the higher diversity in the tropics and so primarily create the LDG[5]. This increase in speciation is generally attributed to higher environmental energy in the tropics, which in turn can hasten evolution[6], such as through shorter generation times, higher mutation rates[7] and foster more complex biotic interactions (e.g., faster coevolution or the 'Red Queen running faster when she is hot'[8]). All these factors make reproductive isolation more likely and favour speciation. However, while some studies on mammals[9], amphibians[10] and butterflies[11] have shown that tropical lineages speciate faster, other studies have found no latitudinal differences[12,13] or higher rates of speciation in temperate areas[14], particularly when focusing on recent rates of speciation[15–17]. Recent speciation rates should be less affected by extinction[18] and more accurate than deep-time estimates, even in the absence of paleontological evidence[19]. They should also reflect any variable that historically influenced speciation and also varied latitudinally, such as environmental energy[20].

Flowering plants are one of the largest eukaryotic radiations (roughly 350,000 species) and show a marked LDG[21]. Despite this pattern, latitudinal differences in

macroevolutionary rates in angiosperms have received comparatively little attention at

a species-level. Previous efforts have shown that angiosperm diversification is faster

in tropical families[22] and in tropical lineages within a limited number of clades[23,24].

The only species-level analysis found no differences in speciation, but did so by

assuming all rate differences in the angiosperm Tree of Life (ToL) were due to

geographic distribution alone[25]. One reason for the lack of attention to angiosperms is

that their vast diversity makes it difficult to obtain well-sampled phylogenetic and

distributional data at a species-level, which is required to estimate speciation rates

accurately.

Results and discussion

Here we use species-level data from the whole angiosperm ToL to assess the

latitudinal variation of speciation rates in plants. We first estimated species-specific

(i.e., tip-based) speciation rates for 29,703 species of flowering plants[26] using

Bayesian Analysis of Macroevolutionary Mixtures (BAMM[27]). BAMM allows for

rate heterogeneity through time and across lineages and accommodates non-random

incomplete taxon sampling. We then obtained latitudinal data for 236,894 angiosperm

species from a previous study[28] and classified species as tropical or temperate using

their median latitude estimated from the Global Biodiversity Information Database

(GBIF, see Methods for details). We collated the BAMM and GBIF datasets to obtain

a dataset (herein 'full') of 28,057 species.

We found that speciation rates consistently increased towards the poles using several

ways of delimiting tropical and temperate species. Using correlation tests that control

for phylogenetic pseudoreplication [24], we found that tropical species had smaller

mean speciation rates (λ) than temperate species (Fig 1a; $\lambda_{temperate}$ = 1.24 species/myr,

$\lambda_{tropical}$ = 0.81 species/myr, p-value = 0.024). Angiosperms in all 10° latitudinal bands

outside the tropics (i.e. absolute median latitude >23.5°) speciated at faster rates than

in any tropical latitudinal band (Fig. 1b). The absolute median latitude occupied by

individual species was also positively correlated with their speciation rate (Fig. 1c;

Spearman's ρ = 0.128, p-value = 0.013). These results were maintained when we: *i*)

analysed only those species that were densely sampled in GBIF; *ii*) compared strictly

tropical and strictly temperate species (i.e., discarding broadly ranging species that

occurred both inside and outside the tropics); and *iii*) discarded the highest latitudinal

bands (>50° N and S), which contained very few species (Table S1).

Figure 1. Temperate species have higher speciation rates (λ) than tropical

species. a) Rank ordered distribution and boxplot (inset) of λ inferred with BAMM

for tropical (red) and temperate (blue) species. * indicates significant difference (p-

value = 0.024) between the two groups, assessed with a Mann-Whitney test

implemented in STRAPP. **b)** λ grouped by latitudinal band of each species. Notches

in boxplots indicate 95% confidence intervals around median, denoted by thick

vertical lines and boxes span the interquartile ranges. **c)** Spearman's ρ correlation

between species absolute median latitude and λ as estimated with STRAPP. Grey

lines are correlations across 1000 samples of the posterior distribution estimated with

BAMM. Inset shows the difference between the empirical and null correlations

estimated with 1000 permutations of evolutionary rates across the phylogeny.

Our full dataset contained a disproportionate amount of temperate species but this did

not explain the consistent association between latitude and speciation rates. The initial

GBIF dataset was consistent with the LDG, containing 42.5% and 57.5% temperate

and tropical species, respectively (Fig S1a-c). However, the data were

disproportionately represented by temperate species (61.7%) after merging GBIF

distributions with the macroevolutionary rates (Fig. S1d-f). We assessed if this

latitudinal bias could affect our results in two ways. First, we generated 100 random

samples of our full dataset (herein 'unbiased' datasets), each with 10,000 species that

maintained the proportion of tropical (57.5%) and temperate (42.5%) species present

in GBIF. Second, we generated another set of 100 random samples (herein 'extreme

tropical' datasets), where the proportions of temperate and tropical species were

29.1% and 70.9% respectively. These were calculated assuming that all species absent

from GBIF were tropical and therefore represent the most extreme scenario of

latitudinal bias. We then assessed whether the λ estimates were affected by the poorer

sampling for tropical species, which may result in average longer branches and lower

λ. We reran BAMM for 10 of the 100 unbiased and extreme tropical datasets and

compared the resulting estimates with the ones from the full dataset (n = 28,057

species). We found that the λ estimates from the full and the subsampled datasets

were very positively correlated (median $\rho = 0.82$, median p-value < 0.001 and median

$\rho = 0.83$, median p-value < 0.001 for the unbiased and the extreme tropical datasets,

respectively), and that this association did not differ between tropical and temperate

species (Figs S2, S3). We also confirmed that the association of λ and latitude was not

a result of the excessive proportion of temperate species in the full dataset by

repeating the STRAPP analyses subsetted to the species present in each of the 100

unbiased and extreme tropical datasets. Our conclusions were identical between the

full and both the unbiased (Fig S4) and extreme tropical datasets (Fig S5). Finally, we

confirmed that our results could not be explained by different levels of sampling for

BAMM between temperate and tropical species. We found that the relationship of

median latitude and BAMM sampling fraction was positive but small (Fig S6).

Moreover, densely sampled clades, which occurred at higher latitudes, tended to have

smaller estimates of λ (Fig. S7), contrary to what we might expect if our results were

a sampling artefact. These analyses collectively exclude the conclusion that our

results stem from lower estimates of speciation rates in more sparsely sampled

tropical clades.

We also confirmed that the speciation rate estimates were robust despite the low

levels of sampling of the angiosperm ToL. Our full dataset contained 8% of the

angiosperm species described in The Plant List[30]. To assess the effect of missing taxa,

we randomly sampled 8% of the species (n = 2,273 species) in our full dataset ten

times and reran BAMM on these 'small' datasets. As above, the λ estimates from the

full and the small datasets were positively correlated (median ρ = 0.66, median p-

value < 0.001), and this association did not differ between tropical and temperate

species (Fig. S8).

The positive association between speciation rates and latitude was consistently

retrieved using two alternative methods for estimating macroevolutionary rates. First,

we used FiSSE (Fast, intuitive State-dependent Speciation-Extinction analysis[31]), a

non-parametric test that determines the effect of a binary trait on speciation rates. We

found a positive association of speciation rate estimated with FiSSE (DR) and

whether species were temperate with our full dataset (median $DR_{temperate}$ = 0.29

species/myr, median $DR_{tropical}$ = 0.18 species/myr, p-value = 0.001). To test whether

the excess of temperate species influenced this result, we repeated the analyses in the

100 unbiased datasets of 10,000 species and found no significant differences

($DR_{temperate}$ = 0.15 species/myr, $DR_{tropical}$ = 0.13 species/myr, p-value = 0.168).

However, simulations suggested that this non-significant result may stem from a lack

of power with FiSSE for our given phylogenetic and trait data (Fig S9, see Methods).

FiSSE, unlike BAMM, also does not account for incomplete sampling in the

speciation rate estimates, so we caution against strong interpretations of the results

obtained with this method with the present levels of missing taxa. Second, we

analysed the association between latitude and speciation rate across clades of different

ages in the angiosperm ToL. Following ref. 26, we defined six 4 million-year-wide

time slices, from 0 to 24 million years (myr), which were used to select clades where

speciation rates could confidently be estimated (i.e., probability of recovering the

crown age of lineages was > 70% following ref. 30) and GBIF coverage was > 50%.

We then fitted models that implemented time-variable and time-constant functions for

speciation and extinction[32], and selected the best fitting model to obtain clade-based

estimates of speciation rates. Mirroring the BAMM results, we found that clades with

a larger proportion of temperate species had higher rates of speciation (Fig S10).

Overall, our findings do not support the long-standing notion that the greater species

diversity of the tropics can be explained by higher rates of speciation. We instead

found higher rates of recent speciation closer to the poles. As temperate biotas have

fewer species, their niche space may be less saturated, thereby increasing

opportunities for lineage divergence[15,33]. Similarly, reproductive isolation may be

elevated at higher latitudes by the greater ecological opportunity generated from

recurrent environmental change and climate instability[34]. Other explanations may be

related to the ecological and life history traits of temperate species. For example,

faster speciation rates at higher latitudes could stem from the higher frequency of

small-seeded species[35]. Small seed size is positively correlated with angiosperm

diversification[26], and temperate species did have smaller seeds in our full dataset (see

Methods, phylANOVA: p value = 0.001, significance assessed with 1,000 random

simulations with phytools[36]). Therefore, our results suggest that latitudinal

differences in speciation rate do not shape the LDG but are shaped by differences in

species diversity and traits[16,37].

Rejecting the role of speciation rate as the cause of the LDG calls for alternative

explanations[25]. From a macroevolutionary perspective, greater tropical species

diversity may arise for at least three reasons[38]: *i*) higher rates of diversification (i.e.,

speciation minus extinction), *ii*) higher rates of net migration, and *iii*) greater potential

for historical colonisation. First, the elevated speciation near the poles could be

coupled with higher extinction rates to produce a scenario where temperate zones act

as both a "cradle" and "grave" of biodiversity[34]. Net diversification might therefore

still be higher in tropical areas despite the absolute rates of speciation and extinction

each being smaller than closer to the poles. However, the lack of reliable extinction

rate estimates from molecular phylogenies in the absence of fossil data[19,39]

complicates a direct test of this hypothesis. Second, the tropical conservatism

hypothesis proposes that lower net migration from the tropics to temperate zones

contributes to the LDG[40]: most lineages originated in the tropics and have been unable

to disperse into harsher environments. Third, if rates of diversification and migration

have no latitudinal variation, the greater diversity of the tropics may stem from their

larger area through time (age-and-area hypothesis[41]).

An alternative to these ideas is that the LDG may simply have nothing to do with

macroevolution. A whole class of hypotheses posits that there are ecological limits to

the number of species that a region can hold, and that these limits vary with latitude[42].

For example, the larger diversity in the tropics could be explained by expanded niche

spaces and greater niche packing[43].

Recent findings have suggested that rates of species origination are highest in species-

poor areas[17,44]. Our results are consistent with these findings and therefore suggest

that latitudinal variation in speciation rates is not a major engine of global diversity

patterns in one of the largest eukaryotic radiations. Using time-heterogeneous

estimates of speciation, we were able to exclude faster rates of evolution, such as

arising from more favourable environmental conditions or more intense biotic

interactions, as responsible for shaping the angiosperm LDG. Future work must now

resolve whether recent rapid speciation nearer the poles is widespread across the ToL

and its underlying causes.

Methods

Latitudinal and phylogenetic datasets

We obtained latitudinal data for 236,894 angiosperms that included the median,

maximum and minimum latitude for each species[28]. We then intersected these data

with speciation rates estimated with BAMM v.2.5.0[27] for 29,703 angiosperms[26]. The

input phylogeny was obtained from a time-calibrated phylogenetic tree for 31,389

land plant species[45,46]. BAMM detects heterogeneity in speciation and extinction rates

through time and across lineages. Non-random incomplete taxon sampling was

incorporated into the estimation process by calculating the number of species sampled

in each family.

The species names in the GBIF and BAMM datasets were standardised with The

Plant List (TPL) using the *Taxonstand*[47] and *Taxonlookup*[48] packages before collation,

resulting in 28,057 species in our full dataset. Tropical and temperate species were

defined as those where the absolute median latitude was below and above 23.5°,

respectively. Species were also grouped into 10° bins according to their median

latitude to define latitudinal bands (starting from a band centered in the Equator from

-5° to 5°, then from 5° to 15, etc). We repeated the analyses using only densely

sampled species (i.e., with five or more data points in the GBIF dataset). Strictly

tropical and temperate species were defined as species with median, maximum and

minimum latitude occurring inside and outside the tropics, respectively. We discarded

widespread species (i.e., neither strictly tropical nor strictly temperate) for analyses

comparing strict tropical and temperate taxa (retaining 20,766 species). Finally, we

discarded data with median latitudes in the highest latitudinal bands (>50° and <-50°),

which contained very few species with extreme estimates of λ (retaining 25,189

species).

BAMM-based analyses

We correlated BAMM speciation rates and latitudinal data using Structured Rate

Permutations on Phylogenies (STRAPP)[29]. STRAPP assesses the significance of the

empirical association of macroevolutionary rates and phenotypic traits by comparing

it to a null distribution generated by permuting the speciation rates across the

phylogeny while maintaining the position of the rate shifts in the phylogenetic tree.

We performed all STRAPP two-tailed tests using the *traitdependentBAMM* function

of the BAMMtools package[49] with 1000 replicates and logging the rates. We assessed

whether *i*) tropical and temperate species had different speciation rates using a Mann

Whitney U test; *ii*) speciation rates for each of the latitudinal bands were different

using the Kruskal-Wallis rank sum statistic; and *iii*) species absolute median latitude

was correlated with its speciation rate using Spearman's rank correlation coefficient.

We analysed the potential effect of the disproportionate amount of temperate species

in our full dataset using BAMM. Subsampled datasets were consistent with the LDG

and either *i*) maintained the proportions of tropical (57.5%) and temperate (42.5%)

taxa observed in the GBIF dataset ('unbiased'); or *ii*) assumed that the species absent

from the GBIF dataset from ref. 28 were all tropical ('extreme tropical') (n = 109,461

species; obtained from subtracting the number of species in The Plant List according

to *TaxonLookup*[48] minus the species in the GBIF dataset). Both subsampled datasets

were generated to assess the effect of the excess of temperate species in our full

dataset, but the extreme tropical dataset represented the most extreme scenario of the

effect of undescribed biodiversity in our analyses. The size of these datasets was fixed

at 10,000 species so that the tropical species present in each subsample, which were

obtained by random draws of 5,750 and 7,091 species for the unbiased and extreme

tropical respectively from a total pool of 10,749 species, were not excessively

redundant. We selected 10 random replicates of the 10,000-species subsampled

datasets and reran the BAMM analyses as detailed in ref. 26. We then assessed

whether the relationship of the full (n=28,057 species) and both the unbiased and

extreme tropical λ estimates was different for tropical and temperate species. To do

this, we predicted the full λ estimates using the subsampled λ estimates, 'tropicality'

(i.e. a binary variable indicating if a species was tropical based on our aforementioned

definition) and the interaction of the λ estimates and tropicality in each of the 10

replicates for the unbiased and extreme tropical datasets. We incorporated the effect

of the phylogeny using the *phylolm* package[50]. We found that the effect size of the

interaction of tropicality and the λ estimates was not different from zero across all the

10 replicates for the unbiased (mean = 0.0009, standard deviation = 0.0015) and the

extreme tropical (mean = 0.0004, standard deviation = 0.0016) datasets. This result

shows that there is no difference in the correlation between λ of the full and

subsampled datasets between temperate and tropical lineages.

We also confirmed that our results could not be explained by latitudinal variation in

the clade sampling fractions of our BAMM analysis. We found a positive relationship

(slope = 0.059, p-value < 0.0001) between the species median latitude and its

sampling fraction (i.e. the percentage of species in the same family present in our

28,057 species phylogenetic tree). This result meant that a 10° increase in median

latitude corresponded with an average of 0.59% increase in sampling fraction.

Furthermore, we found that there was a negative relationship between sampling

fraction and the log-transformed estimated speciation rate (slope = -0.055, p-value <

0.0001). Again, this effect size was small. For example, an increase in sampling

fraction from 20% to 30% corresponded with an average decrease of 0.33

lineages/myr in the estimated speciation rate.

Our full dataset contained λ estimates for 28,057 species, which represent 8% of

described angiosperm species (total n = 346,365 calculated with *Taxonlookup*). To

assess the reliability of λ estimates given the large number of missing taxa, we

randomly generated 10 subsamples ('small' datasets) with 8% of the species in our

full dataset (n = 2,273). As above, we ran BAMM for each of the 10 subsamples and

predicted the full λ estimates using the small λ estimates. Similar to the unbiased and

extreme tropical datasets, we found that the effect size of the interaction of tropicality

and λ was not different from zero across all the 10 replicates (mean = -0.0033,

standard deviation = 0.0189).

Although the reliability of BAMM estimates has recently been questioned[51],

simulations have shown that accurate estimates of speciation can be obtained even in

the absence of fossil data[39,52]. Tip-based estimates of speciation like the ones we use

here should also be robust to the effect of extinction and are increasingly used when

analysing the effect of traits on lineage diversification[20]. Extinction estimates are

thought to be more unreliable[19,53,54] so we do not discuss them here.

FiSSE-based analyses

FiSSE analyses the correlation of a binary trait and diversification and has shown to

have a lower type I error than similar trait-dependent diversification methods[31]. It

calculates the speciation rate associated with each trait state and is based on the

diversification rate metric (DR[55]), which was originally proposed to estimate net

diversification but has since been shown to approximate speciation rates better[20]. The

distributions of DR for each trait value (i.e. tropical and temperate) had long right-

tails and the means were much larger than the medians, so the latter measure better

described the central tendency of the DR distributions. Therefore, following ref. 56

we modified the FiSSE function provided in ref. 31 to compare the median

diversification rate for each trait value instead of the mean.

We also assessed whether the lack of significant differences between tropical and

temperate speciation rates that we found with the 100 unbiased datasets could be

explained by their sample size (n=10,000 species). We generated 100 replicates of the

full dataset of 2,000; 4,000; 6,000; 8,000; 10,000; and 12,000 species and reran FiSSE

for each of them. We then ran an exponential regression with the size of a dataset as a

predictor and the logged p-value of the association of speciation and tropicality as a

response.

Clade-based analyses

We estimated clade-level measures of speciation across the angiosperm phylogenetic

tree. We used a set of 4 million-year-wide time intervals from 0 to 24 myr to define

the ages of the clades. Clades were delimited as the largest non-overlapping

monophyletic groups of four or more species where the sampling was larger than

30%. This criterion ensures that the probability of retrieving the correct crown age of

the clade is above 70%[30]. At least 50% of the species in the clades also had to have

latitudinal data. We used RPANDA (R: Phylogenetic ANalyses of DiversificAtion[32])

to fit six different models of diversification: *i*) constant speciation (λ) and extinction

(μ) fixed at 0; *ii*) constant λ and μ; *iii*) exponential λ and μ fixed at 0; *iv*) exponential λ

and constant μ; *v*) constant λ and exponential μ; and *vi*) exponential λ and μ.

Incomplete sampling within clades was incorporated into the estimates by calculating

the proportion of sampled species within the genera in each clade. Speciation rates

were estimated using model averaging with AIC weights. The correlation of the

clade-level estimates of speciation and the proportion of temperate and tropical

species in each clade was assessed using phylogenetic least squares regression as

implemented in the R package *nlme*[57]. The lambda parameter was optimised using a

maximum likelihood approach. We weighted the regressions by the proportion of

GBIF data points in each clade to account for the effect of different sampling levels of

distribution data. We then confirmed that potential latitudinal differences in sampling

proportions could not explain our results. To do that, we analysed if the sampling

proportions of largely tropical and largely temperate clades (defined as clades where

>50% of the species were tropical and temperate, respectively) across the 6 time

intervals were different using Pearson's chi-squared test. We found that temperate

clades had significantly denser average sampling only in the 16-20 (sampling

proportion $_{tropical}$ = 0.65, sampling proportion $_{temperate}$ = 0.56) and 20-24 (sampling

proportion $_{tropical}$ = 0.69, sampling proportion $_{temperate}$ = 0.57) myr intervals.

Latitudinal variation of seed size

Smaller seeded species have been shown to speciate faster than large seeded species[26].

We assessed whether seed size differed between temperate and tropical lineages as it

has previously been shown to do so[35], and so latitudinal variation in seed size could

explain the differences in speciation between tropical and temperate species. We

obtained seed size measurements for 13,127 species in our full dataset from a

previous study[26]. The resulting dataset was disproportionately temperate (76.04% of

the species were temperate). Average seed sizes were larger in tropical than in

temperate species (mean seed size$_{tropical}$ = 0.149 g; mean seed size$_{temperate}$ = 0.065 g; t =

-298.7; df = 4189.2; p-value < 0.001). This difference remained when phylogeny was

considered (phylANOVA: t = 35.293; p value = 0.001; significance assessed with

1,000 random simulations).

Data availability

Scripts to reproduce the analyses will be deposited upon acceptance in figshare

Author contributions

J.I. conceived the study and performed the analysis; J.I. and A.J.T interpreted the

analysis and wrote the manuscript.

Acknowledgments

We thank the Wellcome Trust (Grant Number 105602/Z/14/Z), Gatsby Charitable

Foundation (Grant Number GAT2962) and Isaac Newton Trust (Grant Number

17.24r) for funding. The authors declare no competing interests.

Declaration of interests

The authors declare no competing interests.

Supplementary Figures and Tables

dataset name	$\lambda_{temperate}$	$\lambda_{tropical}$	p-value	dataset size
more.GBIF.data	1.242	0.802	0.014	26,568
no.widespread	1.337	0.802	0.013	20,766
no.high.latitude	1.207	0.813	0.030	25,819

Table S1. Temperate species have mean higher speciation rates (λ, estimated with

STRAPP) than tropical species across different datasets: "more.GBIF.data" excludes

species with less than five GBIF data points; "no.widespread" excludes species

occurring both in and outside the tropics; "no.high.latitude" excludes species from

poorly sampled latitudinal bands with absolute median latitude ≥50°. Units for λ and

dataset size are lineages/myr and species, respectively.

Figure S1. The full dataset had a disproportionate number of temperate species. **a)**

Proportion of tropical and temperate species in the GBIF dataset (n = 236,894); **b)**

proportion of species in each latitudinal band in the GBIF dataset; **c)** number of

species in each latitudinal band in the GBIF dataset; **d)** proportion of tropical and

temperate species in the full dataset (n = 28,057); **e)** proportion of species in each

latitudinal band in the full dataset; **f)** number of species in each latitudinal band in the

full dataset; in **b), c), e)** and **f)**, the dotted vertical line denotes the 23.5° threshold

between tropical and temperate zones.

Figure S2. BAMM λ estimates were positively correlated in the full and 10 unbiased

datasets with no effect of tropicality on this correlation. **a) - j)** Solid lines are

phylogenetic linear regressions predicting λ in the full tree (n = 28,057 species) with λ

in the unbiased tree (n = 10,000 species) in temperate (shown in blue) and tropical

(shown in red) species. Shaded areas indicate the 95% confidence intervals.

Figure S3. BAMM λ estimates were positively correlated in the full and 10 extreme

tropical datasets with no effect of tropicality on this correlation. **a) - j)** Solid lines are

phylogenetic linear regressions predicting λ in the full tree (n = 28,057 species) with λ

in the extreme tropical tree (n = 10,000 species) in temperate (shown in blue) and

tropical (shown in red) species. Shaded areas indicate the 95% confidence intervals.

Figure S4. Temperate species have higher speciation rates (λ) than tropical species

for 100 subsampled "unbiased" datasets with 42.5% of temperate and 57.5% of

tropical species. **a)** Rank ordered distribution of λ inferred with BAMM for tropical

(red) and temperate (blue) species. **b)** λ grouped by latitudinal band of each species.

Boxes show the average interquartile range across the 100 subsampled datasets, and

lines show the average medians. **c)** Spearman's ρ correlation of species absolute

median latitude and λ as estimated with STRAPP. Solid black line indicates median

correlation ± 95% confidence interval across the posterior distribution. The inset

shows the median (black line) ± 95% confidence interval for the difference between

the empirical and null correlations estimated with 1000 permutations of evolutionary

rates across the phylogeny in each of 100 subsampled datasets.

Figure S5. Temperate species have higher speciation rates (λ) than tropical species

for 100 subsampled "extreme tropical" datasets with 29.1% temperate and 70.9%

tropical species. **a)** Rank ordered distribution of λ inferred with BAMM for tropical

(red) and temperate (blue) species. **b)** λ grouped by latitudinal band of each species.

Boxes and lines are presented as in Fig. S4. **c)** Spearman's ρ correlation of species

absolute median latitude and λ as estimated with STRAPP. Lines and inset shown as

in Fig. S4

Figure S6. Relationship between the species absolute median latitude and the family-

level sampling fraction used in the BAMM analyses. The red line is the slope of the

linear regression (slope = 0.059, p-value < 0.0001).

Figure S7. More densely sampled clades have smaller estimates of speciation rate (λ).

The dotted line is the slope of the linear regression of the log(λ) and the BAMM

sampling fraction (slope = -0.055 , p-value < 0.0001).

475 **Figure S8.** BAMM λ estimates are positively correlated in the full and 10 small

476 datasets and tropicality has no effect on this correlation. **a) - j)** Solid lines are

477 phylogenetic linear regressions predicting λ in the full tree (n = 28,057 species) with λ

478 in the small tree (n = 2,273 species) in temperate (shown in blue) and tropical (shown

479 in red) species. Shaded areas indicate the 95% confidence intervals.

Figure S9. FiSSE power analysis. Each boxplot shows the p-values for the difference

of $DR_{temperate}$ and $DR_{tropical}$ for 100 subsamples of varying size taken from the full

dataset (n = 28,057 species) whilst retaining proportions of tropical and temperate

species observed in the GBIF dataset. Notches in boxplots indicate 95% confidence

intervals around the median, denoted by thick vertical lines. Boxes indicate the

interquartile range (IQR) and whiskers extend to the most extreme data points that are

not further away than ±1.5×IQR. The solid line is an exponential model predicting the

logged mean p-value ± 95% confidence interval (shaded area) from the size of the

dataset. The dotted line indicates p-value = 0.05.

Figure S10. Temperate clades have higher speciation rates in the clade-based

analysis. Correlation of a) the proportion of tropical and b) temperate species in each

clade with the corresponding speciation rate (λ) estimated with RPANDA. The

correlation coefficient is the phylogenetic generalised least squares (PGLS) slope. The

size of the points is scaled to the number of clades in each time interval and their

colours show the statistical significance of the slope.

References

References

1. von Humboldt, A. *Ansichten der Natur mit wissenschaftlichen*

Erlauterungen. (J.G. Cotta, 1808).

2. Wallace, A. R. *Tropical nature and other essays*. (Macmillan, 1878).

3. Pianka, E. R. Latitudinal gradients in species diversity: a review of

concepts. *Am. Nat.* **100**, 33–46 (1966).

4. Hawkins, B. A. Ecology's oldest pattern? *Trends Ecol. Evol.* **16**, 470

(2001).

5. Mittelbach, G. G. *et al.* Evolution and the latitudinal diversity gradient:

speciation, extinction and biogeography. *Ecol. Lett.* **10**, 315–331 (2007).

6. Dowle, E. J., Morgan-Richards, M. & Trewick, S. A. Molecular evolution

and the latitudinal biodiversity gradient. *Heredity* **110**, 501–510 (2013).

7. Rohde, K. Latitudinal gradients in species diversity: the search for the

primary cause. *Oikos* 514–527 (1992).

8. Brown, J. H. Why are there so many species in the tropics? *J. Biogeogr.*

41, 8–22 (2014).

9. Rolland, J., Condamine, F. L., Jiguet, F. & Morlon, H. Faster speciation

and reduced extinction in the tropics contribute to the mammalian

latitudinal diversity gradient. *PLoS Biol.* **12**, e1001775 (2014).

10. Pyron, R. A. & Wiens, J. J. Large-scale phylogenetic analyses reveal the

causes of high tropical amphibian diversity. *Proc. R. Soc. Lond. B Biol.*

Sci. **280**, 20131622 (2013).

11. Condamine, F. L., Sperling, F. A., Wahlberg, N., Rasplus, J.-Y. & Kergoat,

G. J. What causes latitudinal gradients in species diversity? Evolutionary

processes and ecological constraints on swallowtail biodiversity. *Ecol.*

Lett. **15**, 267–277 (2012).

12. Rabosky, D. L. & Huang, H. Minimal effects of latitude on present-day

speciation rates in New World birds. *Proc R Soc B* **282**, 20142889 (2015).

13. Economo, E. P., Narula, N., Friedman, N. R., Weiser, M. D. & Guénard, B.

Macroecology and macroevolution of the latitudinal diversity gradient in

ants. *Nat. Commun.* **9**, 1778 (2018).

14. Weir, J. T. & Schluter, D. The latitudinal gradient in recent speciation and

extinction rates of birds and mammals. *Science* **315**, 1574–1576 (2007).

15. Schluter, D. Speciation, ecological opportunity, and latitude: (American

Society of Naturalists address). *Am. Nat.* **187**, 1–18 (2016).

16. Schluter, D. & Pennell, M. W. Speciation gradients and the distribution of

biodiversity. *Nature* **546**, 48 (2017).

17. Rabosky, D. L. *et al.* An inverse latitudinal gradient in speciation rate for

marine fishes. *Nature* **559**, 392–395 (2018).

18. Nee, S., Holmes, E. C., May, R. M. & Harvey, P. H. Extinction rates can be

estimated from molecular phylogenies. *Philos. Trans. R. Soc. Lond. B.*

Biol. Sci. **344**, 77–82 (1994).

19. Marshall, C. R. Five palaeobiological laws needed to understand the

evolution of the living biota. *Nat. Ecol. Evol.* **1**, 0165 (2017).

20. Title, P. O. & Rabosky, D. L. Tip rates, phylogenies and diversification:

What are we estimating, and how good are the estimates? *Methods Ecol.*

Evol. **0**,

21. Francis, A. P. & Currie, D. J. A globally consistent richness-climate

relationship for angiosperms. *Am. Nat.* **161**, 523–536 (2003).

22. Jansson, R. & Davies, T. J. Global variation in diversification rates of

flowering plants: energy vs. climate change. *Ecol. Lett.* **11**, 173–183

(2008).

23. Svenning, J.-C., Borchsenius, F., Bjorholm, S. & Balslev, H. High tropical

net diversification drives the New World latitudinal gradient in palm

(Arecaceae) species richness. *J. Biogeogr.* **35**, 394–406 (2008).

24. Jansson, R., Rodríguez-Castañeda, G. & Harding, L. E. What can multiple

phylogenies say about the latitudinal diversity gradient? A new look at the

tropical conservatism, out of the tropics, and diversification rate

hypotheses. *Evolution* **67**, 1741–1755 (2013).

25. Antonelli, A. *et al.* An engine for global plant diversity: highest evolutionary

turnover and emigration in the American tropics. *Front. Genet.* **6**, (2015).

26. Igea, J., Miller, E. F., Papadopulos, A. S. & Tanentzap, A. J. Seed size

and its rate of evolution correlate with species diversification across

angiosperms. *PLoS Biol.* **15**, e2002792 (2017).

27. Rabosky, D. L. Automatic detection of key innovations, rate shifts, and

diversity-dependence on phylogenetic trees. *PloS One* **9**, e89543 (2014).

28. Mounce, R., Smith, P. & Brockington, S. Ex situ conservation of plant

diversity in the world's botanic gardens. *Nat. Plants* **3**, 795 (2017).

29. Rabosky, D. L. & Huang, H. A robust semi-parametric test for detecting

trait-dependent diversification. *Syst. Biol.* **65**, 181–193 (2016).

30. Rabosky, D. L. Challenges in the estimation of extinction from molecular

phylogenies: A response to Beaulieu and O'Meara. *Evolution* **70**, 218–228

(2016).

31. Rabosky, D. L. & Goldberg, E. E. FiSSE: A simple nonparametric test for
the effects of a binary character on lineage diversification rates. *Evol. Int.*
J. Org. Evol. **71**, 1432–1442 (2017).

32. Morlon, H. *et al.* RPANDA: an R package for macroevolutionary analyses
on phylogenetic trees. *Methods Ecol. Evol.* **7**, 589–597 (2016).

33. Simpson, G. *The major features of evolution*. (Columbia University Press,
1953).

34. Cutter, A. D. & Gray, J. C. Ephemeral ecological speciation and the
latitudinal biodiversity gradient. *Evolution* **70**, 2171–2185 (2016).

35. Moles, A. T. *et al.* Global patterns in seed size. *Glob. Ecol. Biogeogr.* **16**,
109–116 (2007).

36. Revell, L. J. phytools: An R package for phylogenetic comparative biology
(and other things). *Methods Ecol. Evol.* **3**, 217–223 (2012).

37. Weir, J. T. & Price, T. D. Limits to speciation inferred from times to
secondary sympatry and ages of hybridizing species along a latitudinal
gradient. *Am. Nat.* **177**, 462–469 (2011).

38. Igea, J. & Tanentzap, A. J. Multiple macroevolutionary routes to becoming
a biodiversity hotspot. *Sci. Adv.* **5**, eaau8067 (2019).

39. Mitchell, J. S., Etienne, R. S. & Rabosky, D. L. Inferring diversification rate
variation from phylogenies with fossils. *Syst. Biol.* syy035 (2018).

40. Wiens, J. J. & Donoghue, M. J. Historical biogeography, ecology and
species richness. *Trends Ecol. Evol.* **19**, 639–644 (2004).

41. Fine, P. V. A. & Ree, R. H. Evidence for a time-Integrated species-area
effect on the latitudinal gradient in tree diversity. *Am. Nat.* **168**, 796–804
(2006).

42. Rabosky, D. L. Ecological limits and diversification rate: alternative

paradigms to explain the variation in species richness among clades and

regions. *Ecol. Lett.* **12**, 735–743 (2009).

43. Pellissier, V., Barnagaud, J.-Y., Kissling, W. D., Şekercioğlu, Ç. &

Svenning, J.-C. Niche packing and expansion account for species

richness–productivity relationships in global bird assemblages. *Glob. Ecol.*

Biogeogr. **27**, 604–615 (2018).

44. Quintero, I. & Jetz, W. Global elevational diversity and diversification of

birds. *Nature* **555**, 246–250 (2018).

45. Zanne, A. E. *et al.* Three keys to the radiation of angiosperms into freezing

environments. **506**, 89–92 (2014).

46. Qian, H. & Jin, Y. An updated megaphylogeny of plants, a tool for

generating plant phylogenies and an analysis of phylogenetic community

structure. *J. Plant Ecol.* **9**, 233–239 (2016).

47. Cayuela, L., Cerda, Í. G. la, Albuquerque, F. S. & Golicher, D. J.

taxonstand: An r package for species names standardisation in vegetation

databases. *Methods Ecol. Evol.* **3**, 1078–1083 (2012).

48. Pennell, M. W., FitzJohn, R. G. & Cornwell, W. K. A simple approach for

maximizing the overlap of phylogenetic and comparative data. *Methods*

Ecol. Evol. **7**, 751–758 (2016).

49. Rabosky, D. L. *et al.* BAMMtools: an R package for the analysis of

evolutionary dynamics on phylogenetic trees. *Methods Ecol. Evol.* **5**, 701–

707 (2014).

50. Tung Ho, L. si & Ané, C. A linear-time algorithm for gaussian and non-

gaussian trait evolution models. *Syst. Biol.* **63**, 397–408 (2014).

51. Moore, B. R., Höhna, S., May, M. R., Rannala, B. & Huelsenbeck, J. P.

Critically evaluating the theory and performance of Bayesian analysis of

macroevolutionary mixtures. *Proc. Natl. Acad. Sci.* **113**, 9569–9574

(2016).

52. Rabosky, D. L., Mitchell, J. S. & Chang, J. Is BAMM flawed? Theoretical

and practical concerns in the analysis of multi-rate diversification models.

Syst. Biol. **66**, 477–498 (2017).

53. Rabosky, D. L. Extinction rates should not be estimated from molecular

phylogenies. *Evolution* **64**, 1816–1824 (2010).

54. Mitchell, J. S., Etienne, R. S. & Rabosky, D. L. Inferring diversification rate

variation from phylogenies with fossils. *Syst. Biol.* **68**, 1–18 (2019).

55. Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K. & Mooers, A. O. The

global diversity of birds in space and time. *Nature* **491**, 444 (2012).

56. Faurby, S. & Antonelli, A. Evolutionary and ecological success is

decoupled in mammals. *J. Biogeogr.* **45**, 2227–2237 (2018).

57. Pinheiro, J., Bates, D., DebRoy, S. & Sarkar, D. R Core Team (2014)

nlme: linear and nonlinear mixed effects models. R package version 3.1-

117. *available at https://CRAN.R-project.org/package=nlme.* (2014).