Giraffe Translocation Population Viability Analysis

DEREK E. LEE,[1] Biology Department, Pennsylvania State University, University Park,

Pennsylvania, USA

ELMAR FIENIEG, European Association of Zoos and Aquaria, Amsterdam, The Netherlands

COCK VAN OOSTERHOUT, School of Environmental Sciences, University of East Anglia,

Norwich Research Park, Norwich, UK

ZOE MULLER, School of Biological Sciences, University of Bristol, Bristol, UK

MEGAN STRAUSS, Wild Nature Institute, Concord, New Hampshire, USA

KERRYN D. CARTER, Elephant Connection, Mwandi, Western Province, Zambia

CISKA P.J. SCHEIJEN, Animal, Wildlife and Grassland Sciences, University of the Free State,

Bloemfontein, Free State, South Africa

FRANCOIS DEACON, Animal, Wildlife and Grassland Sciences, University of the Free State,

Bloemfontein, Free State, South Africa

ABSTRACT Most populations of giraffes have declined in recent decades, leading to the recent

decision to upgrade the species to vulnerable status, and some subspecies to endangered.

[1] *Email:* dereklee@psu.edu

Translocations have been used as a conservation tool to re-introduce giraffes to previously

occupied areas or establish new populations, but guidelines for founding populations are lacking.

To provide general guidelines for translocation projects regarding feasibility, we simulated

various scenarios of translocated giraffe populations to identify viable age and sex distributions

of founding populations using Population Viability Analysis (PVA) implemented in Vortex

software. We explored the parameter space for demography (population growth rates: $\lambda = 1.001$,

1.010, 1.024), and the genetic load (number of lethal equivalents: LE = 2.5, 6.29, 12.6),

examining how variation in founding numbers (N = 5 to 80 females) and sex ratios (M:F = 0.1 to

0.5) affected 100-year probability of extinction and genetic diversity. We found that even very

small numbers of founders (N ≤10 females) can appear to be successful in the first decades due

to transient positive population growth, but with moderate population growth rate and moderate

genetic load, long-term population viability (probability of extinction <0.01) was only achieved

with ≥30 females and ≥3 males released. To maintain >95% genetic diversity of the source

population in an isolated population, 50 females and 5 males are recommended to comprise the

founding population. Sensitivity analyses revealed first-year survival and reproductive rate were

the simulation parameters with the greatest proportional influence on probability of extinction

and genetic diversity. These simulations highlight important considerations for translocation

success, and data gaps including true genetic load in wild giraffe populations.

KEYWORDS founding population, giraffe, population model, PVA, translocation, population

viability

Giraffes (*Giraffa camelopardalis*) are endemic African ruminant ungulates, and one of only a

handful of extant terrestrial megaherbivores (Owen-Smith 1988). Most populations of giraffes

have declined in recent decades, leading to the recent decision to upgrade the species to

vulnerable status, and some subspecies as endangered, on the IUCN Red List (Muller et al.

2018). Translocations have been used as conservation tools to re-introduce giraffes to previously

occupied areas or establish new populations (Winter et al. 2019), but quantitative guidelines for

establishing viable populations of translocated giraffes do not exist.

General guidelines for translocations suggest population modelling should be used to assist in the

assessment of project feasibility (IUCN 2013). Population modelling for translocation projects

can offer guidance on age- and sex-classes for founder populations by providing estimates of

population persistence, long-term viability, and genetic diversity (Seddon et al. 2007).

Population viability analysis (PVA) is commonly used to predict the likely future status of a

population and thus offers a quantitative basis for evaluating alternative management strategies

(Boyce 1992, Morris and Doak 2002). The availability of software programs such as Vortex has

facilitated the application of population modelling in re-introduction planning (e.g., Bustamante

1998) and post-release evaluation (e.g., Slotta-Bachmayret et al. 2004).

Our objective here is to identify the minimum size and sex ratio of a founding population that

would ensure long-term population viability and genetic diversity. We simulated various

scenarios of translocated giraffe populations using PVA implemented in Vortex software (Lacy

and Pollack 2017). We used stochastic, single-population, individual-based models to project

future population trajectories and estimate probability of extinction (PE), stochastic rate of

population increase (r), and genetic diversity (GD) at a hypothetically ideal prospective release

site under different founder population release scenarios. Scenarios varied the numbers of

females and males to release (Converse et al. 2013) and included 3 levels of inbreeding genetic

load under demographic rates simulating slow, moderate, and fast population growth rates. We

considered 4 criteria for defining successful translocations: 100-year probability of extinction PE

< 0.05 and PE < 0.01 (Morris and Doak 2002); and 100-year genetic diversity GD $> 80\%$ and

GD $> 95\%$ of the source population (Frankham et al. 2010). We also performed a sensitivity

analysis to determine which demographic and genetic load parameters had the greatest

proportional influence on PE and GD.

METHODS

PVA Scenarios

We constructed population models in program Vortex to simulate a hypothetical translocated

population of simulated individuals using demographic parameters from literature (Lee and

Strauss 2016, Dagg 2014) and publications of the Giraffe International Studbook (Lackey 2009).

Males bred from ages 2 to 25, and females bred from ages 3 to 29, the maximum age of survival

for both sexes was 30 years, females always produced 1 calf, and sex ratio at birth was equal

(Dagg 2014). Within the model parameters, offspring were dependent upon their mother for 1

year, meaning that if the mother died during the calf's first year, then the calf also died (Dagg

2014). All males aged 2 and above were in the breeding pool. Demographic rates are given in

Table 1.

To parameterize our PVA, we based our demographic rates on published observations of means

and variances of age-specific survival and fecundity for wild giraffe populations throughout

Africa (Lee and Strauss 2016), and data for reproductive longevity and inbreeding depression

from the global zoo population (Lackey 2009). We used data from IUCN Red List assessments

to compute mean population growth rates for 7 growing giraffe subpopulations, including some

translocations (Muller et al. 2018). Mean observed population growth rate was 1.024 (range

1.0045 to 1.035). Because the highest observed growth rates were from short time spans, they

were likely due to transient population dynamics and not the asymptotic growth rate. Therefore,

in our PVA we simulated populations with 3 different demographic parameterizations with

different asymptotic population growth rates (λ) that represented slow- (1.001), moderate-

(1.010), and fast-growing (1.024) populations (Lee and Strauss 2016, Muller et al. 2018). The

fastest population growth rate is probably only achievable in translocation destinations that have

few or no large predators (Lee and Strauss 2016, Muller et al. 2018).

Table 1. Demographic rates and variability (SD = standard deviation) used to

102   parameterize population simulations of giraffe translocations reflecting slow,

103   moderate, and fast population growth rates.

Population growth rate	Slow		Moderate		Fast	
λ	1.001		1.010		1.024	
Demographic rates	%	(SD)	%	(SD)	%	(SD)
Mortality Age 0-1	50	(10)	50	(10)	40	(10)
Mortality Age 1-2	20	(6)	20	(6)	20	(6)
Mortality Age >2	5	(3)	5	(3)	5	(3)
Reproduction	45	(10)	50	(10)	50	(10)

Given that few individuals typically are translocated on any given occasion, demographic

stochasticity, Allee effects, and inbreeding depression could all adversely affect long term

viability for translocated populations (O'Grady et al. 2006, Deredec and Courchamp 2007, Van

Houtan et al. 2010). We included stochasticity (demographic rates vary among individuals and

over time) and inbreeding depression (reduced reproduction and calf survival due to inbreeding)

in our simulations because these effects are likely to exist in giraffe populations (O'Grady et al.

2006, Lackey 2009, Lee and Strauss 2016, Lacy and Pollack 2017). We did not include Allee

effects (reduced demographic rates at small population size) because Allee effects are unlikely in

giraffe populations, especially in enclosed reserves where mates can easily locate each other.

We included 3 levels of inbreeding depression. Using data from the giraffe studbook, we

estimated the minimum genetic load as number of total lethal equivalents (LE) per individual

with a regression analysis of natural-log-transformed first-year survival on F-coefficient

according to Morton et al. (1956). The slope of the regression provided an estimate of the

reduction of fitness due to inbreeding and gave the lower bound approximation of the effective

number of LE per gamete, so we doubled the slope to estimate the low value for number of LE

per zygote (Lynch and Walsh 1998). We used Vortex inbreeding values representing a low

genetic load of 2.5 LE per individual (from our regression), a moderate load of 6.29 LE

(O'Grady et al. 2006, Nietlisbach et al. 2018), and a high genetic load of 12.6 LE (double the

moderate load). Fifty percent of LE are due to recessive lethal alleles. The other half of the LE

are due to overdominance (heterozygote superiority), and this genetic load cannot be removed by

selection (Lacy and Pollack 2017). We set demographic temporal correlation between mortality

and reproduction due to environmental variation at 0.5.

We assumed zero translocation-related mortality, so no additional mortality effect above normal

levels due to the process of capturing, relocating, and releasing. If mortality is expected during

the translocation process, our simulation results should be interpreted using the actual number of

successful live releases. We assumed zero post-release dispersal movements because many

translocations will likely be into fenced or otherwise constrained areas, and because we wanted

to keep track of every individual in the translocated population.

We projected 198 PVA scenarios. We simulated populations with various numbers of 2-year-old

females released (5, 10, 20, 30, 40, 50, 60, and 80 females) and different numbers of 2-year-old

males released to vary the sex ratio (SR) at release (SR = males / females, range = 0.1 to 0.5).

We simulated each combination of number of females and males at the 3 levels of asymptotic

growth rate, and 3 levels of genetic load.

For all scenarios we ran 1000 iterations of our stochastic model to project the populations for 100

years. We selected a projection time of 100 years because giraffes are a long-lived species and

we were most interested in the longer-term implications of translocation decisions, particularly

the effect of the number of individuals released. Extinction definition was $N < 2$. We assumed a

hypothetically ideal release site, but included effects of density dependence with a carrying

capacity of $N = 1000$. This was done because we were primarily interested in the effects of

demography and genetics without the confounding effects of habitat limitation, which varies

considerably between translocation sites. Forage quality and availability in all seasons should be

quantified at the release site to determine carrying capacity as part of a comprehensive pre-

translocation assessment. For each scenario we recorded output on PE, and GD (as observed

heterozygosity) in the final extant population. We provide guidelines for founding population

size and sex ratio for successful translocations with success defined 4 ways: probability of

extinction $PE < 0.05$ and $PE < 0.01$; and genetic diversity $GD > 80\%$ and $GD > 90\%$.

We performed a sensitivity analysis to determine which demographic or inbreeding parameter

was most influential to long-term viability by comparing outcomes from simulations that were

identical in every way except that we reduced a single demographic or inbreeding parameter by

approximately 25%. We simulated release of 30 females and 3 males using moderate

demographic rates (Table 1, middle columns) and moderate genetic load (LE = 6.29) as the

reference simulation. We then simulated 6 identical populations except each had a single change:

1) increased age of first reproduction by 1 year for males and females; 2) first year mortality =

63%; 3) second year mortality = 25%; 4) adult mortality = 6%; 5) reproduction = 38%; and 6)

LE = 7.9. The observed differences in PE and GD between the changed simulations and the

reference simulation indicates the sensitivity due to a proportionally similar change in each

demographic or genetic load parameter.

RESULTS

Population projections from all our scenarios followed the same general post-release trajectory.

Overall, there was an initial population decline for 1 year as some released individuals died

before maturity, followed by 27 years of rapid population growth as the founders and their

offspring reproduced, then a 1-year decline in year 29 as the last remaining founding animals

died. This was followed by a brief 3-year period of stable or positive population growth until the

first post-founder generation died and stable age distribution was achieved. Long-term (100-

year) viability depended on population growth rate and inbreeding genetic load, as well as which

success criteria were used, but minimum founding population varied from 10 to 60 females and 1

to 15 males (**Table 2**).

The 100-year PE declined as the number of released females, and sex ratio increased (**Fig. 1**).

Assuming fast population growth demographic rates and minimum inbreeding genetic load, the

100-year PE was below 0.05 when release included at least 10 females and 3 males (**Table 2**).

Under more conservative demographic rates ($\lambda = 1.001$) and realistic inbreeding genetic loads

(LE = 6.3 or 12.6), the minimum release population for 95% viability at 100 years was 30

females and 3–6 males (**Table 2**). Final GD increased with larger numbers of females and males

in the founding population (**Fig. 2**). To meet the success criteria of preserving 80% of GD,

minimum founding population sizes were similar to those required for the criteria of PE < 0.05,

i.e. 30 females and 3–6 males (**Table 2**).

Table 2. Minimum founding population sizes of female (F) and male (M) giraffes

for translocations to achieve 100-year population viability success under 4 criteria

of success, with 3 levels of intrinsic population growth rate and 3 levels of

inbreeding genetic load.

		Inbreeding genetic load (LE = lethal equivalents)		
Criteria	Population growth rate	Low (2.5 LE)	Mod (6.3 LE)	High (12.6 LE)
<0.05 Probability of Extinction	Slow (1.001)	20F 2M	30F 3M	30F 6M
	Mod (1.010)	20F 2M	20F 2M	30F 3M
	Fast (1.024)	10F 3M	10F 5M	20F 2M
<0.01 Probability of Extinction	Slow (1.001)	30F 3M	30F 6M	40F 5M
	Mod (1.010)	20F 2M	30F 3M	30F 9M
	Fast (1.024)	10F 5M	20F 2M	20F 5M
>80% Genetic Diversity	Slow (1.001)	20F 2M	20F 2M	30F 3M
	Mod (1.010)	20F 2M	20F 2M	20F 2M
	Fast (1.024)	10F 1M	10F 3M	20F 2M
>95% Genetic Diversity	Slow (1.001)	50F 15M	60F 12M	60F 18M
	Mod (1.010)	50F 5M	50F 5M	50F 10M
	Fast (1.024)	40F 5M	40F 5M	40F 5M

Figure 1. 100-year probability of extinction decreased as the number of released females

increased in population projections simulating translocations of different numbers of wild

giraffes. Simulations used population growth rates: $\lambda = 1.001, 1.010, 1.024$; and genetic load

number of lethal equivalents: LE = 2.5, 6.29, 12.6.

Figure 2. Gene diversity in final extant populations increased as the number of released females

increased in 100-year population projections simulating 198 translocations of wild giraffes.

Simulations used population growth rates: $\lambda = 1.001, 1.010, 1.024$; and genetic load number of

lethal equivalents: LE = 2.5, 6.29, 12.6.

Sensitivity analysis revealed demographic parameters of first-year survival and adult female

reproductive rate were the most influential on long-term viability metrics of PE and GD (Table

3). Changes in first-year mortality and reproductive rate both increased PE > 10% and reduced

GD > 10%. Increasing the age of first reproduction by 1 year resulted in a 1% increase in PE and

205   2% reduction in GD, and all other parameter changes resulted in < 1% reduction in PE and ≤ 2%

206   reduction in GD (Table 3).

208   **Table 3.** Sensitivity analysis of demographic and genetic load parameters for

209   simulations of giraffe translocations where each parameter was proportionally

210   reduced ~25%. ΔPE is change in probability of extinction relative to the reference

211   simulation, ΔGD is change in genetic diversity relative to the reference simulation.

Parameter	Value	ΔPE	ΔGD
First-year mortality	63%	0.114	-0.106
Reproductive rate	38%	0.102	-0.105
Age of first reproduction	M=3, F=4	0.009	-0.024
Second-year mortality	25%	0.002	-0.020
Adult mortality	6%	0.000	-0.009
Genetic load lethal equivalents	7.90	0.003	-0.006

DISCUSSION

Despite the potentially low chance of success (Fischer and Lindenmeyer 2000), translocations

continue to be used as a major tool in endangered species management. Our analyses evaluated

the likely result of a giraffe translocation under reasonable demographic and genetic assumptions

based on published literature. We considered a population to be minimally demographically

viable when it has a 5% or lower probability of extinction during a 100-year period, and

minimally genetically viable when it has a 20% or lower decrease in genetic diversity compared

to the original wild population over the same period (IUCN 2013). These minimum criteria for

viability of an isolated population were achieved when 20 females and 2 males were released,

assuming demographic parameters for a moderate population growth rate ($\lambda = 1.010$), and a

moderate amount of inbreeding genetic load (lethal equivalents: LE = 6.29). Under more

conservative demographic rates and higher genetic loads, the minimum release population for

95% viability at 100 years was 30 females and 3–6 males. This latter scenario also managed to

preserve at least 80% of the genetic diversity of the source population. Population projections

from our scenarios illuminate the marked impact of population growth and genetic load on the

long-term viability and genetic diversity. Our study thus shows how uncertainty in these

parameters can affect the minimum number of founders and sex ratios of animals required to be

released to successfully establish a new population.

Few studies have thoroughly monitored source and translocated populations for changes in

genetic parameters such as heterozygosity, allelic richness, and rate or level of inbreeding

(Groombridge et al. 2012, Puckett et al. 2014). Ideally, genetic variability should be assessed in

source populations in advance of translocations to use genetic information to guide translocation

plans and to provide a contrast to post-translocation genetic studies (Rocamora and Richardson

2003, Biebach and Keller 2010). This is particularly important if there is reason to assume that

the source population already has low genetic diversity, for example, because it was established

through translocations or because it was a long-isolated population of only a few hundred

individuals. Consideration should also be given to the issues that can arise when using

individuals for reintroduction from captive stocks where genetic adaptation to captivity may have

occurred (Montgomery et al. 2010, Robert 2009), which becomes more likely with more

generations in captivity, but should have been partially mitigated if the population has been

managed to minimize inbreeding (Marsden et al. 2013, Frankham 2010).

During the "founders' years" from year 2 to 23 in all our projections, initial population growth

will likely be positive if 2 or more females are translocated. However, this initial growth is a

transient effect of the young-skewed age distribution of the founding population, and very few

translocations of small (<20 females) founding populations will be viable in the long term in

isolation due to a lack of genetic diversity (**Fig 1**). Increasing the number of females in the initial

release had a strong positive effect on mean annual stochastic rate of population increase, and

final genetic diversity. A 0.01 probability of extinction was achieved assuming moderate

demographic rates and moderate genetic load if the founder population included 30 females and

3 males. In situations where the translocation destination has few or no large predators, and

asymptotic growth rates could approach 1.024, then viable populations could be achieved with

releases of a minimum of 20 females and 2 males that are all unrelated. These minimum

founding populations will result in a 20% loss of genetic diversity, so if genetic diversity

considerations are paramount and additional translocations at a later stage are unlikely, then

much larger founding populations should be attempted.

Note that this is not due to a lack of genetic diversity captured by the founding population, but

due to the rapid loss of genetic diversity in the first years when the population is still small. A

population that is twice as large, loses genetic diversity at half the speed. This highlights the

importance of ensuring that a reintroduced population can grow as fast as possible in this first

stage. Population growth could be facilitated, for example, by providing supplemental water or

food, or veterinary care, especially for individuals with under-represented genes. This may seem

to counteract natural selection, but selection works inefficiently on small populations (Margan et

al. 1998). Instead, maintaining genetic diversity will maximize the material available for

selection once the population has reached a stable size of at least several hundreds of individuals.

Also, the loss of genetic diversity can be mitigated by immigration of a few individuals from

another population every generation, with the exact number of migrants being dependent on the

situation (Wang et al. 2004, Puckett et al. 2014). If there are natural corridors, this exchange may

happen naturally. For fenced or isolated areas, this will require periodic translocations of

individuals. Either way, this carries the risk of spreading diseases.

Rules of thumb for translocations such as we presented here should not be used as a substitute

for site-specific monitoring and analyses, but they can be useful in early decision-making and

feasibility analyses. Ideally, every translocation should be conducted under an adaptive

management framework with continuous pre- and post-translocation monitoring of the source

and destination populations and annual updates to PVA analyses to inform structured decision-

making (McCarthy et al. 2011, Williams and Brown 2014). The use of continuous monitoring,

formal and transparent decision-making, and data-driven adaptive management cycles will

generate the necessary data to estimate demographic parameters, identify prospective subject

animals, predict population viability, and make informed management decisions. The true

demographic rates of any translocated population should be estimated annually and compared

with assumptions made during planning. As our sensitivity analysis showed, if demographic

rates such as juvenile survival or reproductive rates are below critical levels, the translocated

population will be considerably less viable than was predicted.

The ideal female age class for translocations in terms of population growth rate is the youngest

age class that can immediately begin reproducing, but adults are logistically challenging to move

due to their large body size. All of our population projections used the logistically more tractable

age class of 2-year-olds for translocation, as this age class is often chosen for translocations.

Translocating juveniles means there is an inevitable initial post-release decline in population size

as animals are subject to mortality before reaching reproductive age. Sensitivity analyses

indicated ages of first reproduction above our very optimistic values of 3 for females and 2 for

males can adversely affect rate of population increase and long-term viability.

Fewer males than females are necessary in translocations because the male contribution to

population viability is mostly confined to their genetic contributions, and equal offspring sex

ratios will greatly increase the number of adult males within 1 generation. Our projections

assumed all males were in the breeding pool and had equal probability of mating. The details of

giraffe mating success are largely unknown for wild populations, but there is evidence that

dominant males may monopolize mating opportunities potentially resulting in a skewed mating

success among existing males (Pratt and Anderson 1982; 1985). Similarly, female mate choice

could result in a skewed mating success among males (Andersson 1994). Either of these

situations would reduce the pool of breeding males thereby reducing the genetic diversity passed

to subsequent generations, leading to a smaller effective population size, lower genetic diversity,

and possibly higher inbreeding depression.

Future research to inform translocation PVA analyses should focus on two critical data gaps.

First, research should aim to quantify inbreeding depression in wild or semi-wild populations

that have and have not experienced a population bottleneck, and to quantify genetic parameters

in translocated versus source populations over time. Second, research should determine whether

assumptions of panmictic breeding and equal breeding success of all males are accurate. This is

important because skewed mating success will reduce the effective population size, and worsen

the impact of inbreeding, i.e. result in more severe inbreeding depression. Published accounts of

translocation outcomes including annual demographic rates of age-specific survival and

reproduction, genetic diversity, and long-term population viability are also needed as giraffe

translocations are typically not well documented (Winter et al. 2019).

ACKNOWLEDGEMENTS

We thank two anonymous reviewers and the associate editor who improved this manuscript.

LITERATURE CITED

Andersson, M. 1994. Sexual Selection. Princeton University Press, Princeton, New Jersey, USA.

Biebach, I., and, L. F. Keller. 2010. Inbreeding in reintroduced populations: the effects of early reintroduction history and contemporary processes. Conservation Genetics 11:527–538.

Boyce, M. S. 1992. Population viability analysis. Annual Review of Ecology and Systematics 23:481–497. doi: 10.1146/annurev.es.23.110192.002405

Bustamante, J. 1996. Population viability analysis of captive and released bearded vulture populations. Conservation Biology 10:822–831. doi: 10.1046/j.1523-1739.1996.10030822.x

Converse, S. J., C. T. Moore, and D. P. Armstrong. 2013. Demographics of reintroduced populations: estimation, modeling, and decision analysis. The Journal of Wildlife Management 77:1081–1093. doi:10.1002/jwmg.590

Dagg, A. I. 2014. Giraffe: Biology, Behaviour and Conservation. Cambridge University Press, Cambridge, UK.

Deredec, A., and F. Courchamp. 2007. Importance of the Allee effect for reintroductions. Ecoscience 14:440–451. doi: 10.2980/1195-6860(2007)14[440:IOTAEF]2.0.CO;2

Fischer, J., and D.B. Lindenmayer. 2000. An assessment of the published results of animal relocations. Biological Conservation 96:1–11.

Frankham, R., J. D. Ballou, and D. A. Briscoe. 2010. Introduction to conservation genetics. Cambridge University Press, Cambridge, UK.

Groombridge, J. J., C. Raisin, R. Bristol, and D. S. Richardson. 2012. Genetic consequences of reintroductions and insights from population history. Pp. 395-440 in Reintroduction Biology,

Integrating science and management (Eds. J. G. Ewen, D. P. Armstrong, K. A. Parker, and P.

J. Seddon). Wiley-Blackwell, Chichester, UK.

International Union for the Conservation of nature [IUCN]. 2013. Guidelines for reintroductions

and other conservation translocations. IUCN SSC Re-Introduction Specialist Group, Gland,

Switzerland.

Lackey, L. B. 2009. Giraffe studbook *Giraffa camelopardalis* North American Regional/ Global.

Association of Zoos and Aquariums and Disney's Animal Kingdom, Orlando, Florida, USA.

Lacy, R., M. Borbat, and J. Pollak. 2017. VORTEX: a stochastic simulation of the extinction

process. Chicago Zoological Society, Brookfield, Illinois, USA.

Lee, D. E., and, M. K. L. Strauss. 2016. Giraffe Demography and Population Ecology. Reference

Module in Earth Systems and Environmental Sciences. Elsevier, Amsterdam. Available via

ScienceDirect. doi: 10.1016/B978-0-12-409548-9.09721-9.

Lynch, M., and B. Walsh. 1998. Genetics and analysis of quantitative traits. Sinauer, Sunderland,

Massachusetts, USA.

Margan, S. H., R. K. Nurthen, M. E. Montgomery, L. M. Woodworth, E. H. Lowe, D. A.

Briscoe, and R. Frankham. 1998. Single large or several small? Population fragmentation in

the captive management of endangered species. Zoo Biology: Published in affiliation with the

American Zoo and Aquarium Association 17:467–480.

Marsden, C. D., H. Verberkmoes, R. Thomas, R. K. Wayne, and B. K. Mable. 2013. Pedigrees,

MHC and microsatellites: an integrated approach for genetic management of captive African

wild dogs (Lycaon pictus). Conservation Genetics 14:171–183.

Montgomery, M. E., L. M. Woodworth, P. R. England, D. A. Briscoe, and R. Frankham. 2010.

Widespread selective sweeps affecting microsatellites in Drosophila populations adapting to

captivity: implications for captive breeding programs. Biological Conservation 143:1842–

1849.

McCarthy, M. A., D. P. Armstrong, and M. C. Runge. 2012. Adaptive management of

reintroduction. Pp 256-289 in Reintroduction Biology, Integrating science and management

(Eds. J. G. Ewen, D. P. Armstrong, K. A. Parker, and P. J. Seddon). Wiley-Blackwell,

Chichester, UK.

Morris, W. F., and D. F. Doak. 2002. Quantitative conservation biology. Sinauer, Sunderland,

Massachusetts, USA.

Morton, N. E., J. F. Crow, and H. J. Muller. 1956. An estimate of the mutational damage in man

from data on consanguineous matings. Proceedings of the National Academy of Sciences

USA 42:855–863.

Muller, Z., F. Bercovitch, R. Brand, D. Brown, M. Brown, D. Bolger, K. Carter, F. Deacon, J.B.

Doherty, J. Fennessy, S. Fennessy, A. A. Hussein, D. Lee, A. Marais, M. Strauss, A.

Tutchings, and T. Wube. 2018. Giraffa camelopardalis (amended version of 2016

assessment). The IUCN Red List of Threatened Species 2016. doi: 10.2305/IUCN.UK.2016-

3.RLTS.T9194A136266699.en

Nietlisbach, P., S. Muff, J. M. Reid, M. C. Whitlock, and L. F. Keller. 2019. Nonequivalent

lethal equivalents: Models and inbreeding metrics for unbiased estimation of inbreeding load.

Evolutionary applications, 12:266–279.

O'Grady, J. J., B. W. Brook, D. H. Reed, J. D. Ballou, D. W. Tonkyn, and R. Frankham. 2006.

Realistic levels of inbreeding depression strongly affect extinction risk in wild populations.

Biological Conservation 133:42–51. doi: 10.1016/j.biocon.2006.05.016

Owen-Smith, R. N. 1988. Megaherbivores: The influence of very large body size on ecology.

Cambridge University Press, New York, New York, USA.

Pratt, D.M., and V.H. Anderson. 1982. Population, distribution, and behaviour of giraffe in the

Arusha National Park, Tanzania. Journal of Natural History 16:481–489.

Pratt, D.M., and V.H. Anderson. 1985. Giraffe social behaviour. Journal of Natural History

19:771–781.

Puckett, E. E., T. V. Kristensen, C. M. Wilton, S. B. Lyda, K. V. Noyce, P. M. Holahan, D.M.

Leslie, J. Beringer, J.L. Belant, D. White, and L. S. Eggert. 2014. Influence of drift and

admixture on population structure of american black bears (Ursus americanus) in the central

interior highlands, USA, 50 years after translocation. Molecular Ecology 23:2414–2427.

doi:10.1111/mec.12748

Robert, A. 2009. Captive breeding genetics and reintroduction success. Biological Conservation

142:2915–2922.

Rocamora, G., and D.S. Richardson. 2003. Genetic and morphological differentiation between

the two remnant populations of Seychelles white-eye Zosterops modestus. Ibis 145:34–44.

Seddon, P.J., D .P. Armstrong, and R. F. Maloney. 2007. Developing the science of

reintroduction biology. Conservation Biology 21:303–312.

Slotta-Bachmayr, L., R. Boegel, P. Kaczensky, C. Stauffer, and C. Walzer. 2004. Use of

population viability analysis to identify management priorities and success in reintroducing

Przewalski's horses to southwestern Mongolia. Journal of Wildlife Management 68:790–798.

doi: 10.2193/0022-541X(2004)068[0790:UOPVAT]2.0.CO;2

Van Houtan, K. S., J. M. Halley, R. Van Aarde, and S. L. Pimm. 2009. Achieving success with

small, translocated mammal populations. Conservation Letters 2:254–262. doi:

10.1111/j.1755-263X.2009.00081.x

Wang, J. 2004. Application of the one-migrant-per-generation rule to conservation and

management. Conservation Biology 18:332-343.

Williams, B. K., E. D. Brown. 2014. Adaptive management: From more talk to real action.

Environmental Management 53:465–479.

Winter, S., R. T. F. Coimbra, A. Bronec, C. Hay, A. L. Salb, J. Fennessy, and A. Janke. 2019.

Species assignment and conservation genetics of giraffe in the Republic of Malawi.

Conservation Genetics doi:10.1007/s10592-018-01142-4.

413