Elimination of reference mapping bias reveals robust immune related allele-specific expression in crossbred sheep

Mazdak Salavati [1*], **Stephen J. Bush** [1], **Sergio Palma-Vera** [2], **Mary E. B. McCulloch** [1], **David A. Hume** [3] and **Emily L. Clark** [1*]

1. The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush, Midlothian, UK
2. Leibniz Institute for Farm Animal Biology (FBN), Dummerstorf, Germany
3. Mater Research Institute-University of Queensland, Translational Research Institute, Woolloongabba, Qld 4102, Australia

* Correspondence:
Dr Emily L. Clark
Emily.Clark@roslin.ed.ac.uk
Dr Mazdak Salavati
Mazdak.Salavati@roslin.ed.ac.uk

Keywords: Allele-specific expression, mapping bias, RNA-Seq, sheep, transcriptome, WASP, GeneiASE
Running title: Unbiased ovine allele-specific expression
Editorial choice: British English
Article type: Original Research
Manuscript length: 6611 words

Abstract

Pervasive allelic variation at both gene and single nucleotide level (SNV) between individuals is commonly associated with complex traits in humans and animals. Allele-specific expression (ASE) analysis, using RNA-Seq, can provide a detailed annotation of allelic imbalance and infer the existence of cis-acting transcriptional regulation. However, variant detection in RNA-Seq data is compromised by biased mapping of reads to the reference DNA sequence. In this manuscript we describe an unbiased standardised computational pipeline for allele-specific expression analysis using RNA-Seq data, which we have adapted and developed using tools available under open licence. The analysis pipeline we present is designed to minimise reference bias while providing accurate profiling of allele-specific expression across tissues and cell types. Using this methodology, we were able to profile pervasive allelic imbalance across tissues and cell types, at both the gene and SNV level, in Texel x Scottish Blackface sheep, using the sheep gene expression atlas dataset. ASE profiles were pervasive in each sheep and across all tissue types investigated. However, ASE profiles shared across tissues were limited and instead they tended to be highly tissue-specific. These tissue-specific ASE profiles may underlie the expression of economically important traits and could be utilized as weighted SNVs, for example, to improve the accuracy of genomic selection in breeding programmes for sheep. An additional benefit of the pipeline is that it does not require parental genotypes and can therefore be applied to other RNA-Seq datasets for livestock, including those available on the Functional Annotation of Animal Genomes (FAANG) data portal. This study is the first global characterisation of moderate to extreme ASE in tissues and cell types from sheep. We have applied a robust methodology for ASE profiling, to provide both a novel analysis of the multi-dimensional sheep gene expression atlas dataset, and a foundation for identifying the regulatory and expressed elements of the genome that are driving complex traits in livestock.

Introduction

Allele-specific expression (ASE) is the imbalance of allelic expression between parental (diploid) copies at the same locus (Barlow and Bartolomei, 2014). It is most commonly associated with cis-acting regulatory variation that may mediate parent-of-origin, sex- or tissue-specific transcription of one allele relative to the other (Renfree et al., 2009; Hasin-Brumshtein et al., 2014). In a single individual, where there are informative sequence variants (i.e. heterozygote loci) that distinguish the products of two alleles, ASE can be detected by RNA-sequencing (Chamberlain et al., 2015; GTEx Consortium et al., 2017; Cao et al., 2019; Guillocheau et al., 2019). The ratio of allelic read counts obtained from RNA-Seq datasets can be used as a reliable proxy for ASE (i.e. $ASEratio = CountsAllele1/(CountsAllele1 + CountsAllele2)$) (Edsgärd et al., 2016).

Large and complex RNA-Seq datasets give rise to unique and interesting computational challenges, in particular the elimination of reference mapping bias in ASE analysis of diploid genomes. RNA-Seq data are commonly mapped against reference genomes which are typically 'flat', with each position represented only by the reference (most abundant) allele. As such, reads containing heterozygous loci are more likely to be erroneously mapped (Degner et al., 2009; Stevenson et al., 2013; Hodgkinson et al., 2016). This can lead to high false positive ASE locus discovery rates (Degner et al., 2009). Although development of de novo transcript assemblers (Zerbino and Birney, 2008), usage of personalised reference genomes (Rozowsky et al., 2011; Smith et al., 2013) , variant-aware aligners (Xin et al., 2013; Hach et al., 2014) and mapping-free quantification e.g. Kallisto (Bray et al., 2016) have resolved some of these issues, reference allele mapping bias remains a considerable challenge in ASE studies. In the absence of 'trios' of animals or reference population phased haplotype information, which are rare for livestock, correction of mapping bias via synthetic reads with either N masking or alternative mapping bias correction at the heterozygote sites, has proven a robust alternative for ASE discovery (Degner et al., 2009; Mayba et al., 2014; van de Geijn et al., 2015; Miao et al., 2018). In 2015 Van de Geijn et al. benchmarked the WASP software mapping correction strategy against N-masked reads and personal genome mapping. WASP showed consistent correct mapping of reads with multiple alleles and lower false discovery rates in comparison to the other two methods (van de Geijn et al., 2015). The analysis pipeline we present in this manuscript is based on WASP's methodology and is designed to minimise reference bias while providing accurate profiling of allele-specific expression in large and complex RNA-Seq datasets.

We have developed an ASE analysis pipeline using the combination of software available under open licence, WASP (reference mapping bias removal) (van de Geijn et al., 2015), GATK (ASEReadCounter) (McKenna et al., 2010; Van der Auwera et al., 2013) and GeneiASE (Liptak-Stouffer aggregative ASE gene model) (Edsgärd et al., 2016). The GeneiASE model is capable of testing ASE at the gene level using two approaches: i) Static ASE which measures allelic imbalance within a gene (i.e. when ASE variants are located within the boundaries of the gene) and ii) Individual condition dependent ASE (ICD) which measures inducible ASE in a gene under an environmental pressure between 2 timepoints (i.e. in stimulated or unstimulated immune cells).

In addition to ASE at the gene level, we can also measure significant ASE at the single nucleotide level (SNV). ASE has been shown to be enriched within expression quantitative trait loci (eQTL) regions (Montgomery et al., 2010), therefore identifying ASE variants can be useful for understanding the transcriptomic control of complex traits in livestock. Complex trait mapping of ASE loci has been associated with phenotypes such as resistance to Marek's disease in chicken (Meydan et al., 2011) and pigmentation patterns in sheep (García-Gámez et al., 2011).

Understanding ASE is also important because cross-breeding now underlies most
livestock production systems. Knowledge of ASE may provide insights into the molecular basis
of the complex phenomenon of hybrid vigour, as emphasised by recent studies on two Chinese
goat breeds and their F1 hybrids (Cao et al., 2019) and in F1 crosses of two highly inbred
chicken lines (Zhuo et al., 2017). In this study we measure ASE in crossbred sheep. Sheep are
an economically important livestock species in many countries across the globe and
particularly in emerging economies. The identification of prevalent ASE in populations or
breeds, especially in economically relevant phenotypes and tissues, could be used to improve
genomic prediction in sheep breeding programmes, such as those that have been established in
Australia and New Zealand (Daetwyler et al., 2010).
Using the methodology we describe, for mapping bias correction and robust positive
ASE discovery, we were able to profile pervasive allelic imbalance across tissues and cell
types, at both the gene and SNV level, in Texel x Scottish Blackface sheep. We analysed a
subset of total RNA-Seq libraries from liver, spleen, ileum, thymus and bone marrow-
derived macrophages (BMDM) (+/-) lipopolysaccharide (LPS), from six individual adult
crossbred sheep to produce a detailed picture of allelic imbalance in immune-related tissues
and cell types. We chose to focus this analysis on immune related tissues in part because of the
depth of available sequence in those tissues, and in part because they contain abundant immune
cell populations. The diversity of cell populations is reflected in the transcriptional complexity
of immune tissues and cell types in the sheep gene expression atlas dataset (Clark et al., 2017;
Bush et al., 2018). As such this subset of tissues gave us a transcriptionally rich dataset in
which to measure ASE. We also included BMDMs stimulated and unstimulated with LPS to
mimic infection with Gram-negative bacteria, in order to test whether ASE changed in response
to stimulation with LPS in these cells. By measuring ASE in these tissues and cell types from
sheep we were able to: i) provide insight into how pervasive ASE is across tissues at the gene
and SNV level, ii) generate tissue-specific ASE profiles, iii) investigate sex-specific patterns
of ASE, and iv) determine the extent to which ASE changes in response to stimulation with
LPS in an immune cell type. This novel analysis of the multi-dimensional sheep gene
expression atlas dataset provides a foundation for further analysis of the regulatory and
expressed elements of the genome that are driving complex traits in sheep.
130
131 **Methods**
132
133 **Sample preparation and RNA extraction**
134
Data from three male and three female Texel x Scottish Blackface (TxBF) sheep from
the sheep gene expression atlas project (Clark et al., 2017) were used in this study. The dataset
including: 1 cell type (BMDMs (+/-) LPS treatment) and 4 tissues (thymus, spleen, liver and
ileum). Tissue collection, storage and RNA extraction is described in (Clark et al., 2017).
BMDM's were cultured in vitro for 7 days in the presence of macrophage colony-stimulating
factor (CSF1 (10^4 U/ml)) and unstimulated (0 hrs - LPS) and stimulated (7hrs +100ng/ml LPS)
samples of BMDMs were obtained also as described in (Clark et al., 2017). A total of 2 samples
(1 thymus and 1 spleen) did not pass the RNA quality control (RNA integrity number (Mueller
et al., 2004); RIN >7) and were not included in the sheep gene expression atlas. Library
preparation was performed by Edinburgh Genomics (Edinburgh Genomics, Edinburgh, UK).
All total RNA Illumina TruSeq libraries (125bp paired end) were sequenced at a depth of >
100 million reads per sample.
147
148 **Reference mapping bias removal**
149

BAM files from RNA-Seq data, were previously produced by mapping fastq files to the
Oar v3.1 top level DNA fasta track, using HISAT2 (default mismatch penalty MX=6 MN=2)
as described in (Clark et al., 2017). Detailed settings and parameters for all the tools used to
generate the BAM files can be found at (FAANG, 2018). These BAM files were used to locate
reads with heterozygote loci using WASP's find_intersecting.py script (van de Geijn et al.,
2015). The intersection of reads and heterozygote loci in all samples were based on the Ensembl
v92 variant call format (VCF) track (Ensembl v92: ovis_aries_incl_consequences.vcf.gz).
Briefly the Ensembl VCF file was filtered for bi-allelic variants within exonic regions, 5k up
or downstream of exonic regions (5' or 3' UTRs) and intronic regions of all transcripts within
the Oar3.1 sheep assembly (exclusion of indels and intergenic variants). These variants were
used in WASP's find_intersecting.py script in order to extract reads mapped to coordinates
containing variants for each gene. As a result, reads aligned to exonic, 5' or 3' UTRs and
intronic regions were separated into reads intersecting heterozygote loci and reads that did not
intersect heterozygote loci. Synthetic copies of reads intersecting heterozygote loci, were
created with the alternate allele flipped to the remaining options of A, T, C or G (up to 6 loci
/read[2^n] max 64 combinations of synthetic reads) using parameters defined in WASP (van de
Geijn et al., 2015). This was followed by remapping of the synthetic reads using HISAT2
(default mismatch penalty MX=6 MN=2) (Li and Durbin, 2009; Kim et al., 2015) and
eliminating the original reads (and their synthetic copies) which mapped to a different
coordinate in any of its synthetic copies (WASP's filter_reads.py) (van de Geijn et al., 2015).
After merging the retained reads with that did not intersect heterozygote loci, a final BAM file
was produced for ASE read counting step (WASP's remove_dup.py).

172
173 **Allelic read counts and depth filtration**

174
Allele-specific read counting was carried out using the ASEReadCounter module of
GATK v3.8 with parameters -mmq 50 -mbq 25 (McKenna et al., 2010). Multiple pre-
processing steps were performed prior to GeneiASE input as instructed by (Edsgärd et al.,
2016) which included, preparing per chromosome indices, merging the variant set with
corresponding gene coordinates and bi-allelic expression filtering. Loci with < 10 reads
mapped were excluded, as were loci with < 3 reads, or < 1% of the total reads, mapped to both
the reference and alternative allele. This form of filtration will eliminate loci exhibiting mono-
allelic expression (MAE) as previously described (Degner et al., 2009; Stevenson et al., 2013;
Mayba et al., 2014). Producing evidence of MAE using total RNA-Seq datasets produced by
Illumina short read sequences without parent of origin genotypes or imprinting information has
been a controversial issue (DeVeale et al., 2012). Our dataset did not include the trios of
animals or personalised genomes that would be necessary to resolve MAE. As such we decided
to exclude MAE altogether for our analysis using stringent bi-allelic filtration criteria. Similar
bi-allelic filtration criteria have been previously used routinely in ASE studies (Mayba et al.,
2014; Chen et al., 2016a; Edsgärd et al., 2016; GTEx Consortium et al., 2017; Raghupathy et
al., 2018; Cao et al., 2019; Guillocheau et al., 2019; Gutierrez-Arcelus et al., 2019). The
workflow of the analysis pipeline for ASE analysis is detailed in Fig.1.

192
193 **Experimental design for defining allele-specific expression**

194
195 ASE was defined according to the following three categories:

196
197 i) Static ASE: which is inherent allelic imbalance (AI) in each gene calculated by
198 ASE at all heterozygote loci (i.e. $ASE = Counts\,Ref\,Allele/$
199 $(Counts\,Ref\,Allele + Counts\,Alt\,Allele)$ within the boundaries of the gene.

200   The effect size of ASE at gene level was produced by aggregation of the ASE
201   effect size at SNVs within the gene boundaries (the Liptak-Stouffer method,
202   applied by the GeneiASE aggregative model). A null distribution of ASE effect
203   size for genes in each transcriptome was produced by random sub-sampling (n
204   = 1e+5) from a pool of genes having min 2 and max 100 loci within their
205   boundaries. The ASE effect size of each gene (aggregated using Liptak-
206   Stouffer) was then tested against the null distribution of same SNV number, via
207   a modified bi-nomial test (2x1 table). Distribution of p values was examined for
208   uniformity prior to false discovery rate correction (Supplementary Fig.S1, S2
209   and S3) (Benjamini and Hochberg, 1995; Pounds and Cheng, 2006; Barton et
210   al., 2013).
211   ii)   Individual condition-dependent ASE (ICD-ASE): in which the same ASE
212   effect size was calculated for each gene in the treated versus the untreated
213   timepoints of the same sample (i.e. BMDM +/- LPS). The log2ratio
214   ($ASE_{treated}/ASE_{untreated}$) was used in a beta-binomial test (2x2 table) similar to
215   the static mode. The details of this aggregative model have been previously
216   described in the GeneiASE publication (Edsgärd et al., 2016).
217   iii)   Condition dependent ASE at SNV level: in which a contingency table was
218   produced for read counts (ref and alt) for every SNV, present both in treated and
219   untreated conditions (BMDM +/- LPS) (2x2 table) and a Fisher's exact test
220   performed followed by p value multiple testing correction (Benjamini and
221   Hochberg, 1995). The p values from loci showing ASE and shared by the six
222   adult sheep (ID and coordinate) were unified, using the Stouffer method
223   (Dewey, 2016, 2018) and presented as false discovery rate (FDR) for each locus.
224
225   Static ASE was calculated in both tissues and BMDMs (each timepoint was considered
226 separately for BMDMs). Condition dependent ASE analysis was carried out only in BMDMs
227 ± LPS both at gene (ICD-ASE) and SNV (Fisher's exact) level to study LPS inducible ASE.
228
229 **Statistical analysis and thresholds applied**
230
The extraction, transformation and loading of the all datasets and subsequent statistical analysis
was carried out in R v $\geq$ 3.4 unless stated otherwise (R Core Team, 2017). System query
language join statements (Wickham et al., 2019) were used to compare lists of ASE genes or
SNVs between samples. Raw p values resulting from all three types of ASE analysis were
corrected for multiple testing via Benjamini-Hochberg false discovery rate calculations
(Benjamini and Hochberg, 1995). The passing threshold of significance in all analyses was
considered to be FDR < 0.1 (10%) except for the Fisher's exact test association study. Genes
showing ASE in multiple tissues were considered those for which 4 or more of the 6 sheep had
significant ASE.
240
241 **Results**
242
243 **Estimation of heterozygous sites across all individuals**
244
To determine the level of heterozygosity present in the RNA-Seq data we first assessed
the number of bi-allelic heterozygote sites per individual for each of the six sheep (range=
5,673,703-6,438,497) detailed in Fig.2. Individual variation was observed in the SNVs per gene
in each sheep (Fig.2A&C). However, there was no significant difference in the total number of
bi-allelic SNVs captured in the RNA-Seq data across all six individuals or between the male

and female sheep included in the study (Fig.2B). The bi-allelic SNVs captured in the RNA-Seq
data set were annotated using the Ensembl v.92 (Zerbino et al., 2018) reference VCF track. The
distribution of SNVs per gene in the Ensembl track is tail-inflated in comparison to the RNA-
Seq data Fig.2A. This issue could be due to erroneous assignment of SNVs in hypervariable
and repetitive regions, multi-allelic SNVs or simply that there are variants in the Ensembl track
that are not expressed (transcribed). The distribution of SNVs for each individual is shown in
Fig.2C.
257
258 **Reference mapping bias elimination and quality control**
259
We used the WASP ref bias removal script to successfully minimise ref allele mapping
bias in the RNA-Seq samples. The mapping bias was assessed by global distribution of the
allelic ratio i.e. $ref_{counts}/alt_{counts} + ref_{counts}$ in each RNA-Seq sample as shown in Fig.3 (WASP
metrics are included in supplementary Fig.S10,S11 and S12). The ASE discovery rate at the
SNV level on average constituted 5.8% of the heterozygote loci that passed the minimum
filtration criteria in each individual (0.1% of the total expressed). This portion of the
transcriptomic variants belonged to an average of 103 genes in each tissue transcriptome
(approx.1%) or 300 in each individual (Supplementary Fig.S5 and S6). As shown in
Supplementary Fig.S6. expression level varies across tissues but does not affect the
distribution of ASE SNVs.
270
271 **Genes exhibiting tissue-specific and pervasive ASE signatures**
272
We used the static mode of GeneiASE to investigate pervasive and tissue-specific ASE
profiles across all of the available samples. Static ASE represents inherent allelic imbalance
(AI) in each gene calculated by ASE at all heterozygote loci. The number of genes showing
significant static ASE in immune-related tissues across the six sheep are summarised in Table
1. On average approximately 0.5% of the genes in each tissue-specific transcriptome showed
significant ASE (approx. 1% of the filtered set of genes). Pervasive ASE genes were
investigated by applying the minimum 67% shared rule (i.e. a ASE gene was considered
'shared' when it exhibited ASE in a minimum of 4 out of 6 sheep). A list of ASE genes with
significant allelic imbalance (AI) in all tissues, when the effect size was averaged across 6
sheep, was compiled (Fig.4A) (Static ASE measured by GeneiASE's Liptak-Stouffer method).
Six genes exhibited pervasive ASE across tissues (i.e. they were shared across all 4 tissues). In
the order of allelic imbalance effect size they were NAA50 (N(alpha)-acetyltransferase 50,
Histone acetyl transferase) with highest ASE effect size in spleen, UBB (Ubiquitin B, protein
degradation) in thymus, HBP1 (HMG-Box transcription repressor; cell cycle regulation) and
ENSOARG00000016510 both in spleen, C1orf105 (Chromosome 1 Open Reading Frame 105)
in ileum and MTIF2 (mitochondrial translational initiation factor 2) in thymus.
Sets of genes with tissue-specific ASE profiles were also captured (Fig.4B-E). Thymus
had the highest number of tissue-specific ASE genes (n=15) followed by liver (n=12), spleen
(n=5) and ileum (n=4) (Fig.4B-E). Amongst the thymus gene set was CD244 which included
30 heterozygote loci with allelic imbalance, one of which was rs406633825. This missense
allele (Chr1:110308273 C>A; pVal123Phe MAF=0.7, SIFT score=0 deleterious) has
previously been reported in the Texel population characterised by the International Sheep
Genome Consortium (ISGC) (Kijas et al., 2012). The CD244, a non-MHC (major
histocompatibility complex) mediated marker expressed by NK cells and multiple subsets of
CD8+ T cells is known for both pro-inflammatory and inhibitory effects on lymphocytes
(McNerney et al., 2005; Georgoudaki et al., 2015). CD244 exons 2 to 5 are highly conserved
in vertebrates and in mouse a Trypanosoma infection model indicated differential expression

was correlated with multiple copy number variants nearby (Goodhead et al., 2010).The liver-
specific ASE profile included genes involved in amino acid metabolism, cytochrome oxidase
pathways and fibrinogen: FGA (fibrinogen alpha chain), ENSOARG00000003175
(taurochenodeoxycholic 6 alpha-hydroxylase-like, cytochrome P450 family 4 member
[CYP4]), ENSOARG00000001568 (novel gene, complement C4-A-like), CYP3A24
(cytochrome P450 family 3 member) and CA3 (carbonic anhydrase 3). Allelic imbalance in
spleen was present in CACYBP (calcyclin binding protein), DAPK2 (death associated protein
kinase [serine/threonine]) and a novel gene GIMAP8-like (ENSOARG0000001131). The ASE
in GIMAP8 has been previously reported in cattle with a strong paternal parent-of-origin
expression pattern (Chamberlain et al., 2015). The GIMAP/IAN family of proteins, are
involved in survival, selection and homeostasis of lymphocytes (Nitta and Takahama, 2007).
Two genes of functional interest showed evidence of strong tissue-specific ASE in the
spleen: SNAP23 and MYLK. SNAP23 is a key molecule in vesicle transport machinery of the
cell and has been reported to be expressed in sheep spleen. SNAP23 or Synaptosome
Associated Protein 23 is part of protein complex involved in Class 1 MHC mediated antigen
processing and presentation and in neutrophil degranulation (Fabregat et al., 2018). SNAP23
is also vital to lymphocyte development (both B and T) in vitro (Wong et al., 1997; Kaul et al.,
2015). The myosin light chain kinase (MYLK) expression in the splenic trabeculae's smooth
muscle has been demonstrated previously (Jiang et al., 2014; Clark et al., 2017). Overall 199
heterozygote bi-allelic loci were present within the MYLK gene. The variant rs400678033
(Chr1:186347056G>A; .pAla1014Val), a missense SNV in exon 17 of 33 exons in MYLK,
showed consistent allelic imbalance in all spleen samples.
In summary, analysis of ASE across immune related tissues revealed there were a small
number of genes that were shared across tissues. ASE signatures instead tended to be tissue-
specific, within the sub-set of tissues investigated in this study.
325
326 **Individual-specific ASE signatures**
327
In order to investigate whether ASE profiles were either shared across all six sheep or
private to individual sheep we used intersectionality (Fig.5). Each tissue was investigated
separately. A number of private (to each individual) ASE genes were detected for each tissue,
ranging from: 31-123 in ileum (Fig.5A), 24-80 in liver (Fig.5B), 21-83 in spleen (Fig.5C) and
31-66 in thymus (Fig.5D). Some of the shared sets of ASE genes in these tissues were specific
to either male or female sheep, these sex-specific ASE signatures are described in Fig.5. In
ileum no sex-specific set was observed (Fig5A). In contrast to ileum, the ASE profile for liver
included a single gene with female only membership, ENSOARG00000017409 (novel gene;
93% orthology with bovine dicarbonyl and L-xylulose reductase [DCXR]) Fig.5B. In spleen
all female sheep shared significant ASE in PMS1 (DNA mismatch repair system component
during post meiotic segregation), ANKRD10 (Ankyrin Repeat Domain 10) and
ENSOARG00000006103 which was not present in any of the spleen profiles of male sheep
(Fig.5C). In thymus there were 2 sex-specific sets: 16 genes showing ASE only in females and
5 genes only in males. The female-specific thymus gene set included: ARPP21 (cAMP
regulated phosphoprotein 21), CDKL3 (cyclin dependent kinase like 3), CEP19,
ENSOARG00000000710 (novel gene), ENSOARG00000001163 (novel gene),
ENSOARG00000000710, ENSOARG00000001163, ENSOARG00000008981 (novel gene;
T-lymphocyte surface antigen Ly-9-like), ENSOARG00000006215, ENA000000008981,
ENSOARG00000009129, ENSOARG00000011375 (blood vessel epicardial substance
[BVES]), ENSOARG00000015755, ENSOARG00000020354 (novel gene; 53% orthology
with bovine monoacylglycerol acyltransferase [MOGAT1]), ENSOARG00000025005,
ENSOARG00000026030 (novel gene), GPM6A (glycoprotein M6A), RAG1 (recombination

Activating 1), STX8 (syntaxin 8). The male-specific thymus set was comprised of
ENSOARG00000007267 (novel gene; T-cell surface glycoprotein CD1a-like),
ENSOARG00000016841 (novel gene; 98% orthology with bovine ATP synthase membrane
subunit G [ATP5MG]), ENSOARG00000007603, SNX25 (sorting nexin 25) and LDHA
(lactate dehydrogenase A) Fig.5D.
In summary very few ASE genes were shared across all sheep and the majority of ASE
profiles were private to each sheep. Sex-specific ASE signatures were also detected but due to
the small sample size (n=3) in both cases these should be interpreted with caution.
358
359 **ASE in stimulated and unstimulated BMDMs (0hrs vs 7hrs +LPS)**
360
We examined inducible ASE after 7 hrs of exposure to LPS in BMDMs using the ICD
mode of GeneiASE (Edsgärd et al., 2016). A comparison of LPS inducible ICD-ASE genes
and the genes with background static ASE at 0 and 7 hr time points, was also performed. We
first assessed whether differences between 0 and 7 hours could be observed using analysis of
static ASE. Individual-specific ASE profiles and a limited number of shared ASE genes were
also observed in BMDMs. The total number of genes with static ASE in the BMDMs is shown
in Table 2.
Shared static ASE across both time points and independent of LPS induction was only
observed in 5 genes. These genes have a macrophage associated function and include ITGB2
(Yee and Hamerman, 2013), SAA3 (ENSOARG00000009963) (Larson et al., 2003; Deguchi
et al., 2013), CD200R1 (ENSOARG00000019357) (Ocaña-Guzman et al., 2018), DCTN5
(ENSOARG00000017281) (Habermann et al., 2001) and MTIF2 (also seen in the tissue
analysis above) (Overman et al., 2003).
The ICD-ASE in BMDMs +/-LPS captured fewer ASE genes with significant LPS
inducible ASE between the 2 timepoints than the static analysis of ASE. Moreover, there were
large differences in the number of LPS inducible ASE genes in each individual sheep,
indicating significant individual-specific variation in response to LPS stimulation. BMDM
cultured from female 2 showed no LPS inducible response in comparison to male 3 which was
a hyper responder with significant inducible ICD-ASE in 28 genes (including 634 informative
SNVs total). A detailed breakdown of SNVs, aggregated within each gene, with significant
ICD ASE has been summarised in Fig.6.
In summary the ICD-ASE mode of GeneiASE's model was not capable of capturing a
complete picture of differential ASE in the BMDM experiment. Static ASE was present in both
time points, however, there were no inducible ASE genes that were shared across all 6 sheep.
The highly diverse ASE profile of BMDMs was very individual-specific, similar to patterns
observed in tissues (Supplementary Table S1). These individual-specific differences could be
due to individual variation in the innate immune response or experimental variation introduced
during primary cell culture or stimulation with LPS.
389
390 **Condition dependent ASE at the SNV level in BMDMs**
391
To further investigate allelic imbalance at the SNV level without the aggregative gene
model of GeneiASE, Fisher's exact test was used. The filtered read counts for bi-allelic SNVs
shared by all 6 sheep BMDMs were selected (n=646 sites). Allelic read counts of each SNV
were tested using Fisher's exact test between 0hrs and 7hrs (2x2 table). Overall the 6 sheep
shared 646 SNVs with identical allelic genotypes in both time-points of the BMDM RNA-Seq
dataset. These SNVs were tested for association with LPS treatment and only 4 SNVs showed
a strong association (FDR < 1.0e-8) and 12 SNVs had an FDR between 1.0e-02 and 1.0e-08
(Fig.7). The highest F-statistic was at rs430667535 Chr17 Pos:50485358T>C, a synonymous

variant in ubiquitin C (UBC), a polyubiquitin precursor, and also an intronic variant T>C or
A>G in pro-apoptotic BRI3 binding protein (BRI3BP). This variant was shown to have a minor
allele frequency (MAF) of 0.25 in Texel sheep based on the ISCG annotation [ISGC – Ensembl
v.92](Kijas et al., 2012)). The next highest peak was observed on Chr21 under SNVs within
SAA3 gene boundaries (ENSOARG00000009963) at the following coordinates: Highest FDR
peak was observed at SNV rs412192652 (Pos:25826978A>G missense variant [Asn145Asp],
FDR=2.3e-15) surrounded by rs403064928 (Pos:25826884C>A missense variant [Asp113Glu]
in exon 4 of SAA3 FDR=3.6e-07), rs426609498 (Pos:25826845A>G synonymous, FDR=5.5e-
07) and rs405439099 (Pos:25826990G>C 3'UTR variant, MAF 0.6 in Texel sheep, FDR=4.1e-
05). This region contains a strong LD block previously reported by the ISGC COMPOSITE
population (Ensembl v.92) (Supplementary Fig.S4) e.g. rs412192652 and rs405439099
pairwise D' statistics = 1. Two further peaks on Chr3 were observed for rs159926581
(Pos:214731375T>C synonymous, FDR = 3.3e-09) in ribosomal protein L3
(ENSOARG00000016495) and rs159822214 (Pos:112164732G>A 3'UTR variant,
FDR=9.5e-03) in oxysterol binding protein like 8 (OSBPL8). The last inducible ASE
associated signal was on Chr16 rs420037698 (Pos:6887423G >A, FDR= 6.9E-09) in
ENSOARG00000004700, a known synonymous variant in the Texel population (MAF = 0.55).
The SNVs and corresponding genes from Fisher's exact test are summarized in Table 3.
The LPS inducible ASE analysis, of SNVs, using Fisher's exact test revealed a different
picture not captured by the gene level analysis with the GeneiASE model that aggregates SNVs
within each gene (Fig.6 vs Fig.7). The aggregative gene model did not capture any shared ASE
genes in the ICD-ASE mode. Although the 6 sheep shared 646 SNVs and showed highly
significant association with stimulation with the LPS (Fisher's exact method), the aggregation
of ASE effect size from SNV to gene level (ICD-ASE mode) only detected individual-specific
sets of ASE genes in each sheep. This contradicted the results from Fisher's exact test which
detected 4 highly significant LPS inducible shared regions (FDR < 0.01, 16 SNVs) on
chromosomes 3, 16, 17 and 21. For example, the allelic imbalance in the SAA3 genomic
coordinates on chromosome 21 was not detectable in the ICD-ASE model but it was captured
by the Fisher's exact test in all individuals (Fig.7B chromosome 21).
Fisher's exact analysis at SNV level revealed ASE in response to LPS in variants within
CLDN1, ANXA3, BRI3BP, SAA3 and MSR1 (Table3). The anti-inflammatory macrophage
marker CLDN1 (Van den Bossche et al., 2012) and acute-phase inflammation resolution
marker ANXA3 (Yamanegi et al., 2018) have been previously reported with distinct
macrophage functions. The macrophage scavenger receptor 1 (MSR1) has also been shown to
be involved in lipid uptake and migration ability of macrophages (Shi et al., 2019). Three
noncoding RNAs (RF00221 [snoRD43], RF00593[snoU83B] and RF01151[snoU82P] were
amongst genes corresponding to ASE inducible SNVs. These 3 snoRNAs all overlap with the
genomic coordinates of ribosomal protein L3 (ENSOARG00000016495 / RPL3). The
RF00377 [snoU6-53] was also among the ASE positive targets which overlaps the protein-
coding gene CDS2 (CDP-diacylglycerol synthase 2). Using total RNA-Seq (ribosomal RNA
depleted), which includes multiple RNA populations, to generate short read illumina data
makes it difficult to pinpoint the origin of the ASE signal to a specific RNA population.
In summary ASE profiles in BMDMs were highly individual-specific at both gene and
SNV level. Moreover, Fisher's exact SNV level analysis discovered shared ASE SNVs where
the aggregative gene model of ICD-ASE mode did not, indicating for condition dependant ASE
analysis Fisher's exact test is more accurate and robust at SNV level.
446
447 **Discussion**
448
This study is the first to investigate global allele-specific expression across tissues from

sheep using RNA-Seq data. We focused our analysis on immune-related tissues and cell types
from six adult crossbred sheep (TxBF) from the sheep gene expression atlas. ASE profiles were
highly individual-specific in the six sheep analysed in this study. We were able to identify
tissue-specific sets of ASE genes, as well as LPS inducible sets in the BMDM experiment.
Tissue-specific signatures of ASE have been previously reported in similar studies in mouse
(Castel et al., 2015, 2016), goat (Cao et al., 2019) and cattle (Chamberlain et al., 2015).
Several steps were taken in the cattle study (Chamberlain et al., 2015) to mitigate the
ref allele bias, assign parental origin using whole genome sequences and include MAE variants.
The SNV filtration was based on the (Hayes and Daetwyler, 2019) 1000 bull genomes project
to confirm the heterozygote sites. In our pipeline the Ensembl VCF track was used for that
purpose. Chamberlain et al., 2015 use a 0.9 allele frequency cut-off (based on read counts) to
define and include MAE, and as such have a 1 and 0 inflated allelic ratios. In our pipeline no
allelic ratio cut-off is introduced for inclusion as it is difficult to distinguish between
sequencing error and MAE. The minimum read (bi-allelic expression) filtration criteria was
applied to exclude highly sequenced loci (Either count/Total >1%) or sequencing errors
presenting as rare alleles (min either allele count >=3) which consequentially excludes actual
MAE as well as spurious allelic counts. Chamberlain et al. tested 5317 genes (14,495 SNVs)
in spleen and detected 382 ASE genes (with min > 1 SNV per gene, similar to this study).
Although direct comparison would not be appropriate (because we have excluded MAE
variants in our analysis) our analysis of sheep spleen revealed ASE in 86 genes (averaged in 5
sheep) from 8272 filtered genes (averaged in 5 sheep). Similarly, in thymus the cattle study
showed 182 ASE genes from 986 informative genes (9781 SNVs) while from 7961 filtered
genes in sheep thymus 134 ASE genes were captured. The differences in the numbers of genes
exhibiting ASE between the two studies are likely to be a consequence of the filtration criteria
applied, the exclusion of MAE, and species-specific differences between sheep and cow.
Results from a more recent study in goat (Cao et al., 2019) more closely reflect our findings.
They apply similar filtration criteria to our workflow and discovered 144 ASE genes in liver in
comparison to 123 in our sheep liver sets (averaged across 6 sheep). Other recent studies
including those focusing on production relevant tissues, such as muscle (Guillocheau et al.,
2019), have also applied similar stringency in filtration criteria. The filtering criteria we have
used for this analysis is stringent and focused on detecting variants of moderate to extreme
effects. Further analysis of the dataset reducing these criteria might discover additional variants
exhibiting ASE across individuals and tissues, but it would also increase the potential risk of
false positive discovery.
For this analysis we have adapted an ASE analysis workflow with a primary focus on
mapping bias removal prior to allele-specific analysis of the transcriptome. The collection of
scripts for WASP, used for this analysis, or modified versions of them have been utilised by
others for mapping bias removal in reference guided genomic datasets e.g. RNA-Seq
(Mozaffari et al., 2018; Zhou et al., 2018), Chip-Seq (Pelikan et al., 2018) and for methylomic
and epigenetic analysis (Richard Albert et al., 2018).
The ASE analysis pipeline we have adapted for sheep for this study is also adaptable to
other species and tissue types with available RNA-Seq datasets. It could be applied, for
example, to profile allele-specific expression in the RNA-Seq datasets from livestock species
listed on the FAANG data portal (Andersson et al., 2015; Harrison et al., 2018). We used the
Ensembl VCF track to capture information at heterozygote loci however the individual VCF
file from each sheep could also be used in ASE analysis. The latter strategy might enable the
capture of rare variants not included in the publicly available VCF tracks but would also raise
the issue of normalisation/standardization between VCF call sets. The usage of either of these
methods will be limited to the number of loci shared by coordinate and bi-allelic genotype (i.e.
pervasive ASE discovery). Other studies have compared variants at the RNA and DNA level

from the same individual and then removed the DNA variants not present in the RNA-Seq data
from the analysis (Guillocheau et al., 2019). We believe that a strength of the pipeline we
present is that it does not require parental genotypes and can therefore be applied to other RNA-
Seq datasets for livestock where this information is not available.
In our analysis we have not considered either parent-of-origin or breed-of-origin-
specific effects in this analysis. For parent-of-origin or breed-of-origin assignment of these
ASE profiles DNA level genotypes from the parents of the six sheep from the gene expression
atlas (i.e. Texel sire and Scottish Blackface dam) would be required, and these are unfortunately
not available. In this study ASE expression profiles also might be affected by the direction of
the cross (i.e. Texel sire x Scottish Blackface dam). To fully characterise parent-of-origin or
breed-of-origin reciprocal cross experiments would be required. Reciprocal cross studies in
mouse (Huang et al., 2017), chicken (Zhuo et al., 2017) and crossbred cattle (Chen et al.,
2016b) have shed light on the complexity of such pervasive ASE markers and parent-of-origin
effects. Though potentially very interesting these experiments are lengthy and costly to perform
in sheep. Particularly in this case because the reciprocal cross (Scottish Blackface dam x Texel
sire) is rarely used in the UK sheep industry and as a consequence has limited relevance to
production.
Our approach also excludes mono-allelic expression. The minimum filtration criteria
utilised in our workflow along with the reference mapping bias removal step ensures an
unbiased ASE discovery in the transcriptome by excluding the ambiguity surrounding MAE
variants. This form of analysis is based on the principle that absence of evidence (reads) for
either allele of a heterozygote site does not directly amount to evidence of their absence i.e.
MAE. The pattern of ASE (ratio of Alt/Ref+Alt) is dependent on the bi-allelic expression of
loci within the genomic coordinates of the gene or genomic element of interest. For an ASE
effect to be captured by the GeneiASE model, the following criteria must be met: i) Biallelic
expression of the locus; ii) Min depth criteria for each allele (min 3 reads , total 10 reads at that
site and > 1% of total reads containing that allele); iii) The allelic imbalance or departure from
bi-allelic balanced expression being inducible by an environmental trigger (i.e. LPS in ICD
ASE experiment with BMDM data). These stringent criteria secure robust transcriptome-wide
ASE discovery while maximising the usage of read counts from short read RNA-Seq datasets
without considering mono-allelic sites. MAE patterns are impossible to differentiate from
sequencing error or random nonsense mediated decay in total RNA-Seq, unless arbitrary cut-
offs are introduced such as ratio of allelic read counts > 0.9 (Chamberlain et al., 2015) or > 0.7
(Cao et al., 2019). We decided to exclude MAE from this study using read count bi-allelic
expression filtration because it is difficult to distinguish between sequencing error and MAE.
We do appreciate that this form of filtration might lead to a reduced number of ASE genes
discoveries overall and will exclude potentially imprinted loci altogether.
The tissues utilised for ASE analysis in this study (thymus, spleen, liver and ileum) are
highly influential on performance of the immune system. ASE profiles shared across tissues
and cell types were limited and instead they tended to be highly specific. We identified tissue-
specific ASE in several genes in Thymus, for example, that are involved in the T-cell mediated
immune response, including CD47 and CD244. These tissue-specific and cell-type-specific
ASE profiles may underlie the expression of economically important traits such as disease
resistance. Assessment of the connection between economically relevant phenotypes and
tissue-specific ASE profiles could be useful for the improvement of genomics enabled sheep
breeding programmes, particularly those using specialised sire and dam lines (Georges et al.,
2018). Loci exhibiting ASE have been associated with production traits including milk-fat
percentage (Hayes et al., 2010; Suárez-Vega et al., 2017), trypanotolerance in small ruminants
(Kadarmideen et al., 2011; Álvarez et al., 2016), mastitis in goat (Ilie et al., 2018), Johne's
disease in cattle (Mallikarjunappa et al., 2018) and Marek's disease in chicken (Maceachern et

al., 2011; Meydan et al., 2011; Cheng et al., 2015). Although there is no general consensus
currently on the correlation of allelic expression haplotypes and phenotypes under selection in
sheep, this form of ASE analysis could pave the way for functional validation at population
level (e.g. breed or haplotype-specific aseQTL studies in a larger population of sheep).
Examples of population level aseQTL, eQTL and sQTL (QTLs associated with RNA splicing)
already exist for cattle (Wang et al., 2018; Xiang et al., 2018). Knowledge of favourable ASE
in critical genes involved in traits of interest could be used as a performance indicator or
included as weighted SNVs in genomic prediction algorithms to enhance livestock breeding
programmes (Georges et al., 2018). Currently the UK sheep industry is on the cusp of applying
genomic prediction, but suitable genomics enabled breeding programmes for sheep already
exist in New Zealand and Australia (Daetwyler et al., 2010).
561
562 **Conclusions**
563
In this study we characterise extreme to moderate allele-specific expression, at the gene
and SNV level, in immune-related tissues and cells from six adult sheep (TxBF) from the sheep
gene expression atlas dataset. Reference mapping bias removal was an integral component of
the analysis pipeline applied in this study. The correction of reference bias prior to obtaining
the allelic read counts is a critical step toward true ASE discovery. The workflow developed
as part of this manuscript provides an RNA-Seq only dependent tool, without the need for
individual DNA sequences. We note that the stringent filtering process applied would remove
loci where the allelic imbalance was less extreme but might still be of biological significance.
This study is a novel analysis of an existing large-scale complex RNA-Seq dataset from
sheep. Using the pipeline we have adapted for this analysis, we were able to identify ASE
profiles that were pervasive in each sheep and specific to the tissues and cell types investigated.
These tissue and cell-type-specific ASE profiles may underlie the expression of economically
important traits and could be used to identify variants that could be weighted in genomic
prediction algorithms for the improvement of sheep breeding programmes. In summary we
have adapted a robust methodology for ASE profiling, using the sheep gene expression atlas
dataset, and provided a foundation for identifying the regulatory and expressed elements of the
genome that are driving complex traits in livestock.
581
582 **Data Availability**
583
The RNA-Seq sequence data is available via the European Nucleotide Archive (ENA) under
PRJEB19199 (https://www.ebi.ac.uk/ena/data/view/PRJEB19199). The BAM files were
already produced as part of the sheep gene expression atlas following the publicly available
protocol (FAANG, 2018) and as described in (Clark et al. 2017). The BAM files have been
uploaded to ENA under accession numbers ERZ827944, ERZ827949, ERZ827951,
ERZ827955, ERZ827972, ERZ827988, ERZ827995, ERZ827997, ERZ828001, ERZ828016,
ERZ828019, ERZ828036, ERZ828044, ERZ828046, ERZ828050, ERZ828070, ERZ828073,
ERZ828160, ERZ828167, ERZ828168, ERZ828172, ERZ828188, ERZ828192, ERZ828209,
ERZ828215, ERZ828217, ERZ828221, ERZ828240, ERZ828244, ERZ828261, ERZ828268,
ERZ828270, ERZ828274, ERZ828293 and ERZ828297. The Oar v3.1 reference FASTA and
VCF file from Ensembl v92 were used throughout the pipeline. The ASE analysis pipeline
(https://msalavat@bitbucket.org/msalavat/asewrap_public.git) was wrapped using bash
scripting on Edinburgh Compute and Data Facility computing resource Eddie Mark 3
(Edinburgh, 2018). All the raw ASE genes data produced by GeneiASE could be found within
Supplementary_file2.zip.
599

Author's contributions

MS and ELC coordinated and designed the analysis component of the study with assistance from SJB and SPV. MS and SPV designed, optimised and tested the ASE pipeline. DAH acquired the funding for the sheep gene expression atlas project. MEBM, ELC and DAH designed the LPS experiment and generated the data. MEBM performed the LPS stimulation of bone marrow derived macrophages and RNA extraction. SJB performed all bioinformatic analyses prior to analysis with the ASE pipeline. MS performed ASE analysis, visualisation of the results and wrote the manuscript. ELC contributed to manuscript editing and drafting. All authors read and approved the final manuscript.

Acknowledgments

The authors would like to thank University of Edinburgh Computing and Data Support staff (viz. Dr Andy Law and Dr Steve Thorn) for their input towards optimisation of the pipeline and efficient parallel computing. Rachel Young and Lucas Lefevre isolated the bone marrow derived macrophages for the sheep gene expression atlas project and Iseabail Farquhar helped to collect and archive the tissue samples. The authors would also like to thank Professor Kim Summers and Professor Mick Watson for their comments, which helped to improve the manuscript.

Conflict of Interest

The authors have no competing interest regarding the findings presented in this publication.

Funding

The work was supported by a Biotechnology and Biological Sciences Research Council (BBSRC) (http://www.bbsrc.ac.uk) Grant BB/L001209/1 ('Functional Annotation of the Sheep Genome'). Also, BBSRC Institute Strategic Program Grants: 'Farm Animal Genomics' (BBS/E/D/20211550), 'Transcriptomes, Networks and Systems' (BBS/E/D/20211552) and 'Blue Prints for Healthy Animals' (BB/P013732/1). Edinburgh Genomics is partly supported through core grants from BBSRC (BB/J004243/1), NERC (http://www.nerc.ac.uk) (R8/H10/56) and the Medical Research Council (MRC) (https://www.mrc.ac.uk) (MR/K001744/1). SJB was supported by the Roslin Foundation. ELC is supported by the University of Edinburgh Chancellor's Fellowship programme. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

Álvarez, I., Pérez-Pardal, L., Traoré, A., Fernández, I., and Goyache, F. (2016). Lack of specific alleles for the bovine chemokine (C-X-C) receptor type 4 (CXCR4) gene in West African cattle questions its role as a candidate for trypanotolerance. *Infect. Genet. Evol.* 42, 30–3. doi:10.1016/j.meegid.2016.04.029.

Andersson, L., Archibald, A. L., Bottema, C. D., Brauning, R., Burgess, S. C., Burt, D. W., et al. (2015). Coordinated international action to accelerate genome-to-phenome with FAANG, the Functional Annotation of Animal Genomes project. *Genome Biol.* doi:10.1186/s13059-015-0622-4.

Barlow, D. P., and Bartolomei, M. S. (2014). Genomic imprinting in mammals. *Cold Spring*

Harb. Perspect. Biol. 6. doi:10.1101/cshperspect.a018382.

Barton, S. J., Crozier, S. R., Lillycrop, K. A., Godfrey, K. M., and Inskip, H. M. (2013).
Correction of unexpected distributions of P values from analysis of whole genome
arrays by rectifying violation of statistical assumptions. doi:10.1186/1471-2164-14-161.

Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical
and Powerful Approach to Multiple Testing on JSTOR. *Source J. R. Stat. Soc. Ser. B* 57,
289–300. Available at:
https://www.jstor.org/stable/pdf/2346101.pdf?refreqid=excelsior%3A8080132077b9217
878a58e47abbcf2d8 [Accessed October 27, 2015].

Bray, N. L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic
RNA-seq quantification. *Nat. Biotechnol.* 34, 525–527. doi:10.1038/nbt.3519.

Bush, S. J., McCulloch, M. E., Muriuki, C., Salavati, M., Davis, G. M., Farquhar, I. L., et al.
(2018). Comprehensive transcriptional profiling of the gastrointestinal tract of ruminants
from birth to adulthood reveals strong developmental stage specific gene expression.
bioRxiv, 364752. doi:10.1101/364752.

Cao, Y., Xu, H., Li, R., Gao, S., Chen, N., Luo, J., et al. (2019). Genetic Basis of Phenotypic
Differences Between Chinese Yunling Black Goats and Nubian Goats Revealed by
Allele-Specific Expression in Their F1 Hybrids. *Front. Genet.* 10, 145.
doi:10.3389/fgene.2019.00145.

Castel, S. E., Levy-Moonshine, A., Mohammadi, P., Banks, E., and Lappalainen, T. (2015).
Tools and best practices for data processing in allelic expression analysis. *Genome Biol.*
16, 195. doi:10.1186/s13059-015-0762-6.

Castel, S. E., Mohammadi, P., Chung, W. K., Shen, Y., and Lappalainen, T. (2016).
ARTICLE Rare variant phasing and haplotypic expression from RNA sequencing with
phASER. *Nat. Commun.* 7. doi:10.1038/ncomms12817.

Chamberlain, A. J., Vander Jagt, C. J., Hayes, B. J., Khansefid, M., Marett, L. C., Millen, C.
A., et al. (2015). Extensive variation between tissues in allele specific expression in an
outbred mammal. *BMC Genomics* 16, 993. doi:10.1186/s12864-015-2174-0.

Chen, J., Rozowsky, J., Galeev, T. R., Harmanci, A., Kitchen, R., Bedford, J., et al. (2016a).
ARTICLE A uniform survey of allele-specific binding and expression over 1000-
Genomes-Project individuals. doi:10.1038/ncomms11101.

Chen, Z., Hagen, D. E., Wang, J., Elsik, C. G., Ji, T., Siqueira, L. G., et al. (2016b). Global
assessment of imprinted gene expression in the bovine conceptus by next generation
sequencing. *Epigenetics* 11, 501–16. doi:10.1080/15592294.2016.1184805.

Cheng, H. H., Perumbakkam, S., Pyrkosz, A. B., Dunn, J. R., Legarra, A., and Muir, W. M.
(2015). Fine mapping of QTL and genomic prediction using allele-specific expression
SNPs demonstrates that the complex trait of genetic resistance to Marek's disease is
predominantly determined by transcriptional regulation. *BMC Genomics* 16, 816.
doi:10.1186/s12864-015-2016-0.

Clark, E. L., Bush, S. J., McCulloch, M. E. B., Farquhar, I. L., Young, R., Lefevre, L., et al.
(2017). A high resolution atlas of gene expression in the domestic sheep (Ovis aries).
PLOS Genet. 13, e1006997. doi:10.1371/journal.pgen.1006997.

Daetwyler, H. D., Hickey, J. M., Henshall, J. M., Dominik, S., Gredler, B., Van Der Werf, J.
H. J., et al. (2010). Accuracy of estimated genomic breeding values for wool and meat
traits in a multi-breed sheep population. *Anim. Prod. Sci.* doi:10.1071/AN10096.

Degner, J. F., Marioni, J. C., Pai, A. A., Pickrell, J. K., Nkadori, E., Gilad, Y., et al. (2009).
Effect of read-mapping biases on detecting allele-specific expression from RNA-
sequencing data. *Bioinforma. Orig. Pap.* 25, 3207–321210.
doi:10.1093/bioinformatics/btp579.

Deguchi, A., Tomita, T., Omori, T., Komatsu, A., Ohto, U., Takahashi, S., et al. (2013).

Serum amyloid A3 binds MD-2 to activate p38 and NF-κB pathways in a MyD88-
701   dependent manner. *J. Immunol.* 191, 1856–64. doi:10.4049/jimmunol.1201996.
702   DeVeale, B., van der Kooy, D., and Babak, T. (2012). Critical Evaluation of Imprinted Gene
703   Expression by RNA-Seq: A New Perspective. *PLoS Genet.* 8, e1002600.
704   doi:10.1371/journal.pgen.1002600.
705   Dewey, M. (2016). Comparison of various methods for meta–analysis of p–values. Available
706   at: http://www.dewey.myzen.co.uk/meta/comparemet.pdf [Accessed May 8, 2018].
707   Dewey, M. (2018). {metap}: meta-analysis of significance values.
708   Edsgärd, D., Iglesias, M. J., Reilly, S.-J., Hamsten, A., Tornvall, P., Odeberg, J., et al. (2016).
709   GeneiASE: Detection of condition-dependent and static allele-specific expression from
710   RNA-seq data without haplotype information. *Sci. Rep.* 6, 21134.
711   doi:10.1038/srep21134.
712   FAANG (2018). Sheep Atlas Pipeline SOP. Available at:
713   ftp://ftp.faang.ebi.ac.uk/ftp/protocols/analysis/Roslin_SOP_Alignment-based RNA-Seq-
714   Processing_20170917.pdf.
715   Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P., et al.
716   (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res.* 46, D649–D655.
717   doi:10.1093/nar/gkx1132.
718   García-Gámez, E., Reverter, A., Whan, V., McWilliam, S. M., Arranz, J. J., Kijas, J., et al.
719   (2011). Using Regulatory and Epistatic Networks to Extend the Findings of a Genome
720   Scan: Identifying the Gene Drivers of Pigmentation in Merino Sheep. *PLoS One* 6,
721   e21158. doi:10.1371/journal.pone.0021158.
722   Georges, M., Charlier, C., and Hayes, B. (2018). Harnessing genomic information for
723   livestock improvement. *Nat. Rev. Genet.*, 1. doi:10.1038/s41576-018-0082-2.
724   Georgoudaki, A.-M., Khodabandeh, S., Puiac, S., Persson, C. M., Larsson, M. K., Lind, M.,
725   et al. (2015). CD244 is expressed on dendritic cells and regulates their functions.
726   *Immunol. Cell Biol.* 93, 581–590. doi:10.1038/icb.2014.124.
727   Goodhead, I., Archibald, A., Amwayi, P., Brass, A., Gibson, J., Hall, N., et al. (2010). A
728   comprehensive genetic analysis of candidate genes regulating response to Trypanosoma
729   congolense infection in mice. *PLoS Negl. Trop. Dis.* 4, e880.
730   doi:10.1371/journal.pntd.0000880.
731   GTEx Consortium, Gte., Laboratory, Data Analysis &Coordinating Center (LDACC)—
732   Analysis Working Group, Statistical Methods groups—Analysis Working Group,
733   Enhancing GTEx (eGTEx) groups, NIH Common Fund, NIH/NCI, et al. (2017). Genetic
734   effects on gene expression across human tissues. *Nature* 550, 204–213.
735   doi:10.1038/nature24277.
736   Guillocheau, G. M., El Hou, A., Meersseman, C., Esquerré, D., Rebours, E., Letaief, R., et al.
737   (2019). Survey of allele specific expression in bovine muscle. *Sci. Rep.* 9, 4297.
738   doi:10.1038/s41598-019-40781-6.
739   Gutierrez-Arcelus, M., Baglaenko, Y., Arora, J., Hannes, S., Luo, Y., Amariuta, T., et al.
740   (2019). Allele-specific expression changes dynamically during T cell activation in HLA
741   and other autoimmune loci. *bioRxiv*, 599449. doi:10.1101/599449.
742   Habermann, A., Schroer, T. A., Griffiths, G., and Burkhardt, J. K. (2001).
743   Immunolocalization of cytoplasmic dynein and dynactin subunits in cultured
744   macrophages: enrichment on early endocytic organelles. *J. Cell Sci.* 114, 229–240.
745   Available at: http://www.ncbi.nlm.nih.gov/pubmed/11112706 [Accessed March 11,
746   2019].
747   Hach, F., Sarrafi, I., Hormozdiari, F., Alkan, C., Eichler, E. E., and Sahinalp, S. C. (2014).
748   mrsFAST-Ultra: a compact, SNP-aware mapper for high performance sequencing
749   applications. *Nucleic Acids Res.* 42, W494-500. doi:10.1093/nar/gku370.

Harrison, P. W., Fan, J., Richardson, D., Clarke, L., Zerbino, D., Cochrane, G., et al. (2018).
FAANG, establishing metadata standards, validation and best practices for the farmed
and companion animal community. *Anim. Genet.* 49, 520–526. doi:10.1111/age.12736.
Hasin-Brumshtein, Y., Hormozdiari, F., Martin, L., van Nas, A., Eskin, E., Lusis, A. J., et al.
(2014). Allele-specific expression and eQTL analysis in mouse adipose tissue. *BMC*
Genomics 15, 471. doi:10.1186/1471-2164-15-471.
Hayes, B. J., and Daetwyler, H. D. (2019). 1000 Bull Genomes Project to Map Simple and
Complex Genetic Traits in Cattle: Applications and Outcomes. *Annu. Rev. Anim. Biosci.*
7, 89–102. doi:10.1146/annurev-animal-020518-115024.
Hayes, B. J., Pryce, J., Chamberlain, A. J., Bowman, P. J., and Goddard, M. E. (2010).
Genetic architecture of complex traits and accuracy of genomic prediction: coat colour,
milk-fat percentage, and type in Holstein cattle as contrasting model traits. *PLoS Genet.*
6, e1001139. doi:10.1371/journal.pgen.1001139.
Hodgkinson, A., Grenier, J.-C., Gbeha, E., and Awadalla, P. (2016). A haplotype-based
normalization technique for the analysis and detection of allele specific expression.
BMC Bioinformatics. doi:10.1186/s12859-016-1238-8.
Huang, W. C., Ferris, E., Cheng, T., Hörndli, C. S., Gleason, K., Tamminga, C., et al. (2017).
Diverse Non-genetic, Allele-Specific Expression Effects Shape Genetic Architecture at
the Cellular Level in the Mammalian Brain. *Neuron* 93, 1094-1109.e7.
doi:10.1016/j.neuron.2017.01.033.
Ilie, D. E., Kusza, S., Sauer, M., and Gavojdian, D. (2018). Genetic characterization of
indigenous goat breeds in Romania and Hungary with a special focus on genetic
resistance to mastitis and gastrointestinal parasitism based on 40 SNPs. *PLoS One* 13,
e0197051. doi:10.1371/journal.pone.0197051.
Jiang, Y., Xie, M., Chen, W., Talbot, R., Maddox, J. F., Faraut, T., et al. (2014). The sheep
genome illuminates biology of the rumen and lipid metabolism. *Science* 344, 1168–
1173. doi:10.1126/science.1252806.
Kadarmideen, H. N., Watson-Haigh, N. S., and Andronicos, N. M. (2011). Systems biology
of ovine intestinal parasite resistance: disease gene modules and biomarkers. *Mol.*
BioSyst. 7, 235–246. doi:10.1039/C0MB00190B.
Kaul, S., Mittal, S. K., Feigenbaum, L., Kruhlak, M. J., and Roche, P. A. (2015). Expression
of the SNARE protein SNAP-23 is essential for cell survival. *PLoS One* 10, e0118311.
doi:10.1371/journal.pone.0118311.
Kijas, J. W., Lenstra, J. A., Hayes, B., Boitard, S., Porto Neto, L. R., San Cristobal, M., et al.
(2012). Genome-wide analysis of the world's sheep breeds reveals high levels of historic
mixture and strong recent selection. *PLoS Biol.* 10, e1001258.
doi:10.1371/journal.pbio.1001258.
Kim, D., Langmead, B., and Salzberg, S. L. (2015). HISAT: a fast spliced aligner with low
memory requirements. *Nat. Methods* 12, 357–360. doi:10.1038/nmeth.3317.
Larson, M. A., Wei, S. H., Weber, A. T. A., Weber, A. T. A., and McDonald, T. L. (2003).
Induction of human mammary-associated serum amyloid A3 expression by prolactin or
lipopolysaccharide. *Biochem. Biophys. Res. Commun.* 301, 1030–1037.
doi:10.1016/S0006-291X(03)00045-7.
Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* 25, 1754–60. doi:10.1093/bioinformatics/btp324.
Maceachern, S., Muir, W. M., Crosby, S., and Cheng, H. H. (2011). Genome-wide
identification of allele-specific expression (ASE) in response to Marek's disease virus
infection using next generation sequencing. *BMC Proc.* doi:10.1186/1753-6561-5-S4-
S14.
Mallikarjunappa, S., Sargolzaei, M., Brito, L. F., Meade, K. G., Karrow, N. A., and Pant, S.

D. (2018). Short communication: Uncovering quantitative trait loci associated with
resistance to Mycobacterium avium ssp. paratuberculosis infection in Holstein cattle
using a high-density single nucleotide polymorphism panel. *J. Dairy Sci.* 101, 7280–
7286. doi:10.3168/JDS.2018-14388.
Mayba, O., Gilbert, H. N., Liu, J., Haverty, P. M., Jhunjhunwala, S., Jiang, Z., et al. (2014).
MBASED: allele-specific expression detection in cancer tissues and cell lines. *Genome*
Biol. 15, 405. doi:10.1186/s13059-014-0405-3.
McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., et al.
(2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Res.* 20, 1297–303.
doi:10.1101/gr.107524.110.
McNerney, M. E., Lee, K.-M., and Kumar, V. (2005). 2B4 (CD244) is a non-MHC binding
receptor with multiple functions on natural killer cells and CD8+ T cells. *Mol. Immunol.*
42, 489–494. doi:10.1016/J.MOLIMM.2004.07.032.
Meydan, H., Yildiz, M. A., Dodgson, J. B., and Cheng, H. H. (2011). Allele-specific
expression analysis reveals CD79B has a cis-acting regulatory element that responds to
Marek's disease virus infection in chickens. *Poult. Sci.* doi:10.3382/ps.2010-01295.
Miao, Z., Alvarez, M., Pajukanta, P., and Ko, A. (2018). ASElux: an ultra-fast and accurate
allelic reads counter. *Bioinformatics* 34, 1313–1320. doi:10.1093/bioinformatics/btx762.
Montgomery, S. B., Sammeth, M., Gutierrez-Arcelus, M., Lach, R. P., Ingle, C., Nisbett, J.,
et al. (2010). Transcriptome genetics using second generation sequencing in a Caucasian
population. *Nature* 464, 773–777. doi:10.1038/nature08903.
Mozaffari, S. V, Stein, M. M., Magnaye, K. M., Nicolae, D. L., and Ober, C. (2018). Parent
of origin gene expression in a founder population identifies two new candidate imprinted
genes at known imprinted regions. *PLoS One* 13, e0203906.
doi:10.1371/journal.pone.0203906.
Mueller, O., Lightfoot, S., and Schroeder, A. (2004). RNA Integrity Number (RIN) –
Standardization of RNA Quality Control Application. *Agil. Appl. Note, Publ. 5989-*
1165EN, 1–8. doi:10.1101/gr.189621.115.7.
Nitta, T., and Takahama, Y. (2007). The lymphocyte guard-IANs: regulation of lymphocyte
survival by IAN/GIMAP family proteins. *Trends Immunol.* 28, 58–65.
doi:10.1016/j.it.2006.12.002.
Ocaña-Guzman, R., Vázquez-Bolaños, L., and Sada-Ovalle, I. (2018). Receptors That Inhibit
Macrophage Activation: Mechanisms and Signals of Regulation and Tolerance. *J.*
Immunol. Res. 2018, 8695157. doi:10.1155/2018/8695157.
Overman, R. G., Enderle, P. J., Farrow, J. M., Wiley, J. E., and Farwell, M. A. (2003). The
human mitochondrial translation initiation factor 2 gene (MTIF2): Transcriptional
analysis and identification of a pseudogene. *Biochim. Biophys. Acta - Gene Struct. Expr.*
1628, 195–205. doi:10.1016/S0167-4781(03)00144-1.
Pelikan, R. C., Kelly, J. A., Fu, Y., Lareau, C. A., Tessneer, K. L., Wiley, G. B., et al. (2018).
Enhancer histone-QTLs are enriched on autoimmune risk haplotypes and influence gene
expression within chromatin networks. *Nat. Commun.* 9, 2905. doi:10.1038/s41467-018-
05328-9.
Pounds, S., and Cheng, C. (2006). Robust estimation of the false discovery rate. 22, 1979–
1987. doi:10.1093/bioinformatics/btl328.
R Core Team (2017). R: A language and environment for statistical computing. Available at:
https://www.r-project.org/.
Raghupathy, N., Choi, K., Vincent, M. J., Beane, G. L., Sheppard, K. S., Munger, S. C., et al.
(2018). Hierarchical analysis of RNA-seq reads improves the accuracy of allele-specific
expression. *Bioinformatics* 34, 2177–2184. doi:10.1093/bioinformatics/bty078.

Renfree, M. B., Hore, T. A., Shaw, G., Graves, J. A. M., Pask, A. J., Marshall Graves, J. A.,
et al. (2009). Evolution of Genomic Imprinting: Insights from Marsupials and
Monotremes. *Annu. Rev. Genomics Hum. Genet.* 10, 241–262. doi:10.1146/annurev-
genom-082908-150026.

Richard Albert, J., Koike, T., Younesy, H., Thompson, R., Bogutz, A. B., Karimi, M. M., et
al. (2018). Development and application of an integrated allele-specific pipeline for
methylomic and epigenomic analysis (MEA). *BMC Genomics* 19, 463.
doi:10.1186/s12864-018-4835-2.

Rozowsky, J., Abyzov, A., Wang, J., Alves, P., Raha, D., Harmanci, A., et al. (2011).
AlleleSeq: Analysis of allele-specific expression and binding in a network framework.
Mol. Syst. Biol. doi:10.1038/msb.2011.54.

Shi, Q., Zhuang, F., Liu, J.-T., Li, N., Chen, Y.-X., Su, X.-B., et al. (2019). Single-cell
analyses reveal functional classification of dendritic cells and their potential roles in
inflammatory disease. *FASEB J.* 33, 3784–3794. doi:10.1096/fj.201801489R.

Smith, R. M., Webb, A., Papp, A. C., Newman, L. C., Handelman, S. K., Suhy, A., et al.
(2013). Whole transcriptome RNA-Seq allelic expression in human brain. *BMC*
Genomics. doi:10.1186/1471-2164-14-571.

Stevenson, K. R., Coolon, J. D., and Wittkopp, P. J. (2013). Sources of bias in measures of
allele-specific expression derived from RNA-seq data aligned to a single reference
genome. *BMC Genomics* 14, 1. doi:10.1186/1471-2164-14-536.

Suárez-Vega, A., Gutiérrez-Gil, B., Klopp, C., Tosser-Klopp, G., and Arranz, J. J. (2017).
Variant discovery in the sheep milk transcriptome using RNA sequencing. *BMC*
Genomics 18, 170. doi:10.1186/s12864-017-3581-1.

van de Geijn, B., McVicker, G., Gilad, Y., and Pritchard, J. K. (2015). WASP: allele-specific
software for robust molecular quantitative trait locus discovery. *Nat. Methods* 12, 1061–
1063. doi:10.1038/nmeth.3582.

Van den Bossche, J., Laoui, D., Morias, Y., Movahedi, K., Raes, G., De Baetselier, P., et al.
(2012). Claudin-1, Claudin-2 and Claudin-11 Genes Differentially Associate with
Distinct Types of Anti-inflammatory Macrophages In vitro and with Parasite- and
Tumour-elicited Macrophages In vivo. *Scand. J. Immunol.* 75, 588–598.
doi:10.1111/j.1365-3083.2012.02689.x.

Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., Del Angel, G., Levy-
Moonshine, A., et al. (2013). From FastQ data to high confidence variant calls: the
Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinforma.* 43, 11.10.1-
33. doi:10.1002/0471250953.bi1110s43.

Wang, M., Hancock, T. P., Chamberlain, A. J., Vander Jagt, C. J., Pryce, J. E., Cocks, B. G.,
et al. (2018). Putative bovine topological association domains and CTCF binding motifs
can reduce the search space for causative regulatory variants of complex traits. *BMC*
Genomics 19, 395. doi:10.1186/s12864-018-4800-0.

Wickham, H., Francois, R., Henry, L., and Müller, K. (2019). Package 'dplyr'. A Grammar
of Data Manipulation. *R Packag. version 0.8.0.1*, 1–88. Available at:
https://cran.rproject.org/web/packages/dplyr/dplyr.pdf.

Wong, P. P. C., Daneman, N., Volchuk, A., Lassam, N., Wilson, M. C., Klip, A., et al.
(1997). Tissue Distribution of SNAP-23 and Its Subcellular Localization in 3T3-L1
Cells. *Biochem. Biophys. Res. Commun.* 230, 64–68. doi:10.1006/BBRC.1996.5884.

Xiang, R., Hayes, B. J., Vander Jagt, C. J., MacLeod, I. M., Khansefid, M., Bowman, P. J., et
al. (2018). Genome variants associated with RNA splicing variation in bovine are
extensively shared between tissues Ruidong. *BMC Genomics* 19, 521.
doi:10.1186/s12864-018-4902-8.

Xin, H., Lee, D., Hormozdiari, F., Yedkar, S., Mutlu, O., and Alkan, C. (2013). Accelerating

read mapping with FastHASH. *BMC Genomics 2013 141* 14, S13. doi:10.1186/1471-
2164-14-s1-s13.
Yamanegi, K., Yamada, N., Nakasho, K., and Nishiura, H. (2018). Participation of delta
annexin A3 in the ribosomal protein S19 C-terminus-dependent inhibitory mechanism of
the neutrophil C5a receptor through delta lactoferrin. *Pathol. Int.* 68, 109–116.
doi:10.1111/pin.12626.
Yee, N. K., and Hamerman, J. A. (2013). β(2) integrins inhibit TLR responses by regulating
NF-κB pathway and p38 MAPK activation. *Eur. J. Immunol.* 43, 779–92.
doi:10.1002/eji.201242550.
Zerbino, D. R., Achuthan, P., Akanni, W., Amode, M. R., Barrell, D., Bhai, J., et al. (2018).
Ensembl 2018. *Nucleic Acids Res.* 46, D754–D761. doi:10.1093/nar/gkx1098.
Zerbino, D. R., and Birney, E. (2008). Velvet: Algorithms for de novo short read assembly
using de Bruijn graphs. *Genome Res.* doi:10.1101/gr.074492.107.
Zhou, Z.-Y., Hu, Y., Li, A., Li, Y.-J., Zhao, H., Wang, S.-Q., et al. (2018). Genome wide
analyses uncover allele-specific RNA editing in human and mouse. *Nucleic Acids Res.*
46, 8888–8897. doi:10.1093/nar/gky613.
Zhuo, Z., Lamont, S. J., and Abasht, B. (2017). RNA-Seq Analyses Identify Frequent Allele
Specific Expression and No Evidence of Genomic Imprinting in Specific Embryonic
Tissues of Chicken. *Sci. Rep.* 7, 11944. doi:10.1038/s41598-017-12179-9.

919
920 **Tables**
921

922 **Table 1**. Total number of genes with significant static ASE in proportion to genes containing
923 informative SNVs (filtered). Total expressed: Average number of genes being expressed in all
924 4 tissues. Total filtered: Average number of genes (containing heterozygote loci) passing read
925 bi-allelic filtration criteria in 4 tissues. Tissue break-down has been presented as count
926 (%ASE/filtered).
927

Sheep	Thymus	Spleen	Liver	Ileum	Total filtered	Total expressed
Female 1	-	136 (1.31%)	153 (1.47%)	92 (0.88%)	10,379	21,150
Female 2	136 (1.28%)	70 (0.66%)	116 (1.09%)	70 (0.66%)	10,572	21,420
Female 3	151 (1.22%)	75 (0.60%)	105 (0.85%)	140 (1.13%)	12,326	21,689
Avg.	143 (1.25%)	94 (0.85%)	125 (1.13%)	101 (0.89%)	11,092	21,419
Male 1	95 (0.92%)	-	157 (1.52%)	86 (0.83%)	10,282	21,361
Male 2	125 (0.99%)	54 (0.43%)	86 (0.68%)	80 (0.63%)	12,514	21,822
Male 3	106 (1.01%)	71 (0.67%)	110 (1.04%)	82 (0.78%)	10,480	21,545
Avg.	109 (0.97%)	62 (0.55%)	118 (1.08%)	83 (0.74%)	11,092	21,576
Total avg.	126 (1.13%)	78 (0.70%)	121 (1.09%)	92 (0.82%)	11,092	21,497

928
929 **Table 2**. Total number of genes with significant static ASE in BMDMs +/- LPS.
930

Sheep	BMDM 0hrs -LPS	BMDM 7hrs +LPS
Female 1	237	252

Female 2	192	233
Female 3	227	263
Female avg.	219	249
Male 1	205	193
Male 2	300	260
Male 3	260	261
Male avg.	255	238

933 **Table 3**. The variant IDs of LPS inducible ASE SNVs (Fisher's exact) and their respective
934 genes. Data was obtained using Ensembl BioMart query builder. Highly significant SNVs are
935 highlighted in bold (FDR< 1.0e-08)

ID	Chr	Position	Gene ID	Gene Name
rs418350332	1	195297663	ENSOARG00000020472	CLDN1
rs159822214	3	112164732	ENSOARG00000014876	OSBPL8
rs159926581	**3**	**214731375**	**ENSOARG00000016495**	**-**
rs159926581	**3**	**214731375**	**ENSOARG00000022372**	**RF00221**
rs159926581	**3**	**214731375**	**ENSOARG00000024737**	**RF00593**
rs159926581	**3**	**214731375**	**ENSOARG00000025124**	**RF01151**
rs159926581	**3**	**214731375**	**ENSOARG00000025150**	**RF00593**
rs193634916	4	53430934	ENSOARG00000001754	MDFIC
rs418697910	4	53430968	ENSOARG00000001754	MDFIC
rs162298949	6	92979793	ENSOARG00000018710	ANXA3
rs162298949	6	92979793	ENSOARG00000018710	ANXA3
rs160665956	9	76759747	ENSOARG00000001261	SPAG1
rs424104956	13	46654781	ENSOARG00000017292	CDS2
rs424104956	13	46654781	ENSOARG00000017292	CDS2
rs424104956	13	46654781	ENSOARG00000022618	RF00377
rs420037698	**16**	**6887423**	**ENSOARG00000004700**	**-**
rs430667535	**17**	**50485358**	**ENSOARG00000017707**	**BRI3BP**
rs430667535	**17**	**50485358**	**ENSOARG00000018177**	**UBC**
rs403064928	21	25826884	ENSOARG00000009963	SAA3
rs405439099	21	25826990	ENSOARG00000009963	SAA3
rs412192652	**21**	**25826978**	**ENSOARG00000009963**	**SAA3**
rs161531178	26	19663270	ENSOARG00000009626	MSR1

939 **Figure captions**

941
942 **Figure 1**. A flowchart of the allele-specific expression analysis pipeline applied to the sheep
943 gene expression atlas dataset and optimised for WASP and GeneiASE programmes. The
944 remapping was carried out using HISAT2 (Kim et al., 2015) in combination with SAMtools
945 (Li et al., 2009). The Genome Analysis Toolkit v 3.8 was used for the ASE read counting
946 section.
947

948
Figure 2. Distribution of biallelic SNVs expressed per gene in each of the 6 TxBF sheep. The
total number of SNVs was averaged across thymus, liver, ileum and spleen for every animal.
Over 5e+07 SNVs were gathered using Ensembl v.92 VCF track. The total number of SNVs
per genes is averaged across 4 tissue RNA-Seq in each animal (~5.9e+06). **A)** Histogram of
SNVs per gene counts in the reference track (Ensembl in grey) and 6 sheep in red (females)
and blue (males) overlaid. **B).** The overall numbers of genes and SNVs detected in each animal
(averaged over 4 tissues). **C)** Individual histograms from section A with females in red and
males in blue.
957

958
Figure 3. The histogram of a global reference allelic ratio at every locus in the tissues. The
distribution of ref allelic ratio showed a balanced profile without any 0 or 1 inflation which is
observed in the presence of reference mapping bias. The allelic ratio above 0.51 is shown in
blue and below 0.49 in red while balanced bi-allelic expression (0.49-0.51) is coloured in grey.
Ref.dp = read counts for reference allele. Alt.dp = read counts for alternate allele. The y axis is
square root scaled. As discussed in the text SNP that display MAE are not present in any of the
samples analysed, indicating there was no inflation in either 0 or 1 allelic ratio.
966

967
968 **Figure 4**. Genes exhibiting static ASE shared across tissues from all six sheep. The x axis
969 represents the mean allelic imbalance (averaged static ASE across sheep in each tissue). **A)**
970 genes shared by four tissues with significant (FDR < 0.1) static ASE. **B)** ASE genes private to
971 Ileum. **C)** Private to liver. **D)** Private to spleen. **E)** Private to thymus.
972

Figure 5. Intersectionality analysis of genes expressing significant ASE across all six sheep.
In each tissue from left to right the set count of genes (dots connected by lines) illustrates the
number of sheep sharing the gene. The private sets of genes are located at the far right of each
graph (single dots with no line). The intersections are coloured in to illustrate the size of the
set of shared genes (red [common to all 6 sheep], green [shared by 5 or 4 sheep], yellow [only
in females] and purple [only in males]). Detailed lists of genes with ASE shared by at least 4
sheep are presented above each graph for **A)** Ileum, **B)** Liver, **C)** Spleen and **D)** Thymus. Two
sex-specific sets of genes are highlighted: 16 genes showing ASE only in females (in yellow)
and 5 genes only in males (in purple).
983

984
985 **Figure 6**. Intersection analysis of SNVs under genes with significant ICD-ASE in the BMDMs
986 ± LPS. From left to right the set number of genes (dots connected by lines) have been illustrated
987 in order according to the number of sheep sharing the SNV. The private sets of SNVs are
988 located at the far right of each graph (single dots with no line).
989

Figure 7. Scatter plot of the adjusted p values from Fisher's exact test (unified using Stouffer unification) in BMDMs comparing expression from different alleles at 0 vs 7hrs at SNV level (LPS inducible ASE). **A)** The graph shows 646 loci exhibiting LPS inducible allelic imbalance shared across all 6 sheep. **B)** Four loci on chromosomes 3,16,17 and 21 with FDR < 1.0e-08. FDR < 1.0e-02 red line (n=16 SNVs) and FDR < 1.0e-08 blue line (n = 4 SNVs).

1024 **Supplementary material**
1025
1026 A bimodal p value distribution was observed in GeneiASE output as shown in S1 Fig.1:
1027

1028
1029
Figure S1. Distribution of p values resulted from the beta-binomial test (Static ASE
example for spleen in female 1). The uniformity of the p value distribution showed a
bimodal (high frequency close to 1 followed by an extensive lower tail close to 0). R
machine characteristics for printing float numbers were limited to 6 significant numbers
after the decimal point by default which resulted in a high number of absolute zero (0+e0.0)
output-values.
1036

1037
1038
Figure S2. The figure shows the distribution of p values produced by GeneiASE after cut
off introduction (A) and after removal of p value == 1 values (B). The distribution of sorted
FDR calculated by GeneiASE and after correction are shown in C (right and left
respectively). To correct for multiple testing by recalculation of FDR and eliminating the
non-uniformity of p value distribution around 1 the following steps were performed as
shown in S1 Fig.2: i) An arbitrary cut off (1e-7) was placed over all p values smaller than
the said limit including the absolute 0e+0.0; ii) The genes showing p values equal to 1 were
removed from each data frame; iii) The false discovery portion and rate were recalculated
in each sample. iv) Gene showing FDR < 0.1 selected for further analysis.
1048

1049
1050   **Figure S3**. The QQ plot of observed vs expected p values in corrected (left) and GeneiASE's
1051   raw (right) calculated outputs.
1052
1053   **A**

1054
1055   **B**

1056
1057 **Figure S4.** The LD block (A) and LD Manhattan plot (B) of SAA3 transcript
1058 (ENSOART00000010843). The LD block produced as part of International Sheep Genome
1059 Consortium is publicly available on Ensembl using the COMPOSITE population (n = 98).
1060

1061
1062 **Figure S5.** The proportional distribution of ASE genes and SNVs being expressed across
1063 tissues averaged for each sheep. A) Count of genes in 6 sheep expressed in grey,
1064 informative/filtered in blue and ASE positive in red. B) Break down of genes and SNVs
1065 that are expressed, pass minimum filtration criteria and are ASE positive.
1066

1067
1068
1069 **Figure S6**. Scatter plot of allelic depth (read count) in all 4 tissues of the 6 TxBF sheep.
1070 The SNVs passing minimum filtration criteria are coloured in grey and significant ASE
1071 (FDR10%) are marked by red and blue. If reference read count > alternate read count
1072 the point is shown in blue and vice versa in red. Expression level varies across tissues but
1073 does not affect the ASE profile.

Figure S7. Distribution of ASE positive SNVs over the direction of imbalance in the 34 tissues of 6 sheep. Ref.dp > Alt.dp as ASE towards Ref allele and vice versa. There was no significant difference between the count of ASE events given the direction of the allelic imbalance (p-value > 0.05).

Figure S8. The Kmeans cluster analysis of the ASE genes in between static and ICD profiles of BMDMs at 0 and 7hrs samples. Heat map dataframe has been log and centre

1084 scaled prior to cluster analysis. The heatmap shows the ASE genes in 5 groups: 0vs 7 hrs
1085 ICD set, Males 0hrs , Males 7hrs . Females 0hrs and Females 7hrs.
1086
1087 **A**

1105   **B**

1106

1107   **C**

1108
Figure S9. Principle component analysis and Kmeans clustering of ASE gene profiles of 4
tissues and 1 cell type (2 treatment) in 6 sheep. The gene list comprised of all ASE genes
in each tissue or cell type was used to perform PCA which established the presence of a
hierarchical stratification resulting in 3 groups in the dataset. This grouping was used to
cluster the dataset using K means algorithm with 3 centres. **A)** Samples PCA plot of the
first 2 PCs (Matrix 6 sample rows x 3000 gene columns) **B)** Genes PCA plot of the first 2
PC (Matrix 3000 gene rows x 6 sample columns) **C)** Kmeans Clusters overlay on the PC1

vs PC2 scatter plot from section B (Genes PCA plot). Gene name labels were included only
for the genes with overall contribution to the first 2 components > 1.
1118
1119
1120

1121
Figure S10. The read loss proportion during the WASP reference mapping bias removal steps.
The graph shows the input (read counts on the right) and output (read counts on the 2nd left)
RNA-Seq reads from the study before and after WASP processing respectively. Each bar
represents 4 overlaid read proportions labelled by colour and raw read count. The total portion
of reads coloured in grey shows 100% of the reads processed by WASP (input 170 million
reads on average). The green portion shows the non-intersecting reads (and their duplicates)
mapped within the range of Ensemble gene coordinate boundaries (exonic, intronic and UTRs

reads 49 million reads on average). The red portion of the reads represents reads intersecting
bi-allelic heterozygote sites and selected for synthetic read production step (34 million reads
on average). The blue portion of the graph represents the final output of WASP's pipeline
(reference biased and duplicated reads removed from red and green portions respectively). The
output portion (37 million reads on average) was used in GATK ASEReadCounter for allelic
read counts. The last row on the bottom shows the average of these proportions across both
tissue and cell type samples. Reference biased reads count for average 20% of input RNA-Seq
data.

1137
1138
1139 **Figure S11.** Number of SNVs before and after WASP reference bias correction. There is a
1140 significant reduction in number of variants captured (25% on average 1.5e+07) between two
1141 steps. Tissues samples dots mainly present within the upper quantile and BMDMs in the lower.
1142 Average number of variants beforehand = 6,325,570 and afterwards = 4,729,639.

1149
1150
1151

1152
Figure S12. The comparison of reference allele count ratio before and after WASP mapping
bias correction .The distribution of refRatio (refCount/refCount+altcounts) in all the tissue and
cell samples before (in red) and after WASP (in green). The overall counts , left skew (*viz.*
range 0.75-1) and refRatio=1 inflation is significantly reduced after WASP bias correction in
all samples.
Table S1. Mean allelic imbalance (log2(ASE$_{+LPS}$/ASE$_{-LPS}$)) in all 6 sheep. Only 1 gene
(VCAN including 7 SNVs) exhibited inducible ASE which was shared between 3 sheep (in
bold), while two (GCNT1 and SYNE3) showed inducible ASE in two sheep. Diverse and
highly individual ASE response to LPS was observed in each sheep.
1162
1163

Gene	Female1	Female2	Female3	Male1	Male2	Male3
ACOD1	-	-	-	6.8	-	-
ASH2L	-	-	-	-	3.9	-

Gene							
CASP8	-	-	-	-	-	-	9.8
CCDC80	-	-	-	9.1	-	-	-
CLDN1	-	-	-	-	-	-	10.3
CPOX	-	-	-	-	-	-	6.5
EDNRB	-	-	-	-	-	-	11.4
EMP1	-	-	-	-	-	-	23.9
ENSOARG00000001174	-	-	-	-	-	5.0	-
ENSOARG00000001910	-	-	-	-	-	-	7.8
ENSOARG00000002108	-	-	-	-	-	-	8.0
ENSOARG00000002898	-	-	-	-	-	-	5.3
ENSOARG00000004106	-	-	-	-	-	-	7.5
ENSOARG00000006889	-	-	-	31.9	-	-	-
ENSOARG00000007547	-	-	-	-	-	8.2	-
ENSOARG00000007603	-	-	-	-	6.9	-	-
ENSOARG00000014875	-	-	-	-	-	-	5.6
ENSOARG00000016081	6.4	-	-	-	-	-	-
ENSOARG00000018177	-	-	-	11.0	-	-	-
ENSOARG00000021562	-	-	-	-	-	-	9.0
FDX1	-	-	-	-	-	17.3	-
GCNT1	-	-	-	-	16.9	-	30.4
GSDME	-	-	-	-	4.6	-	-
HNMT	-	-	-	-	-	-	15.1
HUNK	-	-	-	-	-	-	4.0
IL1RN	-	-	-	-	-	-	9.7
KANSL1L	-	-	-	-	-	-	4.7
KMO	-	-	-	-	-	-	9.7
MB21D2	-	-	-	-	5.3	-	-
MYO10	-	-	-	-	-	-	7.8
NRP1	-	-	-	9.3	-	-	-
OGFRL1	-	-	-	-	-	-	16.5
PDE4D	-	-	-	-	-	-	11.7
PDK4	-	-	-	-	6.9	-	-
PGM2	-	-	-	-	-	-	9.5
PLK2	-	-	-	-	-	-	6.7
PPIF	6.9	-	-	-	-	-	-
PRNP	-	-	-	-	-	-	8.8
RASSF3	-	-	-	-	-	9.2	-
RGS1	-	-	-	-	-	-	8.4
SAFB2	-	-	-	-	-	-	5.3
SIGLEC15	-	-	-	-	4.8	-	-
SLC35F2	-	-	-	-	-	6.5	-
SYDE1	-	-	-	-	-	-	5.1
SYNE3	-	-	-	7.2	-	-	10.7
TBC1D1	-	-	-	-	6.3	-	-
TFEC	-	-	-	10.3	-	-	-
TRAF1	12.1	-	-	-	-	-	-

VCAN	23.5	-	-	-	14.0	13.9
VLDLR	-	-	-	-	4.7	-

1164
1165