Elimination of reference mapping bias reveals robust immune related allele-specific expression in cross-bred sheep

Mazdak Salavati [1*], Stephen J. Bush [1], Sergio Palma-Vera [2], Mary E.B. McCulloch [1], David A. Hume [3] and Emily L. Clark [1*]

1. The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush, Midlothian, UK
2. Leibniz Institute for Farm Animal Biology (FBN), Dummerstorf, Germany
3. Mater Research Institute-University of Queensland, Translational Research Institute, Woolloongabba, Qld 4102, Australia

* Correspondence:
Dr Emily L. Clark
Emily.Clark@roslin.ed.ac.uk
Dr Mazdak Salavati
Mazdak.Salavati@roslin.ed.ac.uk

Keywords: Allele-specific expression, mapping bias, RNA-Seq, sheep, transcriptome, WASP, GeneiASE
Running title: Unbiased ovine allele-specific expression
Editorial choice: British English
Article type: Original Research
Manuscript length: 6361 words

Abstract

Pervasive allelic variation at both gene and single nucleotide level (SNV) between individuals is commonly associated with complex traits in humans and animals. Allele-specific expression (ASE) analysis, using RNA-Seq, can provide a detailed annotation of allelic imbalance and infer the existence of cis-acting transcriptional regulation. However, variant detection in RNA-Seq data is compromised by biased mapping of reads to the reference DNA sequence. In this manuscript we describe an unbiased standardised computational pipeline for allele-specific expression analysis using RNA-Seq data, which we have adapted and developed using tools available under open licence. The analysis pipeline we present is designed to minimise reference bias while providing accurate profiling of allele-specific expression across tissues and cell types. Using this methodology, we were able to profile pervasive allelic imbalance across tissues and cell types, at both the gene and SNV level, in Texel x Scottish Blackface sheep, using the sheep gene expression atlas dataset. ASE profiles were pervasive in each sheep and across all tissue types investigated. However, ASE profiles shared across tissues were limited and instead they tended to be highly tissue-specific. These tissue-specific ASE profiles may underlie the expression of economically important traits and could be utilized as weighted SNVs, for example, to improve the accuracy of genomic selection in breeding programmes for sheep. An additional benefit of the pipeline is that it does not require parental genotypes and can therefore be applied to other RNA-Seq datasets for livestock, including those available on the Functional Annotation of Animal Genomes (FAANG) data portal. This study is the first global characterisation of moderate to extreme ASE in tissues and cell types from sheep. We have applied a robust methodology for ASE profiling, to provide both a novel analysis of the multi-dimensional sheep gene expression atlas dataset, and a foundation for identifying the regulatory and expressed elements of the genome that are driving complex traits in livestock.

Introduction

Allele-specific expression (ASE) is the imbalance of allelic expression between parental (diploid) copies at the same locus (Barlow and Bartolomei, 2014). It is most commonly associated with cis-acting regulatory variation that may mediate parent-of-origin, sex- or tissue-specific transcription of one allele relative to the other (Renfree et al., 2009; Hasin-Brumshtein et al., 2014). In a single individual, where there are informative sequence variants (i.e. heterozygote loci) that distinguish the products of two alleles, ASE can be detected by RNA-sequencing (Chamberlain et al., 2015; GTEx Consortium et al., 2017; Cao et al., 2019; Guillocheau et al., 2019). The ratio of allelic read counts obtained from RNA-Seq datasets can be used as a reliable proxy for ASE (i.e. $ASE ratio = CountsAllele1/(CountsAllele1 + CountsAllele2)$) (Edsgärd et al., 2016).

Large and complex RNA-Seq datasets give rise to unique and interesting computational challenges, in particular the elimination of reference mapping bias in ASE analysis of diploid genomes. RNA-Seq data are commonly mapped against reference genomes which are typically 'flat', with each position represented only by the reference (most abundant) allele. As such, reads containing heterozygous loci are more likely to be erroneously mapped (Degner et al., 2009; Stevenson et al., 2013; Hodgkinson et al., 2016). This can lead to high false positive ASE locus discovery rates (Degner et al., 2009). Although development of de novo transcript assemblers (Zerbino and Birney, 2008), variant-aware aligners (Xin et al., 2013; Hach et al., 2014) and mapping-free quantification e.g. Kallisto ((Bray et al., 2016)) has removed some of the issues, reference allele mapping bias remains an unresolved challenge in ASE studies. The analysis pipeline we present in the manuscript is designed to minimise reference bias while providing accurate profiling of allele-specific expression in large and complex RNA-Seq datasets.

We have developed an ASE analysis pipeline using the combination of software available under open licence, WASP (reference mapping bias removal) (van de Geijn et al., 2015), GATK (ASEReadCounter) (McKenna et al., 2010; Van der Auwera et al., 2013) and GeneiASE (Liptak-Stouffer aggregative ASE gene model) (Edsgärd et al., 2016). The GeneiASE model is capable of testing ASE at the gene level using two approaches: i) Static ASE which measures allelic imbalance within a gene (i.e. when ASE variants are located within the boundaries of the gene) and ii) Individual condition dependent ASE (ICD) which measures inducible ASE in a gene under an environmental pressure between 2 timepoints (i.e. in stimulated or unstimulated immune cells).

In addition to ASE at the gene level, we can also measure significant ASE at the single nucleotide level (SNV). ASE has been shown to be enriched within expression quantitative trait loci (eQTL) regions (Montgomery et al., 2010), therefore identifying ASE variants can be useful for understanding the transcriptomic control of complex traits in livestock. Complex trait mapping of ASE loci has been associated with phenotypes such as resistance to Marek's disease in chicken (Meydan et al., 2011) and pigmentation patterns in sheep (García-Gámez et al., 2011).

Understanding ASE is also important because cross-breeding now underlies most livestock production systems. Knowledge of ASE may provide insights into the molecular basis of the complex phenomenon of hybrid vigour, as emphasised by recent studies on two Chinese goat breeds and their F1 hybrids (Cao et al., 2019) and in F1 crosses of two highly inbred chicken lines (Zhuo et al., 2017). In this study we measure ASE in cross-bred sheep. Sheep are an economically important livestock species in many countries across the globe and particularly in emerging economies. The identification of prevalent ASE in populations or breeds, especially in economically relevant phenotypes and tissues, could be used to improve genomic prediction in sheep breeding programmes, such as those that have been established in

Australia and New Zealand (Daetwyler et al., 2010).
Using the methodology we describe, for mapping bias correction and robust positive
ASE discovery, we were able to profile pervasive allelic imbalance across tissues and cell
types, at both the gene and SNV level, in Texel x Scottish Blackface sheep. We analysed a
subset of total RNA-Seq libraries from liver, spleen, ileum, thymus and bone marrow-
derived macrophages (BMDM) (+/-) lipopolysaccharide (LPS), from six individual adult
cross-bred sheep to produce a detailed picture of allelic imbalance in immune-related tissues
and cell types. We chose to focus this analysis on immune related tissues in part because of the
depth of available sequence in those tissues, and in part because they contain abundant immune
cell populations. The diversity of cell populations is reflected in the transcriptional complexity
of immune tissues and cell types in the sheep gene expression atlas dataset (Clark et al., 2017;
Bush et al., 2018). As such this subset of tissues gave us a transcriptionally rich dataset in
which to measure ASE. We also included BMDMs stimulated and unstimulated with LPS to
mimic infection with Gram-negative bacteria, in order to test whether ASE changed in response
to stimulation with LPS in these cells. By measuring ASE in these tissues and cell types from
sheep we were able to: i) provide insight into how pervasive ASE is across tissues at the gene
and SNV level, ii) generate tissue-specific ASE profiles, iii) investigate sex-specific patterns
of ASE, and iv) determine the extent to which ASE changes in response to stimulation with
LPS in an immune cell type. This novel analysis of the multi-dimensional sheep gene
expression atlas dataset provides a foundation for further analysis of the regulatory and
expressed elements of the genome that are driving complex traits in sheep.
121
122 **Methods**
123
124 **Sample preparation and RNA extraction**
125
Data from three male and three female Texel x Scottish Blackface (TxBF) sheep from
the sheep gene expression atlas project (Clark et al., 2017) were used in this study. The dataset
including: 2 cell types (BMDMs (+/-) LPS) and 4 tissues (thymus, spleen, liver and ileum).
Tissue collection, storage and RNA extraction is described in (Clark et al., 2017). BMDM's
were cultured in vitro for 7 days in the presence of macrophage colony-stimulating factor
(CSF1 (10^4 U/ml) and unstimulated (0 hrs - LPS) and stimulated (7hrs +100ng/ml LPS)
samples of BMDMs were obtained also as described in (Clark et al., 2017). A total of 2 samples
(1 thymus and 1 spleen) did not pass the RNA quality control (RNA integrity number (Mueller
et al., 2004); RIN <7) and were not included in the sheep gene expression atlas. Library
preparation was performed by Edinburgh Genomics (Edinburgh Genomics, Edinburgh, UK).
All total RNA Illumina TruSeq libraries (125bp paired end) were sequenced at a depth of >
100 million reads per sample.
138
139 **Reference mapping bias removal**
140
BAM files from RNA-Seq data, were previously mapped to Oar v3.1 top level DNA
fasta track, using HISAT2 as described in (Clark et al., 2017). These BAM files were used to
locate reads with heterozygote loci using WASP's find_intersecting.py script (van de Geijn et
al., 2015). The intersection of reads and heterozygote loci in all samples were based on the
Ensembl   v92   variant   call   format   (VCF)   track   (Ensembl   v92:
ovis_aries_incl_consequences.vcf.gz). Synthetic copies of reads containing heterozygote loci,
were created with the alternate allele flipped to the remaining options of A, T, C or G (up to 6
loci /read[2^n] max 64 combinations of synthetic reads) using parameters defined in WASP (van
de Geijn et al., 2015). This was followed by remapping of the synthetic reads using HISAT2

(Li and Durbin, 2009; Kim et al., 2015) and eliminating the original reads (and their synthetic
copies) which mapped to a different coordinate in any of its synthetic copies (WASP's
filter_reads.py) (van de Geijn et al., 2015). After merging the retained reads with non-
intersecting reads, a final BAM file was produced for ASE read counting step (WASP's
remove_dup.py).
155
156   **Allelic read counts and depth filtration**
157
Allele-specific read counting was carried out using the ASEReadCounter module of
GATK v3.8 with parameters -mmq 50 -mbq 25 (McKenna et al., 2010). Multiple pre-
processing steps were performed prior to GeneiASE input as instructed by (Edsgärd et al.,
2016) which included, preparing per chromosome indices, merging the variant set with
corresponding gene coordinates and bi-allelic expression filtering. Loci with < 10 reads
mapped were excluded, as were loci with < 3 reads, or < 1% of the total reads, mapped to both
the reference and alternative allele. This form of filtration will eliminate loci exhibiting mono-
allelic expression (MAE) as previously described (Degner et al., 2009; Stevenson et al., 2013;
Mayba et al., 2014). Similar filtration criteria have been previously used routinely in ASE
studies (Mayba et al., 2014; Chen et al., 2016a; Edsgärd et al., 2016; GTEx Consortium et al.,
2017; Raghupathy et al., 2018; Cao et al., 2019; Guillocheau et al., 2019; Gutierrez-Arcelus et
al., 2019).The workflow of the analysis pipeline for ASE analysis is detailed in Fig.1.
170
171   **Experimental design for defining allele-specific expression**
172
173   ASE was defined according to the following three categories:
174
175   i)   Static ASE: which is inherent allelic imbalance (AI) in each gene calculated by
176   ASE at all heterozygote loci (i.e. $ASE = Counts\,RefAllele/$
177            $(Counts\,RefAllele + Counts\,AltAllele)$ within the boundaries of the gene.
178   The effect size of ASE at gene level was produced by aggregation of the ASE
179   effect size at SNVs within the gene boundaries (the Liptak-Stouffer method,
180   applied by the GeneiASE aggregative model). A null distribution of ASE effect
181   size for genes in each transcriptome was produced by random sub-sampling (n
182   = 1e+5) from a pool of genes having min 2 and max 100 loci within their
183   boundaries. The ASE effect size of each gene (aggregated using Liptak-
184   Stouffer) was then tested against the null distribution of same SNV number, via
185   a modified bi-nomial test (2x1 table). Distribution of p values was examined for
186   uniformity prior to false discovery rate correction (Supplementary file1)
187   (Benjamini and Hochberg, 1995; Pounds and Cheng, 2006; Barton et al., 2013).
188   ii)   Individual condition-dependent ASE (ICD-ASE): in which the same ASE
189   effect size was calculated for each gene in the treated versus the untreated
190   timepoints of the same sample (i.e. BMDM +/- LPS). The log2ratio
191   ($ASE_{treated}/ASE_{untreated}$) was used in a beta-binomial test (2x2 table) similar to
192   the static mode. The details of this aggregative model have been previously
193   described in the GeneiASE publication (Edsgärd et al., 2016).
194   iii)   Condition dependent ASE at SNV level: in which a contingency table was
195   produced for read counts (ref and alt) for every SNV, present both in treated and
196   untreated conditions (BMDM +/- LPS) (2x2 table) and a Fisher's exact test
197   performed followed by p value multiple testing correction (Benjamini and
198   Hochberg, 1995). The p values from loci showing ASE and shared by the six
199   adult sheep (ID and coordinate) were unified, using the Stouffer method

200   (Dewey, 2016, 2018) and presented as false discovery rate (FDR) for each locus.
201
202   Static ASE was calculated in both tissues and BMDMs (each timepoint was considered
203   separately for BMDMs). Condition dependent ASE analysis was carried out only in BMDMs
204   ± LPS both at gene (ICD-ASE) and SNV (Fisher's exact) level to study LPS inducible ASE.
205
Statistical analysis and thresholds applied

207
The extraction, transformation and loading of the all datasets and subsequent statistical analysis
was carried out in R v $\geq$ 3.4 unless stated otherwise (R Core Team, 2017). System query
language join statements (Wickham and Francois, 2015) were used to compare lists of ASE
genes or SNVs between samples. Raw p values resulting from all three types of ASE analysis
were corrected for multiple testing via Benjamini-Hochberg false discovery rate calculations
(Benjamini and Hochberg, 1995). The passing threshold of significance in all analyses was
considered to be FDR < 0.1 (10%) except for the Fisher's exact test association study. Genes
showing ASE in multiple tissues were considered those for which 4 or more of the 6 sheep had
significant ASE.
217
Results

219
Estimation of heterozygous sites across all individuals

221
To determine the level of heterozygosity present in the RNA-Seq data we first assessed
the number of bi-allelic heterozygote sites per individual for each of the six sheep (range=
5,673,703-6,438,497) detailed in Fig.2. Individual variation was observed in the SNVs per gene
in each sheep (Fig.2A&C). However, there was no significant difference in the total number of
bi-allelic SNVs captured in the RNA-Seq data across all six individuals or between the male
and female sheep included in the study (Fig.2B). The bi-allelic SNVs captured in the RNA-Seq
data set were annotated using the Ensembl v.92 (Zerbino et al., 2018) reference VCF track. The
distribution of SNVs per gene in the Ensembl track is tail-inflated in comparison to the RNA-
Seq data Fig.2A. This issue could be due to erroneous assignment of SNVs in hypervariable
and repetitive regions, multi-allelic SNVs or simply that there are variants in the Ensembl track
that are not expressed (transcribed). The distribution of SNVs for each individual is shown in
Fig.2C.
234
Reference mapping bias elimination and quality control

236
We used the WASP ref bias removal script to successfully minimise ref allele mapping
bias in the RNA-Seq samples. The mapping bias was assessed by global distribution of the
allelic ratio i.e. $ref_{counts}/alt_{counts} + ref_{counts}$ in each RNA-Seq sample as shown in Fig.3. The ASE
discovery rate at the SNV level on average constituted 5.8% of the heterozygote loci that passed
the minimum filtration criteria in each individual (0.1% of the total expressed). This portion of
the transcriptomic variants belonged to an average of 103 genes in each tissue transcriptome
(approx.1%) or 300 in each individual (Supplementary file4 – S4 Fig.1 and Fig.2). As shown
in S4 Fig 2. expression level varies across tissues but this does not affect the distribution of
ASE genes.
246
Genes exhibiting tissue-specific and pervasive ASE signatures

248
We used the static mode of GeneiASE to investigate pervasive and tissue-specific ASE

profiles across all of the available samples. The number of genes showing significant static
ASE in immune-related tissues across the six sheep are summarised in Table 1. On average
approximately.0.5% of the genes in each tissue-specific transcriptome showed significant ASE
(approx. 1% of the filtered set of genes). Pervasive ASE genes in all 4 tissues for each of the 6
sheep was investigated by applying the minimum 67% shared rule (i.e. min 4 out of 6 sheep
shared significant ASE for the gene). Genes exhibiting significant ASE shared by all 4 tissues
were then merged and the ASE effect size averaged (Fig.4A). Six genes exhibited pervasive
ASE across tissues (i.e. they were shared across all 4 tissues). In the order of allelic imbalance
effect size they were NAA50 (N(alpha)-acetyltransferase 50, Histone acetyl transferase) with
highest ASE effect size in spleen, UBB (Ubiquitin B, protein degradation) in thymus, HBP1
(HMG-Box transcription repressor; cell cycle regulation) and ENSOARG00000016510 both
in spleen, C1orf105 (Chromosome 1 Open Reading Frame 105) in ileum and MTIF2
(mitochondrial translational initiation factor 2) in thymus.
Sets of genes with tissue-specific ASE profiles were also captured (Fig.4B-E). Thymus
had the highest number of tissue-specific ASE genes (n=15) followed by liver (n=12), spleen
(n=5) and ileum (n=4) (Fig.4B-E). Amongst the thymus gene set was CD244 which included
30 heterozygote loci with allelic imbalance, one of which was rs406633825. This missense
allele (Chr1:110308273 C>A; pVal123Phe MAF=0.7, SIFT score=0 deleterious) has
previously been reported in the Texel population characterised by the International Sheep
Genome Consortium (ISGC) (Kijas et al., 2012). The liver-specific ASE profile included genes
involved in amino acid metabolism, cytochrome oxidase pathways and fibrinogen: FGA
(fibrinogen alpha chain), ENSOARG00000003175 (taurochenodeoxycholic 6 alpha-
hydroxylase-like, cytochrome P450 family 4 member [CYP4]), ENSOARG00000001568
(novel gene, complement C4-A-like), CYP3A24 (cytochrome P450 family 3 member) and
CA3 (carbonic anhydrase 3). Allelic imbalance in spleen was present in CACYBP (calcyclin
binding protein), DAPK2 (death associated protein kinase [serine/threonine]) and a novel gene
GIMAP8-like (ENSOARG0000001131). The ASE in GIMAP8 has been previously reported
in cattle with a strong paternal parent-of-origin expression pattern (Chamberlain et al., 2015).
The GIMAP/IAN family of proteins, which are involved in survival, selection and homeostasis
of lymphocytes (Nitta and Takahama, 2007).
Two genes of functional interest showed evidence of strong tissue-specific ASE in the
spleen: SNAP23 and MYLK. SNAP23 is a key molecule in vesicle transport machinery of the
cell and has been reported to be expressed in sheep spleen. SNAP23 or Synaptosome
Associated Protein 23 is part of protein complex involved in Class 1 MHC mediated antigen
processing and presentation and in neutrophil degranulation (Fabregat et al., 2018). SNAP23
is also vital to lymphocyte development (both B and T) in vitro (Wong et al., 1997; Kaul et al.,
2015). The myosin light chain kinase (MYLK) expression in the splenic trabeculae's smooth
muscle has been demonstrated previously (Jiang et al., 2014; Clark et al., 2017). Overall 199
heterozygote bi-allelic loci were present within the MYLK gene. The variant rs400678033
(Chr1:186347056G>A; .pAla1014Val), a missense SNV in exon 17 of 33 exons in MYLK,
showed consistent allelic imbalance in all spleen samples.
In summary, analysis of ASE across immune related tissues revealed there were a small
number of genes that were shared across tissues. ASE signatures instead tended to be tissue-
specific, within the sub-set of tissues investigated in this study.
294
295 **Individual-specific ASE signatures**
296
In order to investigate whether ASE profiles were either shared across all six sheep or
private to individual sheep we used intersectionality (Fig.5). Each tissue was investigated
separately. A number of private (to each individual) ASE genes were detected for each tissue,

ranging from: 31-123 in ileum (Fig.5A), 24-80 in liver (Fig.5B), 21-83 in spleen (Fig.5C) and
31-66 in thymus (Fig.5D). Some of the shared sets of ASE genes in these tissues were specific
to either male or female sheep, these sex-specific ASE signatures are described in Fig.5. In
ileum no sex-specific set was observed (Fig5A). In contrast to ileum, the ASE profile for liver
included a single gene with female only membership, ENSOARG00000017409 (novel gene;
93% orthology with bovine dicarbonyl and L-xylulose reductase [DCXR]) Fig.5B. In spleen
all female sheep shared significant ASE in PMS1 (DNA mismatch repair system component
during post meiotic segregation), ANKRD10 (Ankyrin Repeat Domain 10) and
ENSOARG00000006103 which was not present in any of the spleen profiles of male sheep
(Fig.5C). In thymus there were 2 sex-specific sets: 16 genes showing ASE only in females and
5 genes only in males. The female-specific thymus gene set included: ARPP21 (cAMP
regulated phosphoprotein 21), CDKL3 (cyclin dependent kinase like 3), CEP19,
ENSOARG00000000710 (novel gene), ENSOARG00000001163 (novel gene),
ENSOARG00000000710, ENSOARG00000001163, ENSOARG00000008981 (novel gene;
T-lymphocyte surface antigen Ly-9-like), ENSOARG00000006215, ENA000000008981,
ENSOARG00000009129, ENSOARG00000011375 (blood vessel epicardial substance
[BVES]), ENSOARG00000015755, ENSOARG00000020354 (novel gene; 53% orthology
with bovine monoacylglycerol acyltransferase [MOGAT1]), ENSOARG00000025005,
ENSOARG00000026030 (novel gene), GPM6A (glycoprotein M6A), RAG1 (recombination
Activating 1), STX8 (syntaxin 8). The male-specific thymus set was comprised of
ENSOARG00000007267 (novel gene; T-cell surface glycoprotein CD1a-like),
ENSOARG00000016841 (novel gene; 98% orthology with bovine ATP synthase membrane
subunit G [ATP5MG]), ENSOARG00000007603, SNX25 (sorting nexin 25) and LDHA
(lactate dehydrogenase A) Fig.5D.
In summary very few ASE genes were shared across all sheep and the majority of ASE
profiles were private to each sheep. Sex-specific ASE signatures were also detected but due to
the small sample size (n=3) in both cases these should be interpreted with caution.
327
328 **ASE in stimulated and unstimulated BMDMs (0hrs vs 7hrs +LPS)**
329
We examined inducible ASE after 7 hrs of exposure to LPS in BMDMs using the ICD
mode of GeneiASE (Edsgärd et al., 2016). A comparison of LPS inducible ICD-ASE genes
and the genes with background static ASE at 0 and 7 hr time points, was also performed. We
first assessed whether differences between 0 and 7 hours could be observed using analysis of
static ASE. Individual-specific ASE profiles and a limited number of shared ASE genes were
also observed in BMDMs. The total number of genes with static ASE in the BMDMs is shown
in Table 2.
Shared static ASE across both time points and independent of LPS induction was only
observed in 5 genes. These genes have a macrophage associated function and include ITGB2
(Yee and Hamerman, 2013), SAA3 (ENSOARG00000009963) (Larson et al., 2003; Deguchi
et al., 2013), CD200R1 (ENSOARG00000019357) (Ocaña-Guzman et al., 2018), DCTN5
(ENSOARG00000017281) (Habermann et al., 2001) and MTIF2 (also seen in the tissue
analysis above) (Overman et al., 2003).
The ICD-ASE in BMDMs +/-LPS captured fewer ASE genes with significant LPS
inducible ASE between the 2 timepoints than the static analysis of ASE. Moreover, there were
large differences in the number of LPS inducible ASE in each individual sheep, indicating
significant individual-specific variation in response to LPS stimulation. BMDM cultured from
female 2 showed no LPS inducible response in comparison to male 3 which was a hyper
responder with significant inducible ICD-ASE in 28 genes (including 634 informative SNVs
total). A detailed breakdown of SNVs, aggregated within each gene, with significant ICD ASE

has been summarised in Fig.6.
In summary the ICD-ASE mode of GeneiASE's model was not capable of capturing a
complete picture of differential ASE in the BMDM experiment. Static ASE was present in both
time points, however, there were no inducible ASE genes that were shared across all 6 sheep.
The highly diverse ASE profile of BMDMs was very individual-specific, similar to patterns
observed in tissues (Supplementary file4. S4 Table 1). These individual-specific differences
could be due to individual variation in the innate immune response or experimental variation
introduced during primary cell culture or stimulation with LPS.
358
359 **Condition dependent ASE at the SNV level in BMDMs**
360
To further investigate allelic imbalance at the SNV level without the aggregative gene
model of GeneiASE, Fisher's exact test was used. The filtered read counts for bi-allelic SNVs
shared by all 6 sheep BMDMs were selected (n=646 sites). Allelic read counts of each SNV
were tested using Fisher's exact test between 0hrs and 7hrs (2x2 table). Overall the 6 sheep
shared 646 SNVs with identical allelic genotypes in both time-points of the BMDM RNA-Seq
dataset. These SNVs were tested for association with LPS treatment and only 4 SNVs showed
a strong association (FDR < 1.0e-8) and 12 SNVs had an FDR between 1.0e-02 and 1.0e-08
(Fig.7). The highest F-statistic was at rs430667535Chr17 Pos:50485358T>C, a synonymous
variant in ubiquitin C (UBC), a polyubiquitin precursor, and also an intronic variant T>C or
A>G in pro-apoptotic BRI3 binding protein (BRI3BP). This variant was shown to have a minor
allele frequency (MAF) of 0.25 in Texel sheep based on the ISCG annotation [ISGC – Ensembl
v.92](Kijas et al., 2012)). The next highest peak was observed on Chr21 under SNVs within
SAA3 gene boundaries (ENSOARG00000009963) at the following coordinates: Highest FDR
peak was observed at SNV rs412192652 (Pos:25826978A>G missense variant [Asn145Asp],
FDR=2.3e-15) surrounded by rs403064928 (Pos:25826884C>A missense variant [Asp113Glu]
in exon 4 of SAA3 FDR=3.6e-07), rs426609498 (Pos:25826845A>G synonymous, FDR=5.5e-
07) and rs405439099 (Pos:25826990G>C 3'UTR variant, MAF 0.6 in Texel sheep, FDR=4.1e-
05). This region contains a strong LD block previously reported by the ISGC COMPOSITE
population (Ensembl v.92) (Supplementary file3) e.g. rs412192652 and rs405439099 pairwise
D' statistics = 1. Two further peaks on Chr3 were observed for rs159926581
(Pos:214731375T>C synonymous, FDR = 3.3e-09) in ribosomal protein L3
(ENSOARG00000016495) and rs159822214 (Pos:112164732G>A 3'UTR variant,
FDR=9.5e-03) in oxysterol binding protein like 8 (OSBPL8). The last inducible ASE
associated signal was on Chr16 rs420037698 (Pos:6887423G >A, FDR= 6.9E-09) in
ENSOARG00000004700, a known synonymous variant in the Texel population (MAF = 0.55).
The SNVs and corresponding genes from Fisher's exact test are summarized in Table 3.
The LPS inducible ASE analysis, of SNVs, using Fisher's exact test revealed a different
picture not captured by the gene level analysis with the GeneiASE model that aggregates SNVs
within each gene (Fig.6 vs Fig7). The aggregative gene model did not capture any shared ASE
genes in the ICD-ASE mode. Although the 6 sheep shared 646 SNVs and showed highly
significant association with stimulation with the LPS (Fisher's exact method), the aggregation
of ASE effect size from SNV to gene level (ICD-ASE mode) only detected individual-specific
sets of ASE genes in each sheep. This contradicted the results from Fisher's exact test which
detected 4 highly significant LPS inducible shared regions (FDR < 0.01, 16 SNVs) on
chromosomes 3, 16, 17 and 21. For example, the allelic imbalance in the SAA3 genomic
coordinates on chromosome 21 was not detectable in the ICD-ASE model but it was captured
by the Fisher's exact test in all individuals (Fig.7B chromosome 21).
Fisher's exact analysis at SNV level revealed ASE in response to LPS in variants within
CLDN1, ANXA3, BRI3BP, SAA3 and MSR1 (Table3). The anti-inflammatory macrophage

marker CLDN1 (Van den Bossche et al., 2012) and acute-phase inflammation resolution
marker ANXA3 (Yamanegi et al., 2018) have been previously reported with distinct
macrophage functions. The macrophage scavenger receptor 1 (MSR1) has also been shown to
be involved in lipid uptake and migration ability of macrophages (Shi et al., 2019). Three
noncoding RNAs (RF00221 [snoRD43], RF00593[snoU83B] and RF01151[snoU82P] were
amongst genes corresponding to ASE inducible SNVs. These 3 snoRNAs all overlap with the
genomic coordinates of ribosomal protein L3 (ENSOARG00000016495 / RPL3). The
RF00377 [snoU6-53] was also among the ASE positive targets which overlaps the protein-
coding gene CDS2 (CDP-diacylglycerol synthase 2). Using bulk RNA-Seq makes it difficult
to differentiate the origin of the ASE signal to specific RNA populations.
In summary ASE profiles in BMDMs were highly individual-specific at both gene and
SNV level. Moreover, Fisher's exact SNV level analysis discovered shared ASE SNVs where
the aggregative gene model of ICD-ASE mode did not, indicating for condition dependant ASE
analysis Fisher's exact test is more accurate and robust at SNV level.
414
415 **Discussion**
416
This study is the first to investigate global allele-specific expression across tissues from
sheep using RNA-Seq data. We focused our analysis on immune-related tissues and cell types
from six adult cross-bred sheep (TxBF) from the sheep gene expression atlas. ASE profiles
were highly individual-specific in the six sheep analysed in this study. We were able to identify
tissue-specific sets of ASE genes, as well as LPS inducible sets in the BMDM experiment.
Tissue-specific signatures of ASE have been previously reported in similar studies in mouse
(Castel et al., 2015, 2016), goat (Cao et al., 2019) and cattle (Chamberlain et al., 2015).
Several steps were taken in the cattle study (Chamberlain et al., 2015) to mitigate the
ref allele bias, assign parental origin using whole genome sequences and include MAE variants.
The SNV filtration was based on the (Hayes and Daetwyler, 2019) 1000 bull genomes project
to confirm the heterozygote sites. In our pipeline the Ensembl VCF track was used for that
purpose. In the cattle study the 1 and 0 inflated ref allelic ratio was still present and the ASE
analysis was done via introduction of 0.9 allelic ratio cut-off. In our pipeline no cut-off is
introduced and because it is difficult to distinguish between sequencing error and MAE, a
minimum filtration criteria was applied to exclude MAE. Chamberlain et al. tested 5317 genes
(14,495 SNVs) in spleen and detected 382 ASE genes (with min > 1 SNV per gene, similar to
this study). Although direct comparison would not be appropriate (because we have excluded
MAE variants in our analysis) our analysis of sheep spleen revealed ASE in 86 genes (average
in 5 sheep) from 8272 filtered genes (average in 5 sheep). Similarly, in thymus the cattle study
showed 182 ASE genes from 986 informative genes (9781 SNVs) while from 7961 filtered
genes in sheep thymus 134 ASE genes were captured. The differences in the numbers of genes
exhibiting ASE between the two studies are likely to be a consequence of the filtration criteria
applied, the exclusion of MAE, and species-specific differences between sheep and cow.
Results from a more recent study in goat (Cao et al., 2019) more closely reflect our findings.
They apply similar filtration criteria to our workflow and discovered 144 ASE genes in liver in
comparison to 123 in our sheep liver sets (averaged across 6 sheep). Other recent studies
including those focusing on production relevant tissues, such as muscle (Guillocheau et al.,
2019), have also applied similar stringency in filtration criteria. The filtering criteria we have
used for this analysis is stringent and focused on detecting variants of moderate to extreme
effects. Further analysis of the dataset reducing these criteria might discover additional variants
exhibiting ASE across individuals and tissues, but it would also increase the potential risk of
false positive discovery.
For this analysis we have adapted an ASE analysis workflow with a primary focus on

mapping bias removal prior to allele-specific analysis of the transcriptome. The collection of
scripts for WASP, used for this analysis, or modified versions of them have been utilised by
others for mapping bias removal in reference guided genomic datasets e.g. RNA-Seq
(Mozaffari et al., 2018; Zhou et al., 2018), Chip-Seq (Pelikan et al., 2018) and for methylomic
and epigenetic analysis (Richard Albert et al., 2018).
The ASE analysis pipeline we have adapted for sheep for this study is also adaptable to
other species and tissue types with available RNA-Seq datasets. It could be applied, for
example, to profile allele-specific expression in the RNA-Seq datasets from livestock species
listed on the FAANG data portal (Andersson et al., 2015; Harrison et al., 2018). We used the
Ensembl VCF track to capture information at heterozygote loci however the individual VCF
file from each sheep could also be used in ASE analysis. The latter strategy might enable the
capture of rare variants not included in the publicly available VCF tracks but would also raise
the issue of normalisation/standardization between VCF call sets. The usage of either of these
methods will be limited to the number of loci shared by coordinate and bi-allelic genotype (i.e.
pervasive ASE discovery). Other studies have compared variants at the RNA and DNA level
from the same individual and then removed the DNA variants not present in the RNA-Seq data
from the analysis (Guillocheau et al., 2019). We believe that a strength of the pipeline we
present is that it does not require parental genotypes and can therefore be applied to other RNA-
Seq datasets for livestock where this information is not available.
In our analysis we have not considered either parent-of-origin or breed-of-origin-
specific effects in this analysis. For parent-of-origin or breed-of-origin assignment of these
ASE profiles DNA level genotypes from the parents of the six sheep from the gene expression
atlas (i.e. Texel sire and Scottish Blackface dam) would be required, and these are unfortunately
not available. In this study ASE expression profiles also might be affected by the direction of
the cross (i.e. Texel sire x Scottish Blackface dam). To fully characterise parent-of-origin or
breed-of-origin reciprocal cross experiments would be required. Reciprocal cross studies in
mouse (Huang et al., 2017), chicken (Zhuo et al., 2017) and cross-bred cattle (Chen et al.,
2016b) have shed light on the complexity of such pervasive ASE markers and parent-of-origin
effects. Though potentially very interesting these experiments are lengthy and costly to perform
in sheep. Particularly in this case because the reciprocal cross (Scottish Blackface dam x Texel
sire) is rarely used in the UK sheep industry and as a consequence has limited relevance to
production.
Our approach also excludes mono-allelic expression. The minimum filtration criteria
utilised in our workflow along with the reference mapping bias removal step ensures an
unbiased ASE discovery in the transcriptome by excluding the ambiguity surrounding MAE
variants. This form of analysis is based on the principle that absence of evidence (reads) for
either allele of a heterozygote site does not directly amount to evidence of their absence i.e.
MAE. The pattern of ASE (ratio of Alt/Ref+Alt) is dependent on the bi-allelic expression of
loci within the genomic coordinates of the gene or genomic element of interest. For an ASE
effect to be captured by the GeneiASE model, the following criteria must be met: i) Biallelic
expression of the locus; ii) Min depth criteria for each allele (min 3 reads , total 10 reads at that
site and > 1% of total reads containing that allele); iii) The allelic imbalance or departure from
bi-allelic balanced expression being inducible by an environmental trigger (i.e. LPS in ICD
ASE experiment with BMDM data). These stringent criteria secure robust transcriptome-wide
ASE discovery while maximising the usage of read counts from short read RNA-Seq datasets
without considering mono-allelic sites. MAE patterns are impossible to differentiate from
sequencing error or random nonsense mediated decay in bulk RNA-Seq, unless arbitrary cut-
offs are introduced such as ratio of allelic read counts > 0.9 (Chamberlain et al., 2015) or > 0.7
(Cao et al., 2019). We decided to exclude MAE from this study for the aforementioned reasons.
The tissues utilised for ASE analysis in this study (thymus, spleen, liver and ileum) are

highly influential on performance of the immune system. ASE profiles shared across tissues
and cell types were limited and instead they tended to be highly specific. We identified tissue-
specific ASE in several genes in Thymus, for example, that are involved in the T-cell mediated
immune response, including CD47 and CD244. These tissue-specific and cell-type-specific
ASE profiles may underlie the expression of economically important traits such as disease
resistance. Assessment of the connection between economically relevant phenotypes and
tissue-specific ASE profiles could be useful for the improvement of genomics enabled sheep
breeding programmes, particularly those using specialised sire and dam lines (Georges et al.,
2018). Loci exhibiting ASE have been associated with production traits including milk-fat
percentage (Hayes et al., 2010; Suárez-Vega et al., 2017), trypanotolerance in small ruminants
(Kadarmideen et al., 2011; Álvarez et al., 2016), mastitis in goat (Ilie et al., 2018), Johne's
disease in cattle (Mallikarjunappa et al., 2018) and Marek's disease in chicken (Maceachern et
al., 2011; Meydan et al., 2011; Cheng et al., 2015). Although there is no general consensus
currently on the correlation of allelic expression haplotypes and phenotypes under selection in
sheep, this form of ASE analysis could pave the way for functional validation at population
level (e.g. breed or haplotype-specific aseQTL studies in a larger population of sheep).
Examples of population level aseQTL, eQTL and sQTL (QTLs associated with RNA splicing)
already exist for cattle (Wang et al., 2018; Xiang et al., 2018). Knowledge of favourable ASE
in critical genes involved in traits of interest could be used as a performance indicator or
included as weighted SNVs in genomic prediction algorithms to enhance livestock breeding
programmes (Georges et al., 2018). Currently the UK sheep industry is on the cusp of applying
genomic prediction, but suitable genomics enabled breeding programmes for sheep already
exist in New Zealand and Australia (Daetwyler et al., 2010).
523
524 **Conclusions**
525
In this study we characterise extreme to moderate allele-specific expression, at the gene
and SNV level, in immune-related tissues and cells from six adult sheep (TxBF) from the sheep
gene expression atlas dataset. Reference mapping bias removal was an integral component of
the analysis pipeline applied in this study. The correction of reference bias prior to obtaining
the allelic read counts is a critical step toward true ASE discovery. The workflow developed
as part of this manuscript provides an RNA-Seq only dependent tool, without the need for
individual DNA sequences. We note that the stringent filtering process applied would remove
loci where the allelic imbalance was less extreme but might still be of biological significance.
This study is a novel analysis of an existing large-scale complex RNA-Seq dataset from
sheep. Using the pipeline we have adapted for this analysis, we were able to identify ASE
profiles that were pervasive in each sheep and specific to the tissues and cell types investigated.
These tissue and cell-type-specific ASE profiles may underlie the expression of economically
important traits and could be used to identify variants that could be weight in genomic
prediction algorithms for the improvement of sheep breeding programmes. In summary we
have adapted a robust methodology for ASE profiling, using the sheep gene expression atlas
dataset, and provided a foundation for identifying the regulatory and expressed elements of the
genome that are driving complex traits in livestock.
543
544 **Data Availability**
545
The RNA-Seq sequence data is available via the European Nucleotide Archive (ENA) under
PRJEB19199 (https://www.ebi.ac.uk/ena/data/view/PRJEB19199). The BAM files were
already produced as part of the sheep gene expression atlas following the publicly available
protocol (FAANG, 2018) and as described in (Clark et al. 2017). The BAM files have been

uploaded to ENA under accession numbers ERZ827944, ERZ827949, ERZ827951,
ERZ827955, ERZ827972, ERZ827988, ERZ827995, ERZ827997, ERZ828001, ERZ828016,
ERZ828019, ERZ828036, ERZ828044, ERZ828046, ERZ828050, ERZ828070, ERZ828073,
ERZ828160, ERZ828167, ERZ828168, ERZ828172, ERZ828188, ERZ828192, ERZ828209,
ERZ828215, ERZ828217, ERZ828221, ERZ828240, ERZ828244, ERZ828261, ERZ828268,
ERZ828270, ERZ828274, ERZ828293 and ERZ828297. The Oar v3.1 reference FASTA and
VCF file from Ensembl v92 were used throughout the pipeline. The ASE analysis pipeline
(https://msalavat@bitbucket.org/msalavat/asewrap_public.git) was wrapped using bash
scripting on Edinburgh Compute and Data Facility computing resource Eddie Mark 3
(Edinburgh, 2018).
560
561 **Author's contributions**
562
MS and ELC coordinated and designed the analysis component of the study with assistance
from SJB and SPV. MS and SPV designed, optimised and tested the ASE pipeline. DAH
acquired the funding for the sheep gene expression atlas project. MEBM, ELC and DAH
designed the LPS experiment and generated the data. MEBM performed the LPS stimulation
of bone marrow derived macrophages and RNA extraction. SJB performed all bioinformatic
analyses prior to analysis with the ASE pipeline. MS performed ASE analysis, visualisation of
the results and wrote the manuscript. ELC contributed to manuscript editing and drafting. All
authors read and approved the final manuscript.
571

572 **Acknowledgments**
573
The authors would like to thank University of Edinburgh Computing and Data Support staff
(viz. Dr Andy Law and Dr Steve Thorn) for their input towards optimisation of the pipeline
and efficient parallel computing. Rachel Young and Lucas Lefevre isolated the bone marrow
derived macrophages for the sheep gene expression atlas project and Iseabail Farquhar helped
to collect and archive the tissue samples. The authors would also like to thank Professor Kim
Summers and Professor Mick Watson for their comments, which helped to improve the
manuscript.
581
582 **Conflict of Interest**
583
584 The authors have no competing interest regarding the findings presented in this publication.
585
586 **Funding**
587
The work was supported by a Biotechnology and Biological Sciences Research Council
(BBSRC) (http://www.bbsrc.ac.uk) Grant BB/L001209/1 ('Functional Annotation of the Sheep
Genome'). Also, BBSRC Institute Strategic Program Grants: 'Farm Animal Genomics'
(BBS/E/D/20211550), 'Transcriptomes, Networks and Systems' (BBS/E/D/20211552) and
'Blue Prints for Healthy Animals' (BB/P013732/1). Edinburgh Genomics is partly supported
through core grants from BBSRC (BB/J004243/1), NERC (http://www.nerc.ac.uk)
(R8/H10/56) and the Medical Research Council (MRC) (https://www.mrc.ac.uk)
(MR/K001744/1). SJB was supported by the Roslin Foundation. ELC is supported by the
University of Edinburgh Chancellor's Fellowship programme. The funders had no role in the
study design, data collection and analysis, decision to publish, or preparation of the manuscript.

598
599

References

Álvarez, I., Pérez-Pardal, L., Traoré, A., Fernández, I., and Goyache, F. (2016). Lack of specific alleles for the bovine chemokine (C-X-C) receptor type 4 (CXCR4) gene in West African cattle questions its role as a candidate for trypanotolerance. *Infect. Genet. Evol.* 42, 30–3. doi:10.1016/j.meegid.2016.04.029.

Andersson, L., Archibald, A. L., Bottema, C. D., Brauning, R., Burgess, S. C., Burt, D. W., et al. (2015). Coordinated international action to accelerate genome-to-phenome with FAANG, the Functional Annotation of Animal Genomes project. *Genome Biol.* doi:10.1186/s13059-015-0622-4.

Barlow, D. P., and Bartolomei, M. S. (2014). Genomic imprinting in mammals. *Cold Spring Harb. Perspect. Biol.* 6. doi:10.1101/cshperspect.a018382.

Barton, S. J., Crozier, S. R., Lillycrop, K. A., Godfrey, K. M., and Inskip, H. M. (2013). Correction of unexpected distributions of P values from analysis of whole genome arrays by rectifying violation of statistical assumptions. doi:10.1186/1471-2164-14-161.

Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing on JSTOR. *Source J. R. Stat. Soc. Ser. B* 57, 289–300. Available at: https://www.jstor.org/stable/pdf/2346101.pdf?refreqid=excelsior%3A8080132077b9217 878a58e47abbcf2d8 [Accessed October 27, 2015].

Bray, N. L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq quantification. *Nat. Biotechnol.* 34, 525–527. doi:10.1038/nbt.3519.

Bush, S. J., McCulloch, M. E., Muriuki, C., Salavati, M., Davis, G. M., Farquhar, I. L., et al. (2018). Comprehensive transcriptional profiling of the gastrointestinal tract of ruminants from birth to adulthood reveals strong developmental stage specific gene expression. *bioRxiv*, 364752. doi:10.1101/364752.

Cao, Y., Xu, H., Li, R., Gao, S., Chen, N., Luo, J., et al. (2019). Genetic Basis of Phenotypic Differences Between Chinese Yunling Black Goats and Nubian Goats Revealed by Allele-Specific Expression in Their F1 Hybrids. *Front. Genet.* 10, 145. doi:10.3389/fgene.2019.00145.

Castel, S. E., Levy-Moonshine, A., Mohammadi, P., Banks, E., and Lappalainen, T. (2015). Tools and best practices for data processing in allelic expression analysis. *Genome Biol.* 16, 195. doi:10.1186/s13059-015-0762-6.

Castel, S. E., Mohammadi, P., Chung, W. K., Shen, Y., and Lappalainen, T. (2016). ARTICLE Rare variant phasing and haplotypic expression from RNA sequencing with phASER. *Nat. Commun.* 7. doi:10.1038/ncomms12817.

Chamberlain, A. J., Vander Jagt, C. J., Hayes, B. J., Khansefid, M., Marett, L. C., Millen, C. A., et al. (2015). Extensive variation between tissues in allele specific expression in an outbred mammal. *BMC Genomics* 16, 993. doi:10.1186/s12864-015-2174-0.

Chen, J., Rozowsky, J., Galeev, T. R., Harmanci, A., Kitchen, R., Bedford, J., et al. (2016a). ARTICLE A uniform survey of allele-specific binding and expression over 1000-Genomes-Project individuals. doi:10.1038/ncomms11101.

Chen, Z., Hagen, D. E., Wang, J., Elsik, C. G., Ji, T., Siqueira, L. G., et al. (2016b). Global assessment of imprinted gene expression in the bovine conceptus by next generation sequencing. *Epigenetics* 11, 501–16. doi:10.1080/15592294.2016.1184805.

Cheng, H. H., Perumbakkam, S., Pyrkosz, A. B., Dunn, J. R., Legarra, A., and Muir, W. M. (2015). Fine mapping of QTL and genomic prediction using allele-specific expression SNPs demonstrates that the complex trait of genetic resistance to Marek's disease is predominantly determined by transcriptional regulation. *BMC Genomics* 16, 816. doi:10.1186/s12864-015-2016-0.

Clark, E. L., Bush, S. J., McCulloch, M. E. B., Farquhar, I. L., Young, R., Lefevre, L., et al.
(2017). A high resolution atlas of gene expression in the domestic sheep (Ovis aries).
PLOS Genet. 13, e1006997. doi:10.1371/journal.pgen.1006997.
Daetwyler, H. D., Hickey, J. M., Henshall, J. M., Dominik, S., Gredler, B., Van Der Werf, J.
H. J., et al. (2010). Accuracy of estimated genomic breeding values for wool and meat
traits in a multi-breed sheep population. *Anim. Prod. Sci.* doi:10.1071/AN10096.
Degner, J. F., Marioni, J. C., Pai, A. A., Pickrell, J. K., Nkadori, E., Gilad, Y., et al. (2009).
Effect of read-mapping biases on detecting allele-specific expression from RNA-
sequencing data. *Bioinforma. Orig. Pap.* 25, 3207–321210.
doi:10.1093/bioinformatics/btp579.
Deguchi, A., Tomita, T., Omori, T., Komatsu, A., Ohto, U., Takahashi, S., et al. (2013).
Serum amyloid A3 binds MD-2 to activate p38 and NF-κB pathways in a MyD88-
dependent manner. *J. Immunol.* 191, 1856–64. doi:10.4049/jimmunol.1201996.
Dewey, M. (2016). Comparison of various methods for meta–analysis of p–values. Available
at: http://www.dewey.myzen.co.uk/meta/comparemet.pdf [Accessed May 8, 2018].
Dewey, M. (2018). {metap}: meta-analysis of significance values.
Edsgärd, D., Iglesias, M. J., Reilly, S.-J., Hamsten, A., Tornvall, P., Odeberg, J., et al. (2016).
GeneiASE: Detection of condition-dependent and static allele-specific expression from
RNA-seq data without haplotype information. *Sci. Rep.* 6, 21134.
doi:10.1038/srep21134.
Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P., et al.
(2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res.* 46, D649–D655.
doi:10.1093/nar/gkx1132.
García-Gámez, E., Reverter, A., Whan, V., McWilliam, S. M., Arranz, J. J., Kijas, J., et al.
(2011). Using Regulatory and Epistatic Networks to Extend the Findings of a Genome
Scan: Identifying the Gene Drivers of Pigmentation in Merino Sheep. *PLoS One* 6,
e21158. doi:10.1371/journal.pone.0021158.
Georges, M., Charlier, C., and Hayes, B. (2018). Harnessing genomic information for
livestock improvement. *Nat. Rev. Genet.*, 1. doi:10.1038/s41576-018-0082-2.
GTEx Consortium, Gte., Laboratory, Data Analysis &Coordinating Center (LDACC)—
Analysis Working Group, Statistical Methods groups—Analysis Working Group,
Enhancing GTEx (eGTEx) groups, NIH Common Fund, NIH/NCI, et al. (2017). Genetic
effects on gene expression across human tissues. *Nature* 550, 204–213.
doi:10.1038/nature24277.
Guillocheau, G. M., El Hou, A., Meersseman, C., Esquerré, D., Rebours, E., Letaief, R., et al.
(2019). Survey of allele specific expression in bovine muscle. *Sci. Rep.* 9, 4297.
doi:10.1038/s41598-019-40781-6.
Gutierrez-Arcelus, M., Baglaenko, Y., Arora, J., Hannes, S., Luo, Y., Amariuta, T., et al.
(2019). Allele-specific expression changes dynamically during T cell activation in HLA
and other autoimmune loci. *bioRxiv*, 599449. doi:10.1101/599449.
Habermann, A., Schroer, T. A., Griffiths, G., and Burkhardt, J. K. (2001).
Immunolocalization of cytoplasmic dynein and dynactin subunits in cultured
macrophages: enrichment on early endocytic organelles. *J. Cell Sci.* 114, 229–240.
Available at: http://www.ncbi.nlm.nih.gov/pubmed/11112706 [Accessed March 11,
2019].
Hach, F., Sarrafi, I., Hormozdiari, F., Alkan, C., Eichler, E. E., and Sahinalp, S. C. (2014).
mrsFAST-Ultra: a compact, SNP-aware mapper for high performance sequencing
applications. *Nucleic Acids Res.* 42, W494-500. doi:10.1093/nar/gku370.
Harrison, P. W., Fan, J., Richardson, D., Clarke, L., Zerbino, D., Cochrane, G., et al. (2018).
FAANG, establishing metadata standards, validation and best practices for the farmed

and companion animal community. *Anim. Genet.* 49, 520–526. doi:10.1111/age.12736.

Hasin-Brumshtein, Y., Hormozdiari, F., Martin, L., van Nas, A., Eskin, E., Lusis, A. J., et al.
(2014). Allele-specific expression and eQTL analysis in mouse adipose tissue. *BMC*
Genomics 15, 471. doi:10.1186/1471-2164-15-471.

Hayes, B. J., and Daetwyler, H. D. (2019). 1000 Bull Genomes Project to Map Simple and
Complex Genetic Traits in Cattle: Applications and Outcomes. *Annu. Rev. Anim. Biosci.*
7, 89–102. doi:10.1146/annurev-animal-020518-115024.

Hayes, B. J., Pryce, J., Chamberlain, A. J., Bowman, P. J., and Goddard, M. E. (2010).
Genetic architecture of complex traits and accuracy of genomic prediction: coat colour,
milk-fat percentage, and type in Holstein cattle as contrasting model traits. *PLoS Genet.*
6, e1001139. doi:10.1371/journal.pgen.1001139.

Hodgkinson, A., Grenier, J.-C., Gbeha, E., and Awadalla, P. (2016). A haplotype-based
normalization technique for the analysis and detection of allele specific expression.
BMC Bioinformatics. doi:10.1186/s12859-016-1238-8.

Huang, W. C., Ferris, E., Cheng, T., Hörndli, C. S., Gleason, K., Tamminga, C., et al. (2017).
Diverse Non-genetic, Allele-Specific Expression Effects Shape Genetic Architecture at
the Cellular Level in the Mammalian Brain. *Neuron* 93, 1094–1109.e7.
doi:10.1016/j.neuron.2017.01.033.

Ilie, D. E., Kusza, S., Sauer, M., and Gavojdian, D. (2018). Genetic characterization of
indigenous goat breeds in Romania and Hungary with a special focus on genetic
resistance to mastitis and gastrointestinal parasitism based on 40 SNPs. *PLoS One* 13,
e0197051. doi:10.1371/journal.pone.0197051.

Jiang, Y., Xie, M., Chen, W., Talbot, R., Maddox, J. F., Faraut, T., et al. (2014). The sheep
genome illuminates biology of the rumen and lipid metabolism. *Science* 344, 1168–
1173. doi:10.1126/science.1252806.

Kadarmideen, H. N., Watson-Haigh, N. S., and Andronicos, N. M. (2011). Systems biology
of ovine intestinal parasite resistance: disease gene modules and biomarkers. *Mol.*
BioSyst. 7, 235–246. doi:10.1039/C0MB00190B.

Kaul, S., Mittal, S. K., Feigenbaum, L., Kruhlak, M. J., and Roche, P. A. (2015). Expression
of the SNARE protein SNAP-23 is essential for cell survival. *PLoS One* 10, e0118311.
doi:10.1371/journal.pone.0118311.

Kijas, J. W., Lenstra, J. A., Hayes, B., Boitard, S., Porto Neto, L. R., San Cristobal, M., et al.
(2012). Genome-wide analysis of the world's sheep breeds reveals high levels of historic
mixture and strong recent selection. *PLoS Biol.* 10, e1001258.
doi:10.1371/journal.pbio.1001258.

Kim, D., Langmead, B., and Salzberg, S. L. (2015). HISAT: a fast spliced aligner with low
memory requirements. *Nat. Methods* 12, 357–360. doi:10.1038/nmeth.3317.

Larson, M. A., Wei, S. H., Weber, A. T. A., Weber, A. T. A., and McDonald, T. L. (2003).
Induction of human mammary-associated serum amyloid A3 expression by prolactin or
lipopolysaccharide. *Biochem. Biophys. Res. Commun.* 301, 1030–1037.
doi:10.1016/S0006-291X(03)00045-7.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* 25, 1754–60. doi:10.1093/bioinformatics/btp324.

Maceachern, S., Muir, W. M., Crosby, S., and Cheng, H. H. (2011). Genome-wide
identification of allele-specific expression (ASE) in response to Marek's disease virus
infection using next generation sequencing. *BMC Proc.* doi:10.1186/1753-6561-5-S4-
S14.

Mallikarjunappa, S., Sargolzaei, M., Brito, L. F., Meade, K. G., Karrow, N. A., and Pant, S.
D. (2018). Short communication: Uncovering quantitative trait loci associated with
resistance to Mycobacterium avium ssp. paratuberculosis infection in Holstein cattle

using a high-density single nucleotide polymorphism panel. *J. Dairy Sci.* 101, 7280–
7286. doi:10.3168/JDS.2018-14388.

Mayba, O., Gilbert, H. N., Liu, J., Haverty, P. M., Jhunjhunwala, S., Jiang, Z., et al. (2014).
MBASED: allele-specific expression detection in cancer tissues and cell lines. *Genome*
Biol. 15, 405. doi:10.1186/s13059-014-0405-3.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., et al.
(2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Res.* 20, 1297–303.
doi:10.1101/gr.107524.110.

Meydan, H., Yildiz, M. A., Dodgson, J. B., and Cheng, H. H. (2011). Allele-specific
expression analysis reveals CD79B has a cis-acting regulatory element that responds to
Marek's disease virus infection in chickens. *Poult. Sci.* doi:10.3382/ps.2010-01295.

Montgomery, S. B., Sammeth, M., Gutierrez-Arcelus, M., Lach, R. P., Ingle, C., Nisbett, J.,
et al. (2010). Transcriptome genetics using second generation sequencing in a Caucasian
population. *Nature* 464, 773–777. doi:10.1038/nature08903.

Mozaffari, S. V, Stein, M. M., Magnaye, K. M., Nicolae, D. L., and Ober, C. (2018). Parent
of origin gene expression in a founder population identifies two new candidate imprinted
genes at known imprinted regions. *PLoS One* 13, e0203906.
doi:10.1371/journal.pone.0203906.

Mueller, O., Lightfoot, S., and Schroeder, A. (2004). RNA Integrity Number (RIN) –
Standardization of RNA Quality Control Application. *Agil. Appl. Note, Publ. 5989-*
1165EN, 1–8. doi:10.1101/gr.189621.115.7.

Nitta, T., and Takahama, Y. (2007). The lymphocyte guard-IANs: regulation of lymphocyte
survival by IAN/GIMAP family proteins. *Trends Immunol.* 28, 58–65.
doi:10.1016/j.it.2006.12.002.

Ocaña-Guzman, R., Vázquez-Bolaños, L., and Sada-Ovalle, I. (2018). Receptors That Inhibit
Macrophage Activation: Mechanisms and Signals of Regulation and Tolerance. *J.*
Immunol. Res. 2018, 8695157. doi:10.1155/2018/8695157.

Overman, R. G., Enderle, P. J., Farrow, J. M., Wiley, J. E., and Farwell, M. A. (2003). The
human mitochondrial translation initiation factor 2 gene (MTIF2): Transcriptional
analysis and identification of a pseudogene. *Biochim. Biophys. Acta - Gene Struct. Expr.*
1628, 195–205. doi:10.1016/S0167-4781(03)00144-1.

Pelikan, R. C., Kelly, J. A., Fu, Y., Lareau, C. A., Tessneer, K. L., Wiley, G. B., et al. (2018).
Enhancer histone-QTLs are enriched on autoimmune risk haplotypes and influence gene
expression within chromatin networks. *Nat. Commun.* 9, 2905. doi:10.1038/s41467-018-
05328-9.

Pounds, S., and Cheng, C. (2006). Robust estimation of the false discovery rate. 22, 1979–
1987. doi:10.1093/bioinformatics/btl328.

R Core Team (2017). R: A language and environment for statistical computing. Available at:
https://www.r-project.org/.

Raghupathy, N., Choi, K., Vincent, M. J., Beane, G. L., Sheppard, K. S., Munger, S. C., et al.
(2018). Hierarchical analysis of RNA-seq reads improves the accuracy of allele-specific
expression. *Bioinformatics* 34, 2177–2184. doi:10.1093/bioinformatics/bty078.

Renfree, M. B., Hore, T. A., Shaw, G., Graves, J. A. M., Pask, A. J., Marshall Graves, J. A.,
et al. (2009). Evolution of Genomic Imprinting: Insights from Marsupials and
Monotremes. *Annu. Rev. Genomics Hum. Genet.* 10, 241–262. doi:10.1146/annurev-
genom-082908-150026.

Richard Albert, J., Koike, T., Younesy, H., Thompson, R., Bogutz, A. B., Karimi, M. M., et
al. (2018). Development and application of an integrated allele-specific pipeline for
methylomic and epigenomic analysis (MEA). *BMC Genomics* 19, 463.

doi:10.1186/s12864-018-4835-2.

Shi, Q., Zhuang, F., Liu, J.-T., Li, N., Chen, Y.-X., Su, X.-B., et al. (2019). Single-cell
analyses reveal functional classification of dendritic cells and their potential roles in
inflammatory disease. *FASEB J.* 33, 3784–3794. doi:10.1096/fj.201801489R.

Stevenson, K. R., Coolon, J. D., and Wittkopp, P. J. (2013). Sources of bias in measures of
allele-specific expression derived from RNA-seq data aligned to a single reference
genome. *BMC Genomics* 14, 1. doi:10.1186/1471-2164-14-536.

Suárez-Vega, A., Gutiérrez-Gil, B., Klopp, C., Tosser-Klopp, G., and Arranz, J. J. (2017).
Variant discovery in the sheep milk transcriptome using RNA sequencing. *BMC*
Genomics 18, 170. doi:10.1186/s12864-017-3581-1.

van de Geijn, B., McVicker, G., Gilad, Y., and Pritchard, J. K. (2015). WASP: allele-specific
software for robust molecular quantitative trait locus discovery. *Nat. Methods* 12, 1061–
1063. doi:10.1038/nmeth.3582.

Van den Bossche, J., Laoui, D., Morias, Y., Movahedi, K., Raes, G., De Baetselier, P., et al.
(2012). Claudin-1, Claudin-2 and Claudin-11 Genes Differentially Associate with
Distinct Types of Anti-inflammatory Macrophages In vitro and with Parasite- and
Tumour-elicited Macrophages In vivo. *Scand. J. Immunol.* 75, 588–598.
doi:10.1111/j.1365-3083.2012.02689.x.

Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., Del Angel, G., Levy-
Moonshine, A., et al. (2013). From FastQ data to high confidence variant calls: the
Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinforma.* 43, 11.10.1-
33. doi:10.1002/0471250953.bi1110s43.

Wang, M., Hancock, T. P., Chamberlain, A. J., Vander Jagt, C. J., Pryce, J. E., Cocks, B. G.,
et al. (2018). Putative bovine topological association domains and CTCF binding motifs
can reduce the search space for causative regulatory variants of complex traits. *BMC*
Genomics 19, 395. doi:10.1186/s12864-018-4800-0.

Wickham, H., and Francois, R. (2015). dplyr: A Grammar of Data Manipulation. *R Packag.*
version 0.4.2., 3.

Wong, P. P. C., Daneman, N., Volchuk, A., Lassam, N., Wilson, M. C., Klip, A., et al.
(1997). Tissue Distribution of SNAP-23 and Its Subcellular Localization in 3T3-L1
Cells. *Biochem. Biophys. Res. Commun.* 230, 64–68. doi:10.1006/BBRC.1996.5884.

Xiang, R., Hayes, B. J., Vander Jagt, C. J., MacLeod, I. M., Khansefid, M., Bowman, P. J., et
al. (2018). Genome variants associated with RNA splicing variation in bovine are
extensively shared between tissues Ruidong. *BMC Genomics* 19, 521.
doi:10.1186/s12864-018-4902-8.

Xin, H., Lee, D., Hormozdiari, F., Yedkar, S., Mutlu, O., and Alkan, C. (2013). Accelerating
read mapping with FastHASH. *BMC Genomics 2013 141* 14, S13. doi:10.1186/1471-
2164-14-s1-s13.

Yamanegi, K., Yamada, N., Nakasho, K., and Nishiura, H. (2018). Participation of delta
annexin A3 in the ribosomal protein S19 C-terminus-dependent inhibitory mechanism of
the neutrophil C5a receptor through delta lactoferrin. *Pathol. Int.* 68, 109–116.
doi:10.1111/pin.12626.

Yee, N. K., and Hamerman, J. A. (2013). β(2) integrins inhibit TLR responses by regulating
NF-κB pathway and p38 MAPK activation. *Eur. J. Immunol.* 43, 779–92.
doi:10.1002/eji.201242550.

Zerbino, D. R., Achuthan, P., Akanni, W., Amode, M. R., Barrell, D., Bhai, J., et al. (2018).
Ensembl 2018. *Nucleic Acids Res.* 46, D754–D761. doi:10.1093/nar/gkx1098.

Zerbino, D. R., and Birney, E. (2008). Velvet: Algorithms for de novo short read assembly
using de Bruijn graphs. *Genome Res.* doi:10.1101/gr.074492.107.

Zhou, Z.-Y., Hu, Y., Li, A., Li, Y.-J., Zhao, H., Wang, S.-Q., et al. (2018). Genome wide

analyses uncover allele-specific RNA editing in human and mouse. *Nucleic Acids Res.*
46, 8888–8897. doi:10.1093/nar/gky613.
Zhuo, Z., Lamont, S. J., and Abasht, B. (2017). RNA-Seq Analyses Identify Frequent Allele
Specific Expression and No Evidence of Genomic Imprinting in Specific Embryonic
Tissues of Chicken. *Sci. Rep.* 7, 11944. doi:10.1038/s41598-017-12179-9.

Tables

Table 1. Total number of genes with significant static ASE in proportion to genes containing
informative SNVs (filtered). Tissue break-down has been presented as count (%ASE/filtered).

Sheep	Thymus	Spleen	Liver	Ileum	Total filtered	Total expressed
Female 1	-	136 (1.31%)	153 (1.47%)	92 (0.88%)	10,379	21,150
Female 2	136 (1.28%)	70 (0.66%)	116 (1.09%)	70 (0.66%)	10,572	21,420
Female 3	151 (1.22%)	75 (0.60%)	105 (0.85%)	140 (1.13%)	12,326	21,689
Avg.	143 (1.25%)	94 (0.85%)	125 (1.13%)	101 (0.89%)	11,092	21,419
Male 1	95 (0.92%)	-	157 (1.52%)	86 (0.83%)	10,282	21,361
Male 2	125 (0.99%)	54 (0.43%)	86 (0.68%)	80 (0.63%)	12,514	21,822
Male 3	106 (1.01%)	71 (0.67%)	110 (1.04%)	82 (0.78%)	10,480	21,545
Avg.	109 (0.97%)	62 (0.55%)	118 (1.08%)	83 (0.74%)	11,092	21,576
Total avg.	126 (1.13%)	78 (0.70%)	121 (1.09%)	92 (0.82%)	11,092	21,497

862　**Table 2**. Total number of genes with significant static ASE in BMDMs +/- LPS.

Sheep	BMDM 0hrs -LPS	BMDM 7hrs +LPS
Female 1	237	252
Female 2	192	233
Female 3	227	263
Female avg.	219	249
Male 1	205	193
Male 2	300	260
Male 3	260	261
Male avg.	255	238

866　**Table 3**. The variant IDs of LPS inducible ASE SNVs (Fisher's exact) and their respective
867　genes. Data was obtained using Ensembl BioMart query builder. Highly significant SNVs are
868　highlighted in bold (FDR< 1.0e-08)

ID	Chr	Start	End	Gene ID	Gene Name
rs418350332	1	195297663	195297663	ENSOARG00000020472	CLDN1
rs159822214	3	112164732	112164732	ENSOARG00000014876	OSBPL8
rs159926581	3	214731375	214731375	ENSOARG00000016495	-

rs159926581	3	214731375	214731375	ENSOARG00000022372	RF00221
rs159926581	3	214731375	214731375	ENSOARG00000024737	RF00593
rs159926581	3	214731375	214731375	ENSOARG00000025124	RF01151
rs159926581	3	214731375	214731375	ENSOARG00000025150	RF00593
rs193634916	4	53430934	53430934	ENSOARG00000001754	MDFIC
rs418697910	4	53430968	53430968	ENSOARG00000001754	MDFIC
rs162298949	6	92979793	92979793	ENSOARG00000018710	ANXA3
rs162298949	6	92979793	92979793	ENSOARG00000018710	ANXA3
rs160665956	9	76759747	76759747	ENSOARG00000001261	SPAG1
rs424104956	13	46654781	46654781	ENSOARG00000017292	CDS2
rs424104956	13	46654781	46654781	ENSOARG00000017292	CDS2
rs424104956	13	46654781	46654781	ENSOARG00000022618	RF00377
rs420037698	16	6887423	6887423	ENSOARG00000004700	-
rs430667535	17	50485358	50485358	ENSOARG00000017707	BRI3BP
rs430667535	17	50485358	50485358	ENSOARG00000018177	UBC
rs403064928	21	25826884	25826884	ENSOARG00000009963	SAA3
rs405439099	21	25826990	25826990	ENSOARG00000009963	SAA3
rs412192652	21	25826978	25826978	ENSOARG00000009963	SAA3
rs161531178	26	19663270	19663270	ENSOARG00000009626	MSR1

Figure captions

Figure 1. A flowchart of the allele-specific expression analysis pipeline applied to the sheep gene expression atlas dataset and optimised for WASP and GeneiASE programmes. The remapping was carried out using HISAT2 (Kim et al., 2015) in combination with SAMtools (Li et al., 2009). The Genome Analysis Toolkit v 3.8 was used for the ASE read counting section.

Figure 2. Distribution of biallelic SNVs expressed per gene in each of the 6 TxBF sheep. The total number of SNVs was averaged across thymus, liver, ileum and spleen for every animal. **A)** Histogram of SNVs per gene counts in the reference track (Ensembl in grey) and 6 sheep in red (females) and blue (males) overlaid. **B)**. The overall numbers of genes and SNVs detected in each animal (averaged over 4 tissues). **C)** Individual histograms from section A with females in red and males in blue.

Figure 3. The histogram of a global reference allelic ratio at every locus in the tissues. The distribution of ref allelic ratio showed a balanced profile without any 0 or 1 inflation which is observed in the presence of reference mapping bias. The allelic ratio above 0.51 is shown in blue and below 0.49 in red while balanced bi-allelic expression (0.49-0.51) is coloured in grey. Ref.dp = read counts for reference allele. Alt.dp = read counts for alternate allele. The y axis is square root scaled. As discussed in the text SNP that display MAE are not present in any of the samples analysed, indicating there was no inflation in either 0 or 1 allelic ratio.

Figure 4. Genes exhibiting static ASE shared across tissues from all six sheep. **A)** genes shared by four tissues with significant (FDR < 0.1) static ASE. **B)** ASE genes private to Ileum. **C)** Private to liver. **D)** Private to spleen. **E)** Private to thymus.

Figure 5. Intersectionality analysis of genes expressing significant ASE across all six sheep.
In each tissue from left to right the set count of genes (dots connected by lines) illustrates the
number of sheep sharing the gene. The private sets of genes are located at the far right of each
graph (single dots with no line). The intersections are coloured in to illustrate the size of the
set of shared genes (red [common to all 6 sheep], green [shared by 5 or 4 sheep], yellow [only
in females] and purple [only in males]). Detailed lists of genes with ASE shared by at least 4
sheep are presented above each graph for **A)** Ileum, **B)** Liver, **C)** Spleen and **D)** Thymus. Two
sex-specific sets of genes are highlighted: 16 genes showing ASE only in females (in yellow)
and 5 genes only in males (in purple).
908
909 **Figure 6**. Intersection analysis of SNVs under genes with significant ICD-ASE in the BMDMs
910 $\pm$ LPS. From left to right the set number of genes (dots connected by lines) have been illustrated
911 in order according to the number of sheep sharing the SNV. The private sets of SNVs are
912 located at the far right of each graph (single dots with no line).
913
914 **Figure 7.** Scatter plot of the adjusted p values from Fisher's exact test (unified using Stouffer
915 unification) in BMDMs comparing expression from different alleles at 0 vs 7hrs at SNV level
916 (LPS inducible ASE). **A)** The graph shows 646 loci exhibiting LPS inducible allelic imbalance
917 shared across all 6 sheep. **B)** Four loci on chromosomes 3,16,17 and 21 with FDR < 1.0e-08.
918 FDR < 1.0e-02 red line (n=16 SNVs) and FDR < 1.0e-08 blue line (n = 4 SNVs).
919
920 **Figure1.**
921

922
923
924
925

926   **Figure2.**

950 **Figure3.**

972   **Figure4.**

1000   **Figure5.**

1001

1002   **Figure6.**
1003
1004

1005
1006
1007   **Figure 7.**
1008

Supplementary material

A bimodal p value distribution was observed in GeneiASE output as shown in S1 Fig.1:

S1 Figure 1. Distribution of p values resulted from the beta-binomial test (Static ASE example for spleen in female 1). The uniformity of the p value distribution showed a bimodal (high frequency close to 1 followed by an extensive lower tail close to 0). R machine characteristics for printing float numbers were limited to 6 significant numbers after the decimal point by default which resulted in a high number of absolute zero (0+e0.0) output-values.

S1 Figure 2. The figure shows the distribution of p values produced by GeneiASE after cut off introduction (top) and after removal of p value == 1 values (middle). The distribution of FDR calculated by GeneiASE and after correction are shown at the bottom (right and left respectively). To correct for multiple testing by recalculation of FDR and eliminating the non-uniformity of p value distribution around 1 the following steps were performed as shown in S1 Fig.2: i) An arbitrary cut off (1e-7) was placed over all p values smaller than the said limit including the absolute 0e+0.0; ii) The genes showing p values equal to 1 were removed from each data frame; iii) The false discovery portion and rate were recalculated in each sample. iv) Gene showing FDR < 0.1 selected for further analysis.

After correction p value **GeneiASE raw p value**

1044
1045 **S1 Figure3**. The QQ plot of observed vs expected p values in corrected (left) and GeneiASE
1046 raw (right) calculated outputs.
1047
1048 Supplementary_file2.zip
1049 The zip file includes all the significant ASE values for genes and SNVs in the study.
1057 Supplementary_file3.pdf
1058 A

B

S3 Figure1. The LD block (A) and LD Manhattan plot (B) of SAA3 transcript (ENSOART00000010843). The LD block produced as part of International Sheep Genome Consortium is publicly available on Ensembl using the COMPOSITE population (n = 98).

Supplementary_file4.pdf

A Gene and SNV counts in 6 sheep
i. Expressed in grey
ii. Passed minimum filter in blue
iii. ASE at FDR10% in red

B

animal	Feature	Expressed	Filtered	ASE
Female1	gene	21150	10379	297
Female1	snv	6151417	125210	5985
Female2	gene	21420	10572	310
Female2	snv	5837076	109205	7095
Female3	gene	21689	12326	417
Female3	snv	5961473	137852	10811
Male1	gene	21361	10282	279
Male1	snv	5862189	125651	6740
Male2	gene	21822	12514	271
Male2	snv	6438497	151451	7110
Male3	gene	21545	10480	279
Male3	snv	5673703	115660	6558

1067
1068 **S4 Figure 1 supp to Figure 2.** The proportional distribution of SNVs being expressed,
1069 passing minimum filtration criteria and showing significant ASE (FDR 10%). A) Count of
1070 genes in 6 sheep expressed in grey, informative/filtered in blue and ASE positive in red.
1071 B) Break down of genes and SNVs that are expressed, pass minimum filtration criteria
1072 and are ASE positive.
1073

1074
1075 **S4 Figure 2 supp to Figure 3**. Scatter plot of allelic depth (read count) in all 4 tissues of
1076 the 6 TxBF sheep. The SNVs passing minimum filtration criteria are coloured in grey and
1077 significant ASE (FDR10%) are marked by red and blue. If reference read count > alternate
1078 read count the point is shown in blue and vice versa in red. Expression level varies across
1079 tissues but does not affect the ASE profile.
1080
1081
1082

1083 **S4 Table 1 supp to Figure 6.** Mean allelic imbalance (log2(ASE$_{+LPS}$/ASE$_{-LPS}$)) in all 6 sheep.
1084 Only 1 gene (VCAN including 7 SNVs) exhibited inducible ASE which was shared between 3
1085 sheep (in bold), while two (GCNT1 and SYNE3) showed inducible ASE in two sheep. Diverse
1086 and highly individual ASE response to LPS was observed in each sheep.
1087
1088

Gene	Female1	Female2	Female3	Male1	Male2	Male3
ACOD1	-	-	-	6.8	-	-
ASH2L	-	-	-	-	3.9	-
CASP8	-	-	-	-	-	9.8
CCDC80	-	-	9.1	-	-	-
CLDN1	-	-	-	-	-	10.3
CPOX	-	-	-	-	-	6.5
EDNRB	-	-	-	-	-	11.4
EMP1	-	-	-	-	-	23.9
ENSOARG00000001174	-	-	-	-	5.0	-
ENSOARG00000001910	-	-	-	-	-	7.8
ENSOARG00000002108	-	-	-	-	-	8.0
ENSOARG00000002898	-	-	-	-	-	5.3
ENSOARG00000004106	-	-	-	-	-	7.5
ENSOARG00000006889	-	-	31.9	-	-	-
ENSOARG00000007547	-	-	-	-	8.2	-
ENSOARG00000007603	-	-	-	6.9	-	-
ENSOARG00000014875	-	-	-	-	-	5.6
ENSOARG00000016081	6.4	-	-	-	-	-
ENSOARG00000018177	-	-	11.0	-	-	-
ENSOARG00000021562	-	-	-	-	-	9.0
FDX1	-	-	-	-	17.3	-
GCNT1	-	-	-	16.9	-	30.4
GSDME	-	-	-	4.6	-	-
HNMT	-	-	-	-	-	15.1
HUNK	-	-	-	-	-	4.0
IL1RN	-	-	-	-	-	9.7
KANSL1L	-	-	-	-	-	4.7
KMO	-	-	-	-	-	9.7
MB21D2	-	-	-	5.3	-	-
MYO10	-	-	-	-	-	7.8
NRP1	-	-	9.3	-	-	-
OGFRL1	-	-	-	-	-	16.5
PDE4D	-	-	-	-	-	11.7
PDK4	-	-	-	6.9	-	-
PGM2	-	-	-	-	-	9.5
PLK2	-	-	-	-	-	6.7
PPIF	6.9	-	-	-	-	-
PRNP	-	-	-	-	-	8.8
RASSF3	-	-	-	-	9.2	-

RGS1	-	-	-	-	-		8.4
SAFB2	-	-	31 -	-	-		5.3
SIGLEC15	-	-	-	4.8	-	-	
SLC35F2	-	-	-	-		6.5	-
SYDE1	-	-	-	-	-		5.1
SYNE3	-	-	7.2	-	-		10.7
TBC1D1	-	-	-	6.3	-	-	
TFEC	-	-	10.3	-	-	-	
TRAF1	12.1	-	-	-	-	-	
VCAN	23.5	-	-	-		14.0	13.9
VLDLR	-	-	-	-		4.7	-

1089
1090   f