Geographic variation in the genetic basis of resistance to leaf rust in locally

adapted ecotypes of the biofuel crop switchgrass (*Panicum virgatum*)

Acer VanWallendael[1,2], Jason Bonnette[3], Thomas E. Juenger[3], Felix B. Fritschi[4], Philip A. Fay[5],

Robert B. Mitchell[6], John Lloyd-Reilley[7], Francis M. Rouquette Jr.[8], Gary Bergstrom[9], and

David Lowry[1,2]

Author Affiliations:

[1]Department of Plant Biology, Michigan State University, East Lansing, MI, USA

[2]Great Lakes Bioenergy Research Center, Michigan State University, East Lansing, MI, USA

[3]Department of Integrative Biology, University of Texas at Austin, Austin, TX, USA

[4]Division of Plant Sciences, University of Missouri, Columbia, MO, USA

[5]USDA-ARS Grassland, Soil and Water Research Laboratory, Temple, Texas, USA

[6]USDA-ARS Wheat, Sorghum, and Forage Research Unit, University of Nebraska, Lincoln, NE,

USA

[7]USDA-NRCS, Kika de la Garza Plant Materials Center, Kingsville, TX, USA

[8]Texas A&M AgriLife Research, Texas A&M AgriLife Research and Extension Center,

Overton, TX, USA

[9]School of Integrative Plant Science, Cornell University, Ithaca, NY, USA

Abstract

Pathogens play an important role in the evolution of plant populations, but genetic

mechanisms underlying disease resistance may differ greatly between geographic areas as well

as over time. Local adaptation is thought to be an important step in plant evolution, and may be

impacted by differential pathogen pressures in concert with abiotic factors. This study uses

locally adapted ecotypes of the native perennial switchgrass (*Panicum virgatum*) to examine the

temporal and spatial variation in the genetic architecture of resistance to fungal pathogens,

namely switchgrass leaf rust (*Puccinia novopanici*). To identify loci underlying variation in

pathogen resistance in switchgrass, we scored rust damage across an outcrossed mapping

population at eight locations across the central United States from southern Texas to Michigan.

We followed rust progression at these sites for three years and mapped quantitative trait loci

(QTLs) using function-valued transformations of rust progression curves. Overall, we mapped 51

QTLs that varied in presence and strength over the three-year period. Two large-effect QTLs

were consistently associated with variation in rust progression in multiple sites and years, and are

therefore potentially the result of the same underlying resistance genes. Interestingly, these two

large-effect QTLs were almost exclusively detected in northern sites. This pattern could be

caused by geographic difference in genetic architecture. The distribution of rust strains or

variation in climatic conditions across the field sites could result in genotype-by-environment

interactions in efficacy of rust resistance loci. Beyond reducing rust damage by 34%, the

beneficial alleles at the two loci also increased biomass by 44%, suggesting a direct benefit by

pleiotropy or indirect benefit through genetic linkage. Our results suggest an important role for

fungal pathogens in the local adaptation of switchgrass and illustrate an influential geographic

component of the genetic architecture of plant disease resistance.

Introduction

Understanding the factors that determine how well a particular population is adapted to

its environment is a major goal of evolutionary biology. Plant populations often exhibit local

adaptation, in which populations are more successful in their local environments than foreign

genotypes in that environment (Leimu & Fisher 2008; Kawecki & Ebert 2004). Traditional

studies of local adaptation have focused on the abiotic factors that contribute to differential

population success (e.g. Clausen et al. 1940). Recent advances in molecular biology have

allowed researchers to directly link abiotic stresses to genetic differences between populations,

showing variation in the genetic basis of local adaptation (McKay et al. 2003; Lowry & Willis

2010; Fournier-Level et al. 2011; Price et al. 2018; He et al. 2018). However, biotic interactions

have been less-studied in this context, but may be essential for explaining variation in local

adaptation (Macel et al. 2007; Grøndahl & Ehlers 2008; Crémieux et al. 2008). Pathogens can

impose strong disruptive selection on plant populations when they are constrained in the areas

they infect, and therefore shape local adaptation (Giraud et al. 2017; Mursinoff & Tack 2017).

Therefore, a full understanding of plant adaptation requires characterization of the genetic basis

of local adaptation to pathogens in conjunction with the abiotic environment.

Molecular coevolution between microbe and host has been documented throughout the

tree of life (Hooper & Gordon 2001; Gagneux et al. 2006; Alfano & Collmer 2004; Dodds et al.

2006) and is an important tool to understand large-scale evolution (Moran et al. 2008;

Woolhouse et al. 2002). The most basic genetic mechanism underlying this dynamic is a gene-

for-gene interaction, whereby infection success is determined by variation at a single locus in

both the pathogen and host (Agrios 1997; Kniskern & Rausher 2006). In the host, resistance (R)

genes usually code for proteins that recognize either the direct product of a single pathogen gene

or some downstream protein in its signaling pathway (Jones & Dangl 2006). Pathogen genes that

have a corresponding R-gene in their host are termed 'avirulence' genes because their presence

means that plant resistance will be effective. Plant R-gene products trigger a cascade of changes

in response to the avirulence gene that may result in the hypersensitive response and systemic

acquired resistance (Kniskern & Rausher 2006). The hypersensitive response (HR) is a local

upregulation of various protective mechanisms, including production of reactive oxygen species

(ROS) and free radicals that may kill a pathogen, but typically at the cost of cell death and

possible loss of fitness and growth (Tian et al. 2003; Jones & Dangl 2006). Systemic acquired

resistance (SAR) is more akin to an innate immune response in vertebrates, whereby pathogen

recognition molecules are increased in expression throughout the plant, improving resistance to

later infections (Jones & Dangl 2006). SAR also shows a fitness cost, but has been demonstrated

to maintain fitness under disease pressure (Traw et al. 2007).

The gene-for-gene coevolutionary model is appealingly simple and can sometimes

explain the pattern of race-specific resistance in many plant species (e.g. van Leur et al. 1989).

However, quantitative resistance may also evolve, whereby resistance is governed by allelic

composition at several to many loci (Geiger & Heun 1989; Agrios 1997; Young 1996). These

mechanisms are typically more complex, involving polygenic adaptations for morphological or

phenological changes, production of antimicrobial compounds, and modification of effector

targets (Niks et al. 2015). Studies have found several examples of both gene-for gene resistance

(Kniskern & Rausher 2006; Bourras et al. 2016), and quantitative resistance (Quesada et al.

2010), but little consensus on the factors that determine which will evolve in a particular system.

Rather, it appears that the expression of host resistance as well as parasite virulence are

environmentally-dependent traits, not stable phenotypes as has been traditionally assumed

(Penczykowski et al. 2016). Since resistance mechanisms are generally investigated under either

stable laboratory conditions or at a single field site, it is challenging to breed for durable

resistance, resistance that prevents disease in many locations and for more than a few years

(Mundt 2014). Addressing the environmental variation in the genetic architecture of plant

disease resistance requires an experiment replicated over both time and space.

The prairie grass switchgrass (*Panicum virgatum*) and its obligate fungal pathogen,

switchgrass leaf rust (*Puccinia novopanici*), are an ideal system to study how loci contributing to

pathogen resistance vary across space. *Panicum virgatum* L. is a long-lived, polyploid, C4,

perennial grass native to North America east of the Rocky Mountains from northern Mexico to

southern Canada (Gleason and Cronquist 1991). It is a common prairie and pasture grass and is

grown as both a forage crop and as a bioenergy feedstock (Casler 2012; Parrish et al. 2012), and

has become an important study system for ecological specialization. *P. virgatum* is split into two

locally adapted ecotypes, upland and lowland (Morris et al. 2011; Lowry et al. 2014; Milano et

al. 2016). The upland ecotype is more common in northern North America, and exhibits small

stature (up to 190 cm) and low pathogen resistance (Casler 2012; Uppalapati et al. 2013; Milano

et al. 2016; Lovell et al. 2016). In contrast, the more southerly lowland ecotype is large (up to

285 cm) and is more resistant to fungal pathogens (Casler 2012; Uppalapati et al. 2013; Milano

et al. 2016; Lovell et al. 2016). While the lowland ecotype produces more biomass, it also has

lower freezing tolerance (Lee et al. 2014; Peixoto & Sage 2016), possibly explaining the rarity of

lowland ecotypes in more northern climates.

Since *P. virgatum* ecotypes differ in their susceptibility to rust infection, this host-

pathogen system is useful for testing the role of local variation in the evolution of resistance.

Switchgrass is infected with at least five species of rust (*Puccinia spp.;* Demers et al. 2017), but

Puccinia novopanici is thought to be dominant in the central US (Gary Bergstrom, *pers. obs.*). *P.*

novopanici is a basidiomycete fungi that infects only living leaf tissue of *P. virgatum*. As such,

these fungi are thought to be extirpated from northern populations every winter. In closely-

related and well-studied wheat rust (*P. triticana*), wind-borne spores blow from warm refugia in

southern Texas across the Great Plains every summer in what is known as the "Puccinia

pathway" (Eversmeyer & Kramer 2000). While this has not been directly examined for

switchgrass rust, it is the dominant explanation for epidemiological patterns, and fits with our

observations. Rust infection is virtually inevitable in switchgrass stands in North America by the

late summer, though damage is less severe in varieties from the lowland ecotype.

Over the range of switchgrass in North America, plant-pathogen interactions occur in a

wide range of abiotic conditions. The environmental dependence of resistance can have

important consequences for disease prediction and breeding for durable resistance (McDonald &

Linde 2002; Michelmore et al. 2013). In *Nicotiana*, for instance, elevated temperature inhibits

plant defense responses, prompting a need for development of heat-resistant R-genes (Zhu et al.

2010). The reasons for altered pathogenicity in different climates can be due to failure of plant

defenses as a result of multiple causes, including stress (Zhu et al. 2010), promotion of beneficial

pathogen conditions (Doke 1983), and variation in pathogen ecology (Weller et al. 2002). While

there have been few studies of the environmental niche of switchgrass rusts, closely related

wheat rusts can offer some clues. In wheat stripe rust (*P. striiformis*), resistance genes are

temperature-dependent (Fu et al. 2009). Further, it is well-established that variation in humidity

can greatly impact the infection by foliar fungal pathogens (Magarey et al. 2005), and may

therefore play a role in plant resistance. Thus, over the geographic range of the switchgrass-rust

interactions, variation in the abiotic environment may play an important role in influencing both

plant defenses and pathogen virulence.

The primary goal of our study was to characterize the genetic architecture of switchgrass

resistance to rust pathogens to better understand the causes of pathogen-mediated local

adaptation. To overcome past limitations due to environmental and temporal variation in host-

pathogen relationships, we measured fungal resistance over three years at a continental scale. We

planted clonally replicated quantitative trait locus (QTL) mapping populations at eight sites

across more than 7000 km of latitude to map QTLs for rust resistance. We first tested whether

rust resistance is controlled by large-effect loci corresponding to specific pathogens in a gene-

for-gene model or by small-effect loci in a polygenic resistance model. The geographic patterns

of QTL presence and strength allowed us to fully summarize the spatial distribution of pathogen

resistance. Therefore we were able to test the degree to which resistance exhibits a genotype-by-

environment interaction, or is homogeneously expressed. In addition, we assessed the

quantitative impact of those QTLs on resistance and other morphological and fitness traits to

determine the evolutionary impacts of pathogen resistance. We hypothesized that resistance

alleles would have evolved in lowland ecotypes and would be correlated with morphological

differences between ecotypes, such as biomass and tiller count.

Methods

Development of mapping populations

To identify loci controlling variation in rust progression, we used a previously developed

a four-way phase-known (pseudo-testcross) population (for full cross details, see Milano et al.

2016). We clonally divided the outbred populations by manually splitting rhizomes at the

Brackenridge Field Laboratory in Austin, TX. In May-July of 2015, the F_0, F_1, and F_2 clones

were potted, moved by truck, and transplanted at eight sites (Figure 1) in Kingsville, TX; Austin,

TX; Temple, TX; Overton, TX; Columbia, MO; Manhattan, KS; Mead, NE; and Hickory

Corners, MI. We assigned plants randomly to a honeycomb design, with 1.56 m between each

plant. To reduce edge effects, we planted a border of lowland plants around the plot that were not

measured experimentally. We watered plants by hand in 2015 to facilitate establishment. To

develop a linkage map for QTL mapping we genotyped 431 second-generation genotypes by

whole genome resequencing (for full details, see Lowry et al. 2019).

Figure 1: Locations of experimental sites in central North America. KING: Kingsville, TX; PKLE: Austin, TX; TMPL: Temple, TX; OVTN: Overton, TX; CLMB: Columbia, MO; MNHT: Manhattan, KS; LINC: Mead, NE; KBSM: Hickory Corners, MI.

Phenotyping

At each site we scored the presence of leaf rust in 2016, 2017, and 2018. We used a method developed for rust on wheat (*Triticum aestivum*; McNeal et al. 1971; Roelfs et al. 1992), which translates well to switchgrass and has been used in previous studies (Uppalapati et al.

 2013). At each site, we scored rust on a 0-10 scale based on the total proportion of the canopy

188 covered in rust spores, a score which we have defined as 'rust damage' for this study (Figure 2).

190
Figure 2: A. Heavily infected single leaf. B. Lightly infected small plant.

In conjuction, we used a simple resistance definition as 1 – rust damage (Simms & Triplett

1994). Other fungal pathogens such as anthracnose and *Bipolaris* were present in plots, but were

generally less common than rust and were not reflected in our ratings.

The effort extended to phenotyping rust damage varied among sites and years due to

logistical challenges. However, for the most part, sampling began three weeks after green-up (the

point at which ~50% of plants had emerged from the soil) and continued weekly until damage

stopped increasing (Figure 3). Over three years, this resulted in over 149,000 rust ratings, which

were used for the QTL analyses. In addition, we measured other morphological and

physiological traits at all sites, including the number of tillers, weekly plant height, date of first

flowering, and end-of-season aboveground biomass (see Milano et al. 2016 and Lowry et al.

2019 *in review* for details of this phenotyping effort).

Figure 3: Rust progression curves in 2016. Black lines show smoothed mean rust values for sampled dates, black dotted lines show fitted logistic curves to sampled data. Green and brown

vertical lines show green-up date and date of first flowering, respectively. (2017 & 2018 in

Supp.)

Rust abundance and composition

Visual pathogen scoring regimes are well known to be subject to statistical artifacts

(Lesaffre et al. 2012) and our data were no exception. Pathogen scores followed a tail-inflated

("U"-shaped) distribution. Therefore, we used nonparametric Wilcoxon signed-rank and

Kruskal-Wallis tests using the functions *wilcox.test* and *kruskal.test* in the *stats* package of R to

test the differences in damage between lineages, years, and sites (R core team 2018). To test for

cytoplasmic effects on rust prevalence, we compared rust scores between F_2 individuals with

maternal cytoplasm from upland and lowland F_1s, also with the aforementioned nonparametric

tests.

Genomic architecture

We first mapped QTLs using traditional QTL mapping on pathogen ratings for each

individual time point. Raw pathogen ratings were processed in R (v3.4; R core team 2018) using

both packages *qtl* and *funqtl* (Broman et al. 2003; Kwak et al. 2016). To examine QTL effects

over time, we scanned for QTLs using Haley-Knott regression for each site by year combination,

with each time point as a separate trait using the functions *scanone* in *qtl* and *geteffects* in *funqtl*

(Broman et al. 2003; Kwak et al. 2016).

We additionally examined QTLs controlling the overall progression of rust by modeling

damage as a function-valued trait (Kwak et al. 2014; 2016). For each individual, we fit a curve to

rust damage ratings using the R package *funqtl* (Kwak et al. 2016). This method has the

advantage of decreasing bias introduced by differences among raters at different sites by using

the parameters of an damage curve as traits, rather than the absolute rating values. In addition, it

overcomes several statistical challenges of QTL mapping of time-valued traits by replacing trait

data with a smoothed approximation, then applying functional principal component analysis

(FPCA) as a dimension-reduction technique (Kwak et al. 2016). QTLs are then mapped for a

small number of principal components (PCs; Kwak et al. 2016). These PCs represent the shape

of the pathogen damage curve of rust, and therefore include both the timing and rate of infection

spread. Previous studies have used area under disease progress curve (AUDPC) measurements to

quantify resistance (Jeger & Viljanen-Rollinson 2001). AUDPC is robust and useful, but may

show bias when infection timing differs among sites (Jeger & Viljanen-Rollinson 2001). FPCA

generally shows the same results as AUDPC, but is less impacted by phenology differences. We

used the first four PCs to map QTLs, and combined their effects by taking the mean LOD

(logarithm of the odds) score, the SLOD score, at each genomic position (Kwak et al. 2016).

Then we conducted 1000 permutations to calculate a penalty for the SLOD score that reduces the

rate of inclusion of extra loci to 5% (Broman & Sen 2009). We examined geographic and

temporal variation by mapping QTLs separately for each site and year, but we also generated a

combined test that summarizes variation in this experiment. For this test, we summed SLOD

scores across multiple sites and years, and concatenated permutations to generate a critical

SLOD cutoff. We produced all plots using *funqtl* and *ggplot2* (Kwak et al. 2016; Wickham &

Wickham 2007).

Finally, we estimated the allele-specific effects of the significant QTLs we discovered in

our study. In the four-way cross design, second-generation offspring can express four possible

allele combinations, L1_L2, U2_L1, U1_L2, or U1_U2. These represent alleles from each of the

four parents (lowland AP13: L1; lowland WBC: L2; upland DAC: U1; and upland VS16: U2).

For each locus, we compared the pathogen scores for the individuals with the "resistant" QTL

alleles to those with the "susceptible" alleles. We additionally made this comparison for

morphological traits including biomass, flowering time, tiller count, and green-up date. We

tested for difference in means using nonparametric Wilcoxon signed-rank tests for pathogen

ratings, and a two-sample *t*-test for all morphological traits.

Results

Rust abundance and composition

Though infection timing varied, rust was present at all sites throughout the study period

(Figure 3). The largest genetic source of variation was between upland and lowland F0

(grandparental) plants, with upland plants experiencing 39.15% more rust damage (W = 1.15e7,

p < 0.0001). Rust damage differed between generations in the cross (χ^2 = 656.98, p < 0.0001),

with F_0 plants showing the least amount of rust, and F_1 plants the greatest (Figure S1). Rust

damage was negatively correlated with green-up date, biomass, flowering time, height, and tiller

269 count (Figure S2). Field sites differed significantly in mean rust damage across years ($\chi^2 =$

270 1.99e5, p < 0.0001), though this was confounded by differing sampling periods across sites.

271 Mean rust damage declined in almost every site each year (Figure S3), decreasing by 19.75% in

272 2017 and 30.74% in 2018. This change correlates with biomass increases of 85.64% in 2017 and

273 46.89% in 2018. Our cross design allowed us to test the phenotypic effect of maternal cytoplasm,

274 the difference between second-generation plants with an upland dam and those with a lowland

275 dam. We compared rust scores between second-generation individuals with maternal cytoplasm

276 from upland and lowland F_1s. There was a cytoplasmic effect (W = 1.72e9, P = 0.0059), but rust

277 scores were only 1.33% higher in plants with lowland cytoplasm (Figure S1).

279 *Genomic architecture*

280   Since we collected data on pathogens over several weeks at all sites, we were able to

281 examine how the genomic architecture of resistance changed over a single season at each site by

282 scanning for QTLs as if each time point were a distinct phenotype. We found extensive variation

283 between sites, but some patterns were shared across several sites. We identified QTLs for

284 resistance at multiple sites on chromosomes 9N and 3N, though they differed in effect direction

285 (Milano et al. 2016). That is, the allele from a lowland grandparent decreased rust damage on

286 chromosome 9N, but increased damage on chromosome 3N. These QTLs had significant effects

287 for ~40-50 days, with the 9N QTL becoming detectable about one week after the 3N QTL

288 (Figure 4 and Supp.). Overall patterns were similar across years, although the collection of fewer

289 time points in 2017 resulted in lower-resolution data.

Columbia, MO 2016

Figure 4: Time-series QTL effects for Columbia, MO in 2016 (remaining sites & years in
Supp.). Blue indicates that the lowland allele decreases rust, red indicates that the upland allele
decreases rust. Color intensity is proportional to LOD score, or the strength of the QTL.

We mapped 51 total QTLs using function-valued traits (Figure 5). Overall, we found the
highest number of QTLs at the most northen site (KBSM), though there was not a clear
geographic pattern in QTL number. Additionally, there was variation between years, with the
greatest number of QTLs in 2016 (24 QTLs), and fewer in 2017 and 2018 (15 and 12,
respectively). We found the same large-effect QTLs using function-valued traits to map overall
QTLs (Figure 5). Across several sites and all years, QTLs on chromosomes 3N and 9N had the
highest LOD scores (Figure 5), indicating close associations between loci and pathogen
resistance, and each explained ~14% of the variation in rust damage.

Figure 5: QTLs for all sites and years. SLOD shows the mean log-odds score for function-valued traits, or the strength of the QTL. Horizontal lines show significance thresholds for each test, peaks above the threshold are shown as solid lines, and peaks below as faded lines. Vertical gray bars highlight large-effect QTLs on chromosomes 3N and 9N. Asterisks show chromosomes that had at least one significant QTL and are colored by year. The combined SLOD plot shows SLOD scores summed over Northern and Southern sites. The bar charts show the phenotypic effect of the large-effect QTLs on rust infection and biomass.

Given their strength and consistency across sites and years, we considered these the most

important QTLs. These large-effect QTLs differed greatly between northern sites and southern

sites. LOD scores were significant at the large-effect sites in northern sites 17 times over the

three years, but only four times in southern sites.

Allelic effects

We calculated allele-specific effects for each QTL in our dataset to quantify the direction

and strength of each QTL. To understand the alleles underlying each QTL, we categorized them

into either "Ecotype-specific" or "Genotype-specific" based on the resistance of each

grandparental allele (Figure 6bc; Milano et al. 2016). For instance, if both upland alleles showed

higher rust damage, and both lowland alleles low rust damage, this QTL would be considered

ecotype-specific. If only one of the grandparental genotypes was resistant or susceptible, we

scored that as genotype-specific. We added an additional category, "Cross-specific," for effects

that were shown in particular upland/lowland combinations (Figure 6a). A majority of loci

showed genotype-specific effects (31 out of 51). Relatively fewer showed ecotype-specific

effects (16 out of 51) and cross-specific effects were rarest (4 out of 51). In addition, we counted

the number of resistance QTLs for each grandparental allele (Figure 6d). Many loci were in the

same direction as the parental divergence, with both lowland alleles causing a reduction in rust

prevalence (23 out of 51), but the same number showed the lowest rust with an upland/lowland

allele combination. A few (5 out of 51) showed opposite pattern, with the least rust with both

upland alleles (Figure 6e).

Figure 6: Pattern of allele-specific effects for significant QTLs. A-C: Examples of three possible patterns of allele-specific effects, cross-specific, genotype-specific, and ecotype-specific. D: Frequency of all QTLs showing the lowest rust score. E: Frequency of each possible pattern of allele-specific effects for all QTLs.

We then calculated the combined effects of the two large-effect QTLs we identified by examining only individuals containing either resistant or susceptible combinations of alleles at these loci. For instance, the allele from the upland VS16 grandparent was most resistant at the 9N locus, but the allele from the lowland WBC3 grandparent was most resistant at the 3N, so the individuals with both of these alleles had the "resistant" combination of alleles. We also chose the individuals with alleles that increased rust, that had the "susceptible" combination of alleles. By comparing these individuals, we estimated the phenotypic impacts of our large-effect QTLs.

The effects of QTLs were most apparent early in the season, when the "resistant" combination

resulted in 34.08% lower rust scores ($W = 56156$, $P = 0.0003$). Later in the summer, the

difference between resistant and susceptible decreased to 6.22%, but variation also decreased (W

$= 41666$, $P < 0.0001$). We repeated this comparison for several other morphological variables.

The resistance alleles increased biomass by 44.3% ($t = 5.44$, $P < 0.0001$, Figure 5), tiller count

by 7.59% ($t = 2.58$, $P = 0.010$), and height by 5.52% ($t = 6.12$, $P < 0.0001$), and contributed to a

2.07% later flowering time ($t = 2.04$, $P = 0.041$; Table 1).

Table 1: Allele-specific effects for the combination of the 3N and 9N QTLs for rust resistance.

"Resistant" represents the mean values for individuals with alleles that decrease rust,

"Susceptible" represents mean values for individuals with alleles that do not confer resistance.

Bold values are significant at α = 0.05.

	Susceptible	Resistant	Δ	Statistic	P
Rust score	**2.161**	**1.425**	**34.08%**	**W = 56156**	**0.0002**
Biomass (g)	**480**	**607**	**26.48%**	***t* = 5.444**	**<0.0001**
Flowering time (Julian Day)	**177**	**180.6**	**2.07%**	***t* = 2.041**	**0.0415**
Tiller Count	**130.9**	**140.9**	**7.59%**	***t* = 2.581**	**0.0100**
Height (cm)	**154.4**	**163**	**5.52%**	***t* = 6.116**	**<0.0001**

Discussion

Our results show that there is a marked difference between the genetic architecture of rust

resistance between populations planted in northern and southern regions. In the north, resistance

appears qualitative, and driven by two large-effect loci that explain ~14% of variation in rust

damage, and clearly influence plant morphological traits. These QTLs were largely stable across

all three years of the study, indicating that they confer durable resistance. In the south, resistance

is more quantitative, driven by many small-effect loci that vary year-to-year.

We found support for a gene-for-gene model in northern populations, but southern

populations were representative of polygenic resistance. This geographic difference leads us to

conclude that there is a strong genotype-by-environment interaction for rust resistance in

switchgrass. Our hypothesis that the majority of resistance QTLs evolved in the lowland

ecotypes was supported, but at many loci, an upland-lowland allele combination produced the

greatest rust resistance. We also found support for the hypothesis that resistance alleles are

correlated with morphological differences, indicating that the two large-effect loci we mapped in

the north may have contributed to ecotypic differentiation between upland and lowland *P.*

virgatum. Overall, our results suggest an important role for two large-effect loci in northern

populations, but primarily minor effect loci underlying variation in resistance at southern field

sites.

Genomic architecture

Locally adapted northern upland and southern lowland ecotypes of *P. virgatum* are

divergent for many traits (Casler et al. 2004, 2007; Lowry et al. 2014; Milano et al. 2016; Lowry

et al. 2018 *in review*), including cold tolerance, biomass, and leaf architecture (Casler 2012

p.30). Ecotypic differences in fungal pathogen resistance have been well documented for *P.*

virgatum and will play a key role in the success of switchgrass as a forage and biofuel crop

(Cornelius & Johnston 1941; Uppalapati et al. 2012; Sykes et al. 2016). Both of the large-effect

resistance QTLs colocalize with previously-identified large-effect QTLs for biomass, height, and

tiller number using the same mapping populations (Lowry et al. 2019 *in review*).

The most striking pattern in QTLs over time and space was between northern and

southern sites. In the north, the two large-effect QTLs are consistently present over time and

across four sites. This pattern would be expected for qualitative gene-for-gene resistance, in

which the two QTL peaks are caused by resistance genes on chromosomes 3N and 9N. However,

in the south, there are many more small-effect QTLs. This pattern is more indicative of

quantitative resistance (Corwin & Kliebenstein 2017). It is possible that the large-effect QTLs in

the northern sites obscured several small-effect QTLs that would have otherwise been above the

critical LOD score, a phenomenon known as the Beavis effect (Beavis 1998; Xu 2003). The

Beavis effect obscures small-effect QTLs and causes effect size overestimation of detected QTLs

(Xu 2003). However, this bias is greatly decreased in sample sizes near to 500, so we expect that

the impact on our study was minimal.

Typically, resistance to a particular pathogen is either quantitative or qualitative, but this

pattern is not generally thought of as being geographically dependent. Studies in wheat have

found that rust resistance can be geographically constrained because resistance is typically strain-

specific (Kolmer 2005). Since rust populations in the central US are dominated by a single

species, *P. novopanici* (Gary Bergstrom *pers. obs.*), resistance differences are inconsistent with

species differences. Rather, we expect that north-south resistance differences may either be

driven by differences in rust strain diversity or a GxE interaction on the resistance loci. Broadly,

we expect higher strain diversity in the south, since this pattern has been documented for wheat

rust in Asia (Ali et al. 2014) and anther smut in *Silene* (Bueker et al. 2016). Southern populations

may have responded to higher stain diversity with more resistance loci. Alternatively, resistance

may be environment-dependent, as a study found for temperature-dependent resistance to wheat

stripe rust (*Puccinia striiformis f. sp. tritici*; Fu et al. 2009). In wheat stripe rust and other

systems, immunity-related proteins often exhibit temperature-dependent activity (Franklin &

Wigge 2014), suggesting a potential mechanism for GxE in disease resistance. Further work in

this system should focus on quantifying the population genetics of *P. novopanici* to better

understand whether resistance is strain-specific.

Previous efforts to map QTLs for rust resistance in switchgrass yielded limited results.

Milano et al. (2016) mapped one QTL on chromosome 8K for a rust prevalence in a population

planted in Austin, Texas. We found small-effect QTLs on this chromosome at three field sites,

but these loci were not consistently detected across years. The differences in QTLs may be due

to temporal differences, but are more likely traceable to inconsistency in phenotyping by

different field technicians. The previous study scored only five tillers per plant and used

principal-component transformation of a 1-4 rating scale (Milano et al. 2016). Our study

improved upon this method as we evaluated rust damage on a whole-plant basis and mapped

QTLs to function-valued transformations of rust progression, allowing for less bias due to tiller

selection and evaluation of the full rust progression curve.

Allelic effects

For the two large-effect QTLs, one showed higher resistance with the upland allele, and

the other with the lowland allele (Figure 5). This is surprising given that the lowland ecotypes

typically display the highest rust resistance (Uppalaptati et al. 2013), so one might expect that all

resistance alleles would come from lowland plants. The presence of upland resistance alleles

may provide further evidence that pathogens differ between the north and south, since upland

plants are much more common in the north.

Previous studies of switchgrass found many fixed differences between ecotypes (Milano

et al. 2016), and considerable variability in effects of loci across geographic locations (Lowry et

al. 2019, *in press*). Our pathogen resistance results were similar, although a surprising number of

loci had genotype-specific effects. This may indicate divergent selection within ecotypes, and

that mechanisms for pathogen resistance differ between populations of the same ecotype, as

would be expected if pathogen strain differences are substantial. A similar pattern was found in

melon (*Cucumis melo*), in which resistance QTLs vary between cultivars due to differences in

Fusarium strains (Perchepied et al. 2005). Since the lowland grandparental genotypes were much

more rust-resistant than the upland, it was surprising that the most rust-resistant plants expressed

resulted from a combination of an upland and a lowland alleles across QTLs.

The two large-effect QTLs decreased rust substantially throughout the season, though the

effect decreased over time. Leaf rust may show differing mechanisms of seedling and adult plant

resistance, so this pattern is not unexpected (German & Kolmer 1992). Similarly, resistance

alleles were associated with higher biomass and other overall morphological traits. This result

runs contrary to the expectation of a growth-defense tradeoff, whereby resistant genotypes

should be slower-growing due to limited resources (Herms & Mattson 1992). However, there

may be many reasons why a negative correlation between growth and defense does not occur

(Kliebenstein 2016; Hahn & Maron 2016). In switchgrass, we find that resistance QTLs

colocalize with biomass and tiller count QTLs (Lowry et al. 2019 *in review*), suggesting either

close linkage or a pleiotropic effect. Importantly for the emerging perennial bioenergy industry,

the correlation between resistance and biomass indicates that breeding switchgrass will be able to

combine positive traits without major trade-offs in biomass across most environmental

conditions. However, both biomass and resistance may instead trade off with freezing tolerance,

which is an important survival trait for switchgrass in high latitudes (Lowry et al. 2019 *in*

review). The major barrier to high-yield lowland traits in northern climates is winter

temperatures, so identifying the molecular basis of upland freezing tolerance in switchgrass is an

important goal. If freezing tolerance is not linked to rust resistance, there is great potential for

improvement of switchgrass for biofuel, especially in northern marginal areas.

Conclusion

Our results show the temporal and geographic variation in the genetic architecture of rust resistance in locally adapted switchgrass, including two large-effect loci that explain both pathogen defense and morphological differences between ecotypes, but show a limited effectiveness in the south. This pattern raises important questions about the drivers of genetic architecture of pathogen resistance and underscores the importance of assaying pathogen resistance across both time and space to capture the inherent variability in the interplay of biotic and abiotic drivers of genetic change. These loci may allow for more effective breeding strategies for rust resistance, if there are not trade-offs with other traits, such as cold tolerance. The role of rust in differentiating these ecotypes illustrates the synergistic role pathogens play in the evolution of different ecotypes and ultimately contributing to genetic variation within species.

References

Agrios GN. 1997. Control of plant diseases. *Plant Pathol.* 5:295–357

Alfano JR, Collmer A. 2004. Type III secretion system effector proteins: double agents in bacterial disease and plant defense. *Annu. Rev. Phytopathol.* 42:385–414

Ali S, Gladieux P, Leconte M, Gautier A, Justesen AF, et al. 2014. Origin, migration routes and worldwide
population genetic structure of the wheat yellow rust pathogen *Puccinia striiformis f. sp. tritici*. *PLoS*
Pathog. 10(1):e1003903

Arnold AE, Herre EA. 2003. Canopy cover and leaf age affect colonization by tropical fungal endophytes:
ecological pattern and process in *Theobroma cacao* (Malvaceae). *Mycologia*. 95(3):388–98

Arnold AE, Lutzoni F. 2007. Diversity and host range of foliar fungal endophytes: are tropical leaves biodiversity
hotspots? *Ecology*. 88(3):541–49

Beavis WD. 1998. QTL analyses: power, precision, and accuracy. *Mol. dissection complex Trait.* 1998:145–62

Broman KW, Sen S. 2009. *A Guide to QTL Mapping with R/qtl*, Vol. 46. Springer

Broman KW, Wu H, Sen Ś, Churchill GA. 2003. R/qtl: QTL mapping in experimental crosses. *Bioinformatics*.
19(7):889–90

Bueker B, Eberlein C, Gladieux P, Schaefer A, Snirc A, et al. 2016. Distribution and population structure of the
anther smut *Microbotryum silenes-acaulis* parasitizing an arctic–alpine plant. *Mol. Ecol.* 25(3):811–24

Casler MD, Vogel KP, Taliaferro CM, Wynia RL. 2004. Latitudinal adaptation of switchgrass populations. *Crop*
Sci. 44(1):293–303

Casler MD. 2012. Switchgrass breeding, genetics, and genomics. In *Switchgrass*, pp. 29–53. Springer

Casler MD, Vogel KP, Taliaferro CM, Ehlke NJ, Berdahl JD, et al. 2007. Latitudinal and longitudinal adaptation of
switchgrass populations. *Crop Sci.* 47(6):2249–60

Clausen J, Keck DD, Hiesey WM. 1940. Experimental studies on the nature of species. I. Effect of varied
environments on western North American plants. *Exp. Stud. Nat. species. I. Eff. varied Environ. West.*
North Am. plants.

Cornelius DR, Johnston CO. 1941. Differences in plant type and reaction to rust among several collections of
Panicum virgatum L. *J. Am. Soc. Agron.*

Corwin JA, Kliebenstein DJ. 2017. Quantitative resistance: more than just perception of a pathogen. *Plant Cell*.
29(4):655–65

Crémieux L, Bischoff A, Šmilauerová M, Lawson CS, Mortimer SR, et al. 2008. Potential contribution of natural
enemies to patterns of local adaptation in plants. *New Phytol.* 180(2):524–33

Demers JE, Liu M, Hambleton S, Castlebury LA. 2017. Rust fungi on *Panicum*. *Mycologia*. 109(1):1–17

Dodds PN, Lawrence GJ, Catanzariti A-M, Teh T, Wang C-I, et al. 2006. Direct protein interaction underlies gene-
for-gene specificity and coevolution of the flax resistance genes and flax rust avirulence genes. *Proc. Natl.*
Acad. Sci. 103(23):8888–93

Doke N. 1983. Involvement of superoxide anion generation in the hypersensitive response of potato tuber tissues to
infection with an incompatible race of *Phytophthora infestans* and to the hyphal wall components. *Physiol.*
Plant Pathol. 23(3):345–57

Eversmeyer MG, Kramer CL. 2000. Epidemiology of wheat leaf and stem rust in the central great plains of the
USA. *Annu. Rev. Phytopathol.* 38(1):491–513

Fournier-Level A, Korte A, Cooper MD, Nordborg M, Schmitt J, Wilczek AM. 2011. A map of local adaptation in
Arabidopsis thaliana. *Science (80-.)*. 334(6052):86–89

Franklin KA, Wigge PA. 2014. *Temperature and plant development*. Wiley Online Library

Fu D, Uauy C, Distelfeld A, Blechl A, Epstein L, et al. 2009. A kinase-START gene confers temperature-dependent
resistance to wheat stripe rust. *Science (80-.)*. 323(5919):1357–60

Gagneux S, DeRiemer K, Van T, Kato-Maeda M, De Jong BC, et al. 2006. Variable host–pathogen compatibility in
Mycobacterium tuberculosis. *Proc. Natl. Acad. Sci.* 103(8):2869–73

German SE, Kolmer JA. 1992. Effect of gene Lr34 in the enhancement of resistance to leaf rust of wheat. *Theor.*
Appl. Genet. 84(1–2):97–105

Giraud T, Koskella B, Laine A. 2017. Introduction: microbial local adaptation: insights from natural populations,
genomics and experimental evolution. *Mol. Ecol.* 26(7):1703–10

Gleason HA, Cronquist A. 1991. *Manual of vascular plants of northeastern United States and adjacent Canada*,
Vol. 834. New York Botanical Garden Bronx, NY

Grøndahl E, Ehlers BK. 2008. Local adaptation to biotic factors: reciprocal transplants of four species associated
with aromatic *Thymus pulegioides* and *T. serpyllum*. *J. Ecol.* 96(5):981–92

He L, Wu W, Zinta G, Yang L, Wang D, et al. 2018. A naturally occurring epiallele associates with leaf senescence
and local climate adaptation in *Arabidopsis* accessions. *Nat. Commun.* 9(1):460

Herms DA, Mattson WJ. 1992. The dilemma of plants: to grow or defend. *Q. Rev. Biol.* 67(3):283–335

Hooper L V, Gordon JI. 2001. Commensal host-bacterial relationships in the gut. *Science (80-.)*. 292(5519):1115–
18

Jeger MJ, Viljanen-Rollinson SLH. 2001. The use of the area under the disease-progress curve (AUDPC) to assess
quantitative disease resistance in crop cultivars. *Theor. Appl. Genet.* 102(1):32–40

Jones JDG, Dangl JL. 2006. The plant immune system. *Nature*. 444(7117):323

Kawecki TJ, Ebert D. 2004. Conceptual issues in local adaptation. *Ecol. Lett.* 7(12):1225–41

Kenaley SC, Quan M, Aime MC, Bergstrom GC. 2018. New insight into the species diversity and life cycles of rust
fungi (Pucciniales) affecting bioenergy switchgrass (*Panicum virgatum*) in the Eastern and Central United
States. *Mycol. Prog.* 17(11):1251–67

Kliebenstein DJ. 2016. False idolatry of the mythical growth versus immunity tradeoff in molecular systems plant
pathology. *Physiol. Mol. Plant Pathol.* 95:55–59

Kniskern JM, Rausher MD. 2006. Major-gene resistance to the rust pathogen *Coleosporium ipomoeae* is common in
natural populations of *Ipomoea purpurea*. *New Phytol.* 171(1):137–44

Kolmer JA. 2005. Tracking wheat rust on a continental scale. *Curr. Opin. Plant Biol.* 8(4):441–49

Kwak I-Y, Moore CR, Spalding EP, Broman KW. 2014. A simple regression-based method to map quantitative trait
loci underlying function-valued phenotypes. *Genetics*. 197(4):1409–16

Kwak I-Y, Moore CR, Spalding EP, Broman KW. 2016. Mapping quantitative trait loci underlying function-valued
traits using functional principal component analysis and multi-trait mapping. *G3 Genes, Genomes, Genet.*
6(1):79–86

Lee DK, Parrish AS, Voigt TB. 2014. Switchgrass and giant *Miscanthus* agronomy. In *Engineering and science of*
biomass feedstock production and provision, pp. 37–59. Springer

Leimu R, Fischer M. 2008. A meta-analysis of local adaptation in plants. *PLoS One*. 3(12):e4010

Lesaffre E, Lawson AB. 2012. *Bayesian biostatistics*. John Wiley & Sons

Lovell JT, Shakirov E V, Schwartz S, Lowry DB, Aspinwall MJ, et al. 2016. Promises and challenges of eco-
physiological genomics in the field: tests of drought responses in switchgrass. *Plant Physiol.* 172(2):734–
48

Lowry DB, Behrman KD, Grabowski P, Morris GP, Kiniry JR, Juenger TE. 2014. Adaptations between ecotypes
and along environmental gradients in *Panicum virgatum*. *Am. Nat.* 183(5):682–92

Lowry DB, Willis JH. 2010. A widespread chromosomal inversion polymorphism contributes to a major life-history
transition, local adaptation, and reproductive isolation. *PLoS Biol.* 8(9):e1000500

Macel M, Lawson CS, Mortimer SR, Šmilauerova M, Bischoff A, et al. 2007. Climate vs. soil factors in local
adaptation of two common plant species. *Ecology*. 88(2):424–33

Magarey RD, Sutton TB, Thayer CL. 2005. A simple generic infection model for foliar fungal plant pathogens.
Phytopathology. 95(1):92–100

McDonald BA, Linde C. 2002. Pathogen population genetics, evolutionary potential, and durable resistance. *Annu.*
Rev. Phytopathol. 40(1):349–79

Mckay JK, Richards JH, Mitchell-Olds T. 2003. Genetics of drought adaptation in *Arabidopsis thaliana*: I.
Pleiotropy contributes to genetic correlations among ecological traits. *Mol. Ecol.* 12(5):1137–51

McNeal FH, Konzak CF, Smith EP, Tate WS, Russell TS. 1971. *A uniform system for recording and processing*
cereal research data. USDA-ARS.

Michelmore RW, Christopoulou M, Caldwell KS. 2013. Impacts of resistance gene genetics, function, and evolution
on a durable future. *Annu. Rev. Phytopathol.* 51:291–319

Milano ER, Lowry DB, Juenger TE. 2016. The genetic basis of upland/lowland ecotype divergence in switchgrass
(Panicum virgatum). *G3 Genes, Genomes, Genet.* 6(11):3561–70

Moran NA, McCutcheon JP, Nakabachi A. 2008. Genomics and evolution of heritable bacterial symbionts. *Annu.*
Rev. Genet. 42:165–90

Morris GP, Grabowski PP, Borevitz JO. 2011. Genomic diversity in switchgrass (*Panicum virgatum*): from the
continental scale to a dune landscape. *Mol. Ecol.* 20(23):4938–52

Mundt CC. 2014. Durable resistance: a key to sustainable management of pathogens and pests. *Infect. Genet. Evol.*
27:446–55

Mursinoff S, Tack AJM. 2017. Spatial variation in soil biota mediates plant adaptation to a foliar pathogen. *New*
Phytol. 214(2):644–54

Parrish DJ, Casler MD, Monti A. 2012. The evolution of switchgrass as an energy crop. In *Switchgrass*, pp. 1–28.
Springer

Peixoto M de M, Sage RF. 2016. Improved experimental protocols to evaluate cold tolerance thresholds in
Miscanthus and switchgrass rhizomes. *Gcb Bioenergy*. 8(2):257–68

Penczykowski RM, Laine A, Koskella B. 2016. Understanding the ecology and evolution of host–parasite
interactions across scales. *Evol. Appl.* 9(1):37–52

Perchepied L, Dogimont C, Pitrat M. 2005. Strain-specific and recessive QTLs involved in the control of partial
resistance to *Fusarium oxysporum f. sp. melonis* race 1.2 in a recombinant inbred line population of melon.
Theor. Appl. Genet. 111(1):65–74

Price N, Moyers BT, Lopez L, Lasky JR, Monroe JG, et al. 2018. Combining population genomics and fitness QTLs
to identify the genetics of local adaptation in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci.* 115(19):5028–33

Price N, Moyers BT, Lopez L, Lasky JR, Monroe JG, et al. 2018. Combining population genomics and fitness QTLs
to identify the genetics of local adaptation in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci.* 115(19):5028–33

Roelfs AP. 1992. *Rust diseases of wheat: concepts and methods of disease management*. Cimmyt

Simms EL, Triplett J. 1994. Costs and Benefits of Plant Responses to Disease : Resistance and Tolerance. *Evolution
(N. Y)*. 48(6):1973–85

Sykes VR, Allen FL, Mielenz JR, Stewart CN, Windham MT, et al. 2016. Reduction of ethanol yield from
switchgrass infected with rust caused by *Puccinia emaculata*. *BioEnergy Res.* 9(1):239–47

Tian D, Traw MB, Chen JQ, Kreitman M, Bergelson J. 2003. Fitness costs of R-gene-mediated resistance in
Arabidopsis thaliana. *Nature*. 423(6935):74

Traw MB, Kniskern JM, Bergelson J. 2007. SAR increases fitness of *Arabidopsis thaliana* in the presence of natural
bacterial pathogens. *Evol. Int. J. Org. Evol.* 61(10):2444–49

Uppalapati SR, Serba DD, Ishiga Y, Szabo LJ, Mittal S, et al. 2013. Characterization of the rust fungus, *Puccinia
emaculata*, and evaluation of genetic variability for rust resistance in switchgrass populations. *Bioenergy
Res.* 6(2):458–68

Van Leur JAG, Ceccarelli S, Grando S. 1989. Diversity for disease resistance in barley landraces from Syria and
Jordan. *Plant Breed.* 103(4):324–35

Wadl PA, Dean D, Li Y, Vito LM, Scheffler BE, et al. 2011. Development and characterization of microsatellites
for switchgrass rust fungus (*Puccinia emaculata*). *Conserv. Genet. Resour.* 3(1):185–88

Weller DM, Raaijmakers JM, Gardener BBM, Thomashow LS. 2002. Microbial populations responsible for specific
soil suppressiveness to plant pathogens. *Annu. Rev. Phytopathol.* 40(1):309–48

Wickham H, Wickham MH. 2007. The ggplot package

Woolhouse MEJ, Webster JP, Domingo E, Charlesworth B, Levin BR. 2002. Biological and biomedical
implications of the co-evolution of pathogens and their hosts. *Nat. Genet.* 32(4):569

Xu S. 2003. Theoretical basis of the Beavis effect. *Genetics*. 165(4):2259–68

Zhang Y, Zalapa JE, Jakubowski AR, Price DL, Acharya A, et al. 2011. Post-glacial evolution of *Panicum
virgatum*: centers of diversity and gene pools revealed by SSR markers and cpDNA sequences. *Genetica*.
139(7):933

Zhu Y, Qian W, Hua J. 2010. Temperature modulates plant defense responses through NB-LRR proteins. *PLoS
Pathog.* 6(4):e1000844