Behavioral syndromes shape evolutionary trajectories via conserved genetic architecture.

Raphael Royauté [a,b]
Ann Hedrick [c,d]
Ned A. Dochtermann [a,e]

[a] Department of Biological Sciences; North Dakota State University
[b] raphael.royaute@gmail.com
[c] Department of Neurobiology, Physiology, and Behavior; University of California, Davis
[d] avhedrick@ucdavis.edu
[e] **corresponding author**: ned.dochtermann@gmail.com

Author to whom correspondence should be addressed:
Ned Dochtermann
1340 Bolley Drive, 201 Stevens Hall
Department of Biological Sciences
North Dakota State University
Fargo, ND 58102

Running Title: Evolutionary conservation of syndromes

Keywords: Behavioral syndromes, evolutionary constraint, **G** matrix, personality, behavioral ecology

Author Contributions: N.A.D. and A.H. conceived the project and supervised the gathering of data, R.R supervised and conducted behavioral trials and analyzed the data. N.A.D., A.H., and R.R. wrote the manuscript.

Abstract

Behaviors are often correlated within broader syndromes, creating the potential for

evolution in one behavior to drive evolutionary changes in other behaviors. Despite

demonstrations that behavioral syndromes are common across taxa, whether this potential

for evolutionary effects is realized has not yet been demonstrated. Here we show that

populations of field crickets (*Gryllus integer*) exhibit a genetically conserved behavioral

syndrome structure despite differences in average behaviors. We found that the

distribution of genetic variation and genetic covariance among behavioral traits was

consistent with genes and cellular mechanisms underpinning behavioral syndromes rather

than correlated selection. Moreover, divergence among populations' average behaviors was

constrained by the genetically conserved behavioral syndrome. Our results demonstrate

that a conserved genetic architecture linking behaviors has shaped the evolutionary

trajectories of populations in disparate environments—illustrating an important way by

which behavioral syndromes result in shared evolutionary fates.

Introduction

Behavior is frequently assumed to have been shaped by selection (Grafen 1984) and thus

populations are expected to differ in a range of behaviors based on local selective

pressures. This implies that behaviors are able to evolve independently, an assumption

increasingly challenged by the ubiquity of behavioral syndromes—correlations among

behaviors (Table 1; Sih et al. 2004a, Sih et al. 2004b)—which have been documented across

taxonomic groups (Garamszegi et al. 2012, 2013) and are comprised of both genetic and

environmental contributions (Dochtermann 2011, Dingemanse and Dochtermann 2014,

Dochtermann et al. 2015).

Given the contribution of genetic correlations to behavioral syndromes, these

syndromes have the potential to constrain the ability of populations to diverge and respond

to local selective pressures (Dochtermann and Dingemanse 2013). Specifically, based on

quantitative genetic theory, if syndromes stem from pleiotropic effects—wherein a single

gene affects multiple behaviors—populations will be constrained to diverge along shared

evolutionary pathways. Further, this divergence is predicted to occur in the direction in

trait space that contains the most variation (Figure 1; Schluter 1996). As a result, if

syndromes have a constraining effect on evolution, the pattern of correlations among traits

will be conserved among populations (Figure 1). Alternatively, if genetic correlations

underpinning syndromes are the result of selection historically favoring particular trait

combinations (i.e. selection-induced linkage disequilibrium (Roff 1997, Saltz et al. 2017)),

the divergence of populations will be generally unconstrained as these genetic correlations

are expected to rapidly break down when selection changes (Roff 1997, Conner 2002, Saltz

et al. 2017).

These two quantitative genetic explanations for behavioral syndromes have explicit

analogs in the behavioral literature: whether syndromes emerge from pleiotropy, tight

genetic linkage, or other shared physiological and cellular effects has been termed the

"constraints hypothesis," as opposed to selection-induced linkage disequilibrium, which

has been termed the "adaptive hypothesis" (Figure 1, *sensu* Bell (2005)). While some

studies have compared phenotypic or among-individual correlations across populations

(e.g. Dingemanse et al. 2007, Pruitt et al. 2010, Dochtermann et al. 2012, Royauté et al.

2014, Michelangeli et al. 2018), only population comparisons of behavioral syndromes at

the additive genetic level allow for properly testing these competing hypotheses.

Unfortunately, data at the additive genetic level has been restricted to a single comparison

of two populations (Bell 2005). Consequently, the overall role of behavioral syndromes in

shaping population divergence is an important gap in our knowledge as support for the

constraints versus adaptive hypotheses remains insufficiently tested.

Knowing whether behavioral syndromes emerge from genetic constraints or

selection induced linkage disequilibrium is important because these mechanisms

differentially affect evolutionary outcomes (Saltz et al. 2017). These potential effects are

broad, from altering responses to environmental changes to speciation dynamics. For

example, in *Anolis* lizards, constraints imposed by genetic correlations on morphological

traits have shaped divergence and what phenotypes can be expressed during adaptive

radiations (McGlothlin et al. 2018). Behavioral syndromes may have an even greater

constraining effect: Dochtermann and Dingemanse (2013) reported that the average

magnitude of genetic correlations between behaviors was sufficient to constrain

evolutionary responses to a greater degree than do correlations between life-history or

morphological traits. Unfortunately, this conclusion was based on data that could not

distinguish between the constraints and adaptive hypotheses. If indeed the constraints

hypothesis underpins behavioral syndromes then syndromes will reduce the rate of

adaptation and reduce the rate at which populations and species diverge. Evaluating

evidence for the constraints and adaptive hypotheses is therefore necessary in order to

understand whether behavioral syndromes are important in behavioral evolution.

Here we evaluated predictions of the adaptive and constraints hypotheses (Figure

1) and tested whether behavioral syndromes have diverged at the genetic level among

populations of the field cricket, *Gryllus integer*. Specifically, according to the constraints

hypothesis we predicted that genetic variation would be expressed in a consistent manner

among populations and genetic correlations would be maintained across generations.

Specific predictions are more difficult to make for the adaptive hypothesis without

knowledge of local selective pressures. This hypothesis can, however, be assessed

indirectly because behavioral divergence is expected to not be constrained and correlations

are expected to rapidly degrade. We evaluated these predictions via estimation and

comparisons of behavioral genetic (co)variance matrices, i.e. **G** (Table 1), estimated for

multiple populations of *G. integer* (Figure 2).

Methods

<u>Cricket collection</u>

G. integer is a particularly appropriate species to evaluate the constraints and adaptive

hypotheses for behavioral syndromes as the species exhibits population differences in

ecologically relevant behaviors that can be assayed in the laboratory (e.g. Hedrick and

Kortet 2006, Kortet and Hedrick 2007, Niemela et al. 2012a). *G. integer* can also readily be

bred in the lab according to quantitative genetic designs making it an ideal model for our

questions of interest (Hedrick 1988).

We collected adult female crickets from four populations throughout the

southwestern and western US: Socorro, NM; Las Cruces, NM; Aguila, AZ; and Dunnigan, CA

(Figure 2) during the summer of 2017. Crickets from these locations are formally

recognized as members of *Gryllus integer* but additional splitting is currently being

considered (D. Weissman, personal communication). These populations are also

geographically distant from each other (Figure 2) and vary in predator and parasitoid

abundance (Hedrick and Kortet 2006).

Around 50 females on average were collected from each population (Table S1) and

taken to animal housing facilities at North Dakota State University. Females were housed

individually in 0.71 L containers and provided with ad libitum food (Purina Chick Starter)

and water (water was provided in glass vials capped with cotton). Each cricket was also

provided with a small piece of cardboard egg carton for shelter. The cricket housing room

was maintained on a 12:12 dark:light cycle reversed such that the room was dark during

daytime hours. The housing room was kept at ~27C.

<u>Breeding design</u>

Females collected from the field (generation P) were allowed to oviposit in water vials

while in their containers. Offspring of these females were designated generation F_0 as sires

were unknown: mating occurred prior to capture and multiple mating is common in the

genus (Simmons 1986). F_0 offspring hatched in their dams' containers and were then

moved to individual housing prior to maturation. We assayed the behavior of 387 F_0

individuals (see below) upon maturation (Table S1). After behavioral trials, F_0 individuals

were assigned to breeding pairs such that individual males were mated to multiple

randomly assigned females from the same population but different dams according to a

standard full-sib, half-sib breeding design (Lynch and Walsh 1998). Matings were

conducted as follows: females were moved from their normal housing containers to a

larger container (34.6 × 21 × 12.4 cm) along with their food dish, water vial, and egg carton

shelter. After the female had been transferred, the assigned male was likewise moved to

the large container, also with its food dish, water vial, and egg carton. The male and female

remained in these containers for 24 hours to allow sufficient time for courtship and

multiple mating. After 24 hours the male and female crickets were returned to their

original containers. If males were to be mated with additional females, they were allowed a

minimum break of 24 hours before repeating the above procedure. These F_0 females were

subsequently allowed to oviposit into water vials within their containers. Resulting F_1

offspring were moved to individual housing prior to maturation and had their behaviors

assayed upon maturation. After behavioral assays, F_1 individuals were likewise paired with

F_1 individuals of the same population but different sires in the same manner as above and

resulting F_2 offspring were moved to individual housing and had their behavior measured

upon maturation. This resulted in the behavioral testing of 395 F_1 individuals and 163 F_2

individuals (Table S1). Across the three generations this represented behavioral testing of

946 individual crickets.

<u>Behavioral testing</u>

All behavioral tests followed standard procedures previously validated in the literature for

Gryllid crickets (Kortet and Hedrick 2007, Kortet et al. 2007, Hedrick and Kortet 2012,

Niemela et al. 2012b, Royauté et al. 2015, Royauté and Dochtermann 2017, Royauté et al.

2019). These assays encompass how individuals vary in risk-taking behavior (Kortet et al.

2007), exploratory propensity (Royauté et al. 2015, Royauté and Dochtermann 2017,

Royauté et al. 2019), and response to predation threat (Royauté and Dochtermann 2017,

Royauté et al. 2019). Based on the known relatedness among individuals we then

estimated **G**—the matrix of additive genetic behavioral variances and covariances—for

each population. Below, we describe these behavioral assays and their ecological relevance.

- Latency to emerge from shelter -

Gryllid crickets, including *G. integer*, use small burrows and natural cracks for refuge from

predators and to which they retreat when under threat. As a result, latency to emerge from

shelter after disturbance can be considered a proxy for risk-taking behavior or "boldness"

(Kortet et al. 2007). Here, we conducted latency tests wherein individuals were transferred

from their home containers to small artificial burrows (40 cm^3) placed within a 34.6 × 21

cm arena. These artificial burrows were capped so that individuals could not immediately

emerge. Crickets were forced to remain in the artificial burrow for two minutes after which

the cap was removed. Crickets were then allowed six minutes and thirty seconds to emerge

from the artificial burrow. During this test we recorded how long it took for an individual

to emerge (in seconds). Individuals that did not emerge were given a maximum latency of

390 seconds.

- Open field exploratory behavior -

Open field tests are a classic behavioral assay across taxa (Walsh and Cummins 1976)

which measure the exploratory propensity of individuals (Réale et al. 2007), including

crickets (Royauté et al. 2015, Royauté and Dochtermann 2017, Royauté et al. 2019).

Individuals that move through more of the arena are considered more thorough explorers

(Réale et al. 2007). Here we used open field tests to measure activity and exploratory

propensity in a 30 × 30 cm plexiglass arena. Individuals were introduced into the arena

under a small container and allowed to rest for 30 seconds after introduction. At the end of

this 30 seconds, the container was removed and the cricket was allowed to explore the

arena for 3 minutes and 40 seconds. The arena was cleaned with isopropyl alcohol between

trials to remove any chemosensory cues from the arena. We used Ethovision XT to record

the total distance the individual moved during the trial (cm), the number of unique zones of

the arena an individual visited during the trial, and the variance in velocity of individuals.

This latter measure indicates whether an individual's speed of exploration was constant

(low velocity variance) or whether individuals had frequent activity bursts punctuated by

long bouts of inactivity (high velocity variance).

- Response to cues of predator presence -

How individuals respond to cues of predator presence often varies within and among

populations and is likely to covary with fitness (Herman and Valone 2000). Crickets

respond to chemical cues of predator presence by either freezing or increasing activity

depending on whether confronted by predator cues of sit-and-wait or active predators

(Storm and Lima 2008, Binz et al. 2014). Here we used a behavioral assay to measure

response to cues of predator presence also previously used with another Gryllid species

(Royauté and Dochtermann 2017, Royauté et al. 2019). Specifically, individuals were

introduced into a 15 cm diameter circular arena (7.5 cm height), the floor of which was

covered with dry filter paper that had been soaked with diluted excreta from leopard

geckos (*Eublepharis macularius*). Crickets respond to exposure to leopard gecko cues by

increasing activity (Royauté and Dochtermann 2017, Royauté et al. 2019). All leopard

geckos were fed a diet of *G. integer* with occasional diet supplementation of mealworms

(i.e. larval *Tenebrio molitor*) and the related decorated cricket (*Gryllodes sigillatus*). Crickets

were introduced to a portion of the arena without predator cue under a small shelter. After

a 30 second rest period, the shelter was removed and the individual allowed to freely move

throughout the arena for 3 minutes and 40 seconds. We then used Ethovision XT to record

the total distance an individual moved during the trial (cm). Total distance moved during

the predator cue trial, the latency to first movement (in seconds), and the variance in

velocity were used in subsequent analyses.

Statistical analyses

- **G** matrix estimation -

We used multi-response mixed effect animal models (Kruuk 2004) implemented using the

`MCMCglmm` package in R (Hadfield 2010) to estimate genetic variances and covariances (i.e.

the **G** matrix). We included the effects of temperature, day and time of testing in the

behavioral arena room along with sex, life-stage and mass of the individual as fixed effects.

We used the individual relatedness matrix (based on the known pedigree) as a random

effect. Traits for which variances and covariances were estimated were: (i) the latency that

an individual emerged from the shelter during the trial (modeled as censored Gaussian),

(ii) the distance moved during the open field trial (Gaussian), (iii) the number of unique

zones an individual visited during the open field trial (Poisson), (iv) the log-transformed

variance in velocity during the open field trial (Gaussian), (v) the square-root transformed

distance an individual moved during the predator cue response trial (Gaussian), (vi) the

latency to initiate movement in the antipredator response trial (Poisson) and (vii) the log-

transformed variance in velocity during the antipredator response trial (Gaussian). The

inclusion of dam ID as a random effect did not improve model fit, indicating negligible or

non-existent maternal effects and was not included in final model runs. Multi-response

models were fit individually by population with each population's variances and

covariances estimated from the posterior of an MCMC chain of 4.8×10^6 iterations, with an

800,000 burn-in period and a thinning interval of 4,000. A prior that was minimally

informative for both variances and covariances was used. All variances and covariances

were estimated at the additive genetic level and on the latent scale (Table S4).

- Evaluating the constraints and adaptive hypotheses -

The location, orientation, and distribution of genetic variation in multivariate space—in

our case seven dimensional space—can differ among groups in a variety of complex ways

(Phillips and Arnold 1999, Blows et al. 2004, Walsh 2007, Roff et al. 2012, Aguirre et al.

2014). We therefore used a suite of statistical approaches to compare the orientation of

genetic variation and evaluated agreement among multiple statistical approaches in

support for either the constraints or adaptive hypothesis.

To determine whether behavioral syndrome structure at the additive genetic level

was shared among populations we used two approaches:

(i)   testing whether populations exhibited shared subspaces (dimensions) of **G**

213   based on Krzanowski's common subspace analysis (Aguirre et al. 2014);

214   (ii)   comparing alignment of dominant eigenvectors among populations (i.e. $\mathbf{g_{max}}$,

215   Table 1 (Schluter 1996))

Krzanowski's common subspace analysis determines whether genetic variation is

expressed in the same dimensions and direction across groups (Aguirre et al. 2014). This

can be thought of as analogous to asking whether the populations shared principal

components (Phillips and Arnold 1999). In two dimensions, this is similar to the directions

the ellipses point in Figure 1, with shared subspaces when the ellipses point in the same

direction. For our data there were seven possible dimensions of overlap but some

dimensions may not possess substantive variation (Table S4). Following Aguirre et al.

(2014) we therefore considered those subspaces that additively contained greater than

90% of the genetic variation, which were then summarized in matrix form (**H**)(Aguirre et

al. 2014). Here, this included three possible shared subspaces ($\mathbf{h_1}$, $\mathbf{h_2}$, and $\mathbf{h_3}$; Table 1) and

we tested whether these subspaces were shared among populations to a degree greater

than expected by chance. Under the constraints hypothesis we expect genetic variation to

be expressed in the same dimensional direction among populations, manifested as shared

genetic subspaces. A lack of shared subspaces would therefore contradict the predictions of

the constraints hypothesis. However, shared subspaces may also be observed under the

adaptive hypothesis if selective pressures are the same across populations.

Because evolutionary trajectories are biased by the dimensional direction (vector)

in which most genetic variation is expressed, we also compared this vector, i.e. g_{max}

(Schluter 1996), among populations. This approach is similar to Krzanowski's common

subspace analysis but focuses on the dimension in which most variation is expressed. Here

we calculated the vector correlation between the g_{max}s of each population and, via

randomization testing, determined whether these correlations significantly differed from 0.

Even if genetic covariances constrain the path of evolutionary change, average

behaviors can change. Therefore, we estimated "$\mathbf{D}$" (Table 1)—a matrix that describes the

phenotypic divergence amongst populations and was here estimated as the (co)variance of

species means (Schluter 1996, McGlothlin et al. 2018). The eigenvectors ($\mathbf{d}$, Table 1) of this

matrix describe the direction in multivariate space in which most divergence has occurred.

If behavioral syndromes have constrained the evolution of populations, we would expect

the direction in which most phenotypic divergence in average behavior has occurred ($\mathbf{d_1}$)

to be correlated with shared subspaces identified by Krzanowski's common subspace

analysis (McGlothlin et al. 2018). Here we calculated the vector correlation between $\mathbf{d_1}$ and

$\mathbf{h_{1:3}}$. Non-zero correlations between $\mathbf{d_1}$ and any shared subspace would be consistent with

the constraints hypothesis.

As with average behaviors, genetic variances across dimensions may differ among

populations even while populations share an overall conserved genetic behavioral

syndrome. We can therefore ask whether genetic variances have been constrained by a

conserved behavioral syndrome. To do so we used methods described by Hine et al. (2009)

to calculate what are known as genetic covariance eigentensors (**E**, Table 1). This approach

starts by calculating the variances and covariances of **G** matrices across populations.

Resulting matrices can subsequently be subjected to further eigen analysis, producing

eigentensor matrices (**E**) and eigenvectors (**e**, Table 1) to describe in which dimension the

Gs of populations differ the most. If a conserved genetic behavioral syndrome, consistent

with the constraints hypothesis, has affected the divergence of **G**s among populations we

would expect correlations between the eigenvectors of eigentensors (**e**) and any shared

subspaces (**h**$_{1:3}$). Put another way, if populations share genetic variation in the same

dimension in which most of the genetic divergence occurred, this would be evidence that

syndromes channel behavioral evolution along a line of least resistance.

For the above approaches we followed the recommendations of Aguirre et al. (2014)

in that all tests were based on the full MCMC posterior distributions and null distributions

for population comparisons were based on randomizations of breeding values. To compare

whether eigenvectors were significantly aligned, we also generated a random distribution

of vector correlations following McGlothlin et al. (2018). The critical values of vector

correlations based on this distribution were 0.93 (P < 0.001), 0.85 (P < 0.01), 0.71 (P <

0.05) and 0.62 (P < 0.1). To assess the significance of eigenvalues of **H** and **E** against

random expectations, we calculated the largest posterior quantiles for which these

distributions did not overlap (Figures S2 and S3 respectively). This threshold serves as a

Bayesian probability in favor of the observed distribution being generated by patterns

other than chance (hereafter, P_{mcmc}). Because there are no clear-cut rules for interpreting

these Bayesian probabilities, we provide the following scale to indicate how we interpreted

support for inferences: P_{mcmc} < 0.7: poor evidence of difference compared to random

expectations; $P_{mcmc} > 0.8$: moderate evidence of difference compared to random

expectations; $P_{mcmc} > 0.9$ strong evidence of difference compared to random expectations;

$P_{mcmc} > 0.95$: very strong evidence of difference compared to random expectations. Other

reported probabilities (hereafter P) were interpreted according to standard criteria.

To further assess support for the adaptive and constraints hypotheses we also

compared genetic correlations across generations. Genetic correlations due to selection-

induced linkage disequilibrium are expected to decline across generations with random

mating. Specifically, Conner (2002) argued that with random mating and in the absence of

physical linkage, the magnitude of genetic correlations should halve every generation.

Because of our breeding design we were able to opportunistically and separately estimate

phenotypic and genetic correlations among behaviors by generation. Here, while mating

was restricted to be within populations, mating was random with regard to behavior and

we would therefore expect both genetic correlations (r_A) and phenotypic correlations (r_P)

to decrease during the duration of the experiment under the adaptive hypothesis. We

therefore generated expected genetic and phenotypic average absolute correlations under

these assumptions (Appendix S1) and compared the observed estimates to these

expectations. According to the constraints hypothesis we would expect correlations to

remain stable across generations while under the adaptive hypothesis we would expect

them to degrade toward a correlation of zero.

Finally, based on the estimated **G** matrices for each population, we calculated

"autonomy" ($\bar{a}$, Table 1) throughout multivariate space following Hansen and Houle

(2008). Autonomy provides an estimate of the "fraction of genetic variation that is

independent of potentially constraining characters"(Hansen and Houle 2008). Put another

way, autonomy estimates the degree to which genetic variation is free to respond to

selection (max $\bar{a}$ = 1) versus constrained by covariance (min $\bar{a}$ = 0). We did not have

predictions as to values for autonomy under either the constraints or adaptive hypotheses.

Instead, these values indicate the potential for future evolutionary constraints for each

population.

Results

Behavioral syndromes were genetically conserved among populations. Based on

Krzanowski's common subspace analysis (**H**, Table 1; Aguirre et al. (2014)), the behavioral

syndrome of *G. integer* was characterized by three dimensions of genetic covariance (**h**$_{1-3}$,

Table 1). These dimensions, and thus the overall syndrome, were shared among

populations, as indicated by all Bayesian probabilities, p_{mcmc}, being < 0.65 (Fig. S1). p_{mcmc}

values that are closer to 1 for this test would indicate departure from random expectations

and would therefore support a lack of shared syndrome structure among populations. The

shared behavioral syndrome was comprised of: *i*) genetic covariation between shelter

emergence time and predator cue responsiveness (**h**$_1$, Table 2); *ii*) a genetic boldness-

activity syndrome in which active individuals were more prone to ignore predator cues and

were quicker to exit from their shelter (**h**$_2$, Table 2); and *iii*) genetic covariance between

activity and shelter emergence (**h**$_3$, Table 2). Each of these three axes explained around

one-third of the observed genetic variance (Table 2).

Following the demonstration of genetic conservation of behavioral syndromes, we

determined whether genetic variation was primarily expressed in the same direction in

multivariate space across populations (**g**$_{max}$ alignment; Table 1). Put another way, given the

general conservation of behavioral syndrome structure at the genetic level, did populations

express most genetic variation in the same combinations of traits? Indeed, the $\mathbf{g_{max}}$s of the

Aguila and Dunnigan and Socorro and Dunnigan populations were strongly correlated with

each other (vector correlation $r > 0.7$, $p < 0.05$) (Figure 2). Moreover, the $\mathbf{g_{max}}$s of each

population were aligned with the shared axes (Figure 2). This alignment demonstrates that

the genetically conserved behavioral syndrome captured the genetic variation expressed in

each population and confirmed that the orientation of genetic variation in multivariate

space was conserved among the populations.

Despite the genetic conservation of behavioral syndrome structure, populations did

exhibit some divergence. Specifically, the populations have diverged in their multivariate

behavioral averages (i.e. "$\mathbf{D}$," Figure 3, Table 1 and S2) and in the magnitude of genetic

variation present in each population (Figure S3). Importantly, however, the direction of

divergence in both means and variances was aligned with the shared behavioral syndrome

(e.g. $r_{d1,h3} = 0.85$, $p < 0.01$; $r_{h1,e11} = 0.92$, $p < 0.01$; Table S2). This alignment demonstrates

that divergence has been constrained by the shared structure of behavioral syndromes.

Behavioral syndromes emerging from either the adaptive or constraints hypotheses

are expected to respond differently to random mating. Specifically, under the adaptive

hypothesis, genetic correlations are expected to erode by 50% every generation. Because

we mated individuals at random, we were able to compare the observed average genetic

and phenotypic correlations (r_A and r_P) with their expected values under the adaptive

hypothesis (see Appendix S1 for details). Contrary to the expectations of the adaptive

hypothesis, but as predicted according to the constraints hypothesis, average genetic and

phenotypic correlations remained stable over the course of three successive laboratory

generations (posterior mean and 95 % credible intervals; $r_{A\ Observed}$ F$_1$ = 0.36 [0.23; 0.52], r_A

$_{Observed}$ F$_2$ = 0.38 [0.23; 0.53], Figure 4).

Finally, for each population, we calculated autonomy (Table 1), which estimates the

degree of constraint on evolutionary outcomes imposed by the genetic architecture

connecting traits. Autonomy varies between 0 and 1, with higher values indicating greater

potential for independent evolution. For *Gryllus integer*, autonomy varied between 0.47 and

0.61 (DUN: $\bar{a}$ = 0.48 [0.31; 0.68], SOC: $\bar{a}$ = 0.47 [0.32; 0.67], AG: $\bar{a}$ = 0.60 [0.44; 0.78], LC: $\bar{a}$ =

0.61 [0.43; 0.76], all populations combined: $\bar{a}$ = 0.57 [0.43; 0.70]). This suggests that the

constraining effect of behavioral syndromes is likely to persist over future generations.

Discussion

Three key results demonstrate conservation of behavioral syndromes at the genetic

level despite differences among populations in average behavior, providing strong support

for the constraints hypothesis. This support for the constraints hypothesis is unexpected

given that a previous study with stickleback (Bell 2005) found that two populations

differed in the magnitude of heritabilities and genetic correlations between two

behaviors—albeit with overlapping confidence intervals—providing support in that case

for the adaptive hypothesis. This conservation of behavioral syndrome structure has also

had the effect of channeling population divergence. Our results therefore suggest that

studying a broader suite of behavioral traits may reveal evolutionary constraints not

apparent from pair-wise correlations.

Our first major result supporting the constraints hypothesis was that the genetic

variation among the four populations was shared along three dimensions. These

dimensions describe the genetic structure of the species' behavioral syndromes and their

being shared demonstrates that the orientation of genetic variation was conserved among

populations. The overall behavioral syndrome consisted of a boldness-activity dimension

($\mathbf{h}_2$, Table 2) frequently described in the literature. This dimension genetically links

activity, exploration and risk-prone behaviors. This dimension has been described at the

phenotypic level (Wilson and Godin 2009, Bókony et al. 2012) but demonstrations at the

genetic level are rare (see Bell (Bell 2005) for one example). The other conserved

dimension ($\mathbf{h}_1$ and $\mathbf{h}_3$) represent potential trade-offs between risk management strategies,

in which individuals either compensate for risk during foraging by being less prone to

resume activity when threatened ($\mathbf{h}_3$, Table 2), or take risks in one context (not moving

away from a predator cue) while avoiding risk in another (taking longer to emerge from

shelter) ($\mathbf{h}_1$, Table 2). Alternatively, dimension $\mathbf{h}_1$ might indicate that individuals with long

latencies are less active. As a result, these individuals may encounter fewer predator cues

resulting in weaker antipredator responses.

Our second major result supporting the constraints hypothesis was that the $\mathbf{g}_{max}$s of

the Aguila and Dunnigan and Socorro and Dunnigan populations were strongly correlated

and all $\mathbf{g}_{max}$s were aligned with the shared behavioral syndrome. This validates that

behavioral syndrome structure is shared among the populations and that the behavioral

syndrome captures the majority of observed genetic variation. Schluter (1996)

demonstrated that morphological divergence among several pairs of populations and

species of vertebrates is constrained by $\mathbf{g}_{max}$. Specifically, evolutionary divergence was

greatest when populations and species shared a common $\mathbf{g}_{max}$ and there was directional

selection for morphological trait combinations in this same direction in phenotypic space

(Schluter 1996). We found that $\mathbf{g}_{max}$ was conserved and that divergence in both average

behavior and genetic (co)variance among the four populations was aligned with $\mathbf{g}_{max}$. This

demonstrates that behavioral syndromes affect population divergence in a manner similar

to that observed for morphology.

Our third result in support of the constraints hypothesis stems from the prediction

that, under the adaptive hypothesis, genetic correlations are expected to decrease by about

50% each generation due to the effects of recombination (Conner 2002). This prediction

assumes an absence of genetic linkage and random mating (Appendix S1). However,

genetic linkage sufficiently strong to resist recombination is also consistent with the

constraints hypothesis—see, for example, the effects of supergenes (Purcell et al. 2014,

Küpper et al. 2016)—and so we consider this assumption appropriate. In contrast to this

prediction of declining correlations, we found that the average genetic correlation did not

change across generations (Figure 4). Similarly, phenotypic correlations did not decrease

according to predictions (Figure 4). Because we were not able to study replicate lines

under random mating, this finding is not conclusive on its own. Instead the result is one

additional line of evidence consistent with the constraints hypothesis and in contradiction

of the adaptive hypothesis.

Importantly, the first two results—shared dimensions of genetic variation and

correlated $\mathbf{g}_{max}$s—could also be observed under the adaptive hypothesis if the selective

pressures each of the populations experienced were the same. We consider this unlikely for

three reasons. First, the degree of geographic separation among populations was extensive,

totaling more than 1500 km in some cases (Figure 2). This degree of geographic separation

makes it unlikely that the populations experienced the exact same selective regime.

Moreover, climate (Table S5) as well as predation and parasitism regimes are highly

variable among the populations (Hedrick and Kortet 2006). Second, if similarity in

selection regimes was the driving force behind these converging patterns of genetic

covariance, we would expect the geographically closest populations to have the greatest

similarity in $\mathbf{g}_{max}$. This was not the case and, in fact, $\mathbf{g}_{max}$ was most similar among

populations that were geographically most separated (Figure 2). Finally, our third main

result directly contradicts the adaptive hypothesis: if trait correlations, like those of

behavioral syndromes, arise due to the adaptive hypothesis and therefore selection-

induced linkage disequilibrium, they are expected to rapidly degrade under random mating

(Roff 1997, Conner 2002). In direct contradiction to this expectation we observed that

correlations did not decrease across generations (Figure 4). Put another way, our first two

results—which showed that the multivariate composition of behavioral syndromes was

shared among populations—are consistent with the predictions of the constraints

hypothesis. Next, our third result—the maintenance of behavioral correlations despite

random mating—demonstrates the failure of predictions made by the adaptive hypothesis.

Our results indicate that the conserved genetic architecture of behavioral

syndromes leads to populations having quantitatively constrained evolutionary trajectories

(Houle 2001) and that these syndromes have limited population divergence. This

quantitative constraint and resulting limitation on divergence is also expected to persist

into the future due to the behavioral syndrome structure imposed by each population's $\mathbf{G}$

matrix. Based on these **G** matrices, we found similar degrees of autonomy (Hansen and

Houle 2008) among populations ranging from 0.47-0.61, a stronger constraint than

observed for life-history or morphological traits (Dochtermann and Dingemanse 2013).

These autonomies indicate that behaviors will rarely evolve independently and the

observed genetic behavioral syndrome will affect future evolution.

Despite the conservation of behavioral syndrome structure at the genetic level

across populations, *Gryllus integer* populations did exhibit divergence in both mean

behaviors and magnitudes of genetic variation present in each population. Both the

divergence in means and variances was strongly aligned with the shape of the shared

behavioral syndrome, demonstrating that the syndrome itself was channeling the

evolutionary divergence of the populations (Table S2). The divergence in means was most

strongly related to differences in latency and distance moved in both the open-field and

antipredator assays (Table 2). Specifically, the Dunnigan, CA and Aguila, AZ populations

exhibited the greatest differences in average behaviors (Figure 4). Similarly, the divergence

in magnitude of genetic variation was driven by the three easternmost populations having

less genetic variation than the Dunnigan, CA population (Figure S1). Whether this

represents a loss of variation due to selection, stochastic effects on the three eastern

populations, or the accumulation of variation for the western population is not currently

clear.

Throughout we have referred to the adaptive and constraints hypothesis as

competing hypotheses for the expression of behavioral syndromes. However, two caveats

to this framing exist: First, according to a Tinbergian framework (Tinbergen 1963), these

hypotheses are not addressing questions at the same level. Within the framework of

Tinbergen's four questions, the constraints hypothesis is a proximate causation question

and reduces to a question about pleiotropy (or other molecular mechanisms) versus

linkage and is agnostic as to selection. In contrast, the adaptive hypothesis is an ultimate

question of function and would be assessed by determining the alignment between **G** and

selection gradients (e.g. following methods described by Berdal and Dochtermann 2019).

This framing does not, however, carry-over to the quantitative genetic literature wherein

selection-induced linkage disequilibrium and molecular mechanisms such as pleiotropy are

considered competing explanations (e.g. Roff 1997). The difference in perspective is partly

due to the history of the fields but is also because the quantitative genetic framework

recognizes the necessary role of continuing correlated selection in maintaining covariances

stemming from linkage disequilibrium. As a second caveat it is important to note that the

hypotheses are not strictly mutually exclusive (Conner et al. 2011, Saltz et al. 2017). It is

possible that some portion of an estimated genetic covariance might stem from pleiotropy

while some other portion stems from linkage disequilibrium (the two could even cancel

each other out). Nonetheless, our results consistently supported the predictions of the

constraints hypothesis across several lines of evidence.

The surprising degree of shared genetic variation in behavioral syndrome reported

here suggests an unrecognized and important role for behavioral syndromes in the

evolution of populations. Behaviors such as those measured here—exploratory behaviors

and responses to predation threat—are frequently assumed to have been under selection

and their responses to selection have been assumed to be unconstrained. In contrast, we

have shown that the genetic contribution to behavioral expression is highly conserved, that

populations share evolutionary fates, and that conserved behavioral variation may be a

driver of population divergence and perhaps even speciation.

479   *Acknowledgements*

We thank Monica Berdal, Katelyn Cannon, Jeremy Dalos, Sarah Felde, Brady Klock, Ishan

Joshi, Hannah Lambert, Jenna LaCoursiere and Alondra Neunsinger for assistance in

conducting behavioral trials and in rearing and care of the crickets and Martori Farms,

David Lightfoot, Scott Bundy, Nico Franz, Sangmi Lee, Cameron Jones, Kenny Chapin, Ti

Eriksson, Meranda Feagins, Charlotte Mulloney, Melody Martinez, Allyson Richins, Mauriel

Rodriguez, Helen Vessels and David Wikman for assistance in collecting the crickets. We

thank J. Bowsher, J. Pruitt, D. Westneat, and members of the Gillam and Dochtermann labs

at NDSU for important discussions and/or comments on earlier versions of this paper. We

also thank Adam Reddiex and two anonymous reviewers for essential critical feedback.

This work was supported by US NSF IOS grants 1557951 and 1558069 to N.A.D. and A.H.

respectively.

491   *References Cited*
492
Aguirre, J., E. Hine, K. McGuigan, and M. Blows. 2014. Comparing **G**: multivariate analysis of

genetic variation in multiple populations. Heredity **112**:21-29.

Bell, A. M. 2005. Behavioural differences between individuals and two populations of

stickleback (*Gasterosteus aculeatus*). Journal of Evolutionary Biology **18**:464-473.

Berdal, M. A., and N. A. Dochtermann. 2019. Adaptive alignment of plasticity with genetic

variation and selection. Journal of Heredity **110**.

Binz, H., R. Bucher, M. H. Entling, and F. Menzel. 2014. Knowing the risk: crickets

distinguish between spider predators of different size and commonness. Ethology

120:99-110.

Blows, M. W., S. F. Chenoweth, and E. Hine. 2004. Orientation of the genetic variance-
covariance matrix and the fitness surface for multiple male sexually selected traits.
American Naturalist **163**:329-340.

Bókony, V., A. Kulcsár, Z. Tóth, and A. Liker. 2012. Personality traits and behavioral
syndromes in differently urbanized populations of house sparrows (Passer
domesticus). Plos One **7**:e36639.

Conner, J. K. 2002. Genetic mechanisms of floral trait correlations in a natural population.
Nature **420**:407-410.

Conner, J. K., K. Karoly, C. Stewart, V. A. Koelling, H. F. Sahli, and F. H. Shaw. 2011. Rapid
Independent Trait Evolution despite a Strong Pleiotropic Genetic Correlation.
American Naturalist **178**:429-441.

Dingemanse, N. J., and N. A. Dochtermann. 2014. Individual behaviour: behavioural ecology
meets quantitative genetics.*in* A. Charmantier, D. Garant, and L. E. B. Kruuk, editors.
Quantitative genetics in the wild. Oxford University Press.

Dingemanse, N. J., J. Wright, A. J. N. Kazem, D. K. Thomas, R. Hickling, and N. Dawnay. 2007.
Behavioural syndromes differ predictably between 12 populations of three-spined
stickleback. Journal of Animal Ecology **76**:1128-1138.

Dochtermann, N. A. 2011. Testing Cheverud's conjecture for behavioral correlations and
behavioral syndromes. Evolution **65**:1814-1820.

Dochtermann, N. A., and N. J. Dingemanse. 2013. Behavioral syndromes as evolutionary
constraints. Behavioral Ecology **24**:806-811.

Dochtermann, N. A., S. H. Jenkins, M. Swartz, and A. C. Hargett. 2012. The roles of
competition and environmental heterogeneity in the maintenance of behavioral
variation. Ecology **93**:1330-1339.

Dochtermann, N. A., T. Schwab, and A. Sih. 2015. The contribution of additive genetic
variation to personality variation: heritability of personality. Proceedings Of The
Royal Society B-Biological Sciences **282**:20142201.

Garamszegi, L. Z., G. Marko, and G. Herczeg. 2012. A meta-analysis of correlated behaviours
with implications for behavioural syndromes: mean effect size, publication bias,
phylogenetic effects and the role of mediator variables. Evolutionary Ecology
26:1213-1235.

Garamszegi, L. Z., G. Marko, and G. Herczeg. 2013. A meta-analysis of correlated behaviors
with implications for behavioral syndromes: relationships between particular
behavioral traits. Behavioral Ecology **24**:1068-1080.

Grafen, A. 1984. Natural selection, kin selection and group selection. Pages 62-84 *in* J. R.
Krebs and N. B. Davies, editors. Behavioural ecology: An evolutionary approach.
Sinauer, Sunderland.

Hadfield, J. D. 2010. MCMC Methods for Multi-Response Generalized Linear Mixed Models:
The MCMCglmm R Package. Journal of Statistical Software **33**:1-22.

Hansen, T. F., and D. Houle. 2008. Measuring and comparing evolvability and constraint in
multivariate characters. Journal of Evolutionary Biology **21**:1201-1219.

Hedrick, A. V. 1988. Female choice and the heritability of attractive male traits: an empirical
study. American Naturalist **132**:267-276.

Hedrick, A. V., and R. Kortet. 2006. Hiding behaviour in two cricket populations that differ
in predation pressure. Animal Behaviour **72**:1111-1118.

Hedrick, A. V., and R. Kortet. 2012. Sex differences in the repeatability of boldness over
metamorphosis. Behavioral Ecology and Sociobiology **66**:407-412.

Herman, C. S., and T. J. Valone. 2000. The effect of mammalian predator scent on the
foraging behavior of *Dipodomys merriami*. Oikos **91**:139-145.

Hine, E., S. F. Chenoweth, H. D. Rundle, and M. W. Blows. 2009. Characterizing the evolution
of genetic variance using genetic covariance tensors. Philosophical Transactions of
the Royal Society B: Biological Sciences **364**:1567-1578.

Houle, D. 2001. Characters as the units of evolutionary change.*in* G. P. Wagner, editor. The
character concept in evolutionary bioloy. Academic Press, New York.

Kortet, R., and A. Hedrick. 2007. A behavioural syndrome in the field cricket Gryllus integer:
intrasexual aggression is correlated with activity in a novel environment. Biological
Journal of the Linnean Society **91**:475-482.

Kortet, R., M. J. Rantala, and A. Hedrick. 2007. Boldness in anti-predator behaviour and
immune defence in field crickets. Evolutionary Ecology Research **9**:185-197.

Kruuk, L. E. B. 2004. Estimating genetic parameters in natural populations using the
'animal model'. Philosophical Transactions of the Royal Society of London Series B-
Biological Sciences **359**:873-890.

Küpper, C., M. Stocks, J. E. Risse, N. dos Remedios, L. L. Farrell, S. B. McRae, T. C. Morgan, N.
Karlionova, P. Pinchuk, and Y. I. Verkuil. 2016. A supergene determines highly
divergent male reproductive morphs in the ruff. Nature genetics **48**:79.

Lynch, M., and B. Walsh. 1998. Genetics and Analysis of Quantitative Traits. Sinauer
Associates, Sunderland, MA.

McGlothlin, J. W., M. E. Kobiela, H. V. Wright, D. L. Mahler, J. J. Kolbe, J. B. Losos, and E. D.
Brodie III. 2018. Adaptive radiation along a deeply conserved genetic line of least
resistance in *Anolis* lizards. Evolution Letters **2**:310-322.

Michelangeli, M., D. G. Chapple, C. T. Goulet, M. G. Bertram, and B. Wong. 2018. Behavioral
syndromes vary among geographically distinct populations in a reptile. Behavioral
Ecology.

Niemela, P. T., N. DiRienzo, and A. V. Hedrick. 2012a. Predator-induced changes in the
boldness of naive field crickets, Gryllus integer, depends on behavioural type.
Animal Behaviour **84**:129-135.

Niemela, P. T., A. Vainikka, A. V. Hedrick, and R. Kortet. 2012b. Integrating behaviour with
life history: boldness of the field cricket, *Gryllus integer*, during ontogeny. Functional
Ecology **26**:450-456.

Phillips, P. C., and S. J. Arnold. 1999. Hierarchical comparison of genetic variance-
covariance matrices. I. Using the Flury hierarchy. Evolution **53**:1506-1515.

Pruitt, J. N., S. E. Riechert, G. Iturralde, M. Vega, B. M. Fitzpatrick, and L. Aviles. 2010.
Population differences in behaviour are explained by shared within-population trait
correlations. Journal of Evolutionary Biology **23**:748-756.

Purcell, J., A. Brelsford, Y. Wurm, N. Perrin, and M. Chapuisat. 2014. Convergent genetic
architecture underlies social organization in ants. Current Biology **24**:2728-2732.

Réale, D., S. M. Reader, D. Sol, P. T. McDougall, and N. J. Dingemanse. 2007. Integrating
animal temperament within ecology and evolution. Biological Reviews **82**:291-318.

Roff, D. A. 1997. Evolutionary Quantitative Genetics. Chapman and Hall, New York.

Roff, D. A., J. M. Prokkola, I. Krams, and M. J. Rantala. 2012. There is more than one way to
skin a **G** matrix. Journal of Evolutionary Biology **25**:1113-1126.

Royauté, R., C. M. Buddle, and C. Vincent. 2014. Interpopulation variations in behavioral
syndromes of a jumping spider from insecticide-treated and insecticide-free
orchards. Ethology **120**:127-139.

Royauté, R., and N. A. Dochtermann. 2017. When the mean no longer matters:
developmental diet affects behavioral variation but not population averages in the
house cricket (*Acheta domesticus*). Behavioral Ecology **28**:337-345.

Royauté, R., C. Garrison, J. Dalos, M. A. Berdal, and N. A. Dochtermann. 2019. Current energy
state interacts with the developmental environment to influence behavioural
plasticity. Animal Behaviour **148**:39-51.

Royauté, R., K. Greenlee, M. Baldwin, and N. A. Dochtermann. 2015. Behaviour, metabolism
and size: phenotypic modularity or integration in *Acheta domesticus*? Animal
Behaviour **110**:163-169.

Saltz, J. B., F. C. Hessel, and M. W. Kelly. 2017. Trait correlations in the genomics era. Trends
in Ecology & Evolution **32**:279-290.

Schluter, D. 1996. Adaptive radiation along genetic lines of least resistance. Evolution
50:1766-1774.

Sih, A., A. Bell, and J. C. Johnson. 2004a. Behavioral syndromes: an ecological and
evolutionary overview. Trends in Ecology & Evolution **19**:372-378.

Sih, A., A. M. Bell, J. C. Johnson, and R. E. Ziemba. 2004b. Behavioral syndromes: An
integrative overview. Quarterly Review of Biology **79**:241-277.

Simmons, L. 1986. Female choice in the field cricket *Gryllus bimaculatus* (De Geer). Animal
Behaviour **34**:1463-1470.

Storm, J. J., and S. L. Lima. 2008. Predator-naïve fall field crickets respond to the chemical
cues of wolf spiders. Canadian Journal of Zoology **86**:1259-1263.

Tinbergen, N. 1963. On aims and methods of ethology. Zeitschrift fur Tierpsychologie
20:410-433.

Walsh, B. 2007. Escape from flatland. Journal of Evolutionary Biology **20**:36-38.

Walsh, R. N., and R. A. Cummins. 1976. The open-field test: a critical review. Psychological
Bulletin **83**:482.

Wilson, A. D., and J.-G. J. Godin. 2009. Boldness and behavioral syndromes in the bluegill

sunfish, Lepomis macrochirus. Behavioral Ecology **20**:231-237.

Table 1. Terms and symbol definitions.

Term or Symbol	Definition
Behavioral syndrome	Among-individual correlations of behavior. For example, individuals that are on average more aggressive also, on average, show a higher degree of exploratory propensity.
$\mathbf{G}$	Additive genetic covariance matrix.
$\mathbf{g_{max}}$	The dominant eigenvector of $\mathbf{G}$, describes the dimension in multivariate trait space with the highest additive genetic variance.
$\mathbf{D}$	Among-population divergence matrix describing patterns of population divergence in average phenotype.
$\mathbf{d}_i$	Eigenvectors of $\mathbf{D}$, $\mathbf{d}_1$ represent the dimension capturing most of the divergence in average phenotype among populations.
$\mathbf{H}$	Common subspaces of genetic variation for all four populations; describes the trait combinations that share the most genetic variation among populations.
$\mathbf{h}_i$	Eigenvectors of $\mathbf{H}$, $\mathbf{h}_1$ is analogous to $\mathbf{g_{max}}$ and describe the major axis of shared genetic variance among populations.
$\mathbf{E}_i$	Eigentensors describing subspaces for which $\mathbf{G}$ varies among population.
$\mathbf{e}_{ij}$	j^{th} eigenvector of the i^{th} eigentensor, describes the trait combinations for which genetic divergence has occurred among populations.
r	Correlation among eigenvectors, values close to 0 indicate independence of eigenvectors, values close to 1 indicate alignment of eigenvectors.
$\bar{a}$	Autonomy of $\mathbf{G}$, indicates the proportion of genetic variation unconstrained by covariance among traits. Values closer to 0 indicate stronger evolutionary constraints and values closer to 1 indicate complete autonomy of genetic variation, meaning that each trait can evolve independently in response to future selection.

Table 2. Eigenvectors of phenotypic divergence (**d**), conserved genetic variation (**h**) and divergence in **G** (**e**). Traits legend: Latency = latency to exit from the shelter, OF.Dist = distance travelled in the open-field test, UZ = number of unique zones explore in the open-field arena, OF.Var.Velo = variance in velocity in the open-field test, AP.Dist = distance travelled in the antipredator response test, AP.Lat.Mov = latency to initiate movement in the antipredator response test, AP.Var.Velo = variance in velocity in the antipredator response test.

Traits	d_1	d_2	h_1	h_2	h_3	E1 (53 %) e_{11}	E2 (31 %) e_{21}	e_{22}
Latency	**-0.32**	**0.90**	**-0.31**	**0.86**	**0.47**	**0.63**	0.08	**0.77**
OF.Dist	**-0.80**	**-0.34**	0.18	**-0.37**	**0.88**	-0.13	**-0.79**	**0.42**
UZ	-0.02	-0.06	0.02	-0.02	0.06	-0.01	-0.08	0.06
OF.Var.Velo	-0.09	-0.06	0.02	-0.02	0.07	-0.02	-0.10	0.06
AP.Dist	**-0.48**	-0.04	**0.93**	**0.35**	0.02	**-0.74**	**-0.59**	**-0.45**
AP.Lat.Mov	-0.05	0.24	-0.10	-0.01	0.02	0.16	0.07	0.15
AP.Var.Velo	-0.07	-0.03	0.05	0.02	0.01	-0.05	-0.05	-0.02
% Variance explained	58.2	31.4	33.1	33.0	32.9	97.4	69.5	30.4

Figure 1. Two contrasting hypotheses can explain the presence of genetic correlations among behavioral traits (i.e. behavioral syndromes): Genetic constraints arising from pleiotropy and shared molecular mechanisms should lead to the expression of the same behavioral syndrome (top left panel). As a result, the vector correlations between major axes of genetic variation ($\mathbf{g_{max}}$) are predicted to be approaching 1 (bottom left panel). Alternatively, selection-induced linkage disequilibrium should lead to differing orientation and strength of behavioral syndromes when selective pressures differ among populations (top right panel). The vector correlation between $\mathbf{g_{max}}$s should therefore be below 1 (bottom right panel).

	Dunnigan (CA)	Aguila (AZ)	Socorro (NM)	Las Cruces (NM)
Dunnigan (CA)	96.4 53.0 0.90**, 0.32, 0.27	Size Shape Orientation $(g_{max} \times h_{1-3})$		
Aguila (AZ)	0.89**	59.5 42.8 0.82*, 0.56, 0.09		
Socorro (NM)	0.59	0.77*	73.4 53.3 0.79*, 0.12, 0.59	
Las Cruces (NM)	0.23	0.24	0.38	51.0 46.7 0.20, 0.81*, 0.55

Figure 2. The genetic architecture of behavioral syndromes is shared in four isolated populations of *Gryllus integer*. Values along the diagonal (peach shading) describe the multivariate structure of behavioral variation: first row the size (total genetic variance), second the shape (percent of variance explained by the major axis of genetic variation, g_{max}), and, third, orientation (vector correlation between g_{max} and conserved genetic subspaces h_{1-3}). Off-diagonal elements represent the correlation between the g_{max} of each population (top row) and the probability that alignment differed from 0: ** $P < 0.01$, * $P < 0.05$.

Figure 3. Evolutionary divergence in the structure of behavioral syndromes occurs along shared axes of genetic variation. A-C) Correlations between pairs of traits that exhibit the greatest variation in divergence (Table S2). Points represent breeding values for each individual within a population centered around the population mean for that trait. >50% of divergence was in latency to emerge from shelter by antipredator response activity D) Population-specific divergence in average behaviors. Population-specific **G** matrices were visualized by transforming estimated breeding values for each trait based on the divergence among populations. Ellipses represent the 95% confidence ellipses for each population centered at the multivariate species mean (DUN: Dunnigan CA, AG: Aguila AZ, SOC: Socorro NM, LC: Las Cruces NM).

Figure 4. A) Additive genetic (r_A) and B) phenotypic correlations (r_P) remained stable over the course of three successive generations compared to theoretical expectations based on selection-induced linkage disequilibrium and random mating ($r_{A\ observed}$ = 0.38, $r_{A\ expected}$ = 0.18, P_{mcmc} for difference from expectations under selection-induced linkage disequilibrium > 0.85; $r_{P\ observed}$ F_0 = 0.32, $r_{P\ observed}$ F_2 = 0.33, $r_{P\ expected}$ F_0 = 0.35, $r_{P\ expected}$ F_2 = 0.26, P_{mcmc} F_2 > 0.80). Error bars correspond to 95% credibility intervals around the posterior mean.