The scaling of genome size and cell size limits maximum rates of photosynthesis with implications for ecological strategies

Adam B. Roddy[1], Guillaume Théroux-Rancourt[2], Tito Abbo[3], Craig R. Brodersen[1], Brook Jensen[4], Guo-Feng Jiang[5,6], R. Alexander Thompson[3], Sara E. Kuebbing[7], Kevin A. Simonin[3]

School of Forestry & Environmental Studies, Yale University, New Haven, CT 06511 USA

Institute of Botany, Universität für Bodenkultur, Vienna, Austria

Department of Biology, San Francisco State University, San Francisco, CA, 94132 USA

Department of Biological Sciences, California State University-Stanislaus, Turlock, CA 95382 USA

State Key Laboratory of Conservation and Utilization of Subtropical Agrobioresources, Guangxi University, Nanning, Guangxi 530004, China

Guangxi Key Laboratory of Forest Ecology and Conservation, College of Forestry, Guangxi University, Nanning, Guangxi 530004, China

Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260 USA

Author for contact:

Email: adam.roddy@yale.edu

Phone: +1.510.224.4432

ORCID: 0000-0002-4423-8729

Abstract

A central challenge in plant ecology is to define the major axes of plant functional variation with direct consequences for fitness. Central to the three main components of plant fitness (growth, survival, and reproduction) is the rate of metabolic conversion of CO_2 into carbon that can be allocated to various structures and functions. Here we (1) argue that a primary constraint on the maximum rate of photosynthesis per unit leaf area is the size and packing density of cells and (2) show that variation in genome size is a strong predictor of cell sizes, packing densities, and the maximum rate of photosynthesis across terrestrial vascular plants. Regardless of the genic content associated with variation in genome size, the simple biophysical constraints of encapsulating the genome define the lower limit of cell size and the upper limit of cell packing densities, as well as the range of possible cell sizes and densities. Genome size, therefore, acts as a first-order constraint on carbon gain and defines the upper limits of allocation to growth, reproduction, and defense. The strong effects of genome size on metabolism, therefore, have broad implications for plant biogeography and for other theories of plant ecology.

Introduction

Quantifying major axes of plant functional variation has given rise to an ever-growing list of traits that impact growth, reproduction, and survival, the three components of individual fitness (Violle et al. 2007). These traits have traditionally been viewed from a reductionist perspective that scales form-function relationships of individual plant organs, e.g. leaves, stems, and roots, to whole organism ecological strategies. As the ultimate source of energy and matter for growth and reproduction, photosynthetic capacity represents a first-order constraint on the emergent properties between whole plant form and function and individual fitness. Here we provide evidence that genome-cellular allometry directly influences interspecific variation in photosynthetic metabolism and provide a mechanistic framework that links genome size and metabolism to other aspects of plant ecology and evolution.

A major limitation on photosynthetic capacity is the ability to deliver resources to, and exporting products from, the sites of metabolic processing (Enquist et al. 1998; West et al. 1999a; Brown et al. 2004). At the level of an individual cell–the fundamental unit of living organisms–rates of resource transport are strongly influenced by cell size because the ratio of cell surface area to cell volume increases exponentially with decreasing cell size. Building on an extensive body of literature we show how variation in the size of the nuclear genome fundamentally limits the sizes and packing densities of cells, and, as a result, the maximum rate of photosynthetic metabolism across vascular plants.

One of the three components of fitness is growth which is ultimately limited by photosynthetic capacity. Relative growth rate (RGR) varies considerably across species and is driven by photosynthetic rate and the resource investment to support photosynthesis as:

$$RGR = A_{mass} \cdot LMR$$

where A_{mass} is the photosynthetic rate per unit leaf biomass and LMR is the leaf mass ratio (the proportion of leaf dry mass to total plant dry mass). A_{mass} is, therefore, frequently

considered a major plant strategy axis (Poorter and Remkes 1990; Poorter et al. 1990;

Reich et al. 1992). However, A_{mass} can be decomposed as:

$$A_{mass} = SLA \cdot A_{area}$$

where SLA is the specific leaf area (leaf area per leaf dry mass) and A_{area} is the net carbon

assimilation rate per unit canopy leaf area. Because of its direct effect on A_{mass}, SLA is often

considered a major predictor of interspecific variation in RGR. A_{area}, on the other hand,

varies orthogonally to SLA (Wright et al. 2004), and, therefore, determines the upper limit

of the relationship between A_{mass}, SLA, and RGR. Maximum potential A_{area} represents, then,

a fundamental limitation on the maximum amount of carbon available for allocation to

growth, reproduction, and survival relative to species ecological strategies. The centrality

of A_{area} to plant ecological strategy suggests two questions:

• First, what are the fundamental features of plant structure that determine maximum

potential A_{area}?

• Second, to what extent do these relationships scale up to affect plant ecological

strategies and evolutionary dynamics?

Here we present a mechanistic framework to address both of these questions, that is based

on a long-recognized but often overlooked relationship: positive scaling between genome

size and cell size. Although the relationship between genome size (i.e. nuclear volume) and

cell size has long been of interest (Sachs 1893), the mechanisms are still not fully

understood (Doyle and Coate 2019). Yet, we show that the allometry between genome size

and cell size directly influences rates of photosynthetic metabolism and argue that the

scaling of genome size and metabolism affects ecological distributions and evolutionary

dynamics. In this way, any factor affecting rates of metabolism is a potential agent of

selection on the genome, if only via selection on genome size.

It is now widely recognized that variation in genome size can have significant

consequences for organismal structure and function, independent of the genes that define

the genotype (Bennett 1971). Positive scaling between genome size and cell size across

terrestrial plants has given rise to numerous studies characterizing the many other

phenotypic correlates of genome size independent of variation in genome structure,
commonly referred to as "nucleotype" effects (Bennett 1971; Cavalier-Smith 1978; 1982;
Bennett and Leitch 2005). Correlates of genome size encompass an incredible diversity of
plant phenotypes, including, for example, the sizes of plant structures, rates of cell division,
rates of physiological processes, and tolerances and responses to abiotic conditions (Table
1).

Our goal is not to recapitulate the many reviews about the nucleotype-phenotype
relationship but, instead, to align these studies more systematically with the field of plant
functional biology. We believe that the diverse impacts of genome-cellular allometry on the
body plan of terrestrial vascular plants strongly influences the coordination between plant
functional traits and, ultimately, whole organism form-function relationships. Here we
summarize previous research, perform new analyses of existing data, and present new data
to show how genome size may, through its impacts on cell size and tissue structure,
determine the biophysical limits of plant metabolic rates, and, therefore, influence other
aspects of ecology and evolution. That genome size may be a key functional trait is not a
new idea (Grime 1998). Yet, despite numerous reports of the phenotypic and ecological
correlates of genome size (Table 1), it has not been fully identified and integrated into the
functional trait literature. Our goal, therefore, is to more directly show how genome size
influences plant traits that impact maximum rates of photosynthetic metabolism.
Metabolism is central to all three aspects of plant fitness, providing the carbon necessary
for allocation to growth, reproduction, and survival. As such, genome size may not itself be
a functional trait but instead may define the limits of variation in numerous other
functional traits.

Genome-cellular allometry limits rates of resource transport and metabolism

Allometry of genome size and cell size

The role of the genome in limiting cell size has been postulated since at least the late 1800s
(Sachs 1893) and was critical in shaping early modern views of the evolution of plant

vascular systems (Bailey and Tupper 1918). At a minimum, a cell must contain its genome,

and there is a strong relationship between the volumes of meristematic cells and genome

size (Šímová and Herben 2012). Cellular expansion from this meristematic minimum size is

cell type-specific (Doyle and Coate 2019). Within a cell type, size can be influenced by

various environmental and developmental factors (Melaragno et al. 1993). Despite this

substantial growth in cell volume during development, there remains a significant effect of

genome size on cell size, particularly for stomatal guard cells (Beaulieu et al. 2008; Knight

and Beaulieu 2008; Lomax et al. 2013; Simonin and Roddy 2018). For example, stomatal

guard cell size and density, which regulate the fluxes of water and CO_2 between the

biosphere and atmosphere, vary within species depending on light, water availability, and

atmospheric CO_2 concentration (Hetherington and Woodward 2003; Franks and Beerling

2009). Furthermore, in the vascular transport network, the sizes of xylem conduits and

their density in the leaf are also affected by variation in genome size (Maherali et al. 2009;

Hao et al. 2013; De Baerdemaeker et al. 2018; Simonin and Roddy 2018). Yet why genome

size and final cell size are correlated within a cell type remains unclear (Doyle and Coate

2019).

We tested whether smaller genomes allow not only for smaller initial and final cell sizes

but also for a greater range in final cell size using published data for terrestrial C3 plants.

We used data for stomatal guard cells because they are the most commonly measured cell

sizes in plants and because their sizes and abundance determine the leaf surface

conductance to CO_2 and water vapor and, therefore, directly control rates of resource

transport for use in photosynthetic metabolism. Sizes of guard cells for angiosperms

(Beaulieu et al. 2008), gymnosperms, and ferns were compiled previously by Simonin and

Roddy (2018), and here we include data for mosses and hornworts from Feild et al. (2015)

and Renzaglia et al. (2017). We assumed that stomatal guard cells are shaped as capsules,

which are composed of a central cylinder with hemispherical ends, such that cell volume

could be estimated from cell length as:

150
$$V = \pi \cdot r^2 \cdot (\frac{4}{3}r + a)$$

where r is the radius of the cylinder and a is the height of the cylinder. We assumed that a
is equal to $2r$. The dumbbell-shaped guard cells common among grasses would likely
violate these assumptions about cell shape and so we have not included in this analysis
data for lineages known to have dumbbell-shaped guard cells. Data for meristematic cell
volume and genome size were taken from Simova and Herben (2012). We used standard
major axis regression (R package *sma*) to fit the mean response and quantile regression (R
package *rq*) to test whether there was greater variation in cell volume among taxa with
smaller genomes (i.e. heteroskedasticity).

For a given genome size, mature guard cells were at least three orders of magnitude more
voluminous than meristematic cells (Figure 1). Genome size was a strong and significant
predictor of meristematic cell volume (log(volume) = 1.24 log(genome size) - 0.23; R^2 =
0.98, P < 0.001; Šímová and Herben 2012). Though it explained less of the variation,
genome size was a significant predictor of final guard cell volume (log(cell volume) = 0.80
log(genome size) + 3.43; R^2 = 0.09, P < 0.001). Quantile regression revealed that the slope
through the 10% quantile was steeper (0.58) than the slope through the 90% quantile (-
0.07), reflecting both the greater variation in final cell volume among species and the
smaller minimum possible cell volume that can be obtained with smaller genomes
(i.e. heteroskedasticity in cell volume with variation in genome size). Thus, smaller
genomes allow for smaller minimum cell size with little or no effect on maximum cell size,
together enabling greater potential variation in final cell size. The greater variation among
species with smaller genomes implies that smaller genomes allow for greater plasticity in
cell sizes and cell packing densities which directly influence maximum rates of leaf surface
conductance to CO_2 and water and ultimately photosynthetic metabolism per unit leaf
surface area (Simonin and Roddy 2018). Further, the greater diversity of cell sizes
observed in plants with small genomes is supporting evidence that the correlation between
genome size and cell size is simply the result of occupying available space within the cell. A
small genome can be housed in either a small or a large cell, but a large genome requires a
cell large enough to contain its volume.

The greater variation in cell volume allowed by smaller genomes across terrestrial C3
plants (Figure 1) further suggests that smaller genomes allow for greater variation in cell

packing densities. For guard cell lengths, stomatal densities, and vein densities, smaller

genomes allowed for greater variation in traits across ferns, gymnosperms and

angiosperms (Simonin and Roddy 2018). Species with smaller genomes in these datasets

are predominantly angiosperms, and these analyses compared distantly related species.

We further tested for greater variation in cell sizes and packing densities with smaller

genomes among closely related species using taxa in *Rhododendron* sect. *Schistanthe* Schltr.

(= sect. *Vireya* Blume). This monophyletic clade has a stepwise phylogeographic history,

having radiated eastward from the Malay Peninsula and reached New Guinea within the

last 15 Ma (Goetsch et al. 2011). We sampled leaves from 19 taxa growing under common

garden conditions at the Rhododendron Species Foundation Botanical Garden in Federal

Way, WA, USA. Genome sizes were measured following standard protocols (Dolezel et al.

2007) at the Benaroya Research Institute in Seattle, WA, USA. For measurements of

stomatal size and density, epidermal impressions were made on fresh leaves using dental

putty (Coltene Whaledent President Light Body), transferred using clear nail polish,

mounted in water, and imaged using a light microscope. Measurements of leaf vein density

were made on leaf sections cleared by soaking in 4% NaOH, 3% sodium hypochlorite,

stained with 1% Safranin O, counterstained with 1% Fast Green, mounted in ethanol, and

imaged with a light microscope. Stomatal traits were averaged across ten images per taxon,

and leaf vein density was averaged across five images per taxon. Quantile regression (R

package 'rq') through the 10th and 90th percentile of the species means were used to

quantify the variation in traits associated with variation in genome size. Consistent with

previous results across vascular plants (Simonin and Roddy 2018), among the closely

related *Rhododendron* sect. *Schistanthe*, there was greater variation in the sizes and packing

densities of veins and stomata among species with smaller genomes (Figure 2).

Furthermore, the steeper quantile slopes for the lower limit of cell size and the upper limit

of packing densities are consistent with the hypothesized constraints imposed by genome

size variation.

Genome size limits maximum photosynthetic metabolism

Previous work has hypothesized that genome size would be linked to maximum

photosynthetic rate but found little support (Knight et al. 2005; Beaulieu et al. 2007). We

hypothesize that one major reason for not finding support is that these previous studies

attempted to predict variation in A_{mass}, which accounts for the construction costs of leaves,

rather than A_{area}, which is the maximum metabolic rate regardless of the construction costs.

As we described above, A_{area} would define the maximum amount of carbon assimilated, but

how the plant allocates the total assimilated carbon–to growth, reproduction, defense,

more durable leaves, etc. (Bazzaz et al. 1987)–would reflect the numerous factors that

influence plant form and other aspects of plant function. Thus, A_{area}, which is orthogonal to

SLA and A_{mass} (Wright et al. 2004), is predicted to be constrained by cell and genome sizes.

Consistent with this prediction, recent work has shown that genome size is a strong

predictor of the sizes and densities of stomatal guard cells and leaf veins across vascular

plants (Simonin and Roddy 2018), and we predicted, therefore, that genome size would, via

its effects on the sizes and packing densities of cells, predict A_{area}.

Data for area-based maximum photosynthetic rate were compiled from the primary

literature (Supplemental Table 1) and merged with the Kew Plant DNA C-Values Database

(Bennett and Leitch). This dataset included 210 species, of which 138 were angiosperms,

46 were gymnosperms, and 26 were ferns. We tested for a relationship between genome

size and A_{area} both for the mean response (standard major axis regression, SMA; R package

'smatr') and for the upper 90% quantile of the data using quantile regression (R package

rq). To test whether the SMA regression was driven by phylogenetic history, we used a

phylogenetically corrected version (R package 'phytools'), which provides a P-value only

for the test against a specific slope, for which we used the non-phylogenetically corrected

SMA slope. For this analysis, we generated a dated supertree using the methods of Simonin

and Roddy (2018), which included generating a family-level supertree using the R package

'brranching' (Chamberlain 2016) and applying node ages from Magallón et al (2015). Of the

128 nodes, 43 with node names had ages. These ages were applied and intermediate

branch lengths smoothed using 'bladj' in Phylocom (Webb et al. 2008). We randomly

resolved polytomies by adding 5 million years to each branch and subtracting an

equivalent amount from the subtending branches to keep the tree ultrametric.

Species with smaller genomes had higher maximum photosynthetic rates across and within

major plant clades (Figure 3). Standard major axis regression on log-transformed data

revealed a significant negative relationship between genome size and A_{area} across all

species (log(A_{area}) = -0.42 log(genome size) + 1.29; R^2 = 0.23, P < 0.001). The phylogenetic

major axis regression was significantly steeper than the non-phylogenetic major axis

regression (log(A_{area}) = -0.75 log(genome size) + 1.76; slope test P < 0.001), suggesting that

the evolution of genome size across land plants is strongly associated with changes in A_{area}

across vascular plants. A similarly negative relationship was apparent among the upper

(90%) quantile, which showed that A_{area} was lower among species with larger genomes.

Despite the large difference in genome size between angiosperms and gymnosperms, the

90% quantile regression slopes were nearly identical for the entire dataset (-0.19) as for

the angiosperms alone (-0.18).

The scaling relationship between A_{area} and genome size follows naturally from the

relationships between genome size and the sizes and densities of veins and stomata.

However, veins and stomata are not the only cells responsible for driving variation in

photosynthetic rates. While the maximum rate of CO_2 diffusion into the leaf is defined by

the sizes and densities of stomata (Franks and Beerling 2009), once inside the leaf, CO_2

must diffuse through the leaf intercellular airspace and into the chloroplasts lining the

interior surfaces of mesophyll cells. The diffusive surfaces of the leaf mesophyll are the

interface between liquid-phase and vapor-phase resource movement, where liquid water

delivered by the veins evaporates and where CO_2 must diffuse into the liquid phase. Thus,

the three-dimensional structure and organization of the mesophyll is predicted to be a

prime target for selection on photosynthetic metabolism (Tholen et al. 2012; Ren et al.

2019) and to be critical to leaf photosynthetic function (Earles et al. 2019), and the limited

evidence on *Arabidopsis thaliana* mutants with altered cell diameter and porosities point in

that direction (Lehmeier et al. 2017). We predict, therefore, that the scaling relationships

between genome size and cell size that coordinate veins and stomata (Simonin and Roddy

2018) extend also to the photosynthesizing cell size, structure, and organization within the

leaf mesophyll. The significant negative scaling relationship between A_{area} and genome size

would emerge from the effects of genome size on the sizes and packing densities in each of

these tissues.

Genome size limits the rate of metabolic up- or down-regulation

Although maximum potential rate of leaf gas exchange is an important parameter determining a species' physiological capacity, the actual rate of leaf gas exchange at any given moment is often substantially lower, depending on a variety of physiological and environmental factors (e.g. light level, atmospheric humidity, leaf temperature, plant water status). Changes in sun angle, shading by passing clouds, and self-shading by fluttering leaves all drive changes in incoming solar radiation, and these rapid dynamics have influenced the evolution of photosynthetic biochemistry (Pearcy 1990). Under naturally varying conditions, leaf gas exchange fluctuates dramatically and rarely reaches its maximum rate, with greater variation occurring at the top of the plant canopy. How frequently a leaf can reach its maximum gas exchange rate and how well optimize its physiological processes to environmental conditions depend on how rapidly the leaf can respond to dynamic, fluctuating conditions.

An emerging consensus is forming that smaller stomata respond more rapidly to fluctuating conditions than larger stomata, allowing leaves with smaller stomata to more closely tune their physiological rates with environmental conditions (Drake et al. 2013; Lawson and Blatt 2014; Lawson and Vialet-Chabrand 2019). Leaf physiological processes change at different rates, with changes in stomatal conductance occurring an order of magnitude more slowly than changes in photosynthesis (McAusland et al. 2016). This difference in response times between physiological processes (e.g. photosynthetic assimilation rate and stomatal conductance) can reduce water use efficiency when stomata are closing and reduce photosynthetic efficiency when stomata are opening (Lawson and Vialet-Chabrand 2019), limiting total photosynthesis by up to 20% (Lawson and Blatt 2014). If stomatal response times are directly limited by the size of stomata then genome-cellular allometry fundamentally limits not only the maximum rate of metabolism but also how quickly metabolism can respond to environmental change. Of the species for which stomatal response times were measured by McAusland et al. (2016) and Drake et al. (2013), twelve were included in the Kew Plant DNA C-Values database. Consistent with previous reports, there was a positive correlation between genome size and guard cell length (Figure 4a), and stomatal response rate exhibited a triangular relationship with

300 genome size such that smaller genomes exhibited both higher maximum stomatal response
301 rates but also a greater variation in stomatal response rate. While the available data on
302 stomatal response rates measured using standard protocols are limited, these preliminary
303 results suggest that genome size indirectly limits the maximum rate of stomatal opening
304 and closing via its effects on the sizes and densities of stomata.

Implications of genome size-metabolism scaling on plant biogeography

Polyploidy thought to increase habitat breadth

Variation in genome size and structure associated with polyploidization has long been
considered to be an important driver of plant evolution and to be associated with
environmental tolerances, habitat breadth, trait variation, and interspecific interactions
(Stebbins 1940; Otto and Whitton 2000; Soltis et al. 2003; Soltis et al. 2014; Michael S
Barker et al. 2016; 2016), all of which influence species biogeography. Describing the types
of polyploids and how they are generated is outside the scope of the current manuscript
and have been thoroughly reviewed elsewhere (e.g. Stebbins Jr 1947; Soltis et al. 2015).
Rather, we focus our discussion here on how and why ploidy–via its relationship with
genome size–may or may not correlate with species distributions and habitat breadth.

Polyploids have been hypothesized to be better adapted to extreme habitats, to have
greater hardiness, and to have greater ecological adaptability [reviewed by (Stebbins 1985:
@Brochmann:2004aa). The possible mechanisms for these effects can be roughly grouped
into two categories: one involving the genetic and genic content of the polyploid genome
and one involving the nucleotypic effects of ploidy and genome size. Because polyploid
genomes have additional genome copies, they have higher absolute genic contents, would
enable neofunctionalization of duplicated genes, and typically have higher heterozygosity,
all of which can promote higher tolerances of environmental conditions. The nucleotypic
effects of ploidal variation, though long recognized (Stebbins 1940), are often confounded
with nucleotypic effects of genome size variation.

While ploidy and genome size are commonly assumed to be synonymous, at broad
phylogenetic scales there is frequently no relationship between genome size and ploidy
(Leitch and Bennett 2004), reflecting the complex history of both ancient and
contemporary whole genome duplications, particularly among the angiosperms (Jiao et al.
2011; Clark and Donoghue 2018; Landis et al. 2018; Ren et al. 2018). Angiosperm genomes
readily rediploidize after polyploidization such that genome size and ploidy are positively
correlated only for narrowly defined phylogenetic groups (i.e. within genera and families,
Figure 5; Leitch and Bennett 2004; Dodsworth et al. 2016). Ferns, by contrast, the other
major group of vascular plants that readily undergoes polyploidization events seem not to
rediploidize as readily (Clark et al. 2016), as evidence by the positive correlation between
genome size and ploidy (Figure 5). If leaf and plant structure and function influence
ecological tolerances and habitat breadth (i.e. if plant structure-function is adaptive), then
the nucleotypic effects of genome size are predicted to influence environmental tolerances.

Smaller genomes enable greater phenotypic plasticity

One long-standing hypothesis is that higher ploidy is related to wider habitat breadth
because polyploids can tolerate greater ecological stress. Higher ploidy is associated with
greater heterozygosity (i.e. greater genetic diversity) and, frequently, higher genic content
due to multiple genome copies, both of which are thought to promote plasticity and enable
polyploids to withstand a greater range of environmental conditions than diploids.
However, several studies testing this hypothesis have not observed polyloids to have
greater habitat breadth (e.g. Stebbins 1985; Martin and Husband 2009; Johnson et al.
2014). Furthermore, these tests frequently find that diploids exhibit greater habitat
breadth than polyploids (Petit and Thompson 1999; Hijmans et al. 2007; Brittingham et al.
2018). One reason is that traits of polyploids are not necessarily more variable than in
diploids (Stebbins 1985; Wei et al. 2018).

We predict that one reason ploidy is not commonly found to correlate with ecological
breadth is because genome size–rather than ploidy per se–drives variation in the absolute
range of potential cell sizes and, by extension, phenotypic plasticity in rates of resource
transport and metabolism. Thus, the phylogenetic scale-dependence of the relationship

between genome size and ploidy could lead to confounding patterns depending on the
phylogenetic scale at which comparisons are made. For example, in the analysis of Rice et
al. (2019), ploidy was determined relative to other closely related species, such that within
genera or families ploidy and genome size are positively correlated, suggesting that the the
bias towards higher abundances of polyploids at higher latitudes may reflect nucelotypic
effects of genome size on cell size and metabolism. The complex, fluctuating process of
polyploidization and rediploidization, which can winnow the genome nonrandomly
(Wendel 2015), would promote the proliferation of beneficial elements associated with
genome duplications (e.g. more gene copies that can neofunctionalize) while reducing the
size of the genome needed to maintain high rates of development and metabolism (Table
1).

We posit here that the nucleotypic effects of genome size, regardless of ploidy, may
influence environmental tolerances. Because smaller genomes allow for greater variation
in cell size (Figure 1), species with smaller genomes may be better able to fine tune their
tissue structure to environmental conditions. This flexibility would allow species with
smaller genomes to better optimize their metabolic rates in order to occupy a wider range
of environmental conditions. Combined with the effects of genome size on rates of cell
division (Van't Hof and Sparrow 1963; Hof 1965; Šímová and Herben 2012), the greater
plasticity in cell size and higher rates of maximum photosynthesis available to species with
small genomes may enable them to better colonize new habitats.

Greater variation in genome size in less productive habitats

In habitats that can support high rates of productivity and primary metabolism, species
with small genomes are expected to predominate. At a broad scale, this prediction holds
because angiosperms, which have, on average, smaller genomes than other vascular plants
are dominant in most ecosystems, particularly those characterized by high productivity.
However, high rates of metabolism, even among the angiosperms, are not always favorable.
Two such habitats are those characterized by extreme water limitation and nutrient
limitation, such as deserts and epiphytic habitats, and by extreme cold, such as high
latitudes. Higher incidences of polyploids have been commonly reported in higher latitudes

and among arctic floras (Brochmann et al. 2004; Rice et al. 2019), but arid habitats have
received less attention.

Arid and epiphytic habitats are characterized by low productivity and may support species
with large genomes. In these habitats, high rates of metabolism are not always favored,
which may relax selection for small genomes. One strategy common in arid and epiphytic
habitats is succulence, which is often associated with Crassulacean acid metabolism (CAM)
photosynthesis. The CAM syndrome limits water loss by sequestering CO_2 uptake and
water loss to nighttime which corresponds with periods of concomitant water loss to
periods of high humidity and little energy available to drive water loss (i.e. at night). As a
result, this strategy is characterized by low rates of resource transport. If metabolism is
one axis of selection on genome size, then we would predict that CAM species may,
therefore, have larger genomes than species employing C3 metabolism. We tested this
hypothesis using the taxonomic distributions of CAM photosynthesis from Smith and
Winter (1996) and genome size data from the Kew Plant DNA C-Values Database (citation).
For C3, we used the broad distribution of angiosperms reported in Simonin and Roddy
(2018), which are representative of extant angiosperm diversity. We scored as CAM the
narrowest taxonomic level in the Kew DNA C-Values Database that was listed as containing
CAM by Smith and Winter (1996). For example, if a genus were listed as containing any
CAM species, all species in the genus were assumed to exhibit CAM photosynthesis. This
strategy was biased against observing differences in genome size between C3 and CAM
species because it necessarily grouped some C3 species as CAM. Nonetheless, genome sizes
were significantly larger among CAM species than among C3 species even after log-
transforming the data to improve normality (t-test: t = 9.72, df = 272.6, P < 0.0001; Figure
6), consistent with the prediction that large genomes may evolve when selection for high
rates of metabolism is weak.

Arid habitats are not exclusively composed of species with large genomes. Rather, they may
harbor a diversity of strategies associated with divergent niches. While many C3 species in
deserts function during a relatively narrow window in the year, either because they are
annuals or because physiological activity in perennial shrubs is restricted to a narrow
range of environmental conditions, when water is available. Thus, because their carbon

gain is limited to such a short time period, they may have small genomes and cells that
enable high rates of metabolism. Desert shrubs have the highest rates of stem hydraulic
conductance measured in C3 plants (Mencuccini 2003), and even among species from
humid tropical forests, dry forest species have higher hydraulic conductance than wet
forest species (Brenes-Arguedas et al. 2013). Thus, less productive habitats may allow for
multiple strategies that encompass a broader range of metabolic rates and, allowing for
greater variation in genome size at the community level than might be observed in
conditions more favorable to high rates of metabolism and productivity.

Smaller genomes increase the probability of invasiveness

The multifaceted effects of genome size on plant structure, function, and ecology (Table 1)
is particularly relevant to the study of invasive species. Identifying the traits that allow an
introduced species to establish, naturalize, and invade into a new environment is a central
aim of invasion biology (Simberloff 2011), with broader implications for plant
biogeographic patterns. Here we distinguish between nonnative species–those that survive
and reproduce in their introduced range–and nonnative invasive species–those that can
disperse and establish far from their original source of introduction (Richardson et al.
2011). This distinction is important because prior studies on the traits of 'invaders' focus
on these different subsets of species, which have slightly different, but overlapping, sets of
traits that determine whether they can survive and reproduce versus invade (Kleunen et al.
2015).

Early theory on the distinguishing traits of invasive plants postulated that "ideal weeds"
should grow rapidly, produce seed continuously and in high number throughout the
growing season, be tolerant to a wide range of environmental conditions, exhibit high trait
plasticity, and be able to reproduce vegetatively from fragments (Baker 1974). On average,
these predictions have been upheld, with nonnative invasive plants tending to exhibit traits
consistent with high fitness (e.g., number of flowers, fruits, or seed or germination rates),
high relative growth rates, high dispersal abilities (smaller seeds), and more efficient
carbon-capture strategies (e.g. high specific leaf area), relative to co-occurring native
species (Leishman et al. 2007; Kleunen et al. 2010; Ordonez et al. 2010; Kuester et al. 2014)

or naturalized but not invasive nonnative species (Rejmánek and Richardson 1996;

Gallagher et al. 2014). Combined, these traits confer a growth advantage, such that plants

with small seeds can disperse further distances, have shorter generation times, and higher

relative growth rates, owing to the greater rates of cell division and higher metabolic rates

provided by smaller genomes (Pandit et al. 2014; Suda et al. 2015). Indeed, even within

species, populations with smaller genomes are more likely to successfully invade new

habitats (Pysek et al. 2018).

Because many of the traits linked with invasiveness can be influenced by both ploidy and

genome size, both have been implicated as underlying features driving invasion (Pandit et

al. 2014; Suda et al. 2015). Because polyploids are thought to be better able to tolerate

environmental fluctuations and to be better able to adapt to new environments, polyploids

tend to be overrepresented among nonnative invasives compared to native angiosperms

(Rejmánek and Richardson 1996; Prentis et al. 2008; Beest et al. 2011; Pandit et al. 2014).

Similarly, nonnative invasive species tend to have smaller genomes than non-invasive

plants (both native and non-native), which is thought to be due to the diverse effects of

genome size on metabolism, rates of development and growth, and seed size (Rejmánek

and Richardson 1996; Bennett et al. 1998; Kubešová et al. 2010; Pandit et al. 2014).

However, the complex, scale-dependent relationship between ploidy and genome size

discussed previously complicate a clear understanding of the effects of ploidy versus

genome size on invasiveness (Rejmánek and Richardson 1996; Pandit et al. 2014). Because

angiosperms, which predominate among nonnative invasives, readily rediploidize and

downsize their genomes subsequent to whole genome duplications (Leitch and Bennett

2004), assessing the relative effects of ploidy versus genome size on invasiveness can be

difficult. For example, the likelihood of being invasive increases with chromosome number

and ploidy but decreases with genome size (Rejmánek and Richardson 1996; Pandit et al.

2014). The multiple paths to polyploidization and the selective retention of only certain

parts of the genome during subsequent genome downsizing (Wendel 2015) could explain

how both higher ploidy and smaller genomes are correlated with invasiveness.

Implications for of genome size variation on the Metabolic Theory of Ecology

As the major source of energy and matter for the biosphere, photosynthetic metabolism represents a first-order control over ecological processes globally. This fundamental link between metabolic and ecological processes has driven the development of the Metabolic Theory of Ecology (MTE) that provides a mechanistic framework for predicting variation in organismal life history attributes, population dynamics, and larger scale ecosystem processes from organismal-level traits related to supplying resources for metabolism (West et al. 1997; Enquist et al. 1998; West et al. 1999a; West et al. 1999b; West et al. 2002; Price et al. 2010). While appealing and seemingly endowed with incredible explanatory power, a number of criticisms of the theory and its assumptions have been consistently raised (Kozłowski and Konarzewski 2004; Kozłowski and Konarzewski 2005; Price et al. 2012). One primary assumption is that the sizes of terminal units in vascular networks (e.g. capillaries in circulatory systems or terminal veins in plant leaves) are invariant. The problems with this assumption have been thoroughly detailed for animal circulatory systems with the allometry of genome size and cell size emerging as a critical factor influencing how body size scales with metabolism (Kozłowski et al. 2003). Furthermore, the allometry of genome size and cell size (Figure 1) and the effects of genome size on maximum metabolic rate (Figure 3) presented here suggest that this assumption is violated in plants, as well. Modifications to the original model that relax some of its assumptions have improved model predictions for plants, particularly by allowing for variation in the packing of xylem conduits (Savage et al. 2010). However, the nucleotypic effects of genome size have yet to be incorporated, although they may further improve models and help to clarify the constraints and major innovations driving botanical form, function, and diversity.

498   Table 1. Brief summary of traits shown previously to correlate with genome size.

Sizes	Reference
Pollen volume	Bennett 1972
Cell mass	Martin 1966
Epidermal cell size	Beaulieu et al. 2008; Knight and Beaulieu 2008
Nuclear volume	Van't Hof and Sparrow 1963; Baetcke et al. 1967
Nuclear dry mass	Bennett et al. 1983; White and Rees 1987
Seed mass	Grotkopp et al. 2004; Knight and Beaulieu 2008
Xylem vessel diameter	Maherali et al. 2009; Hao et al. 2013; De Baerdemaeker et al. 2018
Rates	
Cell division rate, meiosis, mitosis	Van't Hof and Sparrow 1963; Hof 1965; Bennett 1971
Minimum generation time	Bennett 1972
Leaf expansion rate	Grime et al. 1985
Phenology	Grime and Mowforth 1982
Frost tolerance	MacGillivray and Grime 1995

Figure legends

Figure 1. Genome size determines the minimum size of cells, and smaller genomes enable

greater variation in final cell size. Data for meristematic cells (blue triangles) were taken

from Šímová and Herben (2012), and the solid black line is the standard major axis

regression through these points. Data for mature stomatal guard cells (circles and squares)

for ferns (dark green), gymnosperms (pink), and angiosperms (light blue) were taken from

Simonin and Roddy (2018), and data for mosses and hornworts (light green) were taken

from Field et al. (2015) and Renzaglia et al. (2017). The two regression lines through

mature guard cell data represent quantile regressions through the 10th (lower) and 90th

(upper) quantiles of the data.

Figure 2. Variation in the sizes and packing densities of stomatal guard cells and leaf veins

with variation in genome size among *Rhododendron* sect. *Schistanthe* species. The dashed

lines represent regressions through the 90th (upper) and 10th (lower) quantiles. Genome

size limits the lower limit of cell size and the upper limit of cell packing densities, and there

is greater variation in anatomical traits among species with smaller genomes.

Figure 3. Genome size limits the maximum rate of photosynthesis across C3 terrestrial

plants. (a) Untransformed relationship and (b) log-transformed relationship. Dashed black

lines are regressions through the upper 90th quantile. The solid black line is the standard

major axis regression (R^2 = 0.23, P < 0.0001), and the solid red line is the phylogenetically

corrected RMA regression.

Figure 4. Genome size limits the maximum rate of stomatal response (i.e. how fast stomata

can open or close). Data taken from McAusland et al. (2016) and Kew Plant DNA C-values

Database.

Figure 5. Relationship between genome size and ploidy for angiosperms. Each line

represents the SMA regression within a genus. At narrow taxonomic scales, ploidy and

genome size are correlated, but at broad taxonomic scales (i.e. among all angiosperms),

there is no relationship between genome size and ploidy due to rediploidization.

Figure 6. Distributions of genome size for C3 and CAM species show CAM lineages have

significantly larger genomes than C3 lineages. Lineages identified as CAM likely include

many C3 species; see text for details on identification of photosynthetic pathways.

References

Baetcke K, A Sparrow, C Nauman, SS Schwemmer 1967 The relationship of dna content to
nuclear and chromosome volumes and to radiosensitivity (ld50). Proceedings of the
National Academy of Sciences of the United States of America 58:533.

Bailey IW, WW Tupper 1918 Size variation in tracheary cells: I. a comparison between the
secondary xylems of vascular cryptograms, gymnosperms and angiosperms.
Proceedings of the American Academy of Arts and Sciences 54:149–204.

Baker HG 1974 The evolution of weeds. Annual review of ecology and systematics 5:1–24.

Barker MS, N Arrigo, AE Baniaga, Z Li, DA Levin 2016 On the relative abundance of
autopolyploids and allopolyploids. New Phytologist 210:391–398.

Barker MS, BC Husband, JC Pires 2016 Spreading winge and flying high: The evolutionary
importance of polyploidy after a century of study. American Journal of Botany
103:1139–1145.

Bazzaz FA, NR Chiariello, P Coley, LF Pitelka 1987 Allocating resources to reproduction and
defense. BioScience 37:58–67.

Beaulieu JM, IJ Leitch, CA Knight 2007 Genome size evolution in relation to leaf strategy
and metabolic rates revisited. Annals of Botany 99:495–505.

Beaulieu JM, IJ Leitch, S Patel, A Pendharkar, CA Knight 2008 Genome size is a strong
predictor of cell size and stomatal density in angiosperms. New Phytologist
179:975–986.

Beest M te, JJ Le Roux, DM Richardson, AK Brysting, J Suda, M Kubešová, P Pyšek 2011 The
more the better? The role of polyploidy in facilitating plant invasions. Annals of
botany 109:19–45.

Bennett M, J Heslop-Harrison, J Smith, J Ward 1983 DNA density in mitotic and meiotic
metaphase chromosomes of plants and animals. Journal of Cell Science 63:173–179.

Bennett MD 1971 The duration of meiosis. Proceedings of the Royal Society B 178:277–
299.

Bennett MD 1972 Nuclear dna content and minimum generation time in herbaceous plants.
Proceedings of the Royal Society of London Series B Biological Sciences 181:109–
135.

Bennett MD, IJ Leitch 2005 The evolution of the genome. In: Gregory TR, editor. Elsevier.
pp. 89–162.

——— Plant dna c-values database (release 6.0). [accessed 2012].
http://data.kew.org/cvalues/

Bennett MD, IJ Leitch, L Hanson 1998 DNA amounts in two samples of angiosperm weeds.
Annals of Botany 82:121–134.

Brenes-Arguedas T, AB Roddy, TA Kursar 2013 Plant traits in relation to the performance
and distribution of woody species in wet and dry tropical forest types in panama.
Functional Ecology 27:392–402.

Brittingham HA, MH Koski, T-L Ashman 2018 Higher ploidy is associated with reduced
range breadth in the potentilleae tribe. American Journal of Botany 105:700–710.

Brochmann C, A Brysting, I Alsos, L Borgen, H Grundt, A-C Scheen, R Elven 2004 Polyploidy
in arctic plants. Biological Journal of the Linnean Society 82:521–536.

Brown JH, JF Gillooly, AP Allen, VM Savage, GB West 2004 Toward a metabolic theory of
ecology. Ecology 85:1771–1789.

Cavalier-Smith T 1978 Nuclear volume control by nucleoskeletal dna, selection for cell
volume and cell growth rate, and the solution of the dna c-value paradox. Journal of
Cell Science 34:247–278.

——— 1982 Sekeletal dna and the evolution of genome size. Annual Review of Biophysics
and Bioengineering 11:273–302.

Chamberlain S 2016 Brranching: Fetch 'phylogenies' from many sources.

Clark J, O Hidalgo, J Pellicer, H Liu, J Marquardt, Y Robert, M Christenhusz, S Zhang, M Gibby,
IJ Leitch, et al. 2016 Genome evolution of ferns: Evidence for relative stasis of
genome size across the fern phylogeny. New Phytologist 210:1072–1082.

Clark JW, PC Donoghue 2018 Whole-genome duplication and plant macroevolution. Trends
in plant science.

De Baerdemaeker NJF, N Hias, J Van den Bulcke, W Keulemans, K Steppe 2018 The effect of
polyploidization on tree hydraulic functioning. American Journal of Botany
105:161–171.

Dodsworth S, M Chase, AR Leitch 2016 Is post-polyploidization diploidization the key to the
evolutionary success of the angiosperms? Botanical Journal of the Linnean Society
180:1–5.

Dolezel J, J Greilhuber, J Suda 2007 Estimation of nuclear dna content in plants using flow
cytometry. Nature Protocols 2:2233–44.

Doyle JJ, JE Coate 2019 Polyploidy, the nucleotype, and novelty: The impact of genome
doubling on the biology of the cell. International Journal of Plant Sciences 180:1–52.

Drake PL, RH Froend, PJ Franks 2013 Smaller, faster stomata: Scaling of stomatal size, rate
of response, and stomatal conductance. Journal of Experimental Botany 64:495–505.

Earles JM, TN Buckley, CR Brodersen, FA Busch, FJ Cano, B Choat, JR Evans, GD Farquhar, R
Harwood, M Huynh, et al. 2019 Embracing 3D complexity in leaf carbon–water
exchange. Trends in plant science 24:15–24.

Enquist BJ, JH Brown, GB West 1998 Allometric scaling of plant energetics and population
density. Nature 395:163–165.

Field KJ, JG Duckett, DD Cameron, S Pressel 2015 Stomatal density and aperture in non-
vascular land plants are non-responsive to above-ambient atmospheric co2
concentrations. Annals of botany 115:915–922.

Franks PJ, DJ Beerling 2009 Maximum leaf conductance driven by co2 effects on stomatal
size and density over geologic time. Proceedings of the National Academy of
Sciences 106:10343–10347.

Gallagher RV, RP Randall, MR Leishman 2014 Trait differences between naturalized and
invasive plant species independent of residence time and phylogeny. Conservation
Biology 29:360–369.

Goetsch LA, LA Craven, BD Hall 2011 Major speciation accompanied the dispersal of vireya
rhododendrons (ericaceae, rhododendron sect. schistanthe) through the malayan
archipelago: Evidence from nuclear gene sequences. Taxon 60:1015–1028.

Grime J 1998 Plant classification for ecological purposes: Is there a role for genome size?
Annals of Botany 82:117–120.

Grime J, M Mowforth 1982 Variation in genome size—an ecological interpretation. Nature
299:151.

Grime J, J Shacklock, S Band 1985 Nuclear dna contents, shoot phenology and species co-
existence in a limestone grassland community. New Phytologist 100:435–445.

Grotkopp E, M Rejmánek, M Sanderson, TL Rost 2004 Evolution of genome size in pines
(pinus) and its life-history correlates: Supertree analyses. Evolution 58:1705–1729.

Hao G-Y, ME Lucero, SC Sanderson, EH Zacharias, NM Holbrook 2013 Polyploidy enhances
the occupation of heterogeneous environments through hydraulic related trade-offs
inAtriplex canescens(Chenopodiaceae). New Phytologist 197:970–978.

Hetherington A, F Woodward 2003 The role of stomata in sensing and driving
environmental change. Nature 424:901–908.

Hijmans RJ, T Gavrilenko, S Stephenson, J Bamberg, A Salas, DM Spooner 2007 Geographical
and environmental range expansion through polyploidy in wild potatoes (solanum
section petota). Global Ecology and Biogeography 16:485–495.

Hof JV 1965 Relationships between mitotic cycle duration, s period duration and the
average rate of dna synthesis in the root meristem cells of several plants.
Experimental Cell Research 39:48–58.

Jiao Y, NJ Wickett, S Ayyampalayam, AS Chanderbali, L Landherr, PE Ralph, LP Tomsho, Y
Hu, H Liang, PS Soltis, et al. 2011 Ancestral polyploidy in seed plants and
angiosperms. Nature 473:97–100.

Johnson AL, R Govindarajulu, T-L Ashman 2014 Bioclimatic evaluation of geographical
range in fragaria (rosaceae): Consequences of variation in breeding system, ploidy
and species age. Botanical journal of the Linnean Society 176:99–114.

Kleunen M van, W Dawson, N Maurel 2015 Characteristics of successful alien plants.
Molecular Ecology 24:1954–1968.

Kleunen M van, E Weber, M Fischer 2010 A meta-analysis of trait differences between
invasive and non-invasive plant species. Ecology Letters 13:235–245.

Knight CA, JM Beaulieu 2008 Genome size scaling through phenotype space. Annals of
Botany 101:759–766.

Knight CA, NA Molinari, DA Petrov 2005 The large genome constraint hypothesis:
Evolution, ecology and phenotype. Annals of Botany 95:177–190.

Kozłowski J, M Konarzewski 2004 Is west, brown and enquist's model of allometric scaling
mathematically correct and biologically relevant? Functional Ecology 18:283–289.

Kozłowski J, M Konarzewski 2005 West, brown and enquist's model of allometric scaling
again: The same questions remain. Functional Ecology 19:739–743.

Kozłowski J, M Konarzewski, AT Gawelczyk 2003 Cell size as a link between noncoding dna
and metabolic rate scaling. Proceedings of the National Academy of Sciences
100:14080–14085.

Kubešová M, L Moravcova, J Suda, V Jarošík, P Pyšek, others 2010 Naturalized plants have
smaller genomes than their non-invading relatives: A flow cytometric analysis of the
czech alien flora. Preslia 82:81–96.

Kuester A, JK Conner, T Culley, RS Baucom 2014 How weeds emerge: A taxonomic and
trait-based examination using united states data. New Phytologist 202:1055–1068.

Landis JB, DE Soltis, Z Li, HE Marx, MS Barker, DC Tank, PS Soltis 2018 Impact of whole-
genome duplication events on diversification rates in angiosperms. American
Journal of Botany 105:348–363.

Lawson T, MR Blatt 2014 Stomatal size, speed, and responsiveness impact on
photosynthesis and water use efficiency. Plant Physiology 164:1556–1570.

Lawson T, S Vialet-Chabrand 2019 Speedy stomata, photosynthesis and plant water use
efficiency. New Phytologist 221:93–98.

Lehmeier C, R Pajor, MR Lundgren, A Mathers, J Sloan, M Bauch, A Mitchell, C Bellasio, A
Green, D Bouyer, et al. 2017 Cell density and airspace patterning in the leaf can be
manipulated to increase leaf photosynthetic capacity. The Plant Journal 92:981–994.

Leishman MR, T Haslehurst, A Ares, Z Baruch 2007 Leaf trait relationships of native and
invasive plants: Community- and global-scale comparisons. New Phytologist
176:635–643.

Leitch I, M Bennett 2004 Genome downsizing in polyploid plants. Biological journal of the
Linnean Society 82:651–663.

Lomax BH, J Hilton, RM Bateman, GR Upchurch, JA Lake, IJ Leitch, A Cromwell, CA Knight
2013 Reconstructing relative genome size of vascular plants through geological
time. New Phytologist 201:636–644.

MacGillivray CW, J Grime 1995 Genome size predicts frost resistance in british herbaceous
plants: Implications for rates of vegetation response to global warming. Functional
Ecology 9:320–325.

688 Magallón S, S Gómez-Acevedo, LL Sánchez-Reyes, T Hernández-Hernández 2015 A
689   metacalibrated time-tree documents the early rise of flowering plant phylogenetic
690   diversity. New Phytologist 207:437–453.

691 Maherali H, AE Walden, BC Husband 2009 Genome duplication and the evolution of
692   physiological responses to water stress. New Phytologist 184:721–731.

693 Martin PG 1966 Variation in the amounts of nucleic acids in the cells of different species of
694   higher plants. Exp Cell Res 44:84–94.

695 Martin SL, BC Husband 2009 Influence of phylogeny and ploidy on species ranges of north
696   american angiosperms. Journal of Ecology 97:913–922.

697 McAusland L, S Vialet-Chabrand, P Davey, NR Baker, O Brendel, T Lawson 2016 Effects of
698   kinetics of light-induced stomatal responses on photosynthesis and water-use
699   efficiency. New Phytologist 211:1209–1220.

700 Melaragno JE, B Mehrotra, AW Coleman 1993 Relationship between endopolyploidy and
701   cell size in epidermal tissue of arabidopsis. The Plant Cell 5:1661–1668.

702 Mencuccini M 2003 The ecological significance of long-distance water transport: Short-
703   term regulation, long-term acclimation and the hydraulic costs of stature across
704   plant life forms. Plant, Cell & Environment 26:163–182.

705 Ordonez A, IJ Wright, H Olff 2010 Functional differences between native and alien species:
706   A global-scale comparison. Functional Ecology 24:1353–1361.

707 Otto SP, J Whitton 2000 Polyploid incidence and evolution. Annual review of genetics
708   34:401–437.

709 Pandit MK, SM White, MJO Pocock 2014 The contrasting effects of genome size,
710   chromosome number and ploidy level on plant invasiveness: A global analysis. New
711   Phytologist 203:697–703.

712 Pearcy RW 1990 Sunflecks and photosynthesis in plant canopies. Annual Review of Plant
713   Biology 41:421–453.

Petit C, JD Thompson 1999 Species diversity and ecological range in relation to ploidy level
in the flora of the pyrenees. Evolutionary Ecology 13:45–65.

Poorter H, C Remkes 1990 Leaf area ratio and net assimilation rate of 24 wild species
differing in relative growth rate. Oecologia 83:553–559.

Poorter H, C Remkes, H Lambers 1990 Carbon and nitrogen economy of 24 wild species
differing in relative growth rate. Plant physiology 94:621–627.

Prentis PJ, JR Wilson, EE Dormontt, DM Richardson, AJ Lowe 2008 Adaptive evolution in
invasive species. Trends in Plant Science 13:288–294.

Price CA, JF Gilooly, AP Allen, JS Weitz, KJ Niklas 2010 The metabolic theory of ecology:
Prospects and challenges for plant biology. New Phytologist 188:696–710.

Price CA, JS Weitz, VM Savage, J Stegen, A Clarke, DA Coomes, PS Dodds, RS Etienne, AJ
Kerkhoff, K McCulloh, et al. 2012 Testing the metabolic theory of ecology. Chave J,
editor. Ecology Letters 15:1465–1474.

Pysek P, H Skalova, J Cuda, W-Y Guo, J Suda, J Dolezal, O Kauzal, C Lambertini, M Lucanova,
T Mandakova, et al. 2018 Small genome separates native and invasive populations in
an ecologically important cosmopolitan grass. Ecology 99:79–90.

Reich P, M Walters, D Ellsworth 1992 Leaf life-span in relation to leaf, plant, and stand
characteristics among diverse ecosystems. Ecological monographs 62:365–392.

Rejmánek M, DM Richardson 1996 What attributes make some plant species more
invasive? Ecology 77:1655–1661.

Ren R, H Wang, C Guo, N Zhang, L Zeng, Y Chen, H Ma, J Qi 2018 Wide-spread whole genome
duplications contribute to genome complexity and species diversity in angiosperms.
Molecular Plant.

Ren T, SM Weraduwage, TD Sharkey 2019 Prospects for enhancing leaf photosynthetic
capacity by manipulating mesophyll cell morphology. Journal of Experimental
Botany 70:1153–1165.

Renzaglia KS, JC Villarreal, BT Piatkowski, JR Lucas, A Merced 2017 Hornwort stomata:
Architecture and fate shared with 400-million-year-old fossil plants without leaves.
Plant Physiology 174:788–797.

Rice A, P Šmarda, M Novosolov, M Drori, L Glick, N Sabath, S Meiri, J Belmaker, I Mayrose
2019 The global biogeography of polyploid plants. Nature Ecology & Evolution
3:265–273.

Richardson DM, P Pyšek, JT Carlton 2011 A compendium of essential concepts and
terminology in invasion ecology. In: Fifty years of invasion ecology: The legacy of
charles elton. Wiley-Blackwell Oxford, UK. pp. 409–420.

Sachs J von 1893 Physiologische notizen. vi. Über einige beziehungen der specifischen
grösse der pflanzen zu ihren organisation. Flora 77:49–81.

Savage VM, LP Bentley, BJ Enquist, JS Sperry, DD Smith, PB Reich, EI von Allmen 2010
Hydraulic trade-offs and space filling enable better predictions of vascular structure
and function in plants. Proceedings of the National Academy of Sciences 107:22722–
22727.

Simonin KA, AB Roddy 2018 Genome downsizing, physiological novelty, and the global
dominance of flowering plants. PLoS Biology 16:e2003706.

Smith J, K Winter 1996 Crassulacean acid metabolism. In: Winter K; Smith J, editors.
Springer Verlag: Berlin. pp. 427–436.

Soltis DE, PS Soltis, JA Tate 2003 Advances in the study of polyploidy since plant speciation.
New Phytologist 161:173–191.

Soltis DE, CJ Visger, PS Soltis 2014 The polyploidy revolution then...and now: Stebbins
revisited. American Journal of Botany 101:1057–1078.

Soltis PS, DB Marchant, Y Van de Peer, DE Soltis 2015 Polyploidy and genome evolution in
plants. Current opinion in genetics & development 35:119–125.

Stebbins GL 1940 The significance of polyploidy in plant evolution. The American
Naturalist 74:54–66.

Stebbins GL 1985 Polyploidy, hybridization, and the invasion of new habitats. Annals of the
Missouri Botanical Garden:824–832.

Stebbins Jr GL 1947 Types of polyploids: Their classification and significance. In: Advances
in genetics. Vol. 1. Elsevier. pp. 403–429.

Suda J, LA Meyerson, IJ Leitch, P Pysek 2015 The hidden side of plant invasions: The role of
genome size. New Phytologist 205:994–1007.

Šímová I, T Herben 2012 Geometrical constraints in the scaling relationships between
genome size, cell size and cell cycle length in herbaceous plants. Proceedings of the
Royal Society B: Biological Sciences 279:867–875.

Tholen D, C Boom, X-G Zhu 2012 Opinion: Prospects for improving photosynthesis by
altering leaf anatomy. Plant Science 197:92–101.

Van't Hof J, AH Sparrow 1963 A relationship between dna content, nuclear volume, and
minimum mitotic cycle time. Proceedings of the National Academy of Sciences of the
United States of America 49:897.

Violle C, M-L Navas, D Vile, E Kazakou, C Fortunel, I Hummel, E Garnier 2007 Let the
concept of trait be functional! Oikos 116:882–892.

Webb CO, DD Ackerly, SW Kembel 2008 Phylocom: Software for the analysis of
phylogenetic community structure and trait evolution. Bioinformatics 24:2098–
2100.

Wei N, R Cronn, A Liston, T Ashman 2018 Functional trait divergence and trait plasticity
confer polyploid advantage in heterogeneous environments. New Phytologist
221:2286–2297.

Wendel JF 2015 The wondrous cycles of polyploidy in plants. American Journal of Botany
102:1753–1756.

West GB, JH Brown, BJ Enquist 1997 A general model for the origin of allometric scaling
laws in biology. Science 276:122–126.

——— 1999a A general model for the structure and allometry of plant vascular systems.
Nature 400:664–667.

——— 1999b The fourth dimension of life: Fractal geometry and allometric scaling of
organisms. Science 284:1677–1679.

West GB, WH Woodruff, JH Brown 2002 Allometric scaling of metabolic rate from
molecules and mitochondria to cells and mammals. Proceedings of the National
Academy of Sciences 99:2473–2478.

White J, H Rees 1987 Chromosome weights and measures in petunia. Heredity 58:139.

Wright IJ, PB Reich, M Westoby, DD Ackerly, Z Baruch, F Bongers, J Cavender-Bares, T
Chapin, JH Cornelissen, M Diemer, et al. 2004 The worldwide leaf economics
spectrum. Nature 428:821.