Epigenetic bookmarking of H$_2$S exposure in *Caenorhabditis elegans*

Emily M. Fawcett[1,2], Jenna K. Johnson[3], Chris Braden[1], Ellie Garcia[1], and Dana L. Miller[1,2]*

[1]University of Washington School of Medicine, Department of Biochemistry and [2]Graduate Program in Molecular and Cellular Biology, Seattle, WA, and [3]Department of Biology, Luther College, Decorah, IA

*Corresponding author: DLM
phone: 206-685-5025
email: dlm16@uw.edu

short title: Epigenetic bookmarking in *C. elegans*

Abstract

Physiological memories of environmental stress can serve to predict future environmental changes, allowing the organism to initiate protective mechanisms and survive. Although physiological memories, or bookmarks, of environmental stress have been described in a wide range of organisms, from bacteria to plants to humans, the mechanism by which these memories persist in the absence of stress is still largely unknown. We have discovered that *C. elegans* transiently exposed to low doses of hydrogen sulfide (H_2S) survive subsequent exposure to otherwise lethal H_2S concentrations and induce H_2S-responsive transcripts more robustly than naïve controls. H_2S bookmarking can occur at any developmental stage and persists through cell divisions and development but is erased by fasting. We show that maintenance of the H_2S bookmark requires the SET-2 histone methyltransferase and the CoREST-like demethylase complex. We propose a model in which exposure to low doses of H_2S generates a long-lasting, epigenetic memory by modulating H3K4me2 modifications at specific promoters. Understanding the fundamental aspects of H_2S bookmarking in this tractable system can provide mechanistic insight into how environmental exposures are translated into the epigenetic landscape in animals.

Introduction

One strategy animals use to maximize survival in a changing environment is to anticipate changes based on prior experiences. In some situations, physiological adaptations to environmental changes persist long after the initial stimulus has been removed. These sustained cellular memories of previous life experiences are known as environmental bookmarks (reviewed in (Kinoshita and Seki, 2014). Perhaps the best-studied example is in the budding yeast *Saccharomyces cerevisiae*, where cells which have previously been grown in media containing galactose respond to subsequent galactose exposure faster than naïve yeast (Kundu et al., 2007). Similarly, human retinal cells in culture retain markers of high glucose stress long after glucose levels have normalized (Ihnat et al., 2007).

H_2S is common in the environment: in addition to natural sources of H_2S, such as volcanic gasses and anaerobic bacteria in saline marshes, H_2S is produced and emitted from large livestock farms, power plants, oil and natural gas refineries and pipelines, and during the production of glue, plastics, and asphalt (Beauchamp et al., 1984). The effects of long-term exposure to low H_2S are poorly understood, but may contribute to neurological, respiratory, and cardiovascular dysfunction in humans (Kilburn and Warshaw, 1995; Richardson, 1995; Bates et al., 2002). However, exogenous H_2S can also have beneficial effects in mammals, improving outcome in mammalian models of ischemia/reperfusion injury (Bos et al., 2015; Wu et al., 2015; Sen, 2017) and mediating at least some of the beneficial effects of dietary restriction (Hine et al., 2015).

C. elegans is an excellent model to understand the molecular and genetic pathways that mediate the responses to H_2S and thereby the beneficial and/or detrimental physiological effects of H_2S in animals. Just as exposure to high concentrations of H_2S is lethal in mammals, *C. elegans* die when exposed to high concentrations of H_2S (Budde and Roth, 2010). However, *C. elegans* grown in low H_2S are long-lived and thermotolerant (Miller and Roth, 2007) and are better able to maintain proteostasis in hypoxia (Fawcett et al., 2015), suggesting that at least some protective effects of H_2S observed in mammals may be recapitulated in the *C. elegans* model.

We have discovered a novel environmental bookmark in *C. elegans* formed by exposure to hydrogen sulfide (H_2S). We show that transient exposure to low H_2S leads to formation of a bookmark, which enables animals to survive subsequent exposure to otherwise lethal concentrations of H_2S. This bookmark is quite stable, in that it can persist throughout development and far into adulthood. However, we have found that it is possible to reverse the bookmark by short periods of fasting. We further demonstrate that animals that have acquired the H_2S bookmark induce the expression of some H_2S-induced genes more robustly than naïve controls. These data suggest an epigenetic modification at the promoters of these bookmarked genes. Indeed, we demonstrate that mutations in the conserved histone H3 lysine 4 (H3K4) methyltransferase SET-2 and the H3K4me2 demethylase CoREST-like complex abrogate the ability to maintain the persistent effects of H_2S exposure. Our results suggest that exposure to H_2S results in changes in the methylation status of H3K4 at a subset of H_2S-responsive promoters, which facilitates a robust transcriptional response that allows animals to survive subsequent H_2S exposure.

Results

62   *Exposure to low H_2S forms an environmental bookmark.*

       C. elegans grown in low H_2S (50 ppm H_2S in otherwise normal room air) have

64   increased lifespan, increased thermotolerance, and improved maintenance of proteostasis

(Miller and Roth, 2007; Fawcett et al., 2015). These animals raised in low H_2S are also

66   acclimatized, as they can survive when transferred to high H_2S (concentrations higher than

150 ppm), a dose that is lethal to naïve controls (Budde and Roth, 2010). We repeated

68   these experiments, confirming that animals grown in low H_2S survived when transferred to

high sulfide (Fig 1A). Although continuous exposure throughout development is required

70   for H_2S-induced thermotolerance and lifespan extension, shorter exposure to H_2S is

sufficient to protect against hypoxia-induced disruption of proteostasis (Miller and Roth,

72   2007; Fawcett et al., 2015). We therefore asked if shorter exposure to low H_2S was

sufficient to induce acclimatization. We exposed fourth-stage larvae (L4) *C. elegans* to low

74   H_2S for only 6 h and then immediately challenged animals with exposure to high H_2S (150

ppm H_2S in room air) overnight. Although all naïve animals exposed to high H_2S died, 100%

76   of animals that had a short acclimation to low H_2S survived the subsequent exposure to

high H_2S (Fig 1A). This suggests that the physiological changes that occur in low H_2S are

78   relatively rapid and acclimatization does not require continuous growth in low H_2S.

       We next asked if animals that were acclimated to low H_2S could form a persistent,

80   physiological memory of the exposure to H_2S. We exposed first-stage larvae (L1) to low H_2S

for 6h, then returned the animals to room air. When the animals reached L4 we challenged

82   them with exposure to high H_2S overnight. We found that all of the animals survived this

treatment (Fig 1A). Thus, the effects of exposure to low H_2S do not require direct transfer

84 into high H_2S, but instead can be maintained even in the absence of continued exposure to

H_2S. This suggests that there is a process that occurs in addition to acclimatization, which

86 can be achieved simply by activation of HIF-1 transcriptional activity. We refer to this long-

lasting physiological memory of H_2S exposure as H_2S bookmarking to distinguish it from

88 acclimation.

The exposure to low H_2S required to generate the H_2S bookmark is time- and dose-

90 dependent. We initially chose 50 ppm H_2S for the low H_2S exposure based on prior work

demonstrating that this concentration of H_2S causes clear physiological effects, such as

92 increased lifespan and resistance to the hypoxia-induced disturbance of proteostasis

(Miller and Roth, 2007, #51252; Fawcett et al., 2015). However, we found that animals

94 exposed to only 20 ppm H_2S developed the H_2S bookmark and survived subsequent

exposure to high H_2S, though a longer exposure to the low H_2S environment was required

96 than at 50 ppm H_2S (Fig 1B). Conversely, the bookmark was formed more rapidly at higher

concentration of H_2S, as even a 1h exposure to 150 ppm H_2S was sufficient for survival of

98 subsequent exposure. These data indicate that the bookmarking process can be quite rapid,

but that the formation of the bookmark requires a threshold H_2S exposure.

100 Stabilization of the HIF-1 transcription factor has been proposed to underlie the

acclimatization to H_2S (Budde and Roth, 2010). HIF-1 accumulates in animals exposed to

102 H_2S, and constitutive activation of HIF-1, from mutation of either *egl-9* or *vhl-1*, is sufficient

for *C. elegans* to survive exposure to high H_2S (Budde and Roth, 2010). We could not test

104 directly whether *hif-1* is required for acclimatization or bookmarking, because *hif-1* mutant

animals die when exposed to low H_2S (Budde and Roth, 2010). However, we reasoned that

106   if activation of HIF-1 was sufficient for acclimatization then animals exposed to low O_2

(hypoxia) would also survive if transferred to high H_2S. Consistent with this hypothesis, we

108   observed increased survival of animals exposed to high sulfide when transferred directly

from 5000 ppm O_2, a condition where HIF-1 is activated (Fig 1C; Jiang et al., 2001, #24680).

110   This result supports the hypothesis that activation of HIF-1 is sufficient for acclimation.

However, the penetrance of survival was never as high as when animals were transferred

112   from low H_2S. This difference in penetrance could reflect the fact that although HIF-1

induces gene expression in both hypoxia and H_2S, different transcripts accumulate in each

114   condition (Miller et al., 2011). Alternatively, it could be that other factors that contribute to

the process of H_2S bookmarking are not engaged by exposure to hypoxia. We favor this

116   model, because the acclimation induced by exposure to hypoxia is rapidly lost upon return

to room air (Fig 1C). Thus, although the animals had acclimated they had not formed a

118   persistent physiological memory consistent with H_2S bookmarking.

We considered the possibility that the exposure to hypoxia activated stress

120   response pathways that were cross-protective in H_2S. Many well-studied stress response

pathways are activated by and protective against multiple environmental stresses. For

122   example, activation of the insulin-like signaling pathway leads to increased resistance to

thermal, oxidative, and nutritional stress (McColl et al., 2010). Although exposure to H_2S

124   does not activate common stress-response pathways in *C. elegans*, including insulin-like

signaling, TOR, and p53 signaling (Miller and Roth, 2009; Miller et al., 2011), we considered

126   the possibility that protection by H_2S bookmarking may be a result of the activation of such

cross protective stress response pathways. We first tested whether inducing the heat shock

128   response would protect against subsequent H_2S exposure, as the heat-shock response is

cross protective against several other stresses, including salt stress and oxidative stress,

130   and animals grown in H_2S are resistant to thermal stress (Völker et al., 1992; Miller and

Roth, 2007; Verghese et al., 2012). We found that subjecting animals to a non-lethal heat

132   shock did not protect against subsequent exposure to high H_2S (Supplemental Fig 1A).

Furthermore, animals with mutations in candidate cross-protective stress response

134   signaling pathways, including those involved in the response to heat shock, nutrient

deprivation, hypoxic stress, and osmotic stress did not have a defect in the formation or

136   maintenance of H_2S bookmarking (Supplemental Fig 1B). Taken together, these results

suggest that H_2S bookmarking is a specific response to H_2S.

H_2S bookmarking is persistent, but reversible.

140   During development, there are sometimes critical windows during which animals

are most sensitive to certain environmental stimuli. For example, in *C. elegans*, the decision

142   to enter dauer is determined by conditions during the first larval stage (Golden and Riddle,

1984). To determine if there was a specific developmental window in which H_2S

144   bookmarks could be formed, we exposed synchronized cohorts of *C. elegans* at different

developmental stages to low H_2S, and then tested their ability to survive exposure to high

146   H_2S after 48 h in room air. All animals survived subsequent exposure to high H_2S,

regardless of developmental stage at time of adaptation (Fig 2A). We conclude that H_2S

148   bookmarking can occur at any point in the lifetime of the animal.

We next asked how long the H_2S bookmark could persist. To measure this, we

150   acclimated embryos to low H_2S for 8 h, then let them grow in room air for increasing time

before exposure to high H_2S. We found that all animals exposed to H_2S as embryos survived

152   exposure to high H_2S as much as 72h later, when the cohort were reproductive adults (Fig

2B). However, after 72h the survival of bookmarked animals began to decline and by 96h

154   after the initial exposure most animals died when exposed to high H_2S. The exact duration

that the bookmark is maintained is somewhat dose-dependent, in that the decline began

156   sooner if the acclimation period was shortened. These data indicate that the H_2S bookmark

can be maintained throughout embryonic and postembryonic development, but that it

158   eventually spontaneously lost.

The fact that the bookmark spontaneously reversed as the animals aged led us to

160   investigate the possibility that the rate of spontaneous reversal changed with age. To test

this possibility, we exposed synchronized, aging cohorts of animals to low H_2S for 6h and

162   then let the animals age in room air for another 48h before challenging the animals with

exposure to high H_2S. We found that animals were able to form and maintain the H_2S

164   bookmark through day 3 of adulthood. However, starting at day 5 we observed a

progressive decrease in the ability to survive the subsequent exposure to high H_2S (Fig 2C).

166   This result suggests that the H_2S bookmark is formed or maintained less effectively in older

animals. We did not observe a similar decline in acclimatization, as all animals survived

168   when we transferred old animals to high H_2S immediately after exposure to low H_2S.

Together these data suggest that H_2S bookmarking – but not acclimatization – is less

170   efficient with increasing age.

H₂S bookmarking is reversed by fasting

172   While characterizing the H₂S bookmarking phenotype, we noticed that animals on

plates in which the food source had been depleted no longer displayed robust survival in

174   high H₂S. We hypothesized that H₂S bookmarking may be modulated by food availability.

To test this possibility, we adapted embryos to low H₂S for 12 hours on plates seeded with

176   OP50 *E. coli* bacteria. We then removed half of the animals from food for 6-12 hours as

first-stage larvae (L1), then returned these animals to normal food conditions. We

178   challenged both fed and fasted animals with high H₂S when they reached day 1 of

adulthood. The proportion of animals that survived exposure to high H₂S was significantly

180   lower in the fasted animals (Fig 3A). This result suggests that fasting prevents maintenance

of, or actively reverses, the H₂S bookmark. This led us to next determine if fasting also

182   prevented formation of the H₂S bookmark. To test this possibility, we exposed starved L1

to low H₂S for acclimation, then returned the animals to food in the absence of H₂S. In

184   contrast to animals acclimated to low H₂S on food, animals exposed to low H₂S in the

absence of food did not survive subsequent exposure to high H₂S (Fig 3B). These results

186   suggest that both formation and maintenance of the H₂S bookmark is inhibited by fasting.

When *C. elegans* embryos hatch in the absence of food, starved L1 enter a reversible

188   developmental arrest, the L1 diapause (reviewed in (Baugh, 2013)). To confirm that fasting

itself reverses H₂S bookmarking, and not the physiological changes associated with entry

190   into the L1 diapause, we fasted animals at other developmental stages and scored for H₂S

bookmarking retention. We found that fasting at later developmental stages did erase the

192   H₂S bookmark (Fig 3C). Bookmarked animals removed from food for only 6 hours as L4s

displayed a significant decrease in survival when later challenged with high H_2S. This effect

194   was more pronounced when the period of fasting increased. We conclude that fasting, or

the response to food deprivation, effectively reverses the H_2S bookmark.

196   One possibility is that activation of various stress responses, in addition to fasting,

would erase the effect of the H_2S bookmark. We therefore evaluated whether heat shock or

198   hypoxia would reduce the ability of animals with an H_2S bookmark to survive subsequent

exposure to high H_2S. We chose these two stresses because they both have global effects on

200   physiology and metabolism, as fasting does. We found that neither hypoxia nor heat shock

reversed the H_2S bookmark (Fig 3C). Bookmarked animals exposed to hypoxia (1,000 ppm

202   or 5,000 ppm O_2; room air is 210,000 ppm oxygen) or heat shock survived exposure to high

H_2S as well as controls that remained in normal growth conditions. This result supports our

204   assertion that the H_2S bookmark is distinct from other general stress responses. Moreover,

as hypoxia has dramatic effects to decrease protein translation and metabolic activity, this

206   result suggests that maintenance of the H_2S bookmark is not dependent on these processes.

208   *H_2S bookmarking facilitates transcriptional reactivation of H_2S responsive genes*

One possible mechanism of H_2S bookmarking could be that genes induced by

210   exposure to low H_2S persist, and are therefore able to promote survival in subsequent H_2S

exposure. We did not favor this hypothesis as the early transcriptional response to H_2S

212   depends entirely on *hif-1*, which is rapidly degraded in normal conditions (Wang et al.,

1995; Kallio et al., 1999; Miller et al., 2011). To verify that H_2S-induced changes in gene

214   expression were not maintained in animals after exposure to low H_2S, we first measured

expression of the SQRD-1::GFP translational reporter, which is induced by exposure to low

216   H$_2$S (Budde and Roth, 2011). We exposed SQRD-1::GFP animals to low H$_2$S for 8 hours, and

then removed them to house air conditions for 48 hours. Expression of SQRD-1::GFP was

218   significantly increased when animals were exposed to low H$_2$S, but returned to baseline

levels of expression upon return to house air (Fig 4A). To corroborate our results with

220   SQRD-1::GFP, we measured the abundance of 12 H$_2$S-induced transcripts (Miller et al.,

2011). The abundance of these transcripts in bookmarked animals 48 h after exposure to

222   low H$_2$S was not significantly different from naïve controls (Fig 4B). We conclude that

persistent changes in gene expression do not underlie the maintenance of the H$_2$S

224   bookmark.

We next considered the possibility that the H$_2$S bookmark functions to potentiate

226   H$_2$S-induced changes in gene expression upon subsequent re-exposure to H$_2$S. We

measured the expression of SQRD-1::GFP in naïve and bookmarked animals exposed to

228   high H$_2$S for 1 hour. We chose a 1-hour exposure because previous studies showed that

expression of *sqrd-1* is significantly upregulated after 1 hour in low H$_2$S (Miller et al., 2011),

230   and naïve animals survive a 1-hour exposure to high H$_2$S (Budde and Roth, 2011). We

found that the increase in SQRD-1::GFP expression was similar when naïve animals were

232   exposed to low or high H$_2$S for 1 hour. In contrast, expression of SQRD-1::GFP was

significantly higher when animals with the H$_2$S bookmark were exposed to high H$_2$S (Fig

234   4C). This suggests that the H$_2$S bookmark facilitates the expression of H$_2$S-inducible genes.

To corroborate this result, we measured the abundance of transcripts upregulated by

236   exposure to low H$_2$S (Miller et al., 2011). Eight of the 12 genes we tested were increased

more in bookmarked animals than naïve controls when exposed to high H_2S (Fig 4D). We conclude that H_2S bookmarking facilitates the transcriptional re-activation of H_2S-responsive genes, which we refer to as bookmarked genes.

To get a more comprehensive understanding of how H_2S bookmarking affected gene expression we performed RNAseq experiments to measure the abundance of transcripts in naïve and bookmarked animals exposed to high H_2S. We found 16 transcripts that were upregulated in both bookmarked and naïve animals exposed to high H_2S; of these, 12 were also identified in microarray experiments (Table 1; (Miller et al., 2011)). We also observed that 12 gene products were induced only in the naïve animals. None of these were upregulated in our previous microarray experiments. These genes could be induced specifically by exposure to high H_2S, or it could be that they were false negatives (or missing) from the previous microarray experiment. Nevertheless, these studies suggest there is much in common between the transcriptional responses to low and high H_2S.

We observed that eight transcripts were significantly more abundant in the bookmarked animals than in the naïve controls by RNA-seq (Bookmarked genes in Table 1). These results corroborated our qRT-PCR experiments (Fig 4D). In both experiments, gene products for *gst-19*, *nspe-3*, R08E5.1, *sqrd-1*, *cysl-2*, and *nit-1* were more abundant in bookmarked animals than in naïve controls. We also observed 37 transcripts that were induced by exposure to high H_2S only in the bookmarked animals (Bookmark Only in Table 1). Most of these had not been previously shown to be induced by exposure to H_2S. However, this group did include C31C9.2, which was bookmarked in our qRT-PCR experiments (Fig 4D), and *nspe-2*, which was upregulated in microarray experiments

(Miller et al., 2011) but not included in our qRT-PCR panel. It may be that some of these

260   genes are minimally induced by exposure to low H_2S, and do not meet our cut-offs for

determining significance. Together, our data suggest a model where H_2S bookmarking

262   leads to changes in transcriptional accessibility of specific H_2S-inducible genes, which

facilitates robust transcriptional reactivation upon subsequent H_2S exposure.

264   We noted that not all H_2S-responsive genes were bookmarked. In both RNAseq and

qRT-PCR experiments expression of three gene products, *dhs-8, rhy-1,* and *lgc-1*, were

266   induced the same in bookmarked animals and naïve controls exposed to high H_2S. Several

other genes in our RNAseq data were also induced equally in bookmarked and naïve

268   animals. There is no pattern we could detect that predicted whether an H_2S-responsive

gene would be bookmarked for transcriptional reactivation.

Epigenetic factors are required to maintain H_2S bookmarks

272   We hypothesize that there are at least two processes required for H_2S bookmarking:

the original formation of the bookmark, which could be related to acclimation, and

274   maintenance of the bookmark after the end of the exposure to H_2S. We were interested in

the mechanism by which the bookmark was maintained, as the rapid acquisition and

276   reversibility of the H_2S bookmark suggested that it may be mediated by an epigenetic

mechanism (Mirbahai and Chipman, 2014). Alternatively, it is possible that bookmarking

278   could depend upon the persistence of a long-lived protein or small RNA, or through another

uncharacterized mechanism. To distinguish between these possibilities, we performed a

280   candidate screen to identify molecular factors that mediate H_2S bookmarking.

We screened animals with genetic mutations in machinery involved in protein

282   turnover, RNA processing, epigenetic modifications, or factors associated with

environmental bookmarks in other systems (a list of all 250 candidates tested is included

284   as Supplemental Table 1). To identify mutations that specifically disrupted the

maintenance of the H_2S bookmark, we exposed synchronized embryos of each mutant

286   strain to low H_2S for 8 hours, returned them to house air for 48 hours, and then exposed

them to high H_2S for 24 h. We selected candidates that had reduced survival after exposure

288   to high H_2S, as compared to wild-type controls that acclimate and survive the subsequent

exposure to high H_2S. Two of the mutations that caught our attention in this screen were

290   *spr-5* and *set-2* (Fig 5A). For the remainder of this work we focus on these two genes, as

they regulate methylation of lysine 4 on histone 3 (H3K4me) and post-translational

292   modifications of histone proteins occupying promoters can influence transcriptional levels

(reviewed in Filipescu et al., 2014).

294   We were most interested in mutations that disrupted the long-term maintenance of

the H_2S bookmark, but we expected that some of the candidates identified in the primary

296   screen could have defects in initial acclimation to H_2S and/or formation of the H_2S

bookmark. To exclude that *spr-5* and *set-2* were required for acclimation, we exposed each

298   candidate strain to low H_2S for 6 hours, and immediately moved them to high H_2S. We

reasoned that if these animals were able to acclimate to H_2S that they would survive this

300   transition. Both *spr-5* and *spr-2* mutant animals survived this treatment (Fig 5B),

suggesting that acclimation was not abrogated by these mutations.

302   H3K4 can be mono-, di, and tri- methylated. In *C. elegans*, the histone

methyltransferase SET-2 is responsible for the formation of the majority of the bulk

304   H3K4me2 and H3K4me3 (Xiao et al., 2011). However, formation of H3K4me3 by SET-2

requires other components of the ASH-2 methyltransferase complex. The ASH-2 complex in

306   *C. elegans* is composed of three essential subunits: SET-2, ASH-2, and WDR-5.1 (Greer et al.,

2010). To determine if SET-2 was acting as part of the ASH-2 complex in H_2S bookmarking,

308   we tested whether *wdr-5.1(ok1417)* mutant animals were capable of H_2S bookmarking. In

contrast to *set-2(n4589)* mutant animals, we found that *wdr-5.1(ok1417)* mutant animals

310   survived exposure to high H_2S after bookmarking at levels comparable to WT controls (Fig

5A). Similarly, mutations in *rbr-2*, an H3K4me3 demethylase that antagonizes ASH-2

312   methyltransferase complex activity (Greer et al., 2010, #35493), has no effect on H_2S

bookmarking (Fig 5A). Although SET-2 is unable to mediate formation of H3K4me3

314   independently of the ASH-2 complex, generation of H3K4me2 does not require the ASH-2

complex (Xiao et al., 2011). We therefore favor the hypothesis that SET-2 contributes to

316   H_2S bookmarking through its activity to form H3K4me2-modified histones.

        SPR-5 is a component of the histone demethylase CoREST-like complex in *C. elegans*

318   (Eimer et al., 2002). The CoREST complex was first identified in mammals as a corepressor

of the REST transcription factor. Through its histone demethylase activity, the mammalian

320   CoREST complex mediates long-term gene repression that is essential for the maintenance

of cell identity (Andrés et al., 1999). In *C. elegans* the CoREST-like complex has retained its

322   histone demethylase activity and functions in the repression of the *hop-1* gene through

demethylation of H3K4me2 at the *hop-1* gene promoter. For *hop-1* gene repression, SPR-5

324   interacts with the orthologue of mammalian CoREST, SPR-1, and two large proteins with

weak similarity to the mammalian REST transcription factor, SPR-3 and SPR-4 (Eimer et al.,

326   2002). While they have no clear vertebrate homologs, SPR-3 and SPR-4 are predicted to

function in the recruitment of the CoREST-like corepressor complex to gene targets. To

328   determine if SPR-5 is functioning as a part of the CoREST-like complex in H_2S bookmarking,

we examined the effect of a loss-of-function mutation in these other known subunits of the

330   CoREST-like complex. We found that, like *spr-5(by134)* mutant animals, animals with

mutations in *spr-1*, *spr-3*, and *spr-4* all had defects in maintaining the H_2S bookmark after

332   exposure to low H_2S (Fig 5A). Taken together, these results suggest that SPR-5 functions as

a member of the CoREST-like complex to mediate H_2S bookmarking.

334   We hypothesized that the transcriptional reactivation we observed in bookmarked

animals was mediated by changes in H3K4me2 status at bookmarked promoters based on

336   the genetic requirement for *set-2* and the CoREST demethylase complex. This hypothesis

predicts that *set-2(n4589)* and *spr-5(by134)* mutant animals would not be able to

338   effectively reactivate transcription of H_2S-inducible transcripts. We tested this possibility

by measuring transcript abundance in *set-2(n4589)* and *spr-5(by134)* mutant animals

340   exposed to high H_2S. Both *set-2(n4589)* and *spr-5(by134)* mutant animals exposed to low

H_2S as embryos had defects in the transcriptional reactivation of H_2S-inducible genes (Fig

342   6A). One trivial explanation for the lack of transcriptional reactivation of H_2S-inducible

genes in the *set-2* and *spr-5* mutant animals is that they are generally unable to induce gene

344   expression in response to H_2S. We considered this unlikely, as both *set-2(n4589)* and

spr-5(by134) mutant animals survive prolonged exposure to low H_2S as well as wild-type

346   animals. Indeed, we found no difference in the initial transcriptional response to low

sulfide in either *set-2(n4589)* or *spr-5(by134)* mutant animals (Fig 6B). Moreover, both

348   *set-2(n4589)* and *spr-5(by134)* mutant animals upregulated H_2S-responsive genes as

effectively as wild-type controls when exposed to high H_2S after acclimation (Fig 6C). These

350   results indicate that neither *set-2* nor *spr-5* are required for the normal transcriptional

response or acclimatization to H_2S. Instead, our data show that these histone-modifying

352   enzymes are specifically required for the maintenance of the H_2S bookmark and associated

transcriptional reactivation of bookmarked genes.

Discussion

356   We have discovered a novel environmental bookmark to H_2S that enhances survival

when animals re-encounter H_2S in the environment. Our experiments suggest that H_2S

358   exposure leads to formation of a bookmark that has persistent effects on the epigenetic

landscape, allowing for enhanced transcriptional responses and increased survival upon

360   subsequent H_2S exposure.

        To our knowledge, our studies demonstrate the first example of reversal of a stress-

362   induced epigenetic bookmark by fasting. The molecular nature of the interaction between

fasting and H_2S bookmarking is not yet clear. In mammals, changes in diet, including

364   periods of fasting, can lead to dramatic changes in the epigenome, including the reversal of

epigenetic bookmarks (Burdge and Lillycrop, 2010). The ability to erase, or modulate,

366   epigenetic bookmarks on demand has important implications in treatment of human

disease by therapeutic intervention. Offspring of malnourished mothers can carry

368   bookmarks that can lead to diabetes or obesity if food is abundant, which can reduce

fertility (Gluckman et al., 2008). Maternal undernutrition during pregnancy, which results

370   in epigenetic modifications to offspring that are designed to protect against subsequent

famine, is associated with greater susceptibility to cancer in rats (Fernandez-Twinn et al.,

372   2007). Our discovery of epigenetic bookmarking by H_2S exposure provides a uniquely

tractable model to develop strategies to modulate epigenetic marks in animals.

374   We propose that the H_2S bookmark is related to the methylation status of histone

H3K4, based on the genetic requirement for SET-2 and SPR-5 in maintaining the H_2S

376   bookmark. As SET-2 and SPR-5/CoREST both alter methylation status of H3K4me2, one

simple model is that these two enzymes cooperate at the promoters of bookmarked

378   animals to facilitate transcriptional reactivation. However, this interpretation is

complicated by the fact that mutations in *set-2* and *spr-5* do not have entirely overlapping

380   effects. For example, while transcriptional reactivation of *gst-19* and *nspe-3* was abrogated

in both *spr-5* and *set-2* mutant animals, the bookmarking of *nhr-57, nit-1, C31C9.2,* and

382   *R08E5.1* was lost only in *set-2* animals (Fig 5C). Moreover, some bookmarked genes were

not affected in either *set-2(n4589)* or *spr-5(by134)* mutant animals. Future genome-wide

384   studies of histone methylation status and the epigenetic effects of H_2S are necessary to

understand how these two histone-modifying enzymes are coordinated.

386   Our finding that H_2S bookmarking requires histone methylation machinery is

reminiscent of other established examples of epigenetic bookmarks. In plants, multiple

388   rounds of drought stress lead to increased H3K4me3, providing resistance to subsequent

severe droughts (Ding et al., 2012). Additionally, trithorax (TrxG) proteins with histone

390   methyltransferase activity function in the epigenetic bookmarking of low winter

temperatures in plants, which alters the timing of vernalization in subsequent seasons

392   (Buzas et al., 2012; Song et al., 2012). The complexity and pleiotropic nature of

environmental bookmarks has made understanding the mechanistic underpinnings of

394   epigenetic bookmarks difficult, particularly in animals. Although not necessarily a

bookmark, *C. elegans*, that enter into the alternative dauer larval stage have decreased in

396   H3K4me3 and H4 acetylation in the adult animal, resulting in persistent changes in endo-

siRNA levels that can poise genes for activation upon subsequent encounters with

398   environmental stress (Hall et al., 2010).

The genetic factors required to maintain the H_2S bookmark in *C. elegans*, SET-2 and

400   SPR-5, are conserved through humans. Though to our knowledge H_2S has not yet been

shown to have epigenetic effects in mammals, H_2S is a commonly encountered

402   environmental toxin produced by oil refineries and paper mills that has dramatic effects on

neurological, respiratory and cardiovascular function, even at low concentrations (Kilburn

404   and Warshaw, 1995; Richardson, 1995; Bates et al., 2002). We demonstrate that H_2S

concentrations as low as 15 ppm, which is below OSHA limits for industrial exposure, can

406   lead to the establishment of an H_2S bookmark in *C. elegans*. It will be important to learn if

H_2S has similar effects in mammals, as this information could inform future toxic risk

408   assessment to improve worker safety.

410   **Materials and Methods**

C. elegans were maintained on nematode growth media (NGM) with OP50 *E. coli* at 20°C

412   (Brenner, 1974). Worm strains used in this study are listed in Supplementary Table 1.

414   *Constructing defined gaseous environments*

All compressed gas tanks were purchased from Airgas (Seattle, WA) and certified standard

416   to 2% of target gas concentration, with the balance N_2. Tanks with O_2 contained 1,000 or

5,000 ppm, and H_2S source tanks were 5,000 ppm. Gaseous environments were maintained

418   using continuous flow chambers as described in (Fawcett et al., 2012). H_2S was diluted

from a 5,000 ppm stock tank with house air as in (Miller and Roth, 2007). H_2S

420   environments were maintained in a fume hood at 20°C, with matched house air (without

H_2S) continuous flow environments.

H_2S bookmarking assay

424   Embryos were synchronized by allowing gravid adults to lay eggs for 2 h on seeded

NGM plates. First stage larvae (L1) were collected 24 h post egg-lay, L2 after 36 h, and L3s

426   after 48 hours at 20°C on seeded NGM plates. L4 animals were picked from well-fed,

logarithmically growing cultures and moved to seeded NGM plates. Staged animals were

428   immediately exposed to 50 ppm H_2S for indicated amount of time, and then removed to

house air for 48 hours. Animals were then exposed to 150 ppm H_2S overnight (~16 hours)

430   and then scored for survival. Data was reported as % animals alive after 150 ppm H_2S +/-

standard deviation. Bagged animals were censored from the experiment. For differences

432   between genotypes, p-values were calculated by one-way ANOVA using summary statistics

(mean, standard deviation, n).

434   To evaluate the effects of other conditions on bookmarks, 24 h after the adaptation

period animals were moved to hypoxia (1,000 or 5,000 ppm O_2 for 24 h) or heat-shock (37

436   C for 1 h). For fasting, animals were moved to unseeded NGM plates with 25 mg/L

carbenicillin to prevent bacterial growth. After 10 minutes, animals were moved to a new

438   unseeded NGM plate with 25 mg/L carbenicillin to further deplete the food associated with

their cuticle. Palmitic acid (10 mg mL^{-1} in ethanol) was used to form a physical barrier

440   around the edge of each plate to encourage the animals to remain on the surface of the

plate when fasted. After the indicated time period, animals were moved back to NGM plates

442   seeded with live OP50 *E. coli* in room air until 48 h post-adaptation, at which time the

animals were challenged with high H_2S.

qRT-PCR

446   Animals were grown on high-growth plates seeded with NA22 *E. coli* at 20°C. When

animals reached gravid adult, synchronized embryos were obtained by a 5-minute bleach

448   in 1:1:5 water:KOH:hypochloric acid solution. For each strain/condition, ~9,000 embryos

were plated onto a 150 mM NGM plate seeded with live OP50 *E. coli*. Animals were not

450   allowed to starve out the plate at any time during the experiment. The animals were

treated as for the bookmarking assay as above, except that the exposure to 150 ppm H_2S

452   was for one hour. Animals were harvested into 1 mL Trizol solution and immediately

frozen in liquid nitrogen. RNA was isolated from the Trizol preparation as described

454   previously (Chomczynski, 1993). cDNA was made using Invitrogen SuperScript III First

Strand Synthesis System. Primers were as in (Miller et al., 2011), primer sequences

456   available upon request. qPCR was performed using Kappa SYBR FAST qPCR Kit. PCR cycle

was as follows: 95C for 3 min, 95C for 15 sec, 55C for 15 sec x40. 4°C to hold. ΔC_t for each

458   gene product was calculated as in (Miller et al., 2011); briefly, ΔC_t values were calculated by

subtracting each measured C_t from the geometric mean of the control targets that are not

460   altered in response to H_2S (SIR-2.1, HIL-1, IRS-2, and TBA-1). ΔC_t were averaged across

experiments. Student's t-test was used to evaluate differences between ΔC_t values of

462   treated samples and untreated controls. For differences between genotypes, p-values were

calculated with a one-way ANOVA from summary statistics (mean, standard error, n).

464   Reported fold-changes were calculated as $2\text{^}-\Delta\Delta C_t$ where $\Delta\Delta C_t = \Delta C_t$(experimental

condition) - ΔC_t(control condition). Error bars on graphs represent standard error of the

466   mean, which was carried through the fold-change calculation using standard error

propagation ([www.statpages.org)](www.statpages.org).

GFP reporter quantification

470   Animals were synchronized by a 2-hour egg lay of SQRD-1::GFP animals onto seeded NGM

plates. Animals were adapted to H_2S as described above. After 48 hours, animals were

472   exposed to 150 ppm H_2S for 1 hour. Animals were then removed to house air and allowed

to recover for 1 hour to allow for folding of GFP. Roller animals, which contain the SQRD-

474   1::GFP transgene were mounted on an agar pad in a drop of 20 mM sodium azide as

anesthetic. GFP fluorescence was visualized on a Nikon 90i fluorescent microscope with the

476   GFP filter and oil-immersion 20x objective. All images were taken at the same exposure

        time and magnification. Total cell fluorescence was quantified using ImageJ software

478   (Rasband, W.S., ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA,

        http://imagej.nih.gov/ij/, 1997-2014). Student t-tests were used to compare mean cell

480   fluorescence between samples.

482   *Global mRNA measurements with RNAseq*

        Embryos were synchronized by allowing gravid adults to lay eggs for 2-6 hours on seeded

484   NGM plates. Within 24 hours the plates were exposed to 50ppm for 6 hours, and then

        removed to house air until animals reached L4, at which time they were exposed to

486   150ppm H_2S for 1 hour, flash frozen in liquid nitrogen, and immediately stored at -80C.

        RNA was isolated using Trizol, according to manufacturer's instructions, and then

488   isopropanol precipitation. Isolated RNA was sent to Novogene for library preparation using

        Illumina MiSeq for paired-end sequencing. The raw data files were uploaded to the Illumina

490   analysis cloud (BaseSpace) in order to collect raw gene counts. Differential expression of

        genes between conditions was determined using the edgeR analysis package on the raw

492   gene counts (Robinson et al., 2010).

        Acknowledgements

496   We thank our colleagues in the Miller lab that provided constructive comments to improve

        this manuscript and members of the NW Developmental Biology group that provided

498 excellent feedback on this project. We are also thankful to Dr. R. Waterston (Department of

Genome Sciences, University of Washington School of Medicine) for helpful advice and Dr.

500 C. Chatterjee (Department of Chemistry, University of Washington) for insightful

discussions. Mr. Harley Childs helped initiate the candidate screens for factors involved in

502 H_2S bookmarking. This work was supported by NIH grants R00 AG033055 and R01

ES024958 to DLM. DLM is an Ellison Medical Foundation New Scholar in Aging and has

504 been supported by the Glenn Medical Foundation. EMF was supported by NIH

Developmental Biology Predoctoral Training Grant T32 HD007183 from the NICHD. JJ was

506 an Amgen Scholar. Some strains were provided by the CGC, which is funded by NIH Office

of Research Infrastructure Programs (P40 OD010440).

References

510 Andrés ME, Burger C, Peral-Rubio MJ, Battaglioli E, Anderson ME, Grimes J, Dallman J,
      Ballas N, Mandel G. 1999. CoREST: a functional corepressor required for regulation of
512   neural-specific gene expression. *PNAS* **96**: 9873–9878.

Bates MN, Garrett N, Shoemack P. 2002. Investigation of health effects of hydrogen sulfide
514   from a geothermal source. *Arch Environ Health* **57**: 405–411.

Baugh LR. 2013. To grow or not to grow: nutritional control of development during
516   *Caenorhabditis elegans* L1 arrest. *Genetics* **194**: 539–555.

Beauchamp RO, Bus JS, Popp JA, Boreiko CJ, Andjelkovich DA. 1984. A critical review of the
518   literature on hydrogen sulfide toxicity. *Crit Rev Toxicol* **13**: 25–97.

Bos EM, van Goor H, Joles JA, Whiteman M, Leuvenink HG. 2015. Hydrogen sulfide:
520   physiological properties and therapeutic potential in ischaemia. *Br J Pharmacol* **172**:
      1479–1493.

522 Brenner S. 1974. The genetics of *Caenorhabditis elegans*. *Genetics* **77**: 71–94.

Budde MW, Roth MB. 2011. The response of *Caenorhabditis elegans* to hydrogen sulfide and
hydrogen cyanide. *Genetics* **189**: 521–532.

Budde MW, Roth MB. 2010. Hydrogen Sulfide Increases Hypoxia-inducible Factor-1 Activity Independently of von Hippel-Lindau Tumor Suppressor-1 in *C. elegans*. *Molecular Biology of the Cell* **21**: 212–217.

Burdge GC, Lillycrop KA. 2010. Nutrition, epigenetics, and developmental plasticity: implications for understanding human disease. *Annu Rev Nutr* **30**: 315–339.

Buzas DM, Tamada Y, Kurata T. 2012. FLC: a hidden polycomb response element shows up in silence. *Plant Cell Physiol* **53**: 785–793.

Chomczynski P. 1993. A reagent for the single-step simultaneous isolation of RNA, DNA and proteins from cell and tissue samples. *Biotechniques* **15**: 532–4, 536.

Ding Y, Fromm M, Avramova Z. 2012. Multiple exposures to drought 'train' transcriptional responses in Arabidopsis. *Nat Commun* **3**: 740.

Eimer S, Lakowski B, Donhauser R, Baumeister R. 2002. Loss of *spr-5* bypasses the requirement for the *C.elegans* presenilin *sel-12* by derepressing hop-1. *EMBO J* **21**: 5787–
5796.

Fawcett EM, Hoyt JM, Johnson JK, Miller DL. 2015. Hypoxia disrupts proteostasis in
540 *Caenorhabditis elegans*. *Aging Cell* **14**: 92–101.

Fawcett EM, Horsman JW, Miller DL. 2012. Creating Defined Gaseous Environments to
542 Study the Effects of Hypoxia on *C. elegans*. *J. Vis. Exp.* (65), e4088 10.3791/4088.

Fernandez-Twinn DS, Ekizoglou S, Gusterson BA, Luan J, Ozanne SE. 2007. Compensatory
544 mammary growth following protein restriction during pregnancy and lactation increases early-onset mammary tumor incidence in rats. *Carcinogenesis* **28**: 545–552.

546 Filipescu D, Müller S, Almouzni G. 2014. Histone H3 Variants and Their Chaperones During Development and Disease: Contributing to Epigenetic Control. *Annu. Rev. Cell Dev. Biol.*
548 **30**: 615–646.

Gluckman PD, Hanson MA, Cooper C, Thornburg KL. 2008. Effect of *in utero* and early-life
550 conditions on adult health and disease. *N Engl J Med* **359**: 61–73.

Golden JW, Riddle DL. 1984. The Caenorhabditis elegans dauer larva: developmental effects
552 of pheromone, food, and temperature. *Dev Biol* **102**: 368–378.

Greer EL, Maures TJ, Hauswirth AG, Green EM, Leeman DS, Maro GS, Han S, Banko MR,
554 Gozani O, Brunet A. 2010. Members of the H3K4 trimethylation complex regulate lifespan in a germline-dependent manner in *C. elegans*. *Nature* **466**: 383–387.

556   Hall SE, Beverly M, Russ C, Nusbaum C, Sengupta P. 2010. A cellular memory of
         developmental history generates phenotypic diversity in *C. elegans*. *Curr Biol* **20**: 149–
558   155.

       Hine C, Harputlugil E, Zhang Y, Ruckenstuhl C, Lee BC, Brace L, Longchamp A, Treviño-
560   Villarreal JH, Mejia P, Ozaki CK, Wang R, Gladyshev VN, Madeo F, Mair WB, Mitchell JR.
         2015. Endogenous hydrogen sulfide production is essential for dietary restriction
562   benefits. *Cell* **160**: 132–144.

       Ihnat MA, Thorpe JE, Kamat CD, Szabó C, Green DE, Warnke LA, Lacza Z, Cselenyák A, Ross
564   K, Shakir S, Piconi L, Kaltreider RC, Ceriello A. 2007. Reactive oxygen species mediate a
         cellular 'memory' of high glucose stress signalling. *Diabetologia* **50**: 1523–1531.

566   Kallio PJ, Wilson WJ, O'Brien S, Makino Y, Poellinger L. 1999. Regulation of the hypoxia-
         inducible transcription factor 1alpha by the ubiquitin-proteasome pathway. *J Biol Chem*
568   **274**: 6519–6525.

       Kilburn KH, Warshaw RH. 1995. Hydrogen sulfide and reduced-sulfur gases adversely
570   affect neurophysiological functions. *Toxicol Ind Health* **11**: 185–197.

       Kinoshita T, Seki M. 2014. Epigenetic memory for stress response and adaptation in plants.
572   *Plant Cell Physiol* **55**: 1859–1863.

       Kundu S, Horn PJ, Peterson CL. 2007. SWI/SNF is required for transcriptional memory at
574   the yeast GAL gene cluster. *Genes Dev* **21**: 997–1004.

       McColl G, Rogers AN, Alavez S, Hubbard AE, Melov S, Link CD, Bush AI, Kapahi P, Lithgow GJ.
576   2010. Insulin-like signaling determines survival during stress via posttranscriptional
         mechanisms in *C. elegans*. *Cell Metab* **12**: 260–272.

578   Miller DL, Budde MW, Roth MB. 2011. HIF-1 and SKN-1 coordinate the transcriptional
         response to hydrogen sulfide in *Caenorhabditis elegans*. *PLoS One* **6**: e25476.

580   Miller DL, Roth MB. 2009. *C. elegans* Are Protected from Lethal Hypoxia by an Embryonic
         Diapause. *Current Biology* **19**: 1233–1237.

582   Miller DL, Roth MB. 2007. Hydrogen sulfide increases thermotolerance and lifespan in
         Caenorhabditis elegans. *PNAS* **104**: 20618–20622.

584   Mirbahai L, Chipman JK. 2014. Epigenetic memory of environmental organisms: a reflection
         of lifetime stressor exposures. *Mutat Res Genet Toxicol Environ Mutagen* **764-765**: 10–
586   17.

       Richardson DB. 1995. Respiratory effects of chronic hydrogen sulfide exposure. *Am J Ind*
Med **28**: 99–108.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–140.

Sen N. 2017. Functional and Molecular Insights of Hydrogen Sulfide Signaling and Protein Sulfhydration. *J Mol Biol* **429**: 543–561.

Song J, Angel A, Howard M, Dean C. 2012. Vernalization - a cold-induced epigenetic switch. *J Cell Sci* **125**: 3723–3731.

Verghese J, Abrams J, Wang Y, Morano KA. 2012. Biology of the heat shock response and protein chaperones: budding yeast (*Saccharomyces cerevisiae*) as a model system. *Microbiol Mol Biol Rev* **76**: 115–158.

Völker U, Mach H, Schmid R, Hecker M. 1992. Stress proteins and cross-protection by heat shock and salt stress in Bacillus subtilis. *J Gen Microbiol* **138**: 2125–2135.

Wang GL, Jiang BH, Rue EA, Semenza GL. 1995. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O_2 tension. *PNAS* **92**: 5510–5514.

Wu D, Wang J, Li H, Xue M, Ji A, Li Y. 2015. Role of Hydrogen Sulfide in Ischemia-Reperfusion Injury. *Oxid Med Cell Longev* **2015**: 186908.

Xiao Y, Bedet C, Robert VJ, Simonet T, Dunkelbarger S, Rakotomalala C, Soete G, Korswagen HC, Strome S, Palladino F. 2011. Caenorhabditis elegans chromatin-associated proteins
SET-2 and ASH-2 are differentially required for histone H3 Lys 4 methylation in embryos and adult germ cells. *PNAS* **108**: 8305–8310.

Table 1: RNAseq comparison of gene expression in naïve and bookmarked animals.

	Gene	log$_2$ fold change			q-value			log CPM
		Naïve Unexp	Bkmk Unexp	Bkmk Naïve	Naïve Unexp	Bkmk Unexp	Bkmk Naïve	
Bookmarked	gst-19*	5.01	8.96	3.95	1.6E-22	5.0E-56	2.7E-16	7.58
	nspe-3*	4.86	8.00	3.14	5.8E-06	2.5E-15	4.6E-03	4.30
	nspe-4*	4.42	7.43	3.01	1.2E-08	1.7E-22	6.1E-05	4.68
	sqrd-1*	3.86	6.70	2.84	4.4E-20	2.6E-50	1.1E-11	6.96
	nspe-6*	4.92	7.67	2.75	1.9E-09	5.4E-21	1.6E-03	4.91
	R08E5.1*	1.94	4.54	2.59	5.0E-04	4.7E-24	1.0E-08	4.24
	nit-1*	2.73	4.95	2.22	3.6E-12	3.6E-36	2.8E-08	6.72
	cysl-2*	2.00	4.04	2.04	1.1E-05	6.7E-23	8.2E-06	8.85
Not Bookmarked	dhs-8*	3.20	4.33	1.13	1.5E-08	8.6E-16	1.0E+00	4.45
	nhr-62	1.84	2.83	0.99	4.6E-03	1.3E-08	1.0E+00	4.06
	cdo-1	1.40	2.31	0.91	4.8E-04	8.0E-13	2.9E-01	5.95
	M05D6.6	1.50	2.33	0.83	1.0E-04	8.0E-13	6.3E-01	7.25
	rhy-1*	4.44	5.06	0.62	2.6E-28	6.1E-36	1.0E+00	8.21
	lgc-1*	3.30	3.67	0.37	1.0E-02	6.7E-04	1.0E+00	0.45
	gbh-2	1.41	1.46	0.05	2.9E-03	8.1E-04	1.0E+00	5.08
	nhr-57*	4.16	3.43	-0.73	5.0E-28	2.7E-20	1.0E+00	5.65
Naïve Only	hsp-70	4.06	0.54	-3.52	3.3E-05	1.0E+00	7.0E-04	6.22
	R11A5.3	4.04	0.72	-3.32	6.0E-06	1.0E+00	4.6E-04	2.41
	hsp-16.2	3.43	0.79	-2.65	6.3E-04	1.0E+00	3.7E-02	5.23
	nhr-17	2.89	0.33	-2.56	3.6E-12	1.0E+00	2.1E-09	4.88
	cnp-3	2.15	0.49	-1.65	1.9E-04	1.0E+00	2.3E-02	4.01
	lact-4	1.97	0.81	-1.16	1.8E-08	6.3E-01	2.2E-02	5.65
	ugt-8	1.85	1.07	-0.78	1.1E-03	6.8E-01	1.0E+00	4.57
	C06B3.6	1.84	1.16	-0.68	4.3E-04	2.4E-01	1.0E+00	4.64
	egl-9	1.71	1.01	-0.70	7.3E-06	9.8E-02	1.0E+00	5.47
	aip-1	1.69	0.48	-1.21	8.5E-06	1.0E+00	1.7E-02	5.86
	cbs-1	1.67	-0.03	-1.70	9.5E-03	1.0E+00	7.5E-03	5.28
	scp-1	1.37	0.15	-1.23	4.6E-03	1.0E+00	2.9E-02	6.06
Bookmark Only	H12D21.6	3.45	5.40	1.95	1.7E-01	4.9E-07	3.6E-01	-0.31
	B0462.4	2.36	5.39	3.03	4.4E-02	2.6E-14	3.8E-05	2.66
	ptr-22	2.45	4.88	2.43	1.6E-01	2.0E-07	1.4E-01	3.63
	R08F11.4	0.71	4.83	4.12	1.0E+00	7.0E-12	1.3E-08	3.37
	ugt-14	2.33	4.22	1.89	9.1E-02	2.0E-07	4.0E-01	2.55
	R08E5.3	1.47	3.98	2.51	1.2E-02	9.3E-22	1.0E-08	9.26
	C08E8.3	2.51	3.98	1.46	1.5E-02	3.2E-08	8.5E-01	1.60
	nspe-2*	1.57	3.61	2.04	9.3E-02	6.2E-12	1.7E-03	5.00
	gst-30	1.10	3.29	2.19	1.0E+00	1.1E-03	2.7E-01	1.09
	mpst-3	0.88	3.29	2.40	6.0E-01	2.2E-21	3.5E-11	6.40
	ethe-1	0.92	3.15	2.23	1.0E+00	1.4E-09	1.7E-04	3.88
	Y37A1B.5	1.05	2.90	1.85	4.4E-01	8.0E-13	1.1E-04	8.31
	C08E8.10	1.09	2.87	1.78	1.0E+00	7.1E-06	8.5E-02	3.46
	comt-4	2.55	2.74	0.19	2.1E-02	3.5E-03	1.0E+00	0.70
	cysl-3	0.67	2.69	2.02	1.0E+00	1.1E-13	3.2E-07	6.98
	gst-16	-0.03	2.51	2.54	1.0E+00	2.4E-03	3.8E-03	1.80
	M05D6.5	1.14	2.48	1.34	1.4E-02	2.8E-15	7.0E-04	6.50
	C31C9.2*	0.69	2.41	1.72	1.0E+00	3.9E-09	4.6E-04	9.15
	glb-1	1.16	2.38	1.21	1.1E-01	1.4E-09	7.3E-02	5.30
	M162.5	0.52	2.30	1.78	1.0E+00	1.2E-05	7.0E-03	3.95
	cyp-36A1	1.43	2.29	0.86	9.3E-02	6.2E-06	1.0E+00	3.94
	F29C6.1	1.00	2.27	1.27	2.9E-01	8.0E-10	2.2E-02	5.99
	F59B10.4	1.36	2.25	0.89	7.7E-01	1.0E-03	1.0E+00	3.02
	dpy-17	1.23	2.25	1.02	1.0E+00	1.8E-03	1.0E+00	4.10
	nhr-11	0.54	2.16	1.62	1.0E+00	9.3E-05	3.2E-02	3.85
	F26H9.5	0.84	2.14	1.30	1.0E+00	7.0E-08	2.9E-02	8.09
	F56D5.3	0.91	1.98	1.07	1.0E+00	5.2E-03	1.0E+00	3.58
	F54D8.6	0.31	1.83	1.52	1.0E+00	1.8E-07	1.4E-04	7.09
	gst-4	0.37	1.68	1.31	1.0E+00	3.1E-04	3.9E-02	5.10
	Y62E10A.13	0.32	1.63	1.31	1.0E+00	1.2E-03	6.3E-02	8.12
	pck-1	0.78	1.58	0.81	1.0E+00	4.7E-03	1.0E+00	8.81
	C39E9.8	0.73	1.57	0.84	1.0E+00	3.5E-03	1.0E+00	7.33
	C17C3.1	0.74	1.48	0.74	1.0E+00	2.3E-04	1.0E+00	5.83
	acs-2	1.06	1.37	0.30	2.0E-01	5.2E-03	1.0E+00	5.78
	cysl-1	0.34	1.33	0.99	1.0E+00	7.1E-04	1.2E-01	6.73
	lips-10	0.58	1.33	0.74	1.0E+00	4.1E-03	1.0E+00	6.50
	Y11D7A.10	0.19	1.19	1.00	1.0E+00	3.5E-03	8.6E-02	6.40

Table 1: Differentially regulated genes after exposure to high H_2S identified by RNAseq. Bookmarked genes were upregulated significantly more in animals previously exposed to low H_2S than in naïve controls (but were induced by exposure to high H_2S in the naïve animals). Not bookmarked genes were upregulated the same in treated animals and naïve controls. Naïve only genes upregulated only in the previously untreated controls. Bookmark only genes were significantly induced only in animals that had previously been exposed to low H_2S.

A. Physiological memory of H₂S exposure

C. Hypoxia can induce acclimation

B. Bookmarking is time and dose dependent

Fig 1: Exposure to low H₂S induces an environmental bookmark.

A. Survival of animals exposed to high H₂S (black). In the schematic, time in low H₂S is colored yellow, time in room air without H₂S is white, and black indicates when animals were challenged with exposure to high H₂S. TOP ROW: Naïve animals grown in the absence of H₂S (open white). SECOND ROW, animals grown in low H₂S (50 ppm) throughout development. THIRD ROW: Animals exposed to low H₂S for 6 h as L4 and then immediately moved to high H₂S. BOTTOM ROW: Embryos exposed to low H₂S for 6 h, and then raised in room air without H₂S for 48 h. As noted in red below the bottom row, we refer to the initial exposure to low H₂S as acclimation, time in the absence of H₂S as the memory period, and the subsequent exposure to high H₂S as the challenge. Each exposure was repeated at least three times with at least 30 animals in each cohort.

B. Time-dose response to form the H₂S bookmark. Animals were acclimated to each indicated concentrations of H₂S for the indicated time as L1, then challenged with 150 ppm H₂S as L4.

C. Survival of animals exposed to 150 ppm H₂S after previous exposure to hypoxia. Wild-type L4 animals were exposed to 5,000 ppm O₂ for 5h and then either moved directly to high H₂S (hypoxia) or allowed to recover in room air for 3h before exposure to high H₂S (hypoxia + RA). Control animals remained in room air (RA) or were exposed to low H₂S for 5 h (low H₂S). Each point is the average survival from one independent experiment with at least 30 animals in each cohort for each experiment. Lines indicate the mean and standard deviation of the four independent experiments.

A. H₂S bookmarking during development

C. Bookmarking in aging animals.

B. Persistence of the H₂S bookmark

Fig 2: H₂S bookmarks are persistent but reverse spontaneously with age.
A. H₂S bookmarking during development. Animals at each developmental stage were exposed to low H₂S for 4 h, then returned to room air for 48 h before challenge with high H₂S.
B. H₂S bookmark persistence. Animals were exposed to low H₂S for 12 h as embryos and then returned to room air for the indicated period of time before challenge with high H₂S.
C. Bookmarking in aging animals. Animals were exposed to low H₂S for 12 h at the indicated age, then returned to room air for 48 h before challenge with high H₂S. For all panels, each experiment was repeated independently at least 4 times with 30-40 animals in each cohort. Error bars are standard deviation of the mean.

A. Bookmarking is reversed in starved L1.

B. H₂S exposure does not form bookmarks in starved L1

C. Fasting reverses H₂S bookmarking.

Fig 3: Fasting erases H₂S bookmarks.
A. Survival of animals exposed to high H₂S when acclimated as embryos. Embryos were acclimated to low H₂S for 4 h, then allowed to hatch in the presence or absence of food. After 12 h, L1 that hatched without food were moved to plates with food. When animals were L4 they were challenged with high H₂S.
B. Survival of animals acclimated to H₂S in the absence of food. Embryos isolated from naïve adults were allowed to hatch in the absence of food overnight. Starved L1 were moved to plates ± food, then exposed to low H₂S for 12 h. All animals were returned to house air and grown to L4/young adult on plates with food, then challenged with high H₂S.
C. H₂S bookmarking is reversed by fasting, but not other stress responses. As indicated in the schematic, animals were exposed to low H₂S for 4 h as embryos, then returned to room air. During the 48 h recovery time animals were transiently exposed to food deprivation (fasted; green), hypoxia (blue; 1,000 ppm O₂ for 24h), or heat shock (red; 37 C for 1 h). The animals were then challenged with high H₂S as L4/young adult. In all panels, graphs show mean survival ± standard deviation of at least 5 independent experiments with 30-40 animals in each experiment. ***, p < 0.001; **, p < 0.01.

Figure 4

A. SQRD-1::GFP expression

room air   low H$_2$S   low H$_2$S + RA

B. Transcript abundance of H$_2$S-induced genes 48 h after acclimation

C. Reactivation of SQRD-1::GFP expression in bookmarked animals

low H$_2$S (naïve)   high H$_2$S (naïve)   high H$_2$S (bookmarked)

D. Reactivation of H$_2$S-induced genes in bookmarked animals.

Fig 4: H$_2$S bookmarking is associated with transcriptional reactivation of H$_2$S-induced genes.
A. Representative images showing expression of the SQRD-1::GFP translational fusion protein in the absence of H$_2$S (room air; left), after 8 h in low H$_2$S (middle), and then 48 h after return to room air (right). In all images anterior is toward the top.
B. qRT-PCR measurement of abundance of mRNA from known H$_2$S-induced genes. Animals were exposed to low H$_2$S for 12 h and then returned to room air for 48 h. Graph shows mean transcript abundance relative to room air controls. Error bars are standard deviation of at least three experimental replicates. ns: not significant.
C. Representative images showing expression of the SQRD-1::GFP translational reporter in naïve animals exposed to low or high H$_2$S, and in bookmarked animals exposed to high H$_2$S. All animals were first-day adults. Representative images are shown, with measured fluorescence intensity of cohorts of 30-40 animals shown in the graph.
D. Transcriptional reactivation of H$_2$S-inducible genes in bookmarked animals. Transcript abundance of known H$_2$S-inducible genes was measured by qRT-PCR after animals were exposed to high H$_2$S for 1 h. The change in transcript abundance relative to untreated controls is shown for naïve and bookmarked animals. Mean $\pm$ standard deviation is shown. ****, p-value < 0.0001; **, p-value < 0.01; *, p-value < 0.05; ns, not significant.

Figure 5
Fawcett et al.

A. *set-2* and the CoREST-like complex are required to maintain H$_2$S bookmarking

B. *set-2* and *spr-5* are not required for acclimation to H$_2$S

Fig 5: The SET-2 histone demethylase and CoREST histone methyltransferase are required to maintain H$_2$S bookmarking.
A. Bookmarking defects of *set-2* and CoREST-like complex members. Animals were exposed to low H$_2$S for 4 h as embryos, then grown in room air for 48 h. Survival of challenge in high H$_2$S is shown. The *set-2* (purple) and *spr-5* (green) mutant phenotypes were found in a candidate screen of 250 genes (listed in Supplementary Table 1). *spr-5*, *spr-1*, *spr-3*, and *spr-4* are members of the CoREST-like complex, *wdr-5.1* is a component of the ASH complex, and *rbr-2* is a histone demethylase that counteracts the ASH methyltransferase (see references in main text). Graph shows mean ± standard deviation of at least 5 independent experiments with 30-40 animals in each cohort.
B. Acclimation of *set-2* and *spr-5* mutant animals. Animals were exposed to low H$_2$S for 8h as L4 and then moved directly to high H$_2$S overnight. Average survival is shown. Error bars are standard deviation of the mean. Graph shows mean ± standard deviation of at least 5 independent experiments with 30-40 animals in each cohort.

A. Transcriptional reactivation of H₂S-induced genes.

C. Transcriptional response to low H₂S.

B. Transcriptional response to high H₂S after acclimation.

Fig 6: Transcriptional reactivation in bookmarked animals requires *set-2* and *spr-5*.
A. Transcript levels were measured by qRT-PCR for naïve and bookmarked animals exposed to high H₂S for 1 h. The relative fold-change of transcript abundance ($\Delta\Delta C_t$*(bookmarked)*/ $\Delta\Delta C_t$*(naïve)*) is shown ± standard deviation. In all panels, N2 is in blue, *set-2(n4589)* is in purple, and *spr-5(by134)* is in green. In all panels ****, p-value < 0.0001; **, p-value < 0.01; *, p-value < 0.05; if not indicated, not significant.
B. The initial transcriptional response to low H₂S does not require *set-2* or *spr-5*. Transcript abundance for each H₂S-inducible gene was measured in *set-2(n4589)* and *spr-5(by134)* mutant animals after 1 h exposure to low H₂S. Graph shows mean change in transcript abundance, relative to untreated controls ($\Delta\Delta C_t$ ± standard deviation).
C. *set-2* and *spr-5* mutant animals can be preconditioned to H₂S. Animals of each genotype were exposed to low H₂S for 24 h, then moved to high H₂S for 1 h. The transcriptional response of these preconditioned animals is not distinguishable from wild-type controls. Graph shows $\Delta\Delta C_t$ ± standard deviation.

B. Stress response genes are not required for H$_2$S bookmarking

Bookmarking is specific to H$_2$S.

A. Survival of animals exposed to high H$_2$S. L4 animals were exposed to either low H$_2$S for 4 h, 37°C for 1 h (heat shock), or hypoxia (1,000 ppm or 5,000 ppm O$_2$) for 16 h. After treatment, animals were returned to room air. 48 hours later, animals were challenged with high H$_2$S and scored for survival.

B. General stress response pathways are not required for H$_2$S bookmarking. Animals with mutations in genes involved in general stress responses were adapted to low H$_2$S and challenged with high H$_2$S 48 hours later. For both panels, each experiment was repeated at least five times, with 30-40 animals. Graphs shows mean +/- SD. Statistical comparisons were between controls (animals in house air (HA) or the N2 wild-type strain) and treatment groups: ****, p-value <0.0001; ns, not significant.