1

Evolutionary dynamics of culturally transmitted, fertility-reducing traits

Dominik Wodarz[1,2], Shaun Stipp[2], and Natalia L. Komarova[2,1]

1: Department of Ecology and Evolutionary Biology, 321 Steinhaus Hall, University of California, Irvine, CA 92697

2: Department of Mathematics, 340 Rowland Hall, University of California, Irvine, CA 92617

contact: dwodarz@uci.edu; 949-824-2531.

Abstract

Human populations in many countries have undergone a phase of demographic transition, characterized by a significant reduction in fertility at a time of increased resource availability. Typically, the reduction in fertility is preceded by a reduction in mortality and a consequent increase in population density. Various theories have been proposed to account for the demographic transition process, including maladaptation, increased parental investment in fewer offspring, and cultural evolution. Aspects of cultural evolutionary processes in relation to demographic transitions have been studied with mathematical models. The reason for the observation that fertility reduction tends to be preceded by a decline in mortality, however, remains poorly understood. Using a variety of mathematical modeling approaches, we show that the cultural selection of low fertility traits crucially depends on the population death rate: if mortality is relatively high, the trait fails to spread. If mortality is reduced, the trait can spread successfully, thus offering an explanation for the observed trends. Computer simulations can reproduce the central characteristics of the demographic transition process, including significant changes in reproductive behavior within only 1-3 generations. A model tracking the continuous evolution of reproduction rates through "errors" in the cultural transmission process predicts fertility to fall below replacement levels if death rates are sufficiently low. This can potentially explain the very low ideal family sizes in Western Europe.

Introduction

In the 19[th] century, some human populations displayed a demographic transition from relatively high fertility and high mortality towards a significantly reduced fertility and lower mortality [1-4]. This first occurred in more developed parts of the world, such as Europe, the United States, Japan, Australia, and New Zealand, and coincided with an overall increase in resource availability (judged by economic considerations). In Western European countries, fertility has declined below replacement levels since the 1970s and 1980s [5,6], and this also applies to the perceived ideal family size. In German speaking countries the ideal family size has fallen below replacement levels, about 1.7 children [6].

Similarly, it has been observed that fertility reduction tends to be more pronounced in population segments that are economically advantaged than in poorer segments [1]. This is in contrast to trends observed before these demographic transitions, when increased wealth was associated with higher fertility [1,7].

An interesting characteristic of demographic transitions is that the reduction in fertility tends to be preceded by a reduction in the death rate of individuals, and by a consequent temporary population growth phase [4,8,9]. Transitions towards a reduced rate of reproduction following a rise in wealth and resources can be considered paradoxical, especially in the context of evolutionary biology [1,10]. According to

evolutionary thinking, individuals evolve to maximize their reproductive potential, and to

translate increased resource availability into higher reproductive success.

A number of theories have been put forward to account for demographic

transitions towards reduced fertility [1,11]. According to one line of argument, the

transition to reduced fertility might be maladaptive. That is, humans employ behavioral

strategies that used to be advantageous in previous times, but currently result in

behaviors that are not adaptive due to recent significant changes in socio-economic

factors. Another line of argument suggests reduced fertility to be adaptive, because the

current environment favors the production of few offspring with large parental

investment rather than the generation of more offspring with lesser parental investment

per child. This might be a significant consideration in parts of the world where

competition among individuals for jobs and careers is fierce. A third mechanism that

has been invoked to explain the observed decline in fertility is non-genetic, cultural

evolution. According to this argument, behavior that leads to reduced fertility in certain

influential individuals is copied by other individuals in the population, resulting in a

spread of this trait throughout the population. This explanation has the advantage that

cultural evolution can occur on much faster time-scales than genetic evolution, which is

required to account for the relatively fast observed rate of transition to lower fertilities in

human populations.

The dynamics of cultural transmission have been studied extensively with

mathematical models [12-19], and these approaches have also been applied to the

analysis of demographic transitions and the evolution of small family sizes, in particular

by the group of Marc Feldman [20-22]. In the simplest forms, such models are given by

ordinary differential equations that describe the spread of the cultural trait, similar to

infection models [12]. On a more complex level, age-structured models were formulated

[20,21] showing that cultural transmission dynamics can result in demographic

transitions on time scales that are similar to those observed in human data. The spread

of cultural traits that affect fertility, survival, or both was investigated and the effect of

these traits on the demographic structure of the population was investigated. The roles

of vertical, oblique, and horizontal transmission of the trait were examined, and the

models indicated that horizontal and oblique learning could accelerate the spread of the

cultural trait, compared to vertical transmission alone [20]. In an earlier study, it was

shown that cultural niche construction could be an important component that contributed

to the transition to reduced fertilities [21]. According to this mechanism, the first trait to

spread is one of valuing education, which provides an environment that promotes the

spread of a second, fertility-reducing trait. If the trait of valuing education is further

associated with reduced mortality of individuals, the model predicts that the decline in

fertility is preceded by a reduction in the population death rate, as observed in

demographic data. These models were examined further in the context of spatial

population structure, formulated as niche construction models in metapopulations [22].

In these studies, the increased survival / lower death rate of the population was

either considered a culturally transmitted trait (similar to fertility reduction), or it was

coupled to a culturally transmitted trait, e.g. by assuming that appreciation of education

correlates with increased viability. Here, we consider cultural transmission models

where the population death rate is an independent, exogenous variable that can be

modulated (reduced) over time, e.g. as a consequence of sanitary, medical or

technological advances, and investigate the effect of such changes on the ability of a

fertility-reducing trait to spread. We study the spread of the fertility-reducing trait in

spatially structured and non-spatial models, as well as in models with and without age

structure. We first consider a model with two populations, a fast and a slow reproducing

one, and subsequently consider models where the reproduction strategy is a continuous

trait. Because the overall death rate of individuals negatively correlates with the

economic development and the amount of resources available to the population, this

analysis sheds further light onto the correlation between resource availability and the

rate of population growth.

A two-dimensional agent-based model with asexual reproduction

121   We begin by examining the spread of a fertility-reducing cultural trait with the help of a

122   two-dimensional, stochastic agent-based model (ABM) that describes population

123   dynamics on a 2D grid of size nxn. We will refer to this model as ABM1. Two

124   populations are considered which reproduce with a fast and slow rate, respectively.

125 During each time step (representing a generation), the grid is randomly sampled 2M

126 times, where M is the total number of individuals currently present. When an individual

127 is picked, it can undergo either a birth-death update, or a cultural transmission update

128 (with equal probabilities). If the birth-death update is chosen, the individual reproduces

129 with a probability R_f and R_s, depending on whether this is a fast or slow reproducer. A

130 spot is chosen randomly from the eight nearest neighbors. If that target spot is empty,

131 the offspring is placed there, otherwise, the reproduction event is aborted. With an

132 independent probability D, the individual dies; both populations are assumed to have

133 the same death rate.

134 During a cultural transmission update, the individual attempts to change its

135 reproductive strategy with probability C. Given that the cultural transmission update is

136 attempted, the actual change by cultural transmission occurs with the probability defined

137 by the prevalence of individuals with the opposite strategy among the 8 nearest

138 neighbors (including the individual under consideration), see below for more details. The

139 ratio of the constant C to the mean division rate roughly measures prevalence of cultural

140 transmission in the population; it roughly corresponds to the fraction of the individuals in

141 each generation that attempt to change their reproductive strategy based on their

142 neighbors' strategy.

143 A cultural update is performed by gathering the information on the reproductive

144 strategy of the individuals' neighbors, similar to voter models [17,23]. In our model,

145 however, we assume that slow reproducers are more effective at changing a fast

146 reproducer's strategy than vice versa. In particular, when adding up the number of fast

147 and slow reproducers in the neighborhood, there is a probability Q<1 that a fast

148 reproducer is taken into account, while all slow reproducers are always included. This

149 reflects the assumption that the opinion of a slow reproducer is more influential than that

150 of a fast reproducer. This modeling choice is motivated by the assumption that slow

151 reproducers tend to channel the resources available to them into accumulation of wealth

152 and/or social status, and thus they may appear as more attractive models for imitation.

Outcomes of computer simulations for model ABM1

We first discuss the effect of the two components underlying the model: the birth-death

process and the cultural transmission process. If the model is run with only the

reproduction and death processes (without cultural transmission), the only outcome is

the persistence of the fast reproducing population and the competitive exclusion of the

slower reproducing one. This is straightforward competition dynamics behavior, which

underscores the notion that a slower rate of reproduction is a disadvantageous strategy.

If the model is run with only cultural transmission (no reproduction and death, but a

constant population), it essentially becomes a voter model, where "slow" and "fast" are

different opinions held by individuals in the population. As has been described for such

models [17,23], the only eventual outcome is that every individual in the population has

the same opinion. Which of the two opinions wins, depends on drift dynamics and

hence on initial frequencies of the opinions in the population.

When both processes are included, more complex dynamics are observed. Now three

outcomes are possible (Figure 1): the fast reproducers win and exclude the slow

reproducers; the slow reproducers win and exclude the fast reproducers; both

populations coexist in the longer term. Since we study stochastic models, the only

eventual outcome is extinction. In what we call "coexistence", the time until one of the

populations goes extinct is significantly longer than in the parameter regimes we call

"exclusion" (compare Figure 1C to 1A &B). Which outcome is observed depends on

several model parameters, most notably the death rate of agents, D. This is shown in

Figure 2Ai. Each point on this graph depicts the time until one of the populations goes

extinct, depending on the death probability, D. The outcomes are color-coded: purple

depicts fast reproducers remaining, and green slow reproducers. At higher death rates,

the fast reproducers persist and extinction of the slow reproduces occurs at relatively

short time scales. At low death rates, the slow reproducers persist and the fast

reproducers go extinct on a relatively short time scale. At intermediate death rates, the

time to extinction of one of the populations rises sharply, and either population has a

chance to go extinct first. This corresponds to the coexistence regime. Therefore, lower

death rates among individuals in the population create conditions in which the cultural

transmission of the slow reproduction trait is successful, resulting in an overall reduced

level of fertility.

Other parameters further modulate the outcome of the dynamics. Cultural

transmission of the low fertility trait is promoted by lower values of Q, i.e. by a reduced

influence of fast reproducers on choosing the reproduction strategy during the cultural

transmission procedure. Increasing the value of Q results in a lower population death

rate that is required for cultural transmission to be successful (Figure 2Aii). The relative

probability for a cultural transmission event to take place, C, is also an important

determinant of outcome. As expected, higher values of C promote the cultural spread of

the fertility-reducing trait. For lower values of C, lower population death rates are

needed for cultural transmission to be successful (Figure 2Aiii).

Including age structure into the agent-based model

Rather than considering fast versus slow reproducers, we now modify the agent-based

model to consider early versus late reproducers. This model will be referred to as

ABM2. While these two concepts are related, a reduction in fertility due to a later age of

first reproduction might be relevant to current times where segments of the population

with higher degrees of education and more wealth tend to reproduce at later ages. In

the agent-based model, we therefore consider four age classes. Individuals are born

into age-class 1, at which no reproduction is possible. During each time step, all

individuals age by one time unit. After A time units, an individual advances to the next

age class. Reproduction can occur in age classes 2 and 3 for early reproducers, and

only in age class 3 for late reproducers. In either case, reproduction occurs with a

probability R. Age class 4 is a post-reproductive phase, during which the only event that

can occur is death. Death can occur in all age classes, but with increasing probabilities

for successive age classes, i.e. with probabilities $D_4 > D_3 > D_2 > D_1$. This model has the

same properties as the previous one, where we distinguished between slow and fast

reproducers. We again observe three possible outcomes: extinction of either early or

late reproducers, or longer-term coexistence (not shown). The influence of parameters

on outcome is also the same on a qualitative level. Importantly, the death rate of

individuals is again a key determinant of outcome. Since in this model there are different

death rates for the different age classes, we multiplied all of the death rates by a factor

F, which means that all death rates change by the same amount. In Figure 2B, we

plotted the outcome of the dynamics for increasing values of F, which corresponds to

increasingly higher population death rates. Again, each point depicts the time until one

of the two populations goes extinct. We observe the same trend as before. For higher

population death rates, the fast reproducers remain. For low death rates, the slow

reproducers remain. For intermediate death rates, we observe coexistence, indicated by

a sharp rise in the time to extinction, where either fast or slow reproducers have a

chance to persist.

229   Complete mixing versus spatial constraints

The spatial nature of the models explored above makes it difficult to obtain analytical

insights. In this section, we formulate the same processes under the assumption that

individuals mix well. Spatial constraints are most relevant to the cultural transmission

aspect of the model because people are most likely to communicate with and consider

the opinion of neighboring individuals. Mixing, however, can be assumed if populations

are relatively small or information travels more easily through populations.

Fast and slow reproducers. First, we again consider two types of individuals that differ

by their reproduction rates. In the context of well-mixed populations, this can be

described by ordinary differential equations (see Section 1 of the Supplement for more

details of the analysis). Denoting fast reproducers by x_f and slow reproducers by x_s, the

model can be written as follows:

242
$$\dot{x}_f = r_f x_f W - dx_f - \beta x_f \frac{x_s}{x_s + \gamma x_f} + \beta x_s \frac{\gamma x_f}{x_s + \gamma x_f}$$

$$\dot{x}_s = r_s x_s W - dx_s + \beta x_f \frac{x_s}{x_s + \gamma x_f} - \beta x_s \frac{\gamma x_f}{x_s + \gamma x_f},$$

where $W = 1 - \dfrac{x_f + x_s}{K}$ introduces density dependence, limited by the carrying capacity

K. Hence, both populations grow logistically and are in competition with each other, and

die with a rate d (assumed to be the same for slow and fast reproducers). The rate at

which individuals change opinion through cultural transmission (fast reproducers

becoming slow and vice versa) is proportional to the weighted fraction of the individuals

with the opposite opinion in the whole population, multiplied by the rate constant β. (For

comparison, in the spatial setting, the probability to change opinion was proportional to

the weighted fraction of the individuals with the opposite viewpoint among only the

nearest neighbors). The coefficient $\gamma < 1$ indicates the degree to which the influence of

fast reproducers is reduced compared to that of slow reproducers. The equations above

simplify to the following system:

$$\dot{x}_f = r_f x_f W - dx_f - \beta(1-\gamma)\frac{x_f x_s}{x_s + \gamma x_f},$$

$$\dot{x}_s = r_s x_s W - dx_s + \beta(1-\gamma)\frac{x_f x_s}{x_s + \gamma x_f}.$$

255   Another way to model the process of differential strategy switching is to assume simply

256   that cultural transmission is more effective from slow to fast reproducers than vice

257   versa. This can be expressed as

$$\dot{x}_f = r_f x_f W - dx_f - \beta_s x_f \frac{x_s}{x_s + x_f} + \beta_f x_s \frac{x_f}{x_s + x_f}$$

$$\dot{x}_s = r_s x_s W - dx_s + \beta_s x_f \frac{x_s}{x_s + x_f} - \beta_f x_s \frac{x_f}{x_s + x_f},$$

259   where, importantly, $\beta_s > \beta_f$. If we denote $\beta = \beta_s - \beta_f$, then the model can be re-written as

$$\dot{x}_f = r_f x_f W - dx_f - \beta x_f \frac{x_s}{x_s + x_f}$$

$$\dot{x}_s = r_s x_s W - dx_s + \beta x_f \frac{x_s}{x_s + x_f}.$$

The formulation of the cultural transmission dynamics is now equivalent to an infection

model with frequency-dependent infection terms [12,24]. It turns out that for our

analysis, the frequency dependence is not essential, and we can assume that switching

occurs at a rate proportional to the abundance of the opposite type (scaled by the

carrying capacity K), not its fraction in the population. Hence, we also consider the

following version of the model:

$$\dot{x}_f = r_f x_f W - dx_f - \frac{\beta x_f x_s}{K}$$

$$\dot{x}_s = r_s x_s W - dx_s + \frac{\beta x_f x_s}{K}.$$

In the Supplement we demonstrate that all three versions of the model lead to very

similar conclusions. Here we show the results for the latter system, which is the

simplest, and which will be subsequently referred to as the "ODE model". If the

reproduction rates are larger than the death rates ($r_f > d$, $r_s > d$), the model is characterized

by three equilibria:

(i) Fast reproducers win and slow reproducers go extinct, with equilibrium expressions

given by $x_f = K(1 - d/r_f)$, $x_s = 0$. This equilibrium is stable if $d > d_1 \equiv \dfrac{r_1}{1 + (r_1 - r_2)/\beta}$.

(ii) Slow reproducers win and fast reproducers go extinct. The equilibrium expressions

are given by $x_f = 0, x_s = K(1 - d/r_s)$. The solution is stable if $d < d_2 \equiv \dfrac{r_s}{1 + (r_f - r_s)/\beta}$.

(iii) Coexistence of fast and slow reproducers is described by the following equilibrium

expressions: $x_f = \dfrac{K}{\beta}(d - d_2)$, $x_s = \dfrac{K}{\beta}(d_1 - d)$. This is stable if $d_2 < d < d_1$.

Therefore, similar to the corresponding spatial agent-based model, fast reproducers are

only expected to dominate for relatively large population death rates. Lower death rates

result in coexistence, and even lower death rates lead to the exclusive persistence of

slow reproducers.

These results make intuitive sense in the context of the infection dynamics

literature [25,26]. As the death rate of the population is decreased, the population

density increases until it is large enough to allow the slow reproduction meme to spread.

Essentially, a lower death rate increases the basic reproductive ratio of the slow

reproduction trait above unity, allowing invasion. The less the difference between the

strength of cultural trait transmission from fast→slow compared to slow→fast, the lower

the death rate, d, has to be (and hence the higher the population density has to be) to

allow the spread of the slow reproduction trait. Similarly, the larger the fitness difference

between slow and fast reproducers (r_f - r_s), the lower the death rate d has to be to allow

spread of the slow reproduction trait.

Age structure in a well-mixed model. An alternative approach is to consider a non-

spatial, age structured model [27]. We will model the competition dynamics of two types

that differ by their reproductive strategies (see Section 2 of the Supplement for

analytical results). Assume the existence of *N* discrete age groups for the two types,

and denote the abundance of type *s* in age group *i* as $x_i^{(s)}$. Reproduction behavior of

type *s* is described by reproduction rates in each age group, $a_i^{(s)}$. These quantities

comprise the information on the reproductive strategy (how early this type is willing to

start reproducing), and also the biological variation in fecundity (e.g. the assumption that

fecundity may decline with age). We assume that individuals of type *s=2* have a

tendency to reproduce later than individuals of type *s=1*. We can formulate a discrete

time dynamical system for these populations as follows:

$$x_1^{(s)}(t+1) = \sum_{i=1}^{N} a_i^{(s)} x_i^{(s)}(t) W,$$ (1)

$$x_j^{(s)}(t+1) = w_{j-1}^{(s)} x_{j-1}^{(s)}(t)(1 - \beta_j^{(s)} v_j^{(3-s)}(t)) + w_{j-1}^{(3-s)} x_{j-1}^{(3-s)}(t) \beta_j^{(3-s)} v_j^{(s)}, \quad 1 < j \le N$$ (2)

Here, the first equation describes the creation of newborn individuals of type s by older

individuals, where each age group reproduces with intensity $a_i^{(s)}$ and reproduction is

density-dependent, described by term W, e.g. $W = \left(1 + \sum_{s=1}^{2} \sum_{k=1}^{N} x_k^{(s)} / K\right)^{-1}$, with parameter K

having the meaning of carrying capacity. The rest of the equations describe (i) the

population moving from age group to age group, and (ii) the imitation dynamics.

Coefficients $w_{i-1}^{(s)}$ describe the probability for an individual of type s in age group j-1 to

survive until age j. The probability of switching type (imitation) is described by terms

including coefficients β. First we note that expression 3-s simply means "the other

type", as 3-s returns 1 for $s=2$ and it returns 2 for $s=1$. Further, the probability to switch

from type s to type 3-s while transitioning to age group j is given by $\beta_j^{(s)} v_j^{(3-s)}$, where

$$v_j^{(3-s)} = \frac{\sum_{k=j}^{N} x_k^{(3-s)}(t)}{\sum_{k=j}^{N} (x_k^{(3-s)}(t) + x_k^{(s)}(t))},$$

and is proportional to the fraction of individuals of age j and older that belong to "the

other type". We can see that the first term on the right of equation (2) multiplies the

probability that an individual does not switch to the other type, and the second term

multiplies the probability that switching from 3-s to s occurs.

Generally, the behavior of this discrete dynamical system can be quite complex,

and depending on the parameters we see exclusion equilibria (when one type is extinct

and the other reaches a steady positive equilibrium), exclusion periodic solutions (again,

one of the types is extinct and the other enters a steady cycle), and coexistence states

(where both types are present, and may reach a steady state or a cycle). To draw the

parallel between this model and those considered above, we note that parameters $w_{i-1}^{(s)}$

are responsible for survival to the next age group, and thus increasing these quantities

is similar to decreasing the death rate in the previous models. We observe that exactly

as before, large death rates favor the survival and dominance of the early reproducers,

and small death rates lead to the exclusion of early reproducers by late reproducers.

Intuitively, this result can be explained as follows. Assume for simplicity that the

transfer rate, β , does not depend on the age group, and the survival probability, w, is

identical for all age groups and for both types. Fitness of each of the types is comprised

of their net fecundity and their propensity to remain within their type (and not transfer to

the opposite type). In the absence of imitation (transfer), clearly, the fitness of type 1 is

larger than that of type 2. But this can be offset by a larger probability of transfer (if we

assume that $\beta^{(1)}$ is larger than $\beta^{(2)}$ by a sufficient margin). Small death rates (and

therefore large values of w) work against type 1 individuals and benefit type 2

individuals. If w is large, more individuals survive to later stages, and thus type 2 has a

better chance to reproduce. Further, for larger values of w, there will be a larger influx of

individuals transferring from type 1 to type 2: they simply have a longer time to stay

alive and decide to switch. Thus, living longer adds fitness to type 2, such that after a

threshold of w, type 2 becomes stronger and drives type 1 extinct.

Note that many other modeling choices can be made, which are described in the

Supplement. The general trends described here are robust and do not depend on

details on modeling choices. Analytical calculations for certain simple cases are also

presented in the Supplement.

Reproductive strategy as a continuous trait

In this section, we return to the spatial model, but modify the assumptions. So far, we

considered a population of slow and fast reproducers, where slow reproducers could

switch to fast reproducers and vice versa. Switching occurred with a probability that was

proportional to the weighed fraction of the opposing strategy among neighboring

individuals. Rather than considering two discrete reproductive strategies, however, it is

more realistic to assume the probability of reproduction to be a continuous variable. This

model will be called ABM3. We again assume that an individual is chosen for a cultural

transmission event with a probability scaled with C. In this model, however, instead of

adopting (or rejecting) the reproductive probability of the opposite type, the individual

adopts the weighted average of the reproduction probabilities among all neighbors

(including its own reproduction probability). As in the above models, we assume that

slower reproducers are more influential and contribute more to cultural transmission

than faster reproducers. This is implemented during the averaging procedures across

the neighborhood: we weigh the reproduction probability by a factor $Q<1$ if the

reproduction probability of a neighbor is faster than that of the individual under

consideration.

The outcome observed in this model is straightforward. As initial conditions, the

individuals in the system are characterized by different reproduction probabilities. Over

time, the reproduction probabilities converge to a spatially uniform value, the level of

which depends on the initially assigned probabilities. The reason for this eventual

uniformity is the assumption that an individual adopts the average reproduction

probability of the neighborhood during a cultural transmission event.

Next, we introduce "mutations" that can occur during cultural transmission.

Instead of simply adopting the (weighted) average strategy of the neighborhood, with

probability u individuals would modify this strategy by increasing or decreasing it (with

equal probabilities) by a fraction G. We examined the evolution of the average

reproduction probability, R, over time, by running computer simulations, and the

following outcomes were observed (Figure 3): (i) The average probability to reproduce,

R, increases steadily towards the maximum possible value (R+D=1), shown by the

purple, green, and red lines in Figure 3. (Simulations were stopped when R+D=1). (ii)

The average probability to reproduce declines steadily, eventually resulting in

population extinction, shown by the dark blue, light blue, and pink lines in Figure 3.

Extinction occurs because the reproduction rate evolves to levels that are too low to

maintain the population. (iii) The average probability to reproduce converges to an

intermediate level, and fluctuates around this level, shown by the yellow and orange

lines in Figure 3. This level is independent of the starting value of R (not shown).

Parameter values determine which outcome is observed. As before, the population

death probability, D, is a crucial factor (Figure 3). Evolution to maximal reproduction

probabilities, R, is seen for relatively large death rates. Evolution towards low values of

R and hence population extinction is observed for relatively low death rates. This could

be the cultural equivalent to "evolutionary suicide" or "Darwinian extinction" [28].

Evolution towards an intermediate reproduction probability is observed for intermediate

death probabilities, D. A higher probability of cultural transmission, C, and a lower

weight of faster reproducers during the averaging process, Q, further promote evolution

towards declining reproduction rates and population extinction (not shown). Section 3 of

the Supplement further explains the existence of an equilibrium state and explores how

the mean population reproduction rate depends on parameters.

Model with sexual reproduction

So far, all models considered assumed asexual reproduction, which is an obvious

oversimplification when considering human populations. Here, we repeat the analysis of

the last section by introducing sexual reproduction into this modeling framework,

referred to as ABM4. The same model as in the last section will be used, assuming the

probability of reproduction to be a continuous trait, and also assuming that for a cultural

transmission updates, a given individual adopts the weighted average reproduction

probability of the neighborhood with the possibility of "mutations" as defined above.

Sexual reproduction is incorporated in the following way. Two genders are

distinguished, gender 1 and gender 2. Before reproduction can occur, two individuals of

opposing gender have to form an exclusive connection, thus assuming monogamy. The

following events can occur if an individual is chosen for a reproductive update. If the

individual does not have a partner, a connection can be formed with a probability M if an

individual of the opposite gender without a partner is present among the eight nearest

neighbors. The partner is randomly chosen from the neighborhood. If the individual

does have a partner, reproduction happens with a probability R_{av}, which represents the

average reproduction probabilities of the two parents. For simplicity, it is assumed that

once formed, a partnership cannot break, corresponding to life-long monogamy. The

offspring resulting from this partnership are assigned to one of the genders with a 0.5

probability. The reproduction probability of the offspring is given by the average values

of the two parents. The offspring is placed into a randomly chosen empty spot among

the eight nearest neighbors of the parent that was originally picked for reproduction. If

no empty spots exist within the immediate neighborhood, reproduction is not successful.

Potential issues of mate preference for individuals with similar reproduction probabilities

are not taken into account. Death occurs with a probability D, according to the same

rules as described before.

As shown in Figure 4, the outcomes are qualitatively the same as in the

corresponding model without sexual reproduction, described in the previous section. For

relatively low population death rates, the reproduction probability can evolve towards

reduced values until population extinction occurs. For relatively high death rates, the

reproduction probability can evolve towards maximal values. For intermediate death

rates, the population can fluctuate around an intermediate reproduction probability.

Simulating demographic transitions in growing populations

In the modeling approaches described so far, competition is a major driving force of

outcome. In the absence of cultural transmission, faster or earlier reproducing

populations outcompeted the slower or later reproducing ones. Cultural transmission

could reverse the outcome of the competition if individuals were more likely to learn

from slower rather than faster reproducers. Competition, however, is most prevalent if

populations are close to an equilibrium or carrying capacity, which was the case in the

above analysis. The patterns reported above are not observed in exponentially growing

populations, because of the absence of competition (not shown). Because human

populations have been growing, it is important to re-visit the above-explored processes

in the context of unbounded population growth. While the population grows, individuals

can still compete on a local level. Therefore, we consider a growing population that is

subdivided into neighborhoods or demes. In each deme, the dynamics are modeled in

the same way as in the previous sections, imposing a carrying capacity for each deme.

For simplicity, we assume that in each deme, the dynamics are governed by the ODE

model. As initial conditions, a single deme is populated with a majority of fast

reproducers and a minority of slow reproducers. At the end of each time unit, there is a

chance that a new, empty deme is founded into which a fraction of current population

moves. The probability of this occurring is proportional to how full the current deme is. In

addition, the probability to found a new deme is inversely proportional to the number of

existing demes. While the demes are not arranged spatially in this model, founding a

new deme can be thought of as an increase in the density of the population, which gets

more difficult the more demes already exist. Hence, the probability for members of an

individual deme to found a new deme is given by $\dfrac{\alpha(x_f + x_s)}{K(\epsilon N + 1)}$, where N is the number of

currently populated demes, x_f and x_s represent local population sizes of fast and slow

reproducers, K is the local carrying capacity, and α and ε are constants. When a new

deme is founded, a given fraction of both fast and slow reproducers moves into the new

deme. As more demes become populated, the same algorithm is applied to every deme

after each time unit.

In this model, we observe persistence of one type and exclusion of the other,

while the population continues to grow (Figure 5A,B). As before, the fast population

persist for fast overall death rates (Figure 5A), while the slow population persists for low

overall death rates (Figure 5B).

We further used this model to simulate the demographic transition process

(Figure 5C). The simulation was run as before, but at a defined time point in the

simulation, the death rate is continuously and gradually reduced over several time steps

towards a lower, new level. This exogenous reduction is shown by the grey line and is

assumed to correspond to an improvement in various socio-economic factors that

reduce mortality, such as an improvement in disease treatment, sanitary conditions,

technological innovations, etc. The fast reproducing population is shown in purple, the

slow-reproducing population is green, and the total population size is shown by the red

dashed line. Initially, the overall population death rate is relatively high, and the fast

reproducing individuals enjoy a growth advantage. The average reproduction rate is

shown by the black line and is driven by the fast-reproducing population. When the

death rate is reduced, the fertility-reducing cultural trait can spread successfully and

eventually becomes the dominant population. As the death rate declines, we observe a

phase of faster population growth, as observed in data on demographic transitions [9].

Following a time delay after the reduction in the death rate, the reproduction rate also

declines, which is again consistent with data on demographic transitions [9] (this is also

seen in a zoomed-in graph in Figure 5D). The exact timing of events depends on model

parameters. For the purpose of this simulation, we chose parameters such that it takes

about 5 generations to reduce the reproduction rate two-fold. This is an order of

magnitude that is similar to events observed in human populations [1] and shows that

the cultural transmission dynamics underlying our model can lead to sufficiently rapid

changes in fertility. A faster rate of cultural transmission (higher value of β) can lead to

more rapid changes in fertility following the decline in the death rate, while the opposite

holds true for slower rates of cultural transmission.

Discussion and Conclusion

We have used a variety of modeling approaches to investigate the basic dynamics by which a fertility-reducing trait can spread via cultural transmission. A central result was that lower population death rates select for the cultural spread of the low-fertility trait. In a first set of models, the dynamics of two populations were followed, which were fast/slow or early/late reproducers in age structured models. This was done both in the context of nearest neighbor interactions and assuming perfect mixing. Both assumptions have value, depending on the population size and the connectivity of communication networks among individuals; in application to the modern population dynamics, a complete mixing assumption may correspond to the role of media in opinion spread. In a second set of models, the reproduction rate was modeled as a continuous trait, and cultural transmission corresponded to an individual adopting the (weighted) average reproduction rate found within the neighborhood. Assumptions about inaccuracies during cultural transmission allowed us to study the evolution of the average reproduction rate over time. In accordance with the simpler models, faster death rates resulted in evolution towards maximal reproduction rates, while lower death rates resulted in evolution towards ever decreasing reproduction rates, which likely induced in population extinction. Intermediate death rates were found to result in the evolution towards intermediate reproduction rates. These results were true both for asexual and sexual reproduction in the models.

Competition among individuals was a major driving force underlying the

dynamics arising from the model. While in the simpler settings explored here,

competition correlated with populations being close to carrying capacity, we showed

how a deme model can give rise to the same outcomes in populations that continuously

grow without converging to a carrying capacity. Hence, the results described throughout

the paper may hold for growing populations. We demonstrated that, depending on

parameters, the model can reproduce crucial features of the "demographic transition

model" [9].

Our study complements previous mathematical work that analyzed the cultural

spread of small family sizes in relation to demographic transitions [20-22]. Our models

consider a simpler setting involving the basic spread dynamics of the fertility reducing

trait, similar to infection models. The result that lower death rates promote the cultural

spread of the low fertility trait is intuitive if considered in the broader context of infection

dynamics models: lower death rates of individuals increase population density, and this

increases the basic reproductive ratio of the infection [26]. In the context of the

demographic transition towards reduced fertility, this is nevertheless an interesting

result. First, it might provide a simple explanation for the typical observation that a

reduction in fertility is preceded by a reduction in mortality, which our model reproduced.

In addition, the dynamics exhibited by these models suggest that fertility can be

characterized by density dependence, even in the context of increased resource

availability and continuously growing populations. Data indicate that fertility as well as

family size preference are characterized by density dependence, the reasons for which

are not fully understood [29].

An interesting result in our models was that for low population death rates, the

average reproduction rate of the population can continuously decline towards levels that

cannot sustain persistence anymore. In Western European countries, fertility has

declined below replacement levels since the 1970s and 1980s [5,6]. In addition, recent

surveys [6] have revealed that the ideal family size in German speaking countries has

fallen below replacement levels, about 1.7 children, among younger people, indicating

that this trend might continue in the future. These trends could be the result of the type

of cultural evolutionary dynamics explored here.

Some processes in the more complex versions of the models considered here

could also be formulated in slightly different ways. In ABM3 and ABM4, cultural

transmission involves the calculation of the weighted average reproduction rate among

individuals within the immediate neighborhood. The assumption was made that

individuals with a faster reproduction rate than the agent under consideration count less

in this process, irrespective of the magnitude of this difference. Alternatively, it could be

assumed that the reduced weight is proportional to the difference in reproduction rates,

thus taking into account the distance in social hierarchies. While it seems reasonable to

assume that economically more successful individuals carry more weight in cultural

transmission than individuals who are less successful [30,31], the details of this are not

well understood [32,33]. We note that results reported here depend on the assumption

that individuals with lower reproduction rates carry more social weight, an assumption

that has also been made in previous modeling work [21]. Another example of

uncertainties in model construction is the formulation of the sexual reproduction model.

We assumed monogamy, but made some obvious simplifications, as explained in the

results section. There are different assumptions that can be made in models that

describe sexual reproduction, but the most important feature in the current context is

that the reproduction rate of the offspring is not simply a copy of one of the parents, but

represents the average of the two parents. This provides an additional mechanism of

cultural change. Finally, only two types of communication networks have been

considered in the agent based models here, the one where individuals interact with

everyone else in the population, and the one where only interactions among nearest

neighbors are allowed. A large variety of more realistic, random communication

networks can be constructed, but we do not expect the results to differ from the ones

obtained from the two extreme cases of networks considered here.

While some details of the model processes could be formulated in different ways,

we have considered a range of models with different assumptions: with and without

spatial restrictions, in the presence or absence of age structure, with sexual versus

asexual reproduction, with different ways in which cultural transmission changes the

reproduction rate, and in models with different population structure. In all models, the

death rate of the population was identified as a crucial factor that determined whether

cultural transmission of a fertility-reducing trait was successful or not, which could have

implications for understanding the forces that contribute to the occurrence of

demographic transitions and that drive the decline of fertility below replacement levels in

developed countries.

Acknowledgements: We would like to thank Simon Levin for useful discussions that

helped shape this manuscript.

Figure legends:

Figure 1. Time series showing the different outcomes according to ABM1. Individual

realizations are shown. (A) Higher death rates: fast reproducers persist and slow

reproducers go extinct on a short time scale. (B) Lower death rates: slow reproducers

persist and fast reproducers go extinct on a short time scale. (C) Intermediate death

rates: both fast and slow reproducers persist for significantly longer time periods.

Eventually one population goes extinct due to the stochastic nature of the simulation.

Parameters were chosen as follows. R_f=0.005; R_s=0.8R_f; C=0.0008; Q=0.93. For (A),

D=0.001. For (B), D=0.0001. For (C), D=0.00025.

Figure 2. Time to competitive exclusion, as a function of the death rate. (A) Model ABM1. Individual realizations of the computer simulation were run until one of the two populations (fast or slow reproducers) went extinct. This time was recorded with a green dot if the fast reproducer went extinct, and with a purple dot if the slow reproducer went extinct, as a function of the population death rate, D. For low death rates, there are only green dots, corresponding to the slow reproducer persisting and the fast reproducer going extinct relatively fast. For fast death rates, there are only purple dots, corresponding to the fast reproducers persisting and the slow reproducers going extinct in a relatively fast time scale. For intermediate death rates, the time until one of the populations goes extinct becomes sharply longer, and either population can go extinct first. This corresponds to long-term coexistence. For plot (i), parameters were chosen as follows: $R_f=0.005$; $R_s=0.8R_f$; $C=0.0008$; $Q=0.93$. Plots (ii) and (iii) explore parameter dependence of the phenomenon. (ii) A higher value of $Q=0.98$ makes it harder for the slow-reproduction trait to invade, hence requiring lower population death rates. (iii) A lower rate of cultural transmission, $C=0.0004$, makes it harder for the slow-reproduction trait to invade, hence again requiring lower population death rates. (B) Same, but according to ABM2 with age structure. Because each age class is characterized by its own death rate, we multiplied all those death rates by a variable factor F, and plotted the outcome against this parameter. The death rates for the age classes were: $D_1=0.00004$; $D_2=0.00007$; $D_3=0.00009$; $D_4=0.0002$. Other parameters are $R=0.005$; $C=0.0008$; $Q=0.93$; $A=10,000$.

Figure 3. Outcomes of ABM3 with a continuous reproduction strategy and cultural

evolution. The average reproduction probability across the whole population is plotted

over time. Individual simulation results are shown. Simulations were run for different

death rates, decreasing from D1 to D8. For relatively high death rates, the average

reproduction probability increases steadily towards maximal levels. For relatively low

death rates, the average reproduction probability decreases steadily until population

extinction occurs (due to the limited reproduction). For intermediate death rates, the

average reproduction probability comes to oscillate around a steady value, which does

not depend on initial conditions (not shown). Parameters were chosen as follows. Death

rates are given by D1 = 0.002, D2 = 0.001, D3 = 4×10^{-4}, D4 = 3.75×10^{-4}, D5 = 3.6×10^{-4},

D6 = 10^{-4}, D7 = 5×10^{-5}, D8 = 10^{-5}. The reproduction probability of the individuals, R, was

allowed to evolve, starting from R=0.05 for all individuals. C=0.0003; Q=0.965. The

chance to make a mistake during cultural transmission ("mutation") u=0.1. In case of a

mistake, the average reproduction rate was changed by G=2%.

Figure 4. Outcomes of ABM4 with continuous reproduction strategies, cultural

evolution, and sexual reproduction. Simulations are in principle the same as those

presented in Figure 3, but using the model with sexual reproduction. Assumed death

rates decrease from D1 to D8. Results remain robust. Parameters were chosen as

follows. Death rates are given by D1 = 0.002, D2 = 0.001, D3 = 7.5×10^{-4}, D4 = 7×10^{-4}, D5

= 6.5×10^{-4}, D6 = 5×10^{-4}, D7 = 2.5×10^{-4}, D8 = 1×10^{-4}. The reproduction probability of the

individuals, R, was allowed to evolve, starting from R=0.05 among all individuals. The

population death rate, D, is indicated in the graphs. u=0.1, M=0.9; C=0.0003; Q=0.965;

G=0.1; G=2%.

Figure 5. Computer simulations of the deme model, described in the text. (A) The slow-

reproducing population (green) goes extinct and the fast-reproducing population

(purple) continues to grow. Parameter values were chosen as follows: r_f=0.08, r_s=0.04,

d=0.03, β=0.02, K=10, α=0.01, ε=0.01. (B) The fast-reproducing population is going

extinct, and the slow-growing population takes over and continues to grow. The same

parameter values were used, except d=0.003. (C) Simulation of the demographic

transition process. Again, fast and slow reproducers are shown in purple and green,

respectively. The total population size is shown by the dashed red line. The simulation

is started with a death rate d=0.006. In this regime, the fast-reproducing population has

the advantage and is dominant. The cultural spread of the low-fertility trait is not

successful. At a defined time point, the death rate is reduced 1.8 fold every half

generation until it has fallen to a value of d=0.001 (grey line). This creates conditions

under which the cultural transmission of the fertility-reducing trait is successful, and the

population characterized by a slow reproduction rate spreads. This leads to a decline in

the average reproduction rate of the population (black line), which is delayed with

respect to the reduction in the death rate. For the parameter regime considered, the

average reproduction rate is halved within about 2-3 generations, which corresponds to

about 50-100 years (a generation in the model is given by 1/r). The remaining

parameters are given as follows. r_f=0.008, r_s=0.0016, β=0.2, K=10, α=0.01, ε=0.01.

References

1. Borgerhoff Mulder M (1998) The demographic transition: are we any closer to an evolutionary explanation? Trends Ecol Evol 13: 266-270.
2. Caldwell JC (2006) Demographic Transition Theory. Dordrecht, The Netherlands.: Springer.
3. Coale AJ (1989) Demographic transition. In: Eatwell J, Milgate M, Newman P, editors. Social Economics. London.: Palgrave Macmillan UK.
4. Kirk D (1996) Demographic Transition Theory. Population Studies 50: 361-387.
5. Castles FG (2003) Castles, Francis G. "The world turned upside down: below replacement fertility, changing preferences and family-friendly public policy in 21 OECD countries. Journal of European social policy 13: 209-227.
6. Goldstein J, Lutz W, Testa MR (2003) The emergence of sub-replacement family size ideals in Europe. Population Research and Policy Review 22: 479-496.
7. Vining DR (1986) Social versus reproductive success – the central theoretical problem of human sociobiology. Behav Brain Sci 9: 167-260.
8. Thompson V, Roberge MC (2015) An Alternative Visualization of the Demographic Transition Model. Journal of Geography 114: 254-259.
9. Lee R (2003) The Demographic Transition: Three Centuries of Fundamental Change. Journal of Economic Perspectives 17: 167-190.
10. Aarssen LW (2005) Why Is Fertility Lower in Wealthier Countries? The Role of Relaxed Fertility-Selection. Population and Development Review 31: 113-126.
11. Wilson C (1999) Wilson, Chris. "Evolutionary theory and historical fertility change. Population and development review 25: 531-541.
12. Cavalli-Sforza LL, Feldman MW (1981) Cultural Transmission and Evolution: A Quantitative Approach. Princeton, NJ.: Princeton University Press.
13. Creanza N, Kolodny O, Feldman MW (2017) Cultural evolutionary theory: How culture evolves and why it matters. Proc Natl Acad Sci U S A.
14. Feldman MW, Cavalli-Sforza LL (1976) Cultural and biological evolutionary processes, selection for a trait under complex transmission. Theor Popul Biol 9: 238-259.
15. Boyd R, Richerson PJ (1985) Culture and the evolutionary process. . Chicago, IL: University of Chicago Press.
16. Bisin A, Verdier T (2001) The Economics of Cultural Transmission and the Dynamics of Preferences. Journal of Economic theory 97: 298-319.
17. Ehrlich PR, Levin SA (2005) The evolution of norms. PLoS Biol 3: e194.
18. Durrett R, Levin SA (2005) Can stable social groups be maintained by homophilous imitation alone? Journal of Economic Behavior & Organization 57: 267-286.
19. Nakamaru M, Levin SA (2004) Spread of two linked social norms on complex interaction networks. J Theor Biol 230: 57-64.
20. Fogarty L, Creanza N, Feldman MW (2013) The role of cultural transmission in human demographic change: an age-structured model. Theor Popul Biol 88: 68-77.
21. Ihara Y, Feldman MW (2004) Cultural niche construction and the evolution of small family size. Theor Popul Biol 65: 105-111.
22. Borenstein E, Kendal J, Feldman M (2006) Cultural niche construction in a metapopulation. Theor Popul Biol 70: 92-104.
23. Liggett TM (1999) Stochastic interacting systems: contact, voter, and exclusion processes. Berlin: Springer.

24. McCallum H, Barlow N, Hone J (2001) How should pathogen transmission be modelled? Trends in
Ecology & Evolution 16: 295-300.
25. Kermack WO, McKendrick AG (1927) Contributions to the mathematical theory of epidemics – I. Proc
R Soc Med 115A: 700-721.
26. Heesterbeek JA (2002) A brief history of R0 and a recipe for its calculation. Acta Biotheor 50: 189-
204.
27. Charlesworth B (1980) Evolution in age-structured populations. Cambridge, UK: Cambridge
University Press.
28. Webb C (2003) A complete classification of Darwinian extinction in ecological interactions. Am Nat
161: 181-205.
29. Lutz W, Rita Testa M, Penn DJ (2006) Population density is a key factor in declining human fertility.
Population and Environment 28: 69-81.
30. Katz E, Felix P (1955) Personal Influence; the Part Played by People in the Flow of Mass
Communications. Glencoe, IL: Free Press.
31. Grewal R, Mehta R, Kardes FR (2000) The Role of the Social-Identity Function of Attitudes in
Consumer Innovativeness and Opinion Leadership. Journal of Economic Psychology 21: 233-252.
32. Bernardi L, Klärner A (2014) Social networks and fertility. Demographic research 30: 641-670.
33. Nakayama S, Krasner E, Zeno L, Porfiri M (2019) Social information and spontaneous emergence of
leaders in human groups. Journal of The Royal Society Interface 16: 20180938.

Fast reproducer

Slow reproducer

(A)

(i)

(ii)

(iii)

Population death rate, d

(B)

Multiplier of population death rate, F

Evolutionary dynamics of culturally transmitted, fertility-reducing traits

Dominik Wodarz, Shaun Stipp, Natalia L. Komarova

Supplementary information

Contents

1   In the absence of age structured dynamics

1.1   The basic ODE model

Let us describe the dynamics of two competing types of individuals, fast reproducers and slow reproducers. We will denote the population of the former type as $x^{(1)}$ and the population of the latter type as $x^{(2)}$. We have

$$\dot{x}^{(1)} = r_1 x^{(1)} W - d x^{(1)} - \beta x^{(1)} x^{(2)}/K, \qquad (1)$$
$$\dot{x}^{(2)} = r_2 x^{(2)} W - d x^{(2)} + \beta x^{(1)} x^{(2)}/K. \qquad (2)$$

Here, each type reproduces with its own linear reproduction rate, with

$$r_1 > r_2,$$

and the competition is expressed by term W, which for example can take the logistic form,

$$W = 1 - \frac{x^{(1)} + x^{(2)}}{K},$$

where K denotes carrying capacity. Both types die with equal rates, d. There is also a probability of switching from one type to the other, which is proportional to the abundance of the individual of the opposite type. So, the total rate at which type 1 switches to type 2 is given by

$$\beta_1 x^{(1)} \frac{x^{(2)}}{x^{(1)} + x^{(2)}},$$

and the total rate at which type 2 switches to type 1 is given by

$$\beta_2 x^{(2)} \frac{x^{(1)}}{x^{(1)} + x^{(2)}}.$$

If we assume that $\beta_1 > \beta_1$ and denote $\beta = \beta_1 - \beta_2$, we have term

$$\beta x^{(1)} \frac{x^{(2)}}{x^{(1)} + x^{(2)}}$$

with the minus sign in the equation for $x^{(1)}$, and the same term with the plus sign in the equation for $x^{(2)}$. These terms are equivalent in form to frequency dependent infection. It turns out that for our analysis, the frequency dependence is not essential, and we can assume that switching occurs at a rate proportional to the abundance of the opposite type (scaled by the carrying capacity), not its percentage in the population. Therefore we will use equations of the form (1-2). The main assumptions are that

- Type 1 grows faster than type 2, $r_1 > r_2$, and

- There are more individuals switching from type 1 to type 2 than the other way around, that is, $\beta > 0$.

System (1-2) has four steady states.

0. The trivial solution, $x^{(1)} = x^{(2)} = 0$ is unstable as long as $r_1 > d$ and $r_2 > d$. We will assume that both types are viable and the above inequalities hold.

1. Type 1 (fast reproducers) wins:

$$x^{(1)} = K(1 - d/r_1), \quad x^{(2)} = 0.$$

   This solution is stable if

$$d > d_1 \equiv \frac{r_1}{1 + (r_1 - r_2)/\beta}.$$

2. Type 2 (slow reproducers) wins:

$$x^{(1)} = K(1 - d/r_1), \quad x^{(2)} = 0.$$

   This solution is stable if

$$d < d_2 \equiv \frac{r_2}{1 + (r_1 - r_2)/\beta}.$$

   Note that $d_2 < d_1$.

3. Coexistence solution,

$$x^{(1)} = \frac{K}{\beta}(d - d_2), \quad x^{(2)} = \frac{K}{\beta}(d_1 - d).$$

   This solution is positive as long as

$$d_2 < d < d_1. \tag{3}$$

   The characteristic polynomial is given by

$$\lambda^2 + \frac{d(r_1 - r_2)}{\beta}\lambda + \frac{\beta + r_1 - r_2}{\beta}(d_1 - d)(d - d_2),$$

   and the eigenvalues have negative real part as long as the free term is positive, which is only satisfied by condition (3), which means that inequality (3) is the stability condition for the coexistence solution.

1.2   Other ODE models

In the model just considered the conversion process is described by the term

$$\beta x^{(1)} x^{(2)} / K.$$

Alternatively, this term can be formulated as

$$\beta \frac{x^{(1)}x^{(2)}}{x^{(1)} + x^{(2)}},\qquad(4)$$

where the conversion happens proportionally to the current fraction of the individuals of the opposite type. In this case, we have a very similar solution structure. The competitive exclusion solutions are the same as in the previous model, the threshold d values are given by

$$d_1 = \frac{\beta}{1 - r_2/r_1},\quad d_2 = \frac{\beta}{r_1/r_2 - 1},$$

and the coexistence solution is given by

$$x^{(1)} = \frac{K}{\beta}\left(1 - \frac{\beta}{r_1 - r_2}\right)(d - d_2),\quad x^{(2)} = \frac{K}{\beta}\left(1 - \frac{\beta}{r_1 - r_2}\right)(d_1 - d).$$

In a different modeling approach we assume that conversion happens at the same rate for both strategies, but it isn proportional to the weighted fraction of the two strategies in the population. Assuming that strategy 1 is weighed with coefficient $\gamma < 1$, we obtain that the change in numbers for strategy 1 is given by

$$\beta(1 - \gamma)\frac{x^{(1)}x^{(2)}}{\gamma x^{(1)} + x^{(2)}}.\qquad(5)$$

In this case, the competitive exclusion solutions are the same as in the previous model, the threshold d values are given by

$$d_1 = \frac{\beta(1 - \gamma)}{\gamma(1 - r_2/r_1)},\quad d_2 = \frac{\beta(1 - \gamma)}{r_1/r_2 - 1},$$

and the coexistence solution is given by a somewhat different expression,

$$
\begin{aligned}
x^{(1)} &= \frac{K}{\beta + d}\left(\frac{\beta}{r_1 - r_2} + \frac{\beta + d}{\gamma r_2 - r_1}\right)\left(\beta + d + \frac{r_1 - \gamma r_2}{\gamma - 1}\right),\\
x^{(2)} &= \frac{K}{\beta + d}\left(\frac{d\gamma}{\gamma - 1} + \frac{\beta(\beta + d - r_1)}{r_2 - r_1} - \frac{(\beta + d)^2\gamma}{\gamma r_2 - r_1}\right).
\end{aligned}
$$

2 Age structured dynamics

2.1 Model formulation

We will model the competition dynamics of two types that differ by their reproductive strategies. Assume the existence of N discrete age groups for the two types, and denote the abundance of type s in age group i as $x_i^{(s)}$. Reproduction behavior of type s is described by the vector $a_i^{(s)}$, with entries in $[0, 1]$ denoting relative rate of reproduction of this type in age i. Individuals of the first type, $s = 1$, correspond to "fast reproducers", and the second type, $s = 2$, to the "slow reproducers" in the previous section. The latter type generally has a tendency to reproduce later than individuals of type 1. In the approach implemented here, type s is characterized by two integers, $i_{start}^{(s)}$ and $i_{end}^{(s)}$, denoting the first and last age groups where reproduction is possible. We have

$$a_i^{(s)} > 0 \ \text{ if } \ i_{start}^{(s)} \leq i \leq i_{end}^{(s)}, \quad a_i^{(s)} = 0 \text{ otherwise,}$$

where

$$i_{start}^{(1)} < i_{start}^{(2)}.$$

We can formulate a discrete time dynamical system for these populations as follows:

$$x_1^{(s)}(t+1) \ = \sum_{i=1}^{N} a_i^{(s)} x_i^{(s)}(t) W, \tag{6}$$

$$x_j^{(s)}(t+1) \ = w_{j-1}^{(s)} x_{j-1}^{(s)}(t) \left(1 - \beta_j^{(s)} \frac{\sum_{k=j}^{N} x_k^{(3-s)}(t)}{\sum_{k=j}^{N}(x_k^{(3-s)}(t) + x_k^{(s)}(t))} \right)$$

$$+ \quad w_{j-1}^{(3-s)} x^{(3-s)}(t) \beta_j^{(3-s)} \frac{\sum_{k=j}^{N} x_k^{(s)}(t)}{\sum_{k=j}^{N}(x_k^{(3-s)}(t) + x_k^{(s)}(t))}, \quad 1 < j \leq N, \tag{7}$$

where the competition term W can be defined as

$$W \ = \ 1 - \frac{\sum_{s=1}^{2} \sum_{k=1}^{N} x_k^{(s)}}{K} \quad \text{or} \tag{8}$$

$$W \ = \ \left(1 + \frac{\sum_{s=1}^{2} \sum_{k=1}^{N} x_k^{(s)}}{K} \right)^{-1}. \tag{9}$$

Equation (6) describes reproduction. Different age groups reproduce with their own rate $s_i^{(s)}$, and the offspring enters age group 1. Equation (7) describes the population moving from age group to age group. Coefficients $w_{j-1}^{(s)}$ describe the probability for an individual of type s to survive until age j. The probability of switching type is described by terms including coefficient β. First we note that expression $3 - s$ for $s \in \{1, 2\}$ simply returns the type different from type s, because $3 - s$ gives 2 if $s = 1$ and it gives 1 if $s = 2$. The probability to switch from type s to type $3 - s$ while transitioning to age group j is given by

$$\beta_j^{(s)} \frac{\sum_{k=j}^{N} x_k^{(3-s)}(t)}{\sum_{k=j}^{N}(x_k^{(3-s)}(t) + x_k^{(s)}(t))},$$

and is proportional to the fraction of individuals of age j and older that belong to class $3 - s$. With this in mind, we can see that the first term on the right of equation (7) multiplies the probability that an individual does not switch to the other type, and the second term multiplies the probability that switching from $3 - s$ to s occurs. System (6-7) assumes no switching at the first stage. To include switching at the first stage, we replace equation (6) with

$$\begin{aligned}
x_1^{(s)}(t+1) &= \sum_{i=1}^{N} a_i^{(s)} x_i^{(s)}(t) W \left(1 - \beta_1^{(s)} \frac{\sum_{k=1}^{N} x_k^{(3-s)}(t)}{\sum_{k=1}^{N}(x_k^{(3-s)}(t) + x_k^{(s)}(t))} \right) \\
&+ \sum_{i=1}^{N} a_i^{(3-s)} x_i^{(3-s)}(t) W \beta_1^{(3-s)} \frac{\sum_{k=1}^{N} x_k^{(s)}(t)}{\sum_{k=1}^{N}(x_k^{(3-s)}(t) + x_k^{(s)}(t))}. \quad (10)
\end{aligned}$$

2.2 System behavior

System (10, 7) has two exclusion steady states (for $s = 1$ and $s = 2$), which for competition model (9) are given by

$$x_j^{(s)} = K \prod_{k=1}^{j-1} w_k^{(s)} \frac{r \sum_{m=1}^{N} a_m^{(s)} \prod_{k=1}^{m-1} w_k^{(s)} - 1}{\sum_{m=1}^{N} \prod_{k=1}^{m-1} w_k^{(s)}}, \quad 1 \le j \le N, \quad (11)$$

$$x_j^{(3-s)} = 0, \quad 1 \le j \le N. \quad (12)$$

In figure 1 the behavior of a system with $N = 5$ stages is shown. We assumed that for fast reproducers, $i_{start}^{(1)} = 2$, and for slow reproducers, $i^{(2)} = 3$,

while $i_{end}^{(s)} = 5$ for both types. For simplicity we assumed that within the reproductive stages, the values $a_i^{(s)}$ were equal to a constant (independent on type and stage). Further, we assumed that the rates $w_i^{(s)}$ were s- and i-independent, and transfer coefficients $\beta_i^{(s)}$ were i-independent (but dependent on s).

Figure 1: Age structured dynamics according to system (10, 7), numerical simulations. Total populations of individuals of type 1 and type 2 are presented. The steady state values are given on the left as functions of parameter (a) $\beta^{(1)}$ and (b) $w_i^{(s)} = w$ for all $s \in \{1, 2\}, 1 \le i \le N$, the survival probability. Solution types are denoted by a circled number. The parameters are $w = 0.9$ in (a), $\beta^{(1)} = 0.17$ in (b), and $K = 50, \beta_i^{(2)} = 0.1$. The reproductive rate $a_i^{(s)} = 1$ when $2 \le i \le 5$ for $s = 1$ and $3 \le i \le 5$ for $s = 2$. Initially, all populations $x_i^{(s)} = 10$.

In figure 1(a), by fixing all the parameters except for $\beta^{(1)}$, we observed that three different solution types were stable. Solution 1 corresponds to the fast reproducers excluding the slow reproducers and is stable for smaller values transfer away from type 1, $\beta^{(1)}$. Solution 2 corresponds to the slow reproducers excluding the fast reproducers, and corresponds to larger $\beta^{(1)}$. For intermediate values of $\beta^{(1)}$ we observe stable coexistence of both types. Sam-

ple time series of the three solution types (corresponding to three different values of $\beta^{(1)}$) are presented on the right on the figure.

Alternatively, if we fix $\beta^{(1)} > \beta^{(2)}$ and vary the survival probability, w, the same three solution types are observed, 1(b). In particular, we note that low survival rates (that is, high death rates) lead to the dominance of fast reproducers, and high survival rate (low death rates) to the dominance of slow reproducers.

2.3 A two-age system

The simplest nontrivial system that captures the phenomenon of interest is system (10,7) with $N = 2$. Let us assume that $w_j^{(s)} = w$ for both types (that is, mortality is the same for both types). Further, let

$$i_{start}^{(1)} = 1, \quad i_{end}^{(1)} = 2, \quad i_{start}^{(2)} = 2, \quad i_{end}^{(1)} = 2,$$

in other words, type 1 reproduces both in ages 1 and 2, and type 2 only reproduces in age 2. The trivial solution[1] is unstable if $wa_2^{(2)} > 1$ or $wa_2^{(1)} > 1 - a_1^{(1)}$. The following are some of the non-trivial long-term solutions (compare to the equilibria of section 1):

1. Type 1 (fast reproducers) wins – a competitive exclusion steady state:

$$x_1^{(1)} = \frac{K[r(a_1^{(1)} + wa_2^{(1)}) - 1]}{1 + w}, \quad x_2^{(1)} = \frac{Kw[r(a_1^{(1)} + wa_2^{(1)}) - 1]}{1 + w}, \quad x_1^{(2)} = x_2^{(2)} = 0.$$

2. Type 2 (slow reproducers) wins – a competitive exclusion steady state:

$$x_1^{(1)} = x_2^{(1)} = 0, \quad x_1^{(2)} = \frac{K[ra_2^{(2)}w - 1]}{1 + w}, \quad x_2^{(2)} = \frac{Kw[ra_2^{(2)}w - 1]}{1 + w}.$$

3. A coexistence state.

4. Periodic solutions.

---

[1]For the analysis of the trivial solution one has to modify the original system by adding a small constant in the denominators of all the equations, otherwise we have a singularity which is meaningless, because the transfer terms multiplying β must be zero if the population is zero.

Stability of the two exclusion states can be investigated. For simplicity, let us set all nonzero values of fecundity to a constant, $a_j^{(s)} = a$. Further, we will assume that the coefficient of transfer is independent of the age, and is only defined by the type: $\beta_j^{(s)} = \beta^{(s)}$ for $j = 1, 2$, $s = 1, 2$. Let us analyze stability of solution 1 above (fast reproducers win). Stability of the discrete system requires all the eigenvalues of the Jacobian to satisfy $|\lambda| < 1$. The eigenvalues are given by

$$\lambda_{1,2} = \frac{1 \pm \sqrt{1 + 4rw(1+w)^2}}{2r(1+w)^2}, \tag{13}$$

$$\lambda_{3,4} = \frac{\beta^{(1)}(2+w) \pm \sqrt{w[w(2+\beta^{(1)} - 2\beta^{(2)})^2 + 4(1+\beta^{(1)} - \beta^{(2)})(1-\beta^{(2)})]}}{2(1+w)}. \tag{14}$$

The first two eigenvalues do not depend on the transfer rates and correspond to the stability of the type 1 population in the absence of the other population. We can show that $|\lambda_{1,2}| \leq 1$ for all $0 \leq w \leq 1$ and $r \geq 1$. In particular, $\lambda_1 \geq 0$, we have $\lambda_1 = 1$ when $r = 1, w = 0$, it decays with r and w for $r \leq 2$, and for a given $r > 2$, it has a maximum value $(1-w)/2$ when

$$r = \frac{2}{(w-1)^2(w+1)}.$$

Further, $\lambda_2 \in (-1, 0]$ for all values $w \in [0, 1]$ and $r \geq 1$, since $\partial \lambda_2 / \partial r > 0$, and for $r = 1$, $\lambda_2 = 1 - \sqrt{1 + 4w(1+w)^2}/(2(1+w)^2) \in [1/8(1 - \sqrt{17}), 0]$.

The eigenvalues $\lambda_{3,4}$ describe stability against an invasion of type 2 individuals. The solution can become unstable if $\lambda_3 > 1$. This happens when

$$w > w_1 \equiv \frac{(1 - \beta^{(1)})^2}{(\beta^{(2)} - \beta^{(1)})(2 - \beta^{(2)})}.$$

Clearly, if $\beta^{(1)}$ is large (close to 1), the type 1 solution is unstable (because of frequent transfers to type 2). In fact, as long as

$$\beta^{(1)} < \frac{4 - \beta^{(2)} - \sqrt{5(\beta^{(2)})^2 - 16\beta^{(2)} + 12}}{2},$$

the type 1 solution is stable for any values of $w < 1$, because $w_1 > 1$. If however the inequality above is revered (that is, the transfer rate is larger

than a threshold for type 1), the solution becomes unstable for sufficiently large values of w.

Intuitively, fitness of each of the types is comprised of their net fecundity and their propensity to stay (and not transfer to the opposite type). Clearly, the fecundity of type 1 is larger than that of type 2. But this can be offset by a larger probability of transfer (if we assume that $\beta^{(1)}$ is larger than $\beta^{(2)}$ by a sufficient margin). Small death rates (and therefore large values of w) work against type 1 individuals and benefit type 2 individuals. If w is large, more individuals survive to later stages (and thus type 2 has a better chance to reproduce). Further, for larger values of w, there will be a larger influx of individuals transferring from type 1 to type 2: they simply have a longer time to stay alive and decide to switch. Thus, living longer adds fitness to type 2, such that after a threshold of w, type 2 becomes stronger and drives type 1 extinct.

Investigating the stability of type 2 equilibrium, we discover that it is unstable (in this simple 2-age model) for all values of w except for $w = 1$, where it is neutral. Note that for systems with more age stages, this is not the case, and we have a stable type 2 equilibrium (see the previous section). For the 2-age system, for values $w < 1$, but close to 1, instead of equilibrium 1, we observe a stable cycle which contains only type 2 individuals.

3 Birth-death, imitation, and mutation dynamics

3.1 Model formulation and numerical results

Envisage the following process. In a 1D spatial system of a constant size, N, each individual, i, is characterize by a reproduction rate, l_i. During each time unit, N updates are performed, each consisting of two parts, a death-birth (DB) update and a cultural transmission (CT) update. Each update proceeds as follows:

- A DB update: An individual is chosen, randomly and fairly, to be removed (say, this is the individual at location i_1). Then it is replaced by the progeny of one of its two neighbors: the individual at location

$i_1 + 1$ reproduces with probability $l_{i_1+1}/(l_{i_1+1}+l_{i_1-1})$, and the individual at location $i_1 - 1$ reproduces with probability $l_{i_1-1}/(l_{i_1+1} + l_{i_1-1})$. The offspring inherits the reproduction rate of the parent.

- A CT update: this event happens with probability β, which sets the relative time scale of the two types of updates. Pick an individual, randomly and fairly, to perform an imitation update (say this is the individual at location i_2). This individual will change its reproduction rate from l_2 to

$$\tilde{l} = \frac{\sum_{j=i_2-1}^{i_2+1} \alpha_{i_2,j} l_j}{\sum_{j=i_2-1}^{i_2+1} \alpha_{i_2,j}},$$

  where

$$\alpha_{i,j} = \begin{cases} 1, & l_j \leq l_i, \\ s, & l_j > l_i, \end{cases}$$

  and $0 < s < 1$ is a constant that indicates by how much the strategy of fast reproducers is discounted. In other words, a weighted average of all the strategies around the focal individual at i_2 is formed, such that the strategy of those who reproduce faster than the focal individual is discounted with coefficient s. The focal individual adopts the resulting strategy with probability $1 - u$. With probability u, strategy $\tilde{l}$ is increased or decreased (with equal likelihood) by an amount Δl (unless $l < \Delta l$, in which case it can no longer decrease). This process is equivalent to mutations, whereby the phenotype is modified with a certain probability to give rise to variation.

We would like to characterize the equilibrium of this system. First we note that in the absence of mutations ($u = 0$), the state with $l_i = l$ for all i is a equilibrium for any value of l. As a result, the system will converge to one of these neutral equilibria, depending, for example, on the initial condition.

The dynamics change drastically in the presence of mutations, $u > 0$. Now, uniform states are no longer equilibrium states, and the equilibrium reproduction rates will be distributed around some mean value, $\bar{l}$, with the variance that increases with u and Δl. In figure 2(a) we present the time series of the population mean reproduction rates, for 4 different values of Δl, the increment of the reproduction rate. We can see that the population settles to a stochastic equilibrium, where the mean population mean reproduction rate increases with Δl, and convergence time decreases with Δl. Figure 2(b) shows numerically obtained histograms of reproduction rates of populations

Figure 2: The dynamics of a 1D simulation with mutations. (a) The time-series of the population mean reproduction rate, for 4 different values of Δl. (b) Numerically obtained histograms of the population's reproduction rates, taken at generation 50,000, for the same 4 values of Δl. The rest of the parameter are: $N = 100$, $u = 0.04$, $\beta = 1$, $s = 0.9$.

at equilibrium, for the same four values of Δl. We can see that the standard deviation increases with Δl. Similar trends are observed when we vary the mutation rate, u (not shown). 2D simulations that show the same trends are shown in figure 3.

3.2 Analytical considerations

To find the mean equilibrium value of the reproduction rates, we use the following argument. Suppose that the equilibrium distribution[2] of the reproduction rates is given by $\{f_k\}$, such that the probability for an individual to have reproduction rate L_k is given by f_k, with

$$\sum_k f_k L_k = \bar{l}.$$

Under a BD event, suppose an individual at position i_1 with reproduction rate L_1 is picked for replacement, and suppose further than its two neighbors have reproduction rates L_2 and L_3. Then the expected increment in the

---
[2]A similar argument for continuous distributions can be developed.

Figure 3: The dynamics of a 2D simulation with mutations. The population mean reproduction rate is plotted as a function of time, for 3 simulations. The blue line represents a base-line simulation with parameters $u = 0.1, \Delta l/l = 0.02$, the orange line a simulation with an increased mutation rate, $u = 0.3$, and the green line a simulation with an increased $\Delta l/l = 0.04$. The rest of the parameters are as in Fig.3 of the main text, with the death rate 3.75×10^{-4}.

reproduction rate of the focal individual is given by

$$-L_1 + L_2\frac{L_2}{L_2 + L_3} + L_3\frac{L_3}{L_2 + L_3}.$$

Averaging over all the possible reproduction rates, we obtain the expected increment in reproduction rate from a DB update:

$$\Delta L_{DB} = \sum_i \sum_j \sum_k \left(-l_i + \frac{l_j^2}{l_j + l_k} + \frac{l_k^2}{l_j + l_k}\right) f_i f_j f_k. \tag{15}$$

Similarly, we can calculate the expected increment in the reproduction rate resulting from a cultural transmission event:

$$\Delta L_{CT} = \sum_i \sum_j \sum_k \left(-l_i + \frac{l_i + \alpha_{ij}l_j + \alpha_{ik}l_k}{1 + \alpha_{ij} + \alpha_{ik}}\right) f_i f_j f_k. \tag{16}$$

The equation

$$\Delta L_{DB} = -\beta \Delta L_{CT} \tag{17}$$

characterizes the equilibrium. Note that the right hand side of this equation is positive, because the mean increment resulting from CT updates is negative, due to a diminished weight of high reproduction rates in the weighted averages. The left hand side is also positive, because DB updates tend to increase the reproduction rates due to competition among individuals.

Let us assume that the width of the distribution of the equilibrium reproduction rates is defined by the mutation rate (and the increment Δl), and keep it fixed, while varying the mean $\bar{l}$. Note that in equation (16), the expression in the parentheses can be rewritten as

$$\frac{\alpha_{ij}(l_j - l_i) + \alpha_{ik}(l_k - l_j)}{1 + \alpha_{ij} + \alpha_{ik}}.$$

For each location i, let us present $L_i = \bar{l} + \epsilon m_i$, where all m_i are IID with a zero mean and a variance that we denote by $(\sigma/\epsilon)^2$. We can see that $\bar{l}$ cancels from the above expression, and its statistics will only depend on the distribution width. In other words, the mean decrement received by the population reproduction rate as a result of a CT update is defined by the difference between the focal reproduction rate and a weighted average of its neighboring reproduction rates, and does not depend of the absolute value of the rates.

On the contrary, the DB increment defined by equation (15) depends on the magnitude of $\bar{l}$. Intuitively, neighbors compete for filling the empty spot, and the amount of advantage experienced by a neighbor with a higher reproduction rate is proportional to the relative, and not absolute, difference in the rates. Therefore, the increment scales with the relative amount of spread in reproduction rates, and is thus inversely proportional to $\bar{l}$. Again, for each location i, we present $l_i = \bar{l} + \epsilon m_i$, where all m_i are IID with a zero mean and variance $(\sigma/\epsilon)^2$. Then, expanding the expression in parentheses in (15) in terms of ϵ we obtain

$$\left(\frac{m_j + m_k}{2} - m_i \right) \epsilon - \frac{\epsilon}{2} \frac{(m_j - m_k)^2}{m_j + m_k} \sum_{n=1}^{\infty} \left(-\frac{(m_j + m_k)\epsilon}{2\bar{l}} \right)^n.$$

The first term averages to zero, and the second term is given by

$$\frac{\epsilon^2}{4\bar{l}}(m_j - m_k)^2,$$

which upon averaging yields

$$\frac{\sigma^2}{2\bar{\bar{l}}},$$

a quantity inversely proportional to the mean reproduction rate of the population. We further see that it depends on the square of σ in the lowest order.

From the above analysis it follows that the left hand side of equation (17) is a decaying function of $\bar{l}$ which tends to zero as $\bar{l} \to \infty$, and the right hand side of equation (17) is $\bar{l}$-independent. There will be a unique intersection of the two curves as long as β is chosen to be sufficiently low. This intersection defines the equilibrium value of the population mean reproductive rate.

We further note that the quantities Δ_{DB} and $-\Delta_{CT}$ both grow with the distribution width of the reproduction rates, but while $-\Delta_{CT}$ is linear in σ, Δ_{DB} is quadratic in this quantity, and thus grows faster as we increase the width of the distribution of l. Therefore, as u increases and the distribution width increases, the left hand side of equation (17) grows faster than the right hand side, resulting in an increase in the solution, $\bar{l}$.

Figure 4: Finding the equilibrium reproduction rate by solving equation (17), illustrated with example (18-19). The left hand side of equation (17), ΔL_{DB}, is shown as blue lines and the right hand side, $-\beta\Delta L_{CT}$, with yellow lines, as functions of $\bar{l}$. Solid, dashed, and dotted lines correspond to three different values of Δl: $0.05, 0.10, 0.15$. The rest of the parameters are: $s = 0.9, \mu = 0.1, \beta = 1$.

This is illustrated in an example where we assumed that the division

rates are distributed according to the following three-valued distribution with mean $\bar{l}$ and variance $(\Delta l)^2 \mu$:

$i \rightarrow$	1	2	3
L_i	$l - \Delta l$	l	$l + \Delta l$
f_i	$\mu/2$	$1 - \mu$	$\mu/2$

The expressions for ΔL_{DB} and ΔL_{CD} can be obtained explicitly,

$$\Delta L_{DB} = \frac{(\Delta l)^2 \mu}{2\bar{l}} \frac{(\Delta l)^2 \mu - 4\bar{l}^2}{(\Delta l)^2 - 4\bar{l}^2}, \tag{18}$$

$$\Delta L_{CT} = \frac{\Delta l \mu (1 - s)}{6(2 + s)(1 + 2s)} \left((6 - \mu)\mu s - 10s + \mu(3 + \mu) - 8\right). \tag{19}$$

In figure 4, both sides of equation (17) are plotted as functions of $\bar{l}$, and their intersections are marked with vertical lines, for three values of Δl, which represent an increase in the distribution width. We can see that the corresponding solutions $\bar{l}$ become larger for larger distribution widths.

16