Unveiling two new trichome-specific promoters of interest for metabolic engineering in *Nicotiana tabacum*

Mathieu Pottier[$], Raphaëlle Laterre, Astrid Van Wessem, Aldana M. Ramirez, Xavier Herman, Marc Boutry and Charles Hachez[*]

Louvain Institute of Biomolecular Science and Technology, University of Louvain, 1348 Louvain-la-Neuve, Belgium

[$] Current address: InBioS-PhytoSYSTEMS, Laboratory of Plant Physiology, University of Liège, B-4000, Liège, Belgium

* Corresponding author:
Charles Hachez,
charles.hachez@uclouvain.be,
+ 32 10 47 37 96

ORCID:
Mathieu Pottier : 0000-0003-1551-4699
Marc Boutry : 0000-0002-2315-6900
Charles Hachez : 0000-0002-3688-7614

Acknowledgments

The authors are grateful to Joseph Nader for his technical contribution. This work was supported by the Belgian Fund for Scientific Research (Grant ID: MIS – F.4522.17), the Interuniversity Poles of Attraction Program (Belgian State, Scientific, Technical and Cultural Services), and an EU Marie Skłodowska-Curie fellowship (Project ID: 658932) to MP.

Author contributions

MP, RL, CH and MB designed the experiments and analyzed the data. MP, RL, AVW, AR, XH and MB performed experiments. CH, MP and MB wrote the manuscript.

Conflict of Interest: The authors declare that they have no conflict of interest.

Key message

pRbcS-T1 and *pMALD1*, two new trichome-specific promoters of *Nicotiana tabacum*, were identified and their strength and specificity were compared to those of previously described promoters in this species.

Abstract

Nicotiana tabacum has emerged as a suitable host for metabolic engineering of terpenoids and derivatives in tall glandular trichomes, which actively synthesize and secrete specialized metabolites. However, implementation of an entire biosynthetic pathway in glandular trichomes requires the identification of trichome-specific promoters to appropriately drive the expression of the transgenes needed to set up the desired pathway. In this context, RT-qPCR analysis was carried out on wild-type *N. tabacum* plants to compare the expression pattern and gene expression level of *NtRbS-T1* and *NtMALD1*, two newly identified genes expressed in glandular trichomes, with those of *NtCYP71D16, NtCBTS2α, NtCPS2,* and *NtLTP1,* which were reported in the literature to be specifically expressed in glandular trichomes. The latter were previously investigated separately, preventing any accurate comparison of their expression level. We show that *NtRbcS-T1* and *NtMALD1* are specifically expressed in glandular trichomes like *NtCYP71D16, NtCBTS2α,* and *NtCPS2*, while *NtLTP1* was also expressed in other leaf tissues as well as in the stem. Transcriptional fusions of each of the six promoters to the *GUS-VENUS* reporter gene were introduced in *N. tabacum* by *Agrobacterium*-mediated transformation. Most transgenic lines displayed GUS activity in tall glandular trichomes. In some transgenic lines, except for *pNtLTP1:GUS-VENUS*, this expression was specific. In other transgenic lines, GUS expression was extended to other tissues, probably resulting from a position effect during transgene integration. We discuss alternatives to overcome this lack of tissue specificity in some transgenic lines, should some of these promoters be used in the context of metabolic engineering in *N. tabacum*.

Keywords: Rubisco small subunit, Major Allergen Mal D 1.0501, Cembratrien-ol Synthase, Copal-8-ol diphosphate Synthase, Lipid Transfer Protein, Cytochrome P450 oxygenase

Introduction

Trichomes, the epidermal outgrowths covering most of aerial plant tissues are found in a very large number of plant species. Several types of trichomes (unicellular or multicellular, glandular or non-glandular) can be observed in a single plant species. Among those, glandular trichomes are characterized by the presence of cells forming a glandular structure that has the ability to secrete or store large quantities of specialized (also called secondary) metabolites (e.g., phenylpropanoids, flavonoids, acyl sugars, methylketones, and terpenoids),which possess antimicrobial and anti-fungal properties or which act as a defense barrier against herbivorous insects (Schilmiller et al., 2008).

The specialized metabolites secreted by glandular trichomes, which might represent up to 17 % of the leaf dry weight in *Nicotiana tabacum* (tobacco), have been largely exploited over centuries (Wagner et al., 2004). One of their most ancient uses originates from the aromatic properties and fragrance of some of those secretions referred to as essential oils. Besides, many of these specialized metabolites constitute an interesting source of pharmaceuticals and food additives. However, one of the common issues with some specialized metabolites is their low production yield in their natural host plant (Van Agtmael et al., 1999; Yoon et al., 2013). Some of them are highly praised molecules only found in a single plant species or even a single plant cultivar and often at low concentration (e.g., taxol found in *Taxus sp.*, artemisinin in *Artemisia annua* or cannabinoids in *Cannabis sativa*). Therefore, natural resources are often insufficient to reach the global need, while the complex stereochemistry of these compounds often prevents their full chemical synthesis in a cost-effective way.

In order to increase the overall yield, metabolic engineering strategies are undertaken to synthesize such metabolites through homologous or heterologous expression systems (Kirby and Keasling, 2009; Marienhagen and Bott, 2013). Advances in plant biotechnology and increasing knowledge in specialized metabolism also make it possible to exploit plants as production platforms for specific metabolites. One of the main advantages of such a strategy is that plants are photoautotrophic organisms, therefore requiring simple and cheap growth conditions, which accounts for a cost-effective biomass production (Kempinski et al., 2015). Besides, another benefit of using plant hosts is their ability to deal with membrane proteins such as P450 enzymes and posttranslational modifications such as glycosylation, two key features that are frequently limiting in prokaryotic hosts (van Herpen et al., 2010).

Among plant specialized metabolites, terpenoids and derivatives are the most abundant in terms of sheer number and chemical diversity (for review, see Croteau et al. 2000; Bouvier et

al. 2005; Gershenzon and Dudareva 2007). Different plant species have been tested to

perform metabolic engineering. From these trials, *N. tabacum* has emerged as one of the most

suitable plant hosts (Lange et al., 2013; Moses and Pollier, 2013; Wang et al., 2016). *N.*

tabacum synthesizes an important pool of terpenoid precursors (IPP/DMAPP) and, besides

the essential metabolites derived from the isoprenoid biosynthesis pathways, it also produces

a very high amount of a limited range of specialized metabolites (Huchelmann et al., 2017).

This combined to its high biomass, its fast growth rate, its easy genetic transformation and

regeneration make it an interesting host to implement the biosynthesis pathways of terpenoid

compounds and derivatives thereof.

However, engineering terpenoid biosynthetic pathways using ubiquist promoters

frequently leads to severe phenotypes including dwarfism, chlorosis, and decreased seed

production due to the cytotoxicity of these compounds or detrimental impact on the

biosynthesis of essential metabolites (Saxena et al. 2014; Gwak et al. 2017; reviewed in

Huchelmann et al. 2017). To avoid these adverse effects on the plant physiology, a fine

control of the spatiotemporal expression of the transgenes, physically restricting the

biosynthesis of potentially cytotoxic metabolites to specialized organs, is desirable

(Huchelmann et al., 2017). Since glandular trichomes are non-essential organs, which are

involved in the biosynthesis, storage and/or secretion of specialized metabolites and which

naturally evolved to efficiently deal with high concentrations of these metabolites, they make

ideal targets to develop such a metabolic engineering approach. For this purpose,

identification of trichome-specific promoters and therefore of trichome-specific genes is

required.

The promoters of four genes from *Nicotiana* species have previously been shown to be

trichome-specific: CYtochrome P450 oxygenase 71D16 (*NtCYP71D16*), Copal-8-ol

diPhosphate Synthase 2 (*NtCPS2*), Lipid Transfer Protein 1 *(NtLTP1)* of *N. tabacum,* as well

as CemBraTrien-ol Synthase 2α (*NsCBTS2α*) of *Nicotiana sylvestris*. All these genes are

exclusively related to specialized metabolism (Choi et al., 2012; Ennajdaoui et al., 2010;

Sallaud et al., 2012; Wang et al., 2002).

A proteomic comparison was recently performed in *N. tabacum* between proteins extracted

from tall glandular trichomes, which produce large amounts of terpenes, and those extracted

from other plant organs (Laterre et al., 2017). This led to the identification of 47 proteins that

were more abundant in tall glandular trichomes, the most enriched ones being a putative PR-

10 type pathogenesis-related protein, namely Major Allergen Mal D 1.0501 (MALD1) and a

small subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase (RbcS-T1) (Laterre et al., 2017). The transcriptional promoter of a *Nicotiana benthamiana RbcS-T* homolog coupled to a reporter gene was shown to be trichome-specific in leaf tissues (Laterre et al., 2017). In *N. tabacum*, the trichome-specific localization of *NtMALD1* and *NtRbcS-T* transcripts was supported by semi-quantitative RT-PCR (Harada et al., 2010; Laterre et al., 2017).

To limit cytotoxicity issues and/or adverse effects on plant metabolism, elucidating whether gene promoters reported to be active in glandular trichomes are specific to glandular trichomes or display a broader activity in plant tissues is critical if those promoters are to be used in metabolic engineering approaches. The six promoters mentioned above had been investigated separately, preventing one to compare their transcript levels. In addition, for some of them, their cell-type specificity monitored by the GUS reporter gene had not been described in other organs than leaf tissues. Yet, these are key parameters to determine which are the most suitable ones for metabolic engineering in *N. tabacum* tall glandular trichomes.

The present study thus aimed at comparing the expression patterns and expression levels of *NtCYP71D16, NtCBTS2α, NtCPS2, NtLTP1, NtRbS-T1*, and *NtMALD1* promoters in *N. tabacum*. Their transcript levels in trichomes and different organs were compared. Transcriptional fusions of each of the six promoters to *GUS-VENUS* were expressed in transgenic *N. tabacum* plants. Most of the lines obtained with the six reporter constructs displayed GUS activity in the tall glandular trichomes. This expression was trichome-specific in a few lines, except for *pNtLTP1:GUS-VENUS*, corroborating transcripts data. The other lines exhibited GUS activity in other tissues, pointing out to random transgene insertion and positional effects as the most probable causes of such discrepancies. We therefore discuss alternatives to overcome this lack of tissue specificity in some lines, should some of these promoters be used in genetic constructs to drive the expression of specific transgenes.

Results

In a 2D gel analysis of glandular trichome proteins from *N. tabacum*, several spots were identified as trichome-specific proteins, among which RbcS-T1 and MALD1 (Laterre et al., 2017). The RNA levels of *NtRbS-T1* and *NtMALD1* as well as of *NtLTP1*, *NtCYP71D16*, *NtCBTS2α, and NtCPS2*, previously reported as genes specifically expressed in tall glandular trichomes, were compared in trichomes and different other *N. tabacum* organs. To do so, leaves were frozen in liquid nitrogen and carefully scratched with a brush to collect the trichomes. RT-qPCR assays were then performed on RNA extracted from trichomes, roots, trichome-free leaves, and trichome-free stems of six-week-old plants as well as from flowers of 10-week-old plants. Unlike for leaves and stems, trichomes could not be retrieved from flower sepals and petals. For each gene, the relative expression level in trichomes was arbitrarily set to one. The stability of the ubiquitin (*NtUBQ*) and the ATP-synthase β-subunit (*NtATP2*) control genes used to normalize the data is shown in Supplemental Fig. S1. All six studied genes showed a significantly ($p < 0.05$) higher relative expression level in isolated trichomes compared to the levels observed in roots, leaves, stems or flowers (Fig. 1). *NtCYP71D16, NtCBTS2α, NtCPS2*, *NtRbS-T1*, and *NtMALD1* exhibited very low expression in roots, trichome-free leaves and trichome-free stems, while higher transcript levels were found for *NtLTP1* in leaves and stems. The expression was high in flowers for the six genes but, as noted above, trichomes were not removed from these organs.

As most of these genes are involved in the biosynthesis (*NtCYP71D16, NsCBTS2α,* and *NtCPS2*) or transport (*NtLTP1*) of specialized metabolites secreted by mature glands, we wondered whether the leaf developmental stage could impact their expression in trichomes. Thus, glandular trichomes were isolated from leaves at different developmental stages, arbitrarily defined by the leaf length: < 2.5 cm (stage I), between 2.5 cm and 6.5 cm (stage II), between 6.5 cm and 15 cm (stage III), and > 15 cm (stage IV). The transcript levels of the six genes increased with leaf development (Fig. 2). The opposite trend was observed for elongation factor α (*EF1α*), which peaked at stage I, confirming that the observed increasing level of all six genes is not an artifact of the normalization method (Supplemental Fig. S2). While the transcript level of *NtLTP1* appeared stable during leaf development, expression of the other five genes steadily increased until stage III where it reached a plateau (Fig. 2). Among them, *NtRbcS-T1* was the gene for which the transcript level increased the most with leaf development (4-fold increase). Expression of *NtCBTS2α* and *NtCYP71D16* involved in the biosynthesis of cembrenes, the major subgroup of diterpenes produced by *N. tabacum*

glandular trichomes, also exhibited a large increase (3.8- and 3.6-fold, respectively) (Fig. 2). A more moderate increase was found for *NtMALD1* (2.6-fold) and *NtCPS2* (2.4-fold) transcripts, the latter being involved in the biosynthesis of another subgroup of diterpenes, namely labdanes.

The absolute expression levels of all six genes of interest was then determined in isolated trichomes for developmental stage III using an absolute standard curve method (Fig. 3, see Material and methods for details). Several control genes, some of which were used to normalize the relative expression data shown in Figures 1 and 2, were also added to the study for comparison purposes. Among control genes, the absolute expression levels (Fig. 3) were in agreement with previously published data in other Solanaceae species (Lacerda et al., 2015; Lu et al., 2012). Genes involved in cembrene production, *NtCBTS2α* (78.0 copies/pg), and *NtCYP71D16* (67.9 copies/pg), were the most expressed genes at stage III (Fig. 3), while a lower expression was found at this stage for *NtMALD1* (40.8 copies/pg)*, NtLTP1* (28.2 copies/pg)*, NtCPS2* (labdane *diterpenes,* 11.1 copies/pg) and *NtRbS-T1* (5.1 copies/pg).

To further confirm the trichome-specific expression pattern observed by RT-qPCR, we generated transcriptional reporter lines using a *GUS-VENUS* coding sequence. In the 2D gel analysis which led to the identification of trichome-specific proteins, two spots had been identified as trichome-specific RbcS (Laterre et al., 2017). At that time, only the *N. benthamiana* genome sequence was available and a RbcS transcription promoter (*pNbRbcS-T*) corresponding to the minor RbcS spot had been retrieved from this species and characterized (Laterre et al., 2017). Once the sequence of a *N. tabacum* genome became available, we identified *pNtRbcS-T1* (accession: MG493459.1) as the promoter of the gene corresponding to the major RbcS spot (NtRbcS-T1; accession: DV157962). The *pNtMALD1* promoter (accession: MG493458.1), corresponding to the NtMALD1 spot (accession: FS387666) was identified as well. The *GUS-VENUS* coding sequence was fused to *N. tabacum* genomic fragments of 1993 bp and 1974 bp upstream of the translation initiation codon of *NtRbcS-T1* and *NtMALD1*, respectively (Fig. 4). For the other genes, the previously published promoter regions, i.e. 985 bp (*NsCBTS2α*), 849 bp (*NtLTP1*), 1852 bp *(NtCYP71D16),* and 1448 bp (*NtCPS2*) (Choi et al., 2012; Ennajdaoui et al., 2010; Sallaud et al., 2012; Wang et al., 2002) were isolated and similarly fused to the *GUS-VENUS* coding sequence. These constructs were introduced in *N. tabacum* through *Agrobacterium tumefaciens*-mediated transformation. For each construct, 24 to 45 independent T_0 transgenic lines were generated and their GUS activity was monitored in leaves, stems, and roots of six-week-old transgenic plants after 16 h of incubation at 37°C to challenge the cell-type

specificity (Table 1). GUS expression patterns were then confirmed on T_1 lines. Several

observations can be made. i/ For each promoter, a large majority (83-100%) of the lines

displayed GUS activity in tall glandular trichomes (Table 1). This indicates that appropriate

cis-sequences required for expression in tall glandular trichomes are present in the sequences

fused to the reporter gene. ii/ Some lines displayed strict trichome specificity even after

extended GUS assay (see Table I as well as Figure 5 for *pNtRbcS-T1* and *pMALD1* as well as

for *pNtCYP71D16* as a control). As an exception, none of the *NtLTP1* lines displayed

trichome-specific GUS expression (Table I). This agrees with the RT-qPCR data (Fig. 1). iii/

In many lines, however, GUS activity was also found in other organs such as root, stem, leaf

and/or flower (Table 1). In this case, the expression profile was variable according to the line

(examples are displayed in Figure 6), probably as a consequence of the position effect (see

discussion).

Discussion

NtRbS-T1 and NtMALD1 extend the list of trichome-specific genes in Nicotiana tabacum

In this project, the tissue-specific expression of six *N. tabacum* genes, namely *NtLTP1*, *NtCYP71D16*, *NtCBTS2α*, *NtCPS2*, *NtRbS-T1*, and *NtMALD1*, was analyzed. Their trichome-specific expression at the transcript level had not yet been quantified and compared by RT-qPCR. We performed this comparison by RT-qPCR, using tall glandular trichome RNA as well as RNA of different plant organs. Except for *NtLTP1*, all these genes were found to be specifically expressed in tall glandular trichomes in *N. tabacum* (Fig. 1). Apart from *NtLTP1*, whose expression was almost constant during leaf development, that of the other five genes was lower at an early stage of leaf development and reached a maximum at stage III (Fig. 2), presumably when the specialized metabolism is fully operating. This is also true for *RbcS-T1* and this observation is in agreement with the hypothesis that in glandular trichomes, Rubisco recycles the CO_2 released by the specialized metabolism (Pottier et al., 2018).

These expression data may help choose appropriate trichome-specific promoters to drive the expression of a transgene for metabolic engineering purposes. Indeed, although *NtCYP71D16* and *NtCBTS2α* reach higher expression level in trichomes at stage III of leaf development, *NtCPS2* and *NtMALD1* promoters should lead to a more homogenous expression of transgenes among leaves at different developmental stages. *NtLTP1* was an exception since its transcripts were identified in leaf and stem tissues devoid of trichomes. Although the *NtLTP1* promoter was claimed to confer trichome-specific expression, examination of the GUS reporter activity reported by Choi et al. (2012) revealed some activity in other cell types than trichomes and the expression in the stem was not displayed. Our observations are in line with previously published semi-quantitative RT-PCR which showed that *NtLTP1* is expressed in different organs (Harada et al., 2010).

A variety of expression patterns was detected for the different transcriptional reporters

Analysis of *GUS-VENUS* reporter lines revealed that, in most of them, the six promoters drove gene expression in the head cells of tall glandular trichomes of *N. tabacum*. However, expression was rarely exclusively observed in trichomes (Table 1). Moreover, for a given reporter construct, variability in expression patterns was observed between independent lines. Concerning *NtCYP71D16*, Wang et al (2002) noted that after overnight incubation (like in the present work) with the GUS substrate, faint GUS staining was found in guard cells and veins

in some areas of some leaves, and at the cut edges of some leaf blades. No data was reported for the GUS reporter driven by the *NtCBTS-2α* and *NtCPS2* promoters in other organs than leaves (Ennajdaoui et al., 2010; Sallaud et al., 2012). Based on RT-qPCR data, *NtCYP71D16*, *NtCBTS2α*, *NtCPS2*, *NtRbS-T1*, and *NtMALD1* can, however, be considered as specifically expressed in trichomes. A possible reason for the partial discrepancy between RT-qPCR and gene reporter data is that the promoter regions that were used in the constructs are incomplete. This is unlikely to be the case for six promoters, especially for those that are close to 2 kb long (i.e. *pNtRbcS-T1*, *pNtMALD1*, and *pNtCYP71D16*), unless cis regulatory sequences are located downstream of the translation start. In addition, this hypothesis fails to explain the wide diversity in expression patterns observed between lines expressing the same construct. A more likely explanation is a position effect due to the random insertion of the T-DNA in the plant cell genome. Indeed, the genomic environment surrounding the integrated cassette (structure of chromatin, presence of enhancers/silencers near the insertion site) is known to alter the expression level and profile of transgenes (Hernandez-Garcia and Finer, 2014; Kohli et al., 2010). Between independent lines, and thus different insertion sites, those position effects might differ according to the proximal endogenous regulatory elements.

Integration of insulators in the vector, at both ends of the transgene (as well as between different expression cassettes within the construct in case of a multigene construct) could be a way to prevent undesirable effects (Biłas et al., 2016; Hasegawa and Nakatsuji, 2002). Indeed, insulators are sequences that stabilize gene expression by guaranteeing gene autonomy (Biłas et al., 2016; Hasegawa and Nakatsuji, 2002). Another option could be to insert the transgenes in a specific locus, preferentially one promoting high expression of genes (Abdel-ghany et al., 2015). Genome editing technologies such as CRISPR-Cas9, which allows insertions in a desired locus through homologous end joining, could be a way to cope with such an issue (Cao et al., 2016; Khatodia et al., 2016). Finally, a reporter gene coding for a fluorescent protein such as GFP could be included in the genetic construct to facilitate the screening of transformed plants in order to select transgenic lines that only express the reporter in glandular trichomes. If the same promoter is used to drive the expression of both the fluorescent reporter and the transgene(s) of interest, the rationale is that the fluorescence pattern and intensity of the reporter mirror those of the transgene(s) of interest.

It would have been interesting to confirm, in the reporter lines, the quantitative variations of activity between the different promoters. However, because of the position effect, quantitative variations do not necessarily reflect the actual strengths of the promoters. The

positive consequence of this variation is that transgenic plants can be screened to find those
that express the gene of interest at the appropriate level.

Cloning strategy for implementation of an entire metabolic pathway in *Nicotiana*
***tabacum* glandular trichomes**

The use of *N. tabacum* glandular trichomes for metabolic engineering purposes is a very
attractive strategy. However, heterologous expression of an entire metabolic pathway into
such specific plant organ may be a tedious process that poses several technical challenges.
This includes promoting cell type-specific expression at appropriate levels of multiple genes
(coding for enzymes or transporters) that segregate together in the progeny.

For this purpose, the use of several independent T-DNA to generate plants expressing an
entire metabolic pathway is questionable and an approach allowing delivery of multiple
genetic constructs in a unique vector appears to be a far better option. In this way, a single
DNA molecule is transferred into the plant cells, at the same locus, which limits the intergenic
variation in expression due to positional effects in the genome. Such an approach also ensures
that all genes cloned in this molecule are inherited together (Dafny-Yelin and Tzfira, 2007).

However, generating a single T-DNA vector bearing several genes each placed under the
same trichome-specific promoter can lead to a dilution effect since transcription might be
limited by the amount of available transcription factors recruited by cis-elements present in
this promoter (Biłas et al., 2016). To prevent such dilution effect, genes coding for each
enzyme could be placed downstream different trichome-specific promoters. The use of
different promoters will also prevent homologous recombination between similar promoters
in the construct and loss of the sequence in-between.

In addition to that, the level of expression is obviously an essential trait. Beside avoiding
dilution effects of trans elements, using different promoters may be needed to adapt the level
of expression to the efficiency of the enzyme encoded by the transgene. This explains why
identifying different trichome-specific gene promoters is so important. By identifying and
comparing different trichome-specific promoters such as *pNtCYP71D16*, *pNtCBTS2α*,
pNtCPS2, *pNtRbS-T1*, and *pNtMALD1*, our work provides the community with a useful
toolkit to express a gene in glandular trichomes at different levels.

In conclusion, a key and unique feature of glandular trichomes is their ability to synthesize
and secrete large amounts of a limited panel of specialized metabolites. Taking advantage of
the pool of natural precursors to produce specific metabolites in glandular trichomes by
metabolic engineering would therefore be of high biotechnological interest. This requires the

availability of transcriptional promoters specifically active in these structures that could be
used to efficiently drive the expression of the transgenes coding for the enzymes needed to
implement the pathway in a cell-type specific way. In this respect, the identification of the
NtMALD1 and *NtRbS-T1* promoters and their comparison with previously identified trichome-
specific promoters are promising tools for expressing entire biosynthesis pathways in
glandular trichomes of *N. tabacum*. However, use of trichome-specific promoters should
ideally be associated with the addition of insulators at the end of each expression cassette as
well as with site-directed genome edition technologies to ensure cell-type specific expression
at an appropriate level. In the long term, detailed knowledge of the gene network existing in
plant glandular trichomes will generate new leads to tap the largely unexploited potential of
these specialized plant organs to increase plant resistance to pests, or to lead to improved
production of important specialized metabolites via finely tuned metabolic engineering
approaches.
348

Material and methods

Plant material and plant growth conditions

Nicotiana tabacum cv Petit Havana SR1 (Maliga et al., 1973) plants were used in this work. For the *in vitro* cultures, seeds were sterilized by immersion in 1 ml 70% (v/v) ethanol for 1 min and then in 1 ml 50% (v/v) commercial bleach for 2 min. Seeds were then washed three times with 1 ml of sterile MilliQ water and kept at 4°C, in the dark, during 48 h for stratification. Sterilized seeds were sown on solid Murashige and Skoog (MS) medium [4.33 g.l^{-1} MS salts (MP Biochemicals, Solon, OH, USA; www.mpbio.com), 3% (w/v) sucrose, 1% (w/v) agar, pH 5.8 (KOH)] and placed in the growth chamber at 25°C under a 16 h photoperiod (50 μmol photon m^{-2} sec^{-1}). For the soil cultures, seeds were stratified before being sown in potting soil (DCM, Grobbendonk, Belgium; dcm-info.com). Isolated plantlets coming from potting soil or *in vitro* conditions were transferred to Jiffy pots (Gronud, Norway; www.jiffypot.com) before being transferred to bigger pots containing potting soil (DCM). Plants on soil were grown under controlled conditions, in a phytotron set at 25°C and with a 16 h photoperiod (300 μmol photon m^{-2} sec^{-1}).

Tissue isolation, RNA extraction and cDNA synthesis

Trichomes were removed from tissues of 6-week-old plants following the cold-brushing method (Wang et al., 2001). For the analysis of gene expression in trichomes during leaf development, trichomes were isolated from leaves at different developmental stages defined here by leaf length: < 2.5 cm (stage I), between 2.5 cm and 6.5 cm (stage II), between 6.5 cm and 15 cm (stage III), and > 15 cm (stage IV). For the analysis of gene expression in different tissues, we harvested roots, trichomes-free stems, trichomes-free leaves, and leaf trichomes (pool of leaves from stage I to stage III) from 6-week-old plants, and flowers from 10-week-old plants. For each biological replicate (except for isolated trichomes), 100 mg of material was pre-ground in liquid nitrogen using a mortar and pestle. Pre-ground tissues and isolated trichomes were ground in 2 mL Precellys tubes containing 200 μL of ceramic beads Zirmil (0.5 mm, Saint Gobain Zipro, Le Pontet, France) and 500 μL of lysis/2-Mercaptoethanol solution of the Spectrum^{TM} Plant Total RNA Kit (Sigma-Aldrich, St. Louis, Missouri, USA; http://www.sigmaaldrich.com). Samples were subjected to four consecutive 30 s grinding periods at 6,000 rpm using a Precellys 24 (Bertin Technologies, Montigny-le-Bretonneux, France). The homogenates were centrifuged at 1,000 *g* for 3 min (Eppendorf 5430, Hamburg,

Germany). The subsequent steps of the RNA extraction were performed on the supernatants

according to the manufacturer's specifications, except that the 56 °C incubation step was

omitted. RNA was eluted in 50 µl elution buffer and quantified using a spectrophotometer

(Nanodrop® ND-1000, Isogen Life Science, The Netherlands; www.isogen-lifescience.com).

Genomic DNA contamination was eliminated by using the On-Column DNase I Digestion Set

(Sigma-Aldrich, St. Louis, Missouri, USA; www.sigmaaldrich.com). The RNA was finally

flash frozen in liquid nitrogen and stored at -80°C. DNA-free RNA (500 µg) was used for

reverse transcription using the Moloney Murine Leukemia Virus Reverse transcriptase

(Promega, Madison, Wisconsin, USA; be.promega.com) and oligo(dT)$_{18}$. Reverse

transcription mixture was added according to the manufacturer's specifications. After adding

the transcriptase, samples were incubated for 5 min at 25°C, followed by 1 h at 42°C and 5

min at 85°C, placed on ice for 5 min, aliquoted, and stored at -20°C.

Gene expression

Gene-specific RT-qPCR primers listed in Table S1 were designed at the 3'end of the

coding sequence, (size, about 100 bp; melting temperature, 60°C) using OligoPerfect™

Designer (www.thermofisher.com). cDNA (5 µl, 17 fold diluted) was used as a template in 20

µl RT-qPCR reaction, which also contained 10 µl of qPCR master mix plus for SYBR Green I

(Eurogentec, Seraing, Belgium, https://secure.eurogentec.com/eu-home.html) and 5 µl of

primer mix (1.3 µM each). Amplification was performed on an ABI 7500 Real-Time PCR

system (Waltham, Massachusetts, USA; http://www.thermofisher.com). Primer specificity

was confirmed by analysis of the melting curves. For each tissue, primer amplification

efficiency (≥ 95%) was determined using five standards from serial dilutions of a cDNA pool

of the biological replicates used for gene expression analysis. Relative transcript levels were

calculated following the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001) with the geometric

mean of mitochondrial ATP-synthase β-subunit (*ATP2*) and ubiquitin (*UBQ*) used as

references for comparison between different tissues, and of *ATP2*, *UBQ*, and actin (*ACTIN*),

for comparison between different leaf developmental stages. For absolute quantification, PCR

products amplified by gene-specific RT-qPCR primers listed in Supplemental Table S1 were

cloned in pGEM-T Easy vector (Promega, Madison, Wisconsin, USA) prior to their

sequencing. Constructs were linearized by PstI restriction, purified using Nucleospin Extract

II kit (Macherey-Nagel, Düren, Germany) and rigorously quantified using a

spectrophotometer (Nanodrop® ND-1000, Isogen Life Science, The Netherlands;

www.isogen-lifescience.com). For each quantified purified linear plasmid, the copy number

was determined according to the following equation: copy number = (vector amount [g]) ×

6.023×10^{23} [molecules/mole]) / (660 [g/mole/base] × size of the vector+insert

[bases]. Absolute transcript levels were determined through the absolute standard curve

method. Thus, for each studied gene, standards (2.10^6, 2.10^5, 2.10^4, 2.10^3 copies) obtained by

serial dilution of the purified linear plasmids were included in duplicate in q-PCR plates used

to study gene expression during trichome development.

Generation of plants expressing promoter-*GUSVENUS* Fusions

The transcription promoter regions of *NtRbcS-T1* (1993 pb; GenBank accession:

MG493459.1) and *NtMALD1* (1974 pb; GenBank accession: MG493458.1) were identified

blasting the EST corresponding to *NtRbcS-T1* (GenBank accession: DV157962) and

NtMALD1 (GenBank accession: FS387666) coding sequences to the genome of *N.*

tabacum TN90 in the Solgenomics database (http://solgenomics.net). The promoter regions of

NsCBTS2α (985 bp; GenBank accession: HM241151.1), *NtLTP1* (849 bp; GenBank

accession: AB625593.1), *NtCYP71D16* (1852 pb; GenBank accession: AF166332.1), and

NtCPS2 (1448 bp; GenBank accession: HE588139.1) were defined as previously (Choi et al.,

2012; Ennajdaoui et al., 2010; Sallaud et al., 2012; Wang et al., 2002). Promoter regions were

amplified by PCR using as a template genomic DNA prepared from *N. tabacum* or *N.*

sylvestris leaves and the primers listed in Supplemental Table S2. The amplified fragments

were inserted in the pGEM®-T Easy Vector (Promega, Madison, Wisconsin, USA;

www.promega.com) and sequenced. Cloned fragments were cleaved using HindIII (or NotI

for *pNtMALD1* and *pNtRbcS-T1*) and KpnI, prior to their insertion in a pAUX3131 construct

(Navarre et al., 2011), upstream of the *GUSVENUS* coding sequence. The fusion construct

was excised using I-SceI and inserted into the pPZP-RCS2-nptII plant expression vector

(Goderis et al., 2002), also cut with I-SceI. The construct was introduced into *Agrobacterium*

tumefaciens LBA4404 virGN54D (van der Fits et al., 2000) for subsequent *N. tabacum* leaf

disc transformation (Horsch et al., 1986). The regenerated plants were finally transferred to

soil to be analyzed by GUS staining.

GUS histochemical analysis

Histochemical staining of plant tissues for GUS activity was conducted as described

previously (Bienert et al., 2012). To determine the tissue expression frequencies among

independent lines transformed with the same construct (Table 1), GUS staining was

performed for 16h on 6-week-old T0 plants growing in soil. Results were then confirmed in

the corresponding T1 plants of the same age except for flowers tissues, which were analyzed

on 10-week-old plants (Fig. 5 and Fig. 6). Stained tissues were washed with 70% ethanol for

chlorophyll extraction, transferred to 50% glycerol, observed under a light binocular (Carl

Zeiss MicroImaging) and photographed (Moticam 2300).

Statistical analysis

All tests were performed using the R software. For q-PCR, data were analyzed using

kruskal.test (Kruskal–Wallis) function for multiple comparisons. For multiple comparisons,

nparcomp package was used to perform Tukey post-hoc test when significant differences

were detected ($P < 0.05$). Different letters indicate significant differences between samples.

Supplementary data

Supplemental Fig. 1 Expression of *NtATP2* and *NtUBQ* control genes used to normalize the

data in Figure 1.

Supplemental Fig. 2 Expression of *NtEF1α* in trichomes according to leaf developmental

stage.

Supplemental Table S1 List of primers used for RT-qPCR.

Supplemental Table S2 List of primers used to amplify the promoter sequences.

References

References

Abdel-ghany, S. E., Golovkin, M., and Reddy, A. S. N. (2015). "Engineering of Plants for the
Production of Commercially Important Products: Approaches and Accomplishments," in
Plant Biology and Biotechnology, ed. B. Bahadur (New Delhi: Springer India), 551–577.
doi:10.1007/978-81-322-2283-5.

Bienert, M. D., Delannoy, M., Navarre, C., and Boutry, M. (2012). NtSCP1 from Tobacco Is
an Extracellular Serine Carboxypeptidase III That Has an Impact on Cell Elongation.
Plant Physiol. 158, 1220–1229. doi:10.1104/pp.111.192088.

Biłas, R., Szafran, K., Hnatuszko-Konka, K., and Kononowicz, A. K. (2016). Cis-regulatory
elements used to control gene expression in plants. *Plant Cell. Tissue Organ Cult.* 127,
269–287. doi:10.1007/s11240-016-1057-7.

Bouvier, F., Rahier, A., and Camara, B. (2005). Biogenesis, molecular regulation and function
of plant isoprenoids. *Prog. Lipid Res.* 44, 357–429. doi:10.1016/j.plipres.2005.09.003.

Cao, H. X., Wang, W., Le, H. T. T., and Vu, G. T. H. (2016). The power of CRISPR-Cas9-
induced genome editing to speed up plant breeding. *Int. J. Genomics.*
doi:10.1155/2016/5078796.

Choi, Y. E., Lim, S., Kim, H.-J., Han, J. Y., Lee, M.-H., Yang, Y., et al. (2012). Tobacco
NtLTP1, a glandular-specific lipid transfer protein, is required for lipid secretion from
glandular trichomes. *Plant J.* 70, 480–91. doi:10.1111/j.1365-313X.2011.04886.x.

Croteau, R., Kutchan, T. M., and Lewis, N. G. (2000). "Natural Products (Secondary
Metabolites)," in *Biochemistry & Molecular Biology of Plants*, eds. B. Buchanan, W.
Gruissem, and R. Jones (American Society of Plant Physiologists), 1250–1318.
doi:10.1201/b11003-3.

Dafny-Yelin, M., and Tzfira, T. (2007). Delivery of Multiple Transgenes to Plant Cells. *Plant
Physiol.* 145, 1118–1128. doi:10.1104/pp.107.106104.

Ennajdaoui, H., Vachon, G., Giacalone, C., Besse, I., Sallaud, C., Herzog, M., et al. (2010).
Trichome specific expression of the tobacco (Nicotiana sylvestris) cembratrien-ol
synthase genes is controlled by both activating and repressing cis-regions. *Plant Mol.
Biol.* 73, 673–85. doi:10.1007/s11103-010-9648-x.

Gershenzon, J., and Dudareva, N. (2007). The function of terpene natural products in the
natural world. *Nat. Chem. Biol.* 3, 408–414. doi:10.1038/nchembio.2007.5.

Goderis, I. J. W. M., De Bolle, M. F. C., François, I. E. J. A., Wouters, P. F. J., Broekaert, W.
F., and Cammue, B. P. A. (2002). A set of modular plant transformation vectors allowing

flexible insertion of up to six expression units. *Plant Mol. Biol.* 50, 17–27.

doi:10.1023/A:1016052416053.

Gwak, Y. S., Han, J. Y., Adhikari, P. B., Ahn, C. H., and Choi, Y. E. (2017). Heterologous

production of a ginsenoside saponin (compound K) and its precursors in transgenic

tobacco impairs the vegetative and reproductive growth. *Planta* 245, 1105–1119.

doi:10.1007/s00425-017-2668-x.

Harada, E., Kim, J.-A. A., Meyer, A. J., Hell, R., Clemens, S., and Choi, Y.-E. E. (2010).

Expression profiling of tobacco leaf trichomes identifies genes for biotic and abiotic

stresses. *Plant Cell Physiol.* 51, 1627–1637. doi:10.1093/pcp/pcq118.

Hasegawa, K., and Nakatsuji, N. (2002). Insulators prevent transcriptional interference

between two promoters in a double gene construct for transgenesis. *FEBS Lett.* 520, 47–

52. doi:10.1016/S0014-5793(02)02761-8.

Hernandez-Garcia, C. M., and Finer, J. J. (2014). Identification and validation of promoters

and cis-acting regulatory elements. *Plant Sci.* 217–218, 109–119.

doi:10.1016/j.plantsci.2013.12.007.

Horsch, R. B., Klee, H. J., Stachel, S., Winans, S. C., Nester, E. W., Rogers, S. G., et al.

(1986). Analysis of Agrobacterium tumefaciens virulence mutants in leaf discs. *Proc.*

Natl. Acad. Sci. U. S. A. 83, 2571–2575. doi:10.1073/pnas.83.8.2571.

Huchelmann, A., Boutry, M., and Hachez, C. (2017). Plant glandular trichomes: natural cell

factories of high biotechnological interest. *Plant Physiol.* 175, 6–22.

doi:10.1104/pp.17.00727.

Kempinski, C., Jiang, Z., Bell, S., and Chappell, J. (2015). "Metabolic engineering of higher

plants and algae for isoprenoid production.," in *Biotechnology of Isoprenoids*, eds. J.

Schrader and J. Bohlmann (Cham: Springer International Publishing), 161–199.

doi:10.1007/10_2014_290.

Khatodia, S., Bhatotia, K., Passricha, N., Khurana, S. M. P., and Tuteja, N. (2016). The

CRISPR/Cas Genome-Editing Tool: Application in Improvement of Crops. *Front. Plant*

Sci. 7, 1–13. doi:10.3389/fpls.2016.00506.

Kirby, J., and Keasling, J. D. (2009). Biosynthesis of Plant Isoprenoids: Perspectives for

Microbial Engineering. *Annu. Rev. Plant Biol.* 60, 335–55.

doi:10.1146/annurev.arplant.043008.091955.

Kohli, A., Miro, B., and Twyman, R. M. (2010). "Chapter 7 Transgene Integration,

Expression and Stability in Plants: Strategies for Improvements," in *Transgenic Crop*

Plants, ed. C. Kole et al. (Heidelberg: Springer-Verlag, Berlin), 201–237.

doi:10.1007/978-3-642-04812-8.

Lacerda, A. L. M., Fonseca, L. N., Blawid, R., Boiteux, L. S., Ribeiro, S. G., and Brasileiro,
A. C. M. (2015). Reference gene selection for qPCR analysis in tomato-bipartite
begomovirus interaction and validation in additional tomato-virus pathosystems. *PLoS*
One 10, 1–17. doi:10.1371/journal.pone.0136820.

Lange, B. M., Ahkami, A., Markus Lange, B., and Ahkami, A. (2013). Metabolic engineering
of plant monoterpenes, sesquiterpenes and diterpenes-current status and future
opportunities. *Plant Biotechnol. J.* 11, 169–96. doi:10.1111/pbi.12022.

Laterre, R., Pottier, M., Remacle, C., and Boutry, M. (2017). Photosynthetic Trichomes
Contain a Specific Rubisco with a Modified pH-Dependent Activity. *Plant Physiol.* 173,
2110–2120. doi:10.1104/pp.17.00062.

Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using
real-time quantitative PCR and the 2-ΔΔCT method. *Methods* 25, 402–408.
doi:10.1006/meth.2001.1262.

Lu, Y., Xie, L., and Chen, J. (2012). A novel procedure for absolute real-time quantification
of gene expression patterns. *Plant Methods* 8, 1–11. doi:10.1186/1746-4811-8-9.

Maliga, P., Sz-Breznovits, A., and Márton, L. (1973). Streptomycin-resistant plants from
callus culture of haploid tobacco. *Nat. New Biol.* 244, 29–30. Available at:
www.ncbi.nlm.nih.gov/pubmed/4515911.

Marienhagen, J., and Bott, M. (2013). Metabolic engineering of microorganisms for the
synthesis of plant natural products. *J. Biotechnol.* 163, 166–178.
doi:10.1016/j.jbiotec.2012.06.001.

Moses, T., and Pollier, J. (2013). Bioengineering of plant (tri) terpenoids: from metabolic
engineering of plants to synthetic biology in vivo and in vitro. *New Phytol.* 200, 27–43.
doi:10.1111/nph.12325.

Navarre, C., Sallets, A., Gauthy, E., Maîtrejean, M., Magy, B., Nader, J., et al. (2011).
Isolation of heat shock-induced Nicotiana tabacum transcription promoters and their
potential as a tool for plant research and biotechnology. *Transgenic Res.* 20, 799–810.
doi:10.1007/s11248-010-9459-5.

Pottier, M., Gilis, D., and Boutry, M. (2018). The Hidden Face of Rubisco. *Trends Plant Sci.*
23, 382–392. doi:10.1016/j.tplants.2018.02.006.

Sallaud, C., Giacalone, C., Töpfer, R., Goepfert, S., Bakaher, N., Rösti, S., et al. (2012).
Characterization of two genes for the biosynthesis of the labdane diterpene Z-abienol in
tobacco (Nicotiana tabacum) glandular trichomes. *Plant J.* 72, 1–17. doi:10.1111/j.1365-

313X.2012.05068.x.

Saxena, B., Subramaniyan, M., Malhotra, K., Bhavesh, N. S., Potlakayala, S. D., and Kumar,

S. (2014). Metabolic engineering of chloroplasts for artemisinic acid biosynthesis and

impact on plant growth. *J. Biosci.* 39, 33–41. doi:10.1007/s12038-013-9402-z.

Schilmiller, A. L., Last, R. L., and Pichersky, E. (2008). Harnessing plant trichome

biochemistry for the production of useful compounds. *Plant J.* 54, 702–11.

doi:10.1111/j.1365-313X.2008.03432.x.

Van Agtmael, M. A., Eggelte, T. A., and Van Boxtel, C. J. (1999). Artemisinin drugs in the

treatment of malaria: From medicinal herb to registered medication. *Trends Pharmacol.*

Sci. 20, 199–205. doi:10.1016/S0165-6147(99)01302-4.

van der Fits, L., Deakin, E., Hoge, J., and Memelink, J. (2000). The ternary transformation

system: constitutive virG on a compatible plasmid dramatically increases

Agrobacterium-mediated plant transformation. *Plant Mol. Biol.* 43, 495–502.

van Herpen, T. W. J. M., Cankar, K., Nogueira, M., Bosch, D., Bouwmeester, H. J., and

Beekwilder, J. (2010). Nicotiana benthamiana as a production platform for artemisinin

precursors. *PLoS One* 5, e14222. doi:10.1371/journal.pone.0014222.

Wagner, G. J., Wang, E., and Shepherd, R. W. (2004). New approaches for studying and

exploiting an old protuberance, the plant trichome. *Ann. Bot.* 93, 3–11.

doi:10.1093/aob/mch011.

Wang, B., Kashkooli, A. B., Sallets, A., Ting, H.-M. M. H.-M., de Ruijter, N. C. A. A.,

Olofsson, L., et al. (2016). Transient production of artemisinin in Nicotiana benthamiana

is boosted by a specific lipid transfer protein from A. annua. *Metab. Eng.* 38, 159–169.

doi:10.1016/j.ymben.2016.07.004.

Wang, E., Gan, S., and Wagner, G. J. (2002). Isolation and characterization of the CYP71D16

trichome-specific promoter from Nicotiana tabacum L. *J. Exp. Bot.* 53, 1891–1897.

doi:10.1093/jxb/erf054.

Wang, E., Wang, R., DeParasis, J., Loughrin, J. H., Gan, S., and Wagner, G. J. (2001).

Suppression of a P450 hydroxylase gene in plant trichome glands enhances natural-

product-based aphid resistance. *Nat. Biotechnol.* 19, 371–374. doi:10.1038/86770.

Yoon, J. M., Zhao, L., and Shanks, J. V (2013). Metabolic Engineering with Plants for a

Sustainable Biobased Economy. *Annu. Rev. Chem. Biomol. Eng.* 4, 211–37.

doi:10.1146/annurev-chembioeng-061312-103320.

Table 1 Frequency of GUS activity in different organs of transgenic *N. tabacum* lines showing expression in tall glandular trichomes[a].

PROMOTERS	SGT[b]	LEAVES	ROOTS	STEMS	ONLY IN TGT
pNtRbcS-T1	11/20	18/20	10/20	16/20	2/20
pNtMALD1	10/26	20/26	21/26	22/26	1/26
pNsCBTS2α	7/44	29/44	27/44	41/44	2/44
pNtCYP71D16	12/32	21/32	15/32	25/32	3/32
pNtCPS2	21/26	17/26	16/26	24/26	1/26
pNtLTP1	18/23	23/23	22/23	23/23	0/23

[a] The GUS expression frequency was determined as indicated in the Material and methods among independent lines of transgenics plants expressing the *GUS-VENUS* reporter under the control of the indicated promoters.

[b] GUS activity was observed in tall glandular trichomes (TGT), short glandular trichomes (SGT), leaves (besides trichomes), roots, and stems (besides trichomes) of 6-week-old T0 plants.

Figure legends

Fig. 1 Transcript levels in different organs of *N. tabacum*. Normalized transcript levels were determined as indicated in the Material and methods on the indicated organs. Results are shown as mean ± SD of three to five repeats. Different letters indicate significant differences according to a Kruskal-Wallis test ($p < 0.05$) followed by a Tukey post hoc test.

Fig. 2 Transcript levels in trichomes isolated from *N. tabacum* leaves at different developmental stages. Normalized transcript levels were determined as indicated in the Material and methods. St: leaf developmental stage. Stage 1: leaf length < 2.5 cm; stage II: leaf length between 2.5 cm and 6.5 cm; stage III: leaf length between 6.5 cm and 15 cm; stage IV: leaf length > 15 cm. Results are shown as mean ± SD of three repeats. Different letters indicate significant differences according to a Kruskal-Wallis test ($p < 0.05$) followed by a Tukey post hoc test.

Fig. 3 Absolute transcript levels at stage III of leaf development in *N. tabacum*. Absolute transcript levels were determined as indicated in the Material and methods. Results are shown as mean ± SD of three repeats. Different letters indicate significant differences according to a Kruskal-Wallis test ($p < 0.05$) followed by a Tukey post hoc test.

Fig. 4 Molecular constructs used to generate transgenic *N. tabacum* expressing the *GUSVENUS* reporter gene under the control of trichome-specific promoters. The transcription promoter regions of *NtRbcS-T1* (MG493459.1), *NtMALD1* (MG493458.1), *NsCBTS2α* (HM241151.1), *NtLTP1* (AB625593.1), *NtCYP71D16* (AF166332.1), and *NtCPS2* (HE588139.1) were amplified and cloned as described in the Material and methods.

Fig. 5 Specific GUS activity in trichomes of *N. tabacum.* GUS staining was performed in 6-week-old T1 lines.

Fig. 6 Diversity of GUS activity of different transgenic *N. tabacum* lines in leaf (A), stem (B), and root (C) tissues. GUS staining was performed in 6-week-old T1 lines.

Fig. 1 Transcript levels in different organs of *N. tabacum*. Normalized transcript levels were determined as indicated in the Material and methods on the indicated organs. Expression was measured as described in the Material and methods Results are shown as mean ± SD of three to five repeats. Different letters indicate significant differences according to a Kruskal-Wallis test (p < 0.05) followed by a Tukey post hoc test.

Fig. 2 Transcript levels in trichomes isolated from *N. tabacum* leaves at different developmental stages. Normalized transcript levels were determined as indicated in the Material and methods. St: leaf developmental stage. Stage 1: leaf length < 2.5 cm; stage II: leaf length between 2.5 cm and 6.5 cm; stage III: leaf length between 6.5 cm and 15 cm; stage IV: leaf length > 15 cm. Results are shown as mean ± SD of three repeats. Different letters indicate significant differences according to a Kruskal-Wallis test (p < 0.05) followed by a Tukey post hoc test.

Fig. 3 Absolute transcript levels at stage III of leaf development in *N. tabacum*. Absolute transcript levels were determined as indicated in the Material and methods. Results are shown as mean ± SD of three repeats. Different letters indicate significant differences according to a Kruskal-Wallis test ($p < 0.05$) followed by a Tukey post hoc test.

pNtRbcS-T1 (1993 pb)	*GUSVENUS* coding sequence
pNtMALD (1974 pb)	*GUSVENUS* coding sequence
pNtCYP71D16 (1852 pb)	*GUSVENUS* coding sequence
pNsCBTS2α (985 pb)	*GUSVENUS* coding sequence
p*NtLTP1* (849 pb)	*GUSVENUS* coding sequence
p*NtCPS2* (1448 pb)	*GUSVENUS* coding sequence

Fig. 4 Molecular constructs used to generate transgenic *N. tabacum* expressing the *GUSVENUS* reporter gene under the control of trichome-specific promoters. The transcription promoter regions of *NtRbcS-T1* (MG493459.1), *NtMALD1* (MG493458.1), *NsCBTS2α* (HM241151.1), *NtLTP1* (AB625593.1), *NtCYP71D16* (AF166332.1), and *NtCPS2* (HE588139.1) were amplified and cloned as described in the Material and methods.

Fig. 5 Specific GUS activity in trichomes of *N. tabacum.* GUS staining was performed in 6-week-old T1 lines.

Fig. 6 Diversity of GUS activity of different transgenic *N. tabacum* lines in leaf (a), stem (b), and root (c) tissues. GUS staining was performed in 6-week-old T1 lines.