The desmosomal cadherin Desmogon is
necessary for the structural integrity of the
Medaka notochord

Ali Seleit[1,2,#], Karen Gross[1,2], Michaela Woelk[1], Camilla Autorino[1], Lazaro
Centanin[1,#].

1 Animal Physiology and Development, Centre for Organismal Studies (COS)
Heidelberg, Im Neuenheimer Feld 230, Heidelberg 69120, Germany.

2 The Hartmut Hoffmann-Berling International Graduate School of Molecular and
Cellular Biology (HBIGS), University of Heidelberg, Heidelberg, Germany.

authors for correspondence: AS, ali.seleit@cos.uni-heidelberg.de and LC,
lazaro.centanin@cos.uni-heidelberg.de

Running Title: *desmogon* is necessary for notochord integrity

Abstract

The notochord is an embryonic tissue that acts as a precursor to the spine. It is composed of outer sheath cells and inner vacuolated cells. Together they ensure the ability of the notochord to act as a hydrostatic skeleton until ossification begins. To date, there is still a paucity in our understanding of how the notochord cell types are specified and the molecular players controlling both their formation and maintenance remain poorly understood. Here we report that *desmogon,* a desmosomal cadherin, is essential for proper vacuolated cell shape and therefore correct notochord morphology. We trace *desmogon+* precursors and uncover an early developmental heterogeneity that dictates the balance of vacuolated and sheath cell formation. We demonstrate that the growth of vacuolated cells occurs cell autonomously and reveal the presence of distinct injury sensing mechanisms in the notochord. Additionally, using a small-scale F0 CRISPR screen we implicate uncharacterized genes in notochordal integrity.

Introduction

The notochord is the defining characteristic that unites all chordates (Lim et al. 2017; Satoh et al. 2012; Stemple 2005; Stemple et al. 1996; Corallo et al. 2018). Developmentally, it derives from the dorsal organizer region in vertebrate embryos (Stemple 2005; Ellis et al. 2013; Corallo et al. 2018). Subsequently, its constituent cells adopt a mesodermal fate and undergo convergent-extension movements (Satoh et al. 2012; Tada & Heisenberg 2012; Corallo et al. 2018; Stemple 2005; Stemple 2004). This results in a tube-like structure that runs along the anterior-posterior axis while simultaneously delimiting the dorso-ventral axis (Stemple 2005). In addition to providing structural support to embryos by acting as the major skeletal element during embryonic development (Stemple et al. 1996; Stemple 2005), it also has important signalling roles (Yamada et al. 1991; Yamada et al. 1993; Pourquie et al. 1993; Hebrok et al. 1998; Fouquet et al. 1997; Corallo et al. 2018; Stemple et al. 1996; Stemple 2005; Satoh et al. 2012). Indeed, patterning of adjacent tissues by the notochord is essential for correct morphogenesis to occur (Yamada et al. 1991; Yamada et al. 1993; Pourquie et al. 1993; Hebrok et al. 1998; Fouquet et al. 1997; Talbot et al. 1995; Stemple et al. 1996; Fleming 2004; Corallo et al. 2018). In vertebrates the notochord is a transient structure and a precursor to spine formation, it is eventually almost entirely replaced by vertebrae (Fleming 2004; Stemple 2005; Corallo et al. 2018; Lopez-Baez et al. 2018; Gray et al. 2014). Recently, it has been shown that correct spine patterning relies on segmentation cues present in the embryonic notochord (Wopat et al. 2018; Lleras Forero et al. 2018). This and other observations has led to the understanding that correct notochord morphogenesis is essential for normal spine formation (Fleming 2004; Lim et al. 2017; Wopat et al. 2018; Lleras Forero et al. 2018; Gray et al. 2014).

The notochord of all vertebrates studied to date (Ellis et al. 2014; Corallo et al. 2018) is formed of two cell-types: an outer sheath epidermal-like cell population and an inner vacuolated cell core (Stemple 2005; Corallo et al. 2018). The outer sheath cells cover the notochord tube and secrete ECM components that help in building and maintaining the peri-notochordal membrane

(Ellis et al. 2013; Corallo et al. 2018; Yamamoto et al. 2010; Lim et al. 2017), while the inner-

cells have large lysosomally-derived vacuoles that can withstand high hydrostatic pressure

(Ellis et al. 2013; Ellis et al. 2014). This ability to act as a hydrostatic skeleton is particularly

important in teleost fish as the strength and flexibility of the notochord is essential for proper

locomotion – which in turn is necessary for survival – as embryogenesis concludes (Stemple

et al. 1996; Ellis et al. 2014; Jiang & Smith 2007). A number of mutants affecting overall

notochord formation and differentiation have previously been described in ENU screens

(Stemple et al. 1996; Talbot et al. 1995). More recently, new players have been uncovered

that are important for vacuolated cell formation and maintenance (Lim et al. 2017; Garcia et

al. 2017). However, many decades after the first description of the notochordal cell types, there

is still a paucity in our understanding of the molecular players that control the correct formation

and maintenance of vacuolated and sheath cells and by extension the structural integrity of

the notochord (Ellis et al. 2013).

Here we uncover new regulators of correct notochord morphogenesis in vertebrates.

Desmogleins are a conserved family of calcium-binding cadherin transmembrane proteins.

They localize to cellular membranes and are important for establishing strong cell-cell contacts

and maintaining tissue integrity (Garrod & Chidgey 2008a). Structurally, they are part of the

intercellular desmosome junctions (Garrod & Chidgey 2008a) and are expressed in tissues

that undergo significant mechanical strain (Delva et al. 2009). We characterize the role of a

fish-specific Desmoglein, hereafter referred to as *desmogon*, that is expressed in the

notochord of Medaka and is to our knowledge the first Desmoglein reported to be present in

chordate notochords. The loss of *desmogon* causes vacuolated cell defects and leads to

structural deformities in the notochord. We generated a Tg(*desmogon*:EGFP) to address the

origin and growth dynamics of vacuolated cells by 4D-imaging, a process revealed to be

incremental and locally uncoordinated; vacuolated cells behave like autonomous units.

Interestingly, *desmogon* also labels early disc-shaped notochord precursors that constitute a

bipotent population. When analysed in 4D at the single-cell level, however, we reveal that each

progenitor is unipotent and generates either one vacuolated cell or a number of sheath cells.

In addition, exploiting the stable labelling of vacuolated cells in our Tg(*desmogon*:EGFP), we

uncover two distinct types of regeneration responses in Medaka notochords, one spatially

localized and the other global, that depend on the type of injury sustained. Finally, we use

Tg(*desmogon*:EGFP) to carry out a small scale reverse-genetics screen on highly conserved

and uncharacterized genes enriched in the notochord (Briggs et al. 2018; Farrell et al. 2018).

Using this fast and straight-forward methodology we were able to implicate new players in

correct notochord morphology in a vertebrate model. This work and approach, we believe,

could lead to a deepening of our understanding of the origin of spine and vertebral defects.

Results

***desmogon* is a fish-specific desmosomal cadherin expressed in the notochord**

While searching for a stable marker for neuromast border cells (Seleit et al. 2017) we serendipitously came across a novel uncharacterized *desmog-2-like* gene (ENSORLG00000017110), which we named *desmogon*. The 5.3Kb long transcript of *desmogon* is distributed over 14 exons and encodes a protein with at least 3 desmosomal cadherin domains and one cytoplasmic cadherin domain (Supplementary Figure 1). Based on the amino acid sequence the expected sub-cellular localization is plasma membrane and it is predicted to function as a component of the inter-cellular desmosome junctions. A list of all known orthologues of *desmogon* suggests that this gene is fish-specific, as it is absent in all other sequenced chordates (materials and methods for details) (Supplementary Table 1 and 2). Among fish, the *desmogon* locus is conserved in the vast majority of teleost branches although interestingly, it seems to have been lost in Zebrafish and Tetraodon (as evidenced by the syntenic conservation of the surrounding genomic region) (Supplementary Figure 1). *In situ* hybridization showed *desmogon* to be highly expressed in the developing notochord of Medaka (Figure 1A). To gain a better understanding of the dynamic spatial expression of *desmogon* we generated the Tg(*desmog*:EGFP) (Figure 1B-F) by using a 2.2kb proximal promoter region that contained strong peaks of H3K4 methylation (Supplementary Figure 1). Confocal analysis of mosaic, injected *desmogon*:EGFP and of Tg(*desmogon*:EGFP) medaka embryos revealed EGFP expression in the developing notochord throughout embryogenesis (Figure 1B-F, Supplementary Movies 1 & 2), this expression persists in adult fish in a segmented pattern along the spine (data not shown). Within the notochord, *Desmogon* labels vacuolated cells *(*Figure 1 C, F, arrows in Figure 1E, 1F) and a proportion of covering sheath cells *(*yellow asterisks in Figure 1E, 1F). The expression of a Desmoglein family member in vacuolated and sheath cells suggests the presence of desmosomes in both cell-types, therefore we followed an electron microscopy (EM) approach to characterise the notochord of 10 dpf *wild-type* medaka larvae at the sub-cellular level. Previous studies reported the existence of caveolae in the cellular membrane of vacuolated cells in the zebrafish notochord

(Nixon et al., 2007; Lim et. al, 2017), which we confirmed is also present in medaka (Figure

1G, G'). Additionally, we observed the occurrence of desmosomes mediating the physical

association of neighboring vacuolated cells (Figure 1G, G'), presumably to enhance their inter-

cellular adhesion capacities. Desmosomes were also found connecting sheath to vacuolated

cells (Figure 1H, I), and sheath to sheath cells (Figure 1I). Altogether, our results reveal the

expression of an uncharacterised Desmoglein family member in the two cell types of the

notochord that concurrently display desmosomes on their cellular membranes.

Vacuolated cells grow as autonomous units

The Tg(*desmog*:EGFP) allowed us to follow the axial extension of the notochord as the embryo

develops. Notochord expansion was mediated by the growth of vacuolated cells that not only

increase in size but also change their circular morphology into a more oblique shape as the

notochord matures (Figure 2A, B) (n>10). Using single plane illumination microscopy (SPIM)

(Krzic et al., 2012) we observed that vacuolated cells grow anisotropically over time (Figure

2C-E, and Supplementary Figure 2) in a process that appears to be irreversible (N>50 cells).

Our 4D data also suggests that *desmogon+* vacuolated cells in medaka are post-mitotic, since

we have not detected dividing vacuolated cells in any of our imaging (N=3 embryos at 4 dpf,

and N=2 embryos at 3dpf image for 24h, N>50 cells) (Supplementary movie 3). Remarkably,

our analysis revealed that vacuolated cells increment their size in a locally uncoordinated

manner, where even neighbouring vacuolated cells grow at different rates (Figure 3C-E',

Supplementary Movie 4). These results demonstrate that vacuolated cells grow as

autonomous units. We next wondered whether the Medaka notochord grows in size just by the

increase in volume of vacuolated cells, or if in addition new vacuolated cells are being

incorporated. We observed *desmogon+* precursor cells differentiating into vacuolated cells

(Figure 2D-E', Supplementary Movies 5 & 6) in a process that occurred along the notochord.

The formation of new vacuolated cells was more frequently observed at the posterior tip of the

extending medaka notochord (Figure 2E-E'', Supplementary Movie 6) (Total N=8 cells in 2

independent embryos). Our data indicates that during axial extension the number of

vacuolated cells increases in the notochord, however, this is not the only cell-type that needs

to be amplified during notochord extension. We therefore aimed to better resolve the cellular

processes leading to the incorporation of new vacuolated and sheath cells as the notochord

grows.

Behavioural heterogeneity of notochord precursor cells

In addition to labelling vacuolated and sheath cells in medaka, the (*desmogon*:EGFP) shows

expression in early, undifferentiated notochord precursors known as disc-shaped-progenitors

(Figure 1B, Figure 3A and Supplementary Movie 7). The population of disc-shaped-progenitors

was previously shown to generate both differentiated cell types (Yamamoto et al., 2010), and

therefore constitutes a bi-potent population. While analysing sparsely labelled notochords of

medaka larvae injected with *desmogon*:EGFP (Figure 3B-C), we noticed that clusters tended

to contain either vacuolated or sheath cells (N=41 sheath cells clusters, N=39 vacuolated cell

clusters, N=80/93 cell-type specific cluster and N=13/93 clusters containing both cell types,

N=93 clusters in 26 mosaic larvae). These results suggest that under physiological conditions

disc-shaped progenitors might be bi-potent as a population, but fate-restricted as individual

cells. To gain a finer spatial/temporal resolution on the formation of vacuolated and sheath

cells in the notochord, we decided to inject the *desmogon*:EGFP construct into zebrafish

embryos. Even though a *desmogon* orthologue is not present in this species, we could observe

labelled vacuolated and sheath cells in *desmogon*:EGFP zebrafish injected embryos

(Supplementary Figure 3, and Supplementary Movie 8), indicating that the transcriptional

machinery driving expression of medaka *desmogon* is conserved in distantly related teleosts.

Live-imaging on embryos with EGFP+ clones in the notochord revealed that the growth of

vacuolated cells is incremental but also locally uncoordinated in Zebrafish (n=6 embryos n>10

vacuolated cells) (Supplementary movie 8). Focusing on earlier stages of notochord

development, we were able to clearly observe vacuolated and sheath cell formation from

undifferentiated *desmogon*+ notochord precursor cells. These disc-shaped precursors were

located primarily but not exclusively at the growing posterior tip of the notochord. We observed

two distinct, mutually exclusive cellular behaviours in *desmogon+* precursors. Either they

directly generate a single vacuolated cell (Supplementary Movie 9, Supplementary Figure 3,

magenta asterisk) (N=9 cells), or they undergo a dorso-ventral symmetric division leading to

the formation of sheath cells (Supplementary Movie 9, Supplementary Figure 3, yellow

asterisk) (n>10 cells). This behavioural heterogeneity does not seem to be temporally

controlled or governed by morphogenetic gradients, since both cellular outcomes can occur

simultaneously in neighbouring precursors. Our data shows that disc-shaped *desmogon+*

precursors can directly differentiate into vacuolated cells that will not undergo mitosis

throughout our imaging (n>10 cells in=6 embryos), while the formation of sheath cells

necessitates a symmetric division that could be followed by further amplifying rounds of

divisions (n>10 cells in n=6 embryos). Interestingly, we consistently observed that disc-shaped

precursors contained a detectable vacuole which was expanded during vacuolated cell

differentiation, and lost during the maturation of sheath cells. Indeed, directly after the dorso-

ventral symmetric division, a proportion of newly formed sheath cells do contain small vacuoles

(n>10 cells in n=6 embryos) (Supplementary Movie 10), suggesting that this feature could

reflect an earlier developmental state that must be repressed during the maturation of sheath

cells. Although phylogenetically distantly related within the teleost branch, the results from

Zebrafish mirrored our findings in Medaka, particularly the non-mitotic nature of desmogon+

vacuolated cells. Our results indicate that in both species disc-shaped precursors are

exhausted by the end of notochordal development, which raises the question of whether and

how vacuolated cells can be replaced after injury of mature Medaka notochords.

201
202   **Vacuolated cell loss triggers a robust and localized regeneration response in medaka**
203
Spatially targeted and precise multi-photon laser ablation of 6-10 vacuolated cells of 5-6 dpf

Tg*(desmogon:*EGFP*)* embryos resulted in the specific loss of cells in the area of injury (Figure

4A, Supplementary movie 11 and 12). Both sheath and vacuolated cells outside the ablated

zone retain a normal morphology and the overall integrity of the notochord is unaffected. We

next assessed whether a regeneration response can occur in the ablated area. Indeed, 48-

hours post injury we observed the appearance of small *desmogon+* vacuolated cells

specifically in the area of injury (Figure 4B, Supplementary movie 13) (n=8 embryos). Interestingly, the overall morphology, presence of a small vacuole, and EGFP+ expression of these cells was highly reminiscent of the earliest vacuolated sheath cells we observed during development in Zebrafish notochords, suggesting that the source of new vacuolated cells is the same in both species (Garcia et al. 2017). The small vacuolated cells grow in size over time as assessed at 5 days post injury (Figure 4C, Supplementary movie 14) (n=8 embryos), this growth followed the same autonomous rationale we observed under physiological conditions in Medaka. Overall, our results indicate that Medaka notochords can mount robust regeneration in response to vacuolated cell loss. This occurs in a highly spatially localized manner that is restricted to the initial injury site.

Peri-notochordal membrane injury triggers a global regeneration response

The perinotochordal membrane is a thick ECM layer that ensheathes the notochord and helps to maintain its integrity. To test the effect of a sudden loss of hydrostatic pressure within the notochordal tube, we ablated the lower lining of the peri-notochordal membrane (Figure 4D, Supplementary movie 15). Two days post-injury we observed *desmogon+* cells leaking outside of the notochord tube, indicating that the lower lining of the notochordal membrane failed to be repaired. This in turn led to a loss of notochord morphology and the appearance of small vacuolated cells no longer became restricted to the injury site but were rather spread over the entire tube (Figure 4E). Interestingly, perinotochordal injury and the concomitant lack of repair triggered a global regenerative response even in the absence of vacuolated cell ablation anterior and posterior to the targeted site. Five days post-injury the leakage of notochord vacuolated cells continued and the morphology of the notochord was heavily perturbed. We observed a cluster of *desmogonGFP+* vacuolated cells ventral to the notochord tube at the site of the original injury, and small vacuolated cells persisted anterior and posterior to the injury site (Figure 4F) (n=3). Overall, we conclude that injury to the peri-notochordal membrane cannot be repaired efficiently in Medaka. This leads to cell leakage and a perturbed notochord morphology, which in turn triggers a global regeneration response that is not spatially restricted to the initial injury site.

Desmogon crispants and stable mutants exhibit notochordal lesions of collapsed vacuolated cells

To test if _desmogon_ has any functional role during notochord morphogenesis we designed 3 gRNAs targeting different exons along its coding sequence. CRISPR/Cas9 injection into Tg(_desmogon:_EGFP) resulted in lesions containing collapsed vacuolated cells along the length notochord, which were absent in embryos injected with control _oca2_ gRNAs (Figure 5 A-C, Supplementary Movies 16, 17) (Lischik et al. 2018). The vast majority of injected embryos survived until stage 42 and no obvious pleiotropic effects were detected, suggesting the effects observed were notochord specific. Importantly, a proportion of F0 injected embryos that survived to adulthood showed strong signs of bending and defective spine formation (data not shown). The described phenotypes for the F0 injected _desmogon_ crispants were consistently recapitulated in _desmogon_ mutants (Figure 5D, Supplementary Figure 5, Supplementary Movies 18, 19). To gain a better understanding of the structural phenotypes we decided to perform electron microscopy (EM) on _desmogon_ mutants. Longitudinal sections on wild-type notochords were characterised by the typical highly ordered array of vacuolated cells (Figure 5E). This contrasts with the structural disorganisation present in lesioned areas of the _desmogon_ mutant notochords (Figure 5F-H). EM data also showed the presence of vacuolated cells of appreciably different sizes (Figure 5F, arrows), evidence of vacuolated cell collapse (Figure 5F-G'), and invasion of sheath cells into the central notochord tube that indicates the possible triggering of a regenerative response in the lesioned area (Figure 5F asterisk and Figure 5H, Supplementary Figure 4). Stronger _desmogon_ phenotypes included the loss of notochord integrity and shape as revealed by kinking and buckling along the notochord tube (Figure 6A-C). A closer analysis of the phenotypes affecting embryos with collapsed notochords revealed the presence of larger lesions that were consistently filled with _desmogon_+ small vacuolated cells (Supplementary Movie 18 & 19). In addition to being a marker for vacuolated cells, we therefore believe that _desmogon_ has a functional role in proper notochord development and shape. Its loss leads to vacuolated cell collapse and the appearance of lesions that contain small vacuolated cells, this can in turn lead to gross morphological defects in the notochord.

Targeted CRISPR screen uncovers novel regulators of notochord integrity

Realizing the concurrence of F0 CRISPR phenotypes with the ones observed for the stable

mutant line, as recently reported for other systems (Wu et al. 2018; Lischik et al. 2018;

Trubiroha et al. 2018), we decided to use the Tg*(desmogon:*EGFP*)* as a fast and straight-

forward read-out for notochord defects. We searched for putative regulators of notochord

integrity by using the recently generated single-cell transcriptome data from Zebrafish (Briggs

et al. 2018; Farrell et al. 2018). We selected a number of well annotated and poorly

characterized genes that were conserved across vertebrates and highly expressed during

Zebrafish notochord morphogenesis (*arrdc3a, kcnk6, pmp22b, si:dkey- 261h17.1 and vgll2b*).

We then performed a small-scale F0 CRISPR screen (Wu et al. 2018) in Tg*(desmogon:*EGFP*)*

embryos by targeting each selected gene with 2 gRNAs (for details of selection criteria see

materials and methods). For *vgll2b* this resulted in 55% of injected embryos showing

morphological defects in notochord shape and integrity including twisting and bending of the

notochord tube (Figure 6D, F and Table 1). Targeting *arrdc3a* resulted in 30% of injected

embryos showing disruption in notochordal integrity, with strong phenotypes including buckling

and kinking (Figure 6E, F, and Table 1). Targeting *kcnk6, si:dkey- 261h17.1, pmp22b* with the

same approach resulted in strong phenotypes on the notochord but additionally all three

showed significant pleiotropic effects (Stemple et al. 1996), including general growth

retardation, shorter body axes and gross morphological defects (Table 1). It is therefore more

difficult to assign causal phenotypes to these genes. It is clear, however, that they directly or

indirectly affect correct notochord morphology. In conclusion, using a straight-forward reverse-

genetics approach we have uncovered novel and conserved regulators of notochord

morphogenesis and maintenance in vertebrates.

Discussion

Desmogleins are a conserved family of desmosomal cadherins that localize to the plasma membrane of cells (Garrod & Chidgey 2008b; Delva et al. 2009). They are a constituent part of desmosomes and are important mediators of strong inter-cellular adhesion. Indeed, desmosomes have been shown to be expressed in cell types that operate under significant mechanical strain (Garrod & Chidgey 2008b; Delva et al. 2009). Loss of desmosomal function leads to disruption of tissue integrity (Garrod & Chidgey 2008a). The notochord is a tissue that is constantly assailed by strong mechanical stresses (Lim et al. 2017; Garcia et al. 2017; Corallo et al. 2018). It is able to withstand those pressures primarily because of its structural organization: large vacuolated cells on the inside of the tube are surrounded by a strong peri-notochordal membrane formed by sheath cells (Lim et al. 2017; Yamamoto et al. 2010; Koehl et al. 2000; Adams et al. 1990). Our study on *desmogon,* a fish-specific desmosomal cadherin, is the first showing a desmoglein family member expressed and functional in vertebrate notochords.

desmogon is necessary for correct notochord morphology in Medaka

The structural demands on teleost notochords are particularly high given that larvae need to swim and feed as soon as embryogenesis concludes (Jiang & Smith 2007; Stemple et al. 1996; Ellis et al. 2014). Significantly, this happens before the ossification and formation of spines (Corallo et al. 2018; Lleras Forero et al. 2018; Wopat et al. 2018; Fleming 2004; Gray et al. 2014). This could explain the specific allocation of this desmoglein family member expression to the notochord. To test whether *desmogon* has a functional role in the Medaka notochord we targeted it by CRISPR/Cas9 and observed flattened vacuolated cells and lesions along the length of the notochord. This could either be a sign of collapsed vacuolated cells (Lim et al., 2017) or a failure of vacuolated cells to properly form. Our EM data on lesioned mutant notochords strongly suggests that the phenotype results from the local collapse of vacuolated cells. Interestingly, in areas that contained larger lesions (both in injected fish and in stable

mutants), we consistently observed the appearance of small vacuolated cells. This is highly

reminiscent of results we report from the regeneration experiments and suggests that the

collapse of *desmogon* mutant vacuolated cells can trigger a regenerative response. Indeed

EM results on lesioned notochords shows evidence of invading sheath cells. Our results

indicate that *desmogon* is necessary for the maintenance and/or growth, but not the formation,

of vacuolated cells. Additionally, we observed a strong correlation between the size of the

mutant lesion and the structural integrity of the notochord, bigger lesions invariably led to

buckling and kinking of the notochord tube. Our results are in line with previous observations

that have shown that correct vacuolated cell morphology is essential for the notochord to

withstand the high mechanical stresses it faces (Lim et al. 2017; Ellis et al. 2013; Garcia et al.

2017; Fleming 2004; Adams et al. 1990). Failure of properly building up and maintaining the

high hydrostatic pressure inside the notochord tube leads to bending and buckling along the

length of the notochord (Ellis et al. 2013; Corallo et al. 2018). Our results demonstrate that the

presence of *desmogon* in the Medaka notochord is necessary for correct vacuolated cell shape

and by extension proper notochord morphology and integrity.

The fact that *desmogon* is present in the vast majority of teleost branches argues that it has a

conserved role in notochord maintenance in fish. However, there are two intriguing exceptions,

desmogon has no detectable orthologues in Tetraodon or in Zebrafish. It seems likely that both

fish have lost *desmogon* through the course of their evolution, this interpretation is supported

by the fact that we can identify syntenic genomic regions that do not contain *desmogon.* It

remains a formal possibility, however, that an orthologue with a highly divergent nucleotide

sequence exists in Zebrafish and Tetraodon. It would be interesting to see whether other

desmosomal cadherins have taken over the notochordal role of *desmogon* in those species.

Indeed, other desmosomal cadherin family members have been implicated in notochord

integrity in Zebrafish (Goonesinghe et al. 2012), although mutants display earlier gastrulation

defects that complicate the proper characterisation of notochordal phenotypes. Interestingly,

injecting our medaka *desmogon* partial promoter driving GFP in Zebrafish results in vacuolated

and sheath cell labelled clones. This suggests that the core transcriptional machinery driving

tissue-specific expression of *desmogon* remains in place and active in Zebrafish. It would be

of interest to investigate when the deployment of desmosomal cadherins in notochords arose

during evolution and how wide-spread its usage is among the different chordate clades.

354
355 ***Tg(desmogon:EGFP)*as a screening tool for genes involved in proper notochord**
356 **morphology**
357
It has been reported that most biomedical research today focuses heavily on a limited number

of genes, leaving behind potentially important genes understudied (Stoeger et al. 2018).

Making use of our newly generated transgenic line in combination with the recently published

and publicly available single cell transcriptomics data from vertebrate embryos (Briggs et al.

2018; Farrell et al. 2018), we attempted to address this imbalance. To do so we performed a

small scale F0 CRISPR screen, the efficacy of which has been recently demonstrated in

Zebrafish (Wu et al. 2018) and confirmed in other fish species (own observations and personal

communications). Indeed, recent work in Medaka (Lischik et al. 2018), in addition to our results

from *desmogon* mutants, argues for the use of F0 injected embryos as a method to analyse

tissue-specific phenotypes. Briefly, we focused on conserved well-annotated genes that were

highly and differentially expressed in developing notochords. We addressed whether they

might be involved in proper notochord morphology by using the *desmogon:GFP* line as a fast

and straight-forward read-out for notochord shape and integrity. Broadly, the targeted genes

fell into two categories. *kcnk6, si:dkey- 261h17.1, pmp22b* constituted the first category and

showed strong pleiotropic effects in addition to notochordal defects. Delineating the cause of

the pleiotropy is difficult given the essential signalling role of the notochord. (Yamada et al.

1991; Yamada et al. 1993; Pourquie et al. 1993; Hebrok et al. 1998; Fouquet et al. 1997;

Corallo et al. 2018; Stemple et al. 1996). The observed defects could either arise from the

inability of the notochord to correctly pattern adjacent tissue or from notochord independent

roles for these genes during embryogenesis. This makes assigning causal phenotypes more

difficult. The second group of targeted genes contained the conserved putative co-

transcriptional factor *vgll2b* and the highly conserved membrane bound arrestin *arrdc3a.* Both

genes showed specific defects in notochord morphology and structure when targeted and no

overt pleiotropic phenotypes. This argues for functional roles for these genes that are likely to

be notochord specific. The precise cellular defects caused by these two genes remain unclear

although in both cases vacuolated cells appeared morphologically normal. It is clear, though,

that targeted notochords appear unable to withstand the high mechanical strain and buckle

under pressure. It remains a challenge for the future to decipher the genetic networks of these

genes and to assess whether their roles are functionally conserved in higher vertebrates. All

in all, using a simple, targeted, reverse-genetics approach, in combination with publicly

available data from single cell transcriptomics (Briggs et al. 2018; Farrell et al. 2018), we

believe we have implicated new players in correct vertebrate notochord integrity. This

methodology can be easily adapted to other contexts and promises to aid in the study of

neglected genes with potentially important functions.

394 Notochord vacuolated cells during development

We made use of the newly generated *Tg(desmogon:EGFP)* line to address the formation and

growth dynamics of notochord vacuolated cells during development. It has previously been

reported that one vacuole exists per vacuolated cell under homeostatic conditions (Ellis et al.

2014) and therefore vacuole growth can be used as a proxy for cellular volume growth. Our

results confirm that vacuolated cells grow in volume over time as has been reported before

(Ellis et al. 2013; Ellis et al. 2014) and reveal that they do so anisotropically, changing their

morphology in the process from more roundish to more oblique shapes. This might be due to

the increased cellular packing as the notochord expands. While it has been previously shown

that notochords extend over time and that this supports axis elongation in vertebrate embryos

(Ellis et al. 2014; Garcia et al. 2017; Ellis et al. 2013), it is still unclear exactly how this growth

and expansion is coordinated. A possible mechanism could be that a morphogen gradient

synchronizes the growth of neighbouring cells in an orderly fashion. Our dynamic data, in both

medaka and zebrafish embryos, argues against the presence of such a signal. We reveal that

the growth of vacuolated cells in Medaka and Zebrafish is an incremental one-way process

that is locally uncoordinated; neighbouring vacuolated cells grow at different rates. This

strongly suggests cell-autonomous mechanisms of vacuolated cell growth, how this is

coordinated globally to eventually reach an equivalent size remains unclear and constitutes an

interesting avenue for future research. It could be that the final size reached is close to the

physiological limit as has been previously suggested (Ellis et al. 2014).

In addition to incrementing their size cell-autonomously we report that the number of

vacuolated cells increases as the notochord grows. However, in line with previous

observations from Zebrafish (Garcia et al. 2017), we have observed no cell division of

vacuolated cells. To characterize the initial steps of vacuolated cell formation during notochord

development we employed a highly temporally resolved 4-D approach. We observed that

vacuolated cells arise from disc-shaped precursors as previously reported (Melby et al. 1996;

Dale & Topczewski 2011). Sparse labelling of these *desmogon*+ precursors in developing

Zebrafish notochords combined with long-term live-imaging revealed a hitherto unrecognized

in vivo behavioural heterogeneity of the disc-shaped precursors. Either they directly

differentiate into vacuolated cells (forgoing any division), or they generate sheath cells by

undergoing a dorso-ventral symmetric division, that could be followed by further amplifying

rounds of mitosis. When considering the position along the AP axis, the developmental time

and the levels of EGFP+ expression, we were unable to reliably predict the output of a

precursor cell. It seems plausible therefore that the decision to form a vacuolated or sheath

cell is not predetermined nor controlled by tissue-level morphogens, but rather depends on

sensing the needs of the growing notochord tube in a local manner. Indeed, it has been

previously reported that Notch-Jag1 signalling is important in the balance between vacuolated

and sheath cell formation (Yamamoto et al. 2010), suggesting that fate acquisition could be

resolved locally among neighbouring cells. Complementarily, it is possible that local

mechanical forces that arise during the expansion of the tube could operate on inherently

plastic disc-shaped precursors and contribute to adopting vacuolated or sheath cell identity.

437
438

Notochord vacuolated cells in regeneration

It has been recently shown that Medaka hearts and retinas have a vastly reduced regenerative potential as compared to Zebrafish (Lust & Wittbrodt 2018; Ito et al. 2014; Lai et al. 2017). However, we have previously reported that Medaka neuromasts are able to regenerate efficiently (Seleit, et al. 2017), and the same happens after mechanical amputations on the caudal fin (Katogi et al. 2004) and injuries to the liver (Van Wettere et al. 2013). It therefore seems that Medaka has a highly variable tissue-specific regenerative capacity (Kang et al. 2016). We wondered whether Medaka can regenerate lost notochord vacuolated cells. It has been shown in Zebrafish that loss of *caveolae* mutant vacuolated cells in response to mechanical strain triggers an efficient regeneration response (Garcia et al. 2017). This is mediated by sheath cells that invade the inside of the notochord tube and trans-differentiate into vacuolated cells (Garcia et al. 2017). Localized laser-ablation of vacuolated cells in Medaka leads to a very similar process as in Zebrafish where small vacuolated cells specifically invade the injury site. Interestingly, these invading cells start growing in size cell-autonomously mirroring the un-coordinated nature of growth that we observe for vacuolated cells during development. The appearance of small vacuolated cells during regeneration is another reminiscent feature of what we report during notochord development. Indeed, a proportion of sheath cells formed from dividing precursors do initially contain small vacuoles that highly resemble the cells participating in the regenerative response. This suggests two things; a) the presence of sub-populations of sheath cells that display distinct behaviours both during development and regeneration (Lopez-Baez et al. 2018), b) sheath cells participating in the regenerative response most likely re-acquire a small vacuole and could thus be reverting to an earlier state in their developmental history. Reactivation of developmental programs has been a hallmark of efficient regeneration in a variety of models (Tanaka & Galliot 2009; Rodrigo Albors et al. 2015; Kaloulis et al. 2004) and it seems likely that this takes place in the notochord of Medaka after vacuolated cell loss.

The highly specific spatially localized response to vacuolated cell injury we observe in Medaka

indicates that there are mechanisms in place that can sense injured tissue without the need to

activate a global regeneration program (LoCascio et al. 2017). Indeed, it has been shown that

the release of vacuolated cell contents upon apoptosis can trigger a local regenerative reaction

from neighbouring sheath cells (Garcia et al. 2017). A similar process can therefore be

occurring in Medaka. Interestingly, when we ablated a small part of the peri-notochordal

membrane this led to vacuolated cell leakage at the injury site. As the vacuolated cell leakage

continued we witnessed the activation of a global regenerative response anterior and posterior

to the original injury site. This could be observed by the presence of small vacuolated cells

along the entire length of the notochord. It therefore seems likely that sheath cells can also

respond to injury without specific vacuolated cell death, suggesting that sheath cells might be

able to sense other stresses like tissue tension. It has recently been reported in Zebrafish that

a *wilms+* subpopulation of sheath cells gets activated in response to a coarse needle injury to

the notochord (Lopez-Baez et al. 2018). This sub-population mainly forms scar-tissue that acts

as a stopper to maintain notochordal integrity (Lopez-Baez et al. 2018). In Medaka, leakage

of vacuolated cells at the site of injury indicates that the peri-notochordal membrane was not

efficiently repaired. It is tempting to speculate that this is due to the absence or delayed

activation of the wilms+ subpopulation of sheath cells. Overall, our results strongly argue for

the presence of distinct injury-sensing mechanisms in sheath cells (dependent and

independent of vacuolated cell death). We also report the existence of two regenerative

responses in Medaka notochords, one spatially localized and the other global, that depend on

the type of injury sustained. Our data positions vacuole re-acquisition by sheath cells as the

key step for replenishing vacuolated cells regardless of the type of injury sustained. Identifying

the molecular trajectories sheath cells traverse will allow a better understanding of the existing

heterogeneities among and plasticity of sheath cells both during development and

regeneration. This will in turn allow a more targeted exploitation of the potential of sheath cells

in treating notochordal and by extension spinal cord defects in vertebrates.

Figure Legends

Figure 1 d*esmogon* is a desmosomal cadherin that labels the notochord throughout embryogenesis in Medaka.

(A) *in-situ* hybridization on *desmogon* in stage 33 medaka embryos reveals strong enrichment in the notochord. Scalebar=20 microns (B) mosaic injection of desmogon:EGFP in medaka embryos stage 30, labels the notochord . Scale bar=100um (C-D) Transgenic line *Tg(desmog:EGFP)* labels the notochord in Medaka at stage 42 embryos. Scalebar=100 microns (E-F) Maximum projection of mosaic injected (E) and *Tg(desmog:EGFP)* (F) labelling notochord vacuolated cells and a proportion of covering sheath cells. Magenta arrows indicate vacuolated cells and yellow asterisks sheath cells. Scalebar= 50 microns (G) Longitudinal EM section in between vacuolated cells connected by desmosomes (black asterisk) Scale bar= 1 micron (G') Desmosomal structure and caveolae present between vacuolated cells Scale bar = 0.2 micron. (H) Longitudinal EM section showing desmosomal connections between sheath cells and vacuolated cells scale bar= 1 micron. (I) Cross-section EM showing desmosomal connections between two sheath cells in addition to desmosomes between sheath cells and vacuolated cells scale bar= 1 micron.

Supplementary movie1 *Tg(desmog:EGFP)* stage 42 Medaka embryo. Z-stack shows labelled vacuolated cells throughout the notochord and a proportion of sheath cells labelled. Scalebar=100 microns.

Supplementary movie2 *Tg(desmog:EGFP)* stage 42 Medaka embryo. 3D projection shows the specific labelling of the notochord tube by 2.2kb *desmogon* promoter region. Scalebar=100 microns.

Figure 2 Live-imaging of *Tg(desmog:EGFP)* reveals Medaka notochord growth dynamics.

(A) Single plane of 3-4dpf *Tg(desmog:EGFP)* embryo showing vacuolated cells, notice more circular shape and small size compared to (B) 5-6dpf *Tg*Desmogon::EGFP, bigger and oblique vacuolated cells. n>10. Scalebar=50 microns. (C-C''') SPIM time-lapse imaging *of desmogon+* vacuolated cell growth over time highlighted with yellow and magenta asterisks (n>10 vacuolated cells in 3 embryos at 4-5dpf and n>10 vacuolated cells in 2 embryos at 3-4dpf). Scalebar=20microns. (D-D''') Differentiation of desmogon+ precursors into vacuolated cells near mid-point of notochord. Arrow labels cell of interest. (E-E') Differentiation of a number of desmogon+ precursors into vacuolated cells near tip of the tail and growing notochord (magenta asterisks). Scalebar=20 microns. Time in hours. Total n=8 sheath cells in 2 independent embryos.

Supplementary movie3 *Tg(desmog:EGFP)* 4-5dpf embryo.Time-lapse imaging of developing notochord. No observable divisions of *desmogon+* vacuolated cells. n=3 embryos at 4-5dpf and n=2 embryos at 3-4dpf. Scalebar=20 microns. Time in hours.

Supplementary movie4 *Tg(desmog:EGFP)* 4-5dpf embryo. Time-lapse imaging of growing *desmogon+* vacuolated cells. Notice the anisotropic and autonomous nature of vacuolated cell growth, n>10 vacuolated cells in 3 embryos at 4-5dpf and n>10 vacuolated cells in 2 embryos at 3-4dpf. Scalebar= 20microns. Time in hours.

Supplementary movie5 *Tg(desmog:EGFP)* 4-5dpf embryo. *desmogon+* precursor differentiates into vacuolated cells that grow in size over time near notochord mid-point.Time in hours. SPIM-imaging. Scalebar=20 microns.

Supplementary movie6 *Tg(desmog:EGFP)* 4-5dpf embryo near extending posterior end of the tail. *desmogon+* precursors differentiate into vacuolated cells that grow in size over time.Time in hours. SPIM-imaging. Scalebar=20 microns.

Supplementary movie7 3-D reconstruction of *desmog:EGFP injected 3*dpf embryo near extending posterior end of the tail. Notice the labelling of undifferentiated disc-shaped progenitors.

Supplementary movie8 Clones of *desmogon+* cells in 24hpf Zebrafish embryo developing notochord. Both sheath cell clones and vacuolated cell clones are present. Notice the growth of vacuolated cells is locally uncoordinated in Zebrafish as is the case in Medaka. Scalebar=30 microns. Time in hours.

Supplementary movie9 Clones of *desmogon+* disc shaped precursors in 24hpf Zebrafish embryo developing notochord. Direct differentiation of the anterior disc-shaped precursor into a vacuolated cell. Posterior disc shaped precursor produces two sheath cells after a dorso-ventral division Scalebar=30 microns. Time in hours.

Supplementary movie10 Clones of *desmogon+* disc shaped precursors in 24hpf Zebrafish embryo. Notice the presence of small vacuoles in newly formed sheath cells, that disappear over time. Scalebar=30 microns. Time in hours.

Figure 3 Bi-potent disc-shaped precursor population is fate-restricted at single cell level
(A) Mosaic injected medaka embryo *desmog:EGFP* showing the presence of disc-shaped precursors at the posterior tip of the growing notochord tube. (B-C) independent clones of sheath cells and vacuolated cells occur in mosaic injected fish. (D) Disc-shaped precursors at the growing tip of the medaka notochord are fate-restricted and either generate vacuolated cells or sheath cells. Scale bars are 20um in A, 1mm in B, and 100um in C.

Figure 4 Local and global regeneration dynamics after notochord injury
(A)*Tg(desmog:EGFP)* 6dpf embryo post ablation. Ablation area indicated by magenta lines.

Vacuolated cells in the ablated zone are missing. Vacuolated cells outside ablation zone are intact and morphologically normal. Scalebar= 100 microns. (B) Same embryo 48 hours post injury. Notice the appearance of small vacuolated cells specifically in the ablated zone. Regeneration of injured area is initiated. Vacuolated cells in uninjured area appear normal. Scalebar= 100 microns. (C) Same embryo 5 days post injury vacuolated cells grow in size, notochord is intact. Scalebar= 100 microns. n=8 embryos. (D) Ablation of vacuolated cells and lower lining of notochord tube. (E) 48 hours post injury, magenta arrow highlights position of injury, notice failure to regenerate and leakage of *desmogon+* cells outside of notochord. Overall notochord morphology grossly altered both anterior and posterior to injury site. Notice appearance of small vacuolated cells outside original injury site. Scalebar= 100 microns. (F) 5 days post injury magenta arrow highlights growing leakage of *desmogon+* vacuolated cells outside of the notochord. Failure to repair and correctly regenerate lower tube lining is evident. Gross morphological defects apparent over entire length of notochord. Notice the appearance of small vacuolated cells throughout the notochord. Scalebar= 100 microns. n=3 embryos.

Supplementary movie11 *Tg(desmog:EGFP)* 6dpf embryo. Pre-Ablation stack through notochord. Scalebar= 100 microns.

Supplementary movie12 *Tg(desmog:EGFP)* 6dpf embryo. Post-ablation stack through notochord. Notice the loss of vacuolated cells anteriorly at the site of injury. Scalebar= 100 microns.

Supplementary movie13 *Tg(desmog:EGFP)* embryo. 48hours post-injury stack through notochord. Small vacuolated cells appear specifically in the site of injury. Scalebar= 100 microns.

Supplementary movie14 *Tg(desmog:EGFP)* embryo. 5 days post-injury stack through notochord. Vacuolated cells grow in size specifically in the site of injury. Scalebar= 100

microns.

Supplementary movie15 *Tg(desmog:EGFP)* embryo. Post-ablation stack through notochord.

Vacuolated cells and lower lining of notochord ablated. Scalebar= 100 microns.

Figure 5 *desmogon* mutants exhibit notochordal lesions and vacuolated cell collapse.

(A)Control CRISPR injected *Tg(desmog:EGFP)* with o*ca2* gRNA1,2 & Cas9. Single plane.

Scalebar=100 microns. (B, C) *desmogon* gRNA1, 2, 3 & Cas9 injected into

*Tg*Desmogon:E:GFP results in local collapse of vacuolated cells and lesions in the notochord.

Single plane. Scalebar=100 microns. (D) Stable desmogon CRISPR mutant line recapitulates

phenotypes observed in the injected generation Scalebar=100microns. (E) Longitudinal EM

section through wild-type stage 42 medaka notochords with highly ordered vacuolated cell

arrangement (F) Longitudinal EM section through lesioned *desmogon* mutant stage 42

notochords, notice the structural disorganization, vacuolated cells with varying sizes (yellow

arrows), invading sheath cells (black asterisk) and evidence of collapsed vacuolated cells. (G-

G') Longitudinal EM section on *desmogon* mutant notochords reveals the presence of

collapsed vacuolated cells. (H) Small vacuolated cells are present in Longitudinal sections of

EM in *desmogon* mutants and can be clearly distinguished from the neighboring vacuolated

cells and sheath cell.

Supplementary movie16 *Tg(desmog:EGFP)* 6-7dpf *oca2* gRNA1,2+ Cas9 injection. Z-stack

through notochord. Notice normal notochord and vacuolated cell morphology. Scalebar= 100

microns.

Supplementary movie17 *Tg(desmog:EGFP)* 6-7dpf Desmogon gRNA1,2,3 & Cas9 injection.

Z-stack through notochord. Local collapses of vacuolated cells are evident and vacuolated cell

morphology is highly perturbed. Scalebar= 100 microns.

Supplementary movie18 *Tg(desmog:EGFP)* 6-7dpf desmogon gRNA1,2,3 & Cas9 injection.

Z-stack through notochord. Strong phenotypes show abnormal notochord lesions. Notice the

presence of small vacuolated cells and cellular debris. Overall a disorganized notochord

structure is evident. Scalebar= 50 microns.

Supplementary movie19 *Tg(desmog:EGFP)* 6-7dpf desmogon F1 mutant. Z-stack through

notochord. Strong phenotypes show abnormal notochord lesions. Phenotype resembles what

is observed in the injected generation. Scalebar= 30 microns.

Figure 6 Gross morphological defects in the notochord of *desmogon, vgll2b and*

***arrdc3a* CRISPR injected embryos**

(A) Overall morphology of control *oca2* gRNA & Cas9 injected notochord in

Tg(desmogon:EGFP)(B, C) Strong phenotypes in *desmogon* CRISPR injections results in

gross morphological defects in notochord integrity and notochord buckling and bending.

maximum projections. Scalebar=100 microns. Embryos with notochord phenotypes 65/160.

Over 90% of embryos survive to stage 42. (D, F) *vgll12b* gRNA1,2 & Cas9 injected into

Tg(desmog:EGFP) results in morphological defects in the notochord. Notice magenta arrow

where notochord is twisted, overall notochord bending observed. Scale bar=100microns.

Embryos with phenotypes in notochord 66/120. 79% of embryos survive to stage 42. (E, F)

arrdc3a gRNA1,2 & Cas9 injected into *Tg(desmog:EGFP)* results in notochord bending.

Scalebar=100microns. Embryos with phenotypes 29/97. 77% of embryos survive to stage

42.

Table 1 List of genes targeted by CRISPR and quantification of phenotypes in F0 injections.

Supplementary Table 1 List of all available species with known orthologues to Medaka

d*esmogon* ranked from the highest identity score to the lowest using the 1-1 and 1-many

ortholouges comparison. The gene seems to be fish-specific. (ENSEMBLE).

Supplementary Table 2 List of all available species without any othrologues to Medaka

desmogon. Notice the absence of an orthologous gene in Zebrafish and the Tetraodon

(highlighted in magenta). (ENSEMBLE).

Supplementary Table 3 List of all available species with known orthologues to Medaka vgl12b

ranked from the highest identity score to the lowest. The gene is highly conserved in

vertebrates (ENSEMBLE).

Supplementary Table 4 List of all available species with known orthologues to Medaka

arddc3a ranked from the highest identity score to the lowest. The gene is highly conserved in

vertebrates(ENSEMBLE).

Supplementary Figure 1 (A) Pfam predicted *desmogon* protein domains. In green, 3 cadherin

domains and in red, a cytoplasmic cadherin domain. Multiple sequence alignment of 12

selected species to uncharacterized Medaka protein(H2MRM9). Identity score reveals weak

amino acid sequence conservation with closest hits (B) Comparative genomic alignment of

uncharacterized medaka transcript(ENSORLG00000017110) using GENOMICUS shows

conservation of locus in the vast majority of teleost branches, notice the loss of locus in

Zebrafish and the Tetraodon. Locus of interest is highlighted by black desmogon label.

Scheme modified from Genomicus to highlight syntenic geneomic region (C) Choosing of

desmogon partial promoter. 2.2kb promoter region (highlighted in blue). H3K27ac, H3K4me1

and H3K4me2 peaks from UCSC genome browser Medaka blastula stage data at 2.2kb

upstream of predicted *desmogon* TSS.

Supplementary Figure 2

(A)Area measurement on 12 paired vacuolated cells at 4 different time-point over a 20-hour

period reveals the autonomous nature of vacuolated cell growth. Neighbouring vacuolated

cells share the same colour code and grow at different rates. Area was calculated on maximum

projections using standard Fiji software.

Supplementary Figure 3

(A, A') Time-lapse recording of clones of vacuolated cells (purple asterisks) and sheath cells

(yellow asterisks) labelled after injection of desmogonGFP plasmid into Zebrafish embryos.

Notice the growth of vacuolated cells over time is asynchronous. (B,B') Time-lapse recording

of notochord disc-shaped precursor cells labelled in injected zebrafish embryos. One labelled

precursor directly trans-differentiates into vacuolated cell (purple asterisks) while the other

undergoes a dorso-ventral division giving rise to two sheath cells (yellow asterisks). Scale

bar= 30 microns. Time in hours.

Supplementary Figure 4 (A) Longitudinal EM section in *desmogon* mutant notochord.

Notice the sheath cell invasion to the center of the lesioned tube Scale bar= 5 microns. (B)

High magnification on sheath cell nucleus from (A). Scale bar= 1 micron.

707
Supplementary Figure 5 (A) Single Z plane projection in a Tg(desmogon:EGFP) stage 42

embryo, notice the organization and size of vacuolated cells. (B) 3 consecutive Z plane

projections in an F1 CRISPR *desmogon* mutant stage 42 embryo, notice the appearance of

lesions full of small vacuolated cells (C) Multiple sequence alignment of 9 selected species to

uncharacterized Medaka protein *vgll2b*. Identity score reveals very strong amino acid

sequence conservation in vertebrates. (D) Multiple sequence alignment of 10 selected

species to uncharacterized Medaka protein *arrdc3a*. Identity score reveals very strong amino

acid sequence conservation across vertebrates.

Materials and Methods

Fish stocks and generation of transgenic lines

Medaka (*Oryzias latipes*) and Zebrafish (*Danio rerio*) stocks were kept in a fish facility built according to the local animal welfare standards (Tierschutzgesetz §11, Abs. 1, Nr. 1) as described before (Seleit et. al, 2017). Animal experiments were performed in accordance with European Union animal welfare guidelines.(Tierschutzgesetz 111, Abs. 1, Nr. 1, Haltungserlaubnis AZ35–9185.64 and AZ35–9185.64/BH KIT).

The strain used in this study is: Cab (medaka Southern wild type population). Tg(*desmogon*:EGFP) transgenic line was generated for this study by I-SceI mediated insertion, as previously described(Rembold et al. 2006). Briefly, a 2.2kb promoter region upstream of the desmogon CDS (but including the ATG of the first exon) was amplified from Medaka genomic DNA using forward primer (TCGCTGCTTGTTGTGTAGGT) and reverse primer (CATTGGCGCAGTGATTTGAA). The amplified fragment was subsequently cloned by A-tailing into a PGEMTeasy vector (Promega). From there it was sub-cloned into a vector with I-SceI sites and eGFP. This was then injected into Medaka embryos to obtain the transgenic line Tg(*desmogon*:eGFP).

Imaging and Image analysis

Embryos were prepared for live-imaging as previously described (Seleit, et al. 2017a; Seleit, et al. 2017b) 20 mg/ml as a 20x stock solution of tricaine (Sigma-Aldrich, A5040-25G) was utilized to anaesthetise embryos. Glass-bottomed dishes (MatTek Corporation, Ashland, MA 01721, USA) contained the embryos that were in turn covered with low melting agarose (0,6% in ERM). Embryos were screened using an Olympus MVX10 binocular coupled to a Leica DFC500 camera. For image acquisition we used confocal microscopes Leica TCS SPE and Leica TCS SP5 II. Developing notochords were imaged for approximately 1 day using EMBL MuVi-SPIM (Krzic, Gunther, Saunders, Streichan & Hufnagel 2012b) with two illumination

objectives (10x Nikon Plan Fluorite Objective, 0.30 NA) and two detection objectives (16X Nikon CFI LWD Plan Fluorite Objective, 0.80 NA). Embryos were placed in glass capillaries using 0,6% low melting agarose at room temperature. Standard Fiji software was used for all image analysis. Stitching was performed using 2D and 3D plugins on Fiji. For mosaic analysis, a *desmogon*:EGFP plasmid was injected embryos into 2 to 4 cell-stage embryos, which were grown until day 9 pf. For the quantification, we selected embryos bearing 1 and up to 4 clusters in the notochord, where clusters were defined as single cells or groups of cell that were at least one somite away from each other.

Multi-photon Laser Ablations

A multi-photon laser was used in combination with a Leica TCS SP5 II microscope to perform ablations on vacuolated cells of the notochord and the peri-notochordal membrane in Tg(desmogon:eGFP). 'Point ablations' were chosen along the perimeter of each vacuolated cell. Laser of 880 nm wavelength was used and the power used ranged from 30-35% for 500ms (the time parameter was adjusted in the biological replicates when necessary). The targeted area was immediately imaged post-ablation to checked for signs of cellular bursting, debris and the release of GFP signal.

CRISPR gRNAs and Cas9

Lab-made Cas9 mRNA was transcribed by mMachine Sp6 Transcription Kit (Thermo Fisher). The gRNAs for all targeted genes were designed in-silico using CCtop (Stemmer, Thumberger, et al. 2015). gRNA synthesis was done as previously reported (Stemmer, Thumberger, et al. 2015). All genes were targeted with 2 gRNAs against the CDS except *desmogon* where 3 gRNAs were designed and injected. The following is a list of gRNA sequences used in this study for each gene.

desmogon gRNA1 (CCCAUAUGGUGUGUUCAGCGUGG) gRNA2
(UUGGUGCCAGCUCUUCAGCGGGG) gRNA3 (AUCUAUGACUAUGAAGGUCGAGG)

vgll2b gRNA 1 (UGGGCCCCCAGACAUUCCUUCGG)

gRNA2(GGGUGCGCCCGUUUCACAGUGGG)

arrdc3a gRNA1(AAGUUCGUUGGACGGAAUCGAGG)

gRNA2(AUCCCGCAUGGUGGUCCCAAAGG)

kcnk6 gRNA1 (GUUCUCCAGCAUCGAGCGGCCGG) gRNA2

(UGUAUUGUAGUCGCCGGGCGAGG)

si:dkey- 261h17.1 gRNA1 (GAGCUUGCUGUCACAAGCCUUGG)

gRNA2(CUGAGUCACUUCAAUCUGCCUGG)

pmp22b gRNA1(CCUGCUGCACAUUGCUGCACUGG) gRNA2

(CAGCUCUUCACCUUGCAGAAGGG)

Tg(*desmogon*:eGFP) were injected at the 1 cell stage with a solution containing 15 ng/µl of

each gRNA for each gene individually and 150 ng/µl of CAS9 mRNA. The embryos were

screened iteratively over the course of development for gross morphological phenotypes and

for phenotypes in the notochord. For control injections two gRNAs targeting the *Oca2* locus

and Cas9 (Lischik et al. 2018) were injected into Tg(*desmogon*:eGFP) embryos. The stable

desmogon F2 mutants exhibited a small deletion in the region of guide RNA2.

Whole-mount *in-situ* hybridization

A 604bp *in situ* probe for *desmogon* was generated from total cDNA of stage 33 medaka

embryos by PCR using the following primers fwd: TTCTGCGAGATCAGGCTCAC rev:

AAGGCCCCTCCTCTGTAACT and subsequently A-tailed and cloned into a PGEMTeasy

vector (Promega). Sense and anti-sense probes were generated using Sp6 and T7

polymerases (Invitrogen) and the In situ hybridization protocol was followed as previously

reported in (Stemmer, Schuhmacher, et al. 2015). Hybridisations were performed overnight at

65°C then samples were incubated with an antibody against anti-digoxigenin conjugated with

AP Fab fragments (1:2000; Roche, 11093274910). Staining was done using NBT/BCIP

(Roche).

Electron Microscopy sample preparation and imaging

10 dpf fish from Medaka wt Cab strain and stable Desmogon mutants were placed in a fixative

consisting of 2.5% glutaraldehyde and 4% paraformaldehyde in 0.1M PHEM buffer. The

fluorescing part of the notochord in mutant fish and equally small pieces of the wt fish was cut

out in the fixative and fixation was continued for 30 min at room temperature and at 4˚C

overnight. The samples were further fixed in 1% osmium in 0.1M PHEM buffer, washed in

water, and incubated in 1% uranylacetate in water overnight at 4˚C. Dehydration was done in

10 min steps in an acetone gradient followed by stepwise Spurr resin infiltration at room

temperature and polymerization at 60˚C.The blocks were trimmed to get longitudinal sections

of the notochord. 70nm thick sections were obtained using a leica UC6 ultramicrotome (Leica

Microsystems, Vienna) and the sections were collected on Formvar-coated, copper slot grids

and thereafter post-stained with uranyl acetate and Reynold's lead citrate. Imaging was done

using a JEOL JEM-1400 electron microscope (JEOL, Tokyo) operating at 80 kV and equipped

with a 4K TemCam F416 (Tietz Video and Image Processing Systems GmBH, Gautig).

Transcriptomics data and candidate picking

The targeted genes were selected from single cell transcriptomics data generated by the Klein

Lab (Briggs et al., 2018) based on a number of different criteria. First, high enrichment during

early development of the notochord in zebrafish. Precisely, candidates were picked from 10,

14, 18 and 24 hpf data. Second, the selected genes were checked for conservation across

vertebrates. Third the genes had to be well annotated but poorly characterized in both Medaka

and Zebrafish. Lastly, a compact and short CDS, simple exon-intron structure/number and

limited alternative splicing was favoured to improve chances of efficient mutagenesis with the

CRISPR machinery.

Bioinformatics tools
830
ENSEMBL(www.ensembl.org) was used to obtain all the coding sequences and upstream

promoter regions for all genes analysed in this study. Supplementary Table 1 and 2 were

generated using standard ensembl software using the 1-1 and 1-many orthologue

comparisons. Genomic sequence alignments for synteny were performed on ensembl or using

GENOMICUS (www.genomicus.biologie.ens.fr/). Protein information and amino acid

sequence alignments were done using Uniprot (www.uniprot.org). UCSC

(https://genome.ucsc.edu/) genome browser was used for the identification of the promoter

region of desmogon. Pfam and PRINTS were used for information on protein domains and

conserved motifs (https://pfam.xfam.org/) (http://130.88.97.239/PRINTS/index.php). Single

cell transcriptomics data was visualized using

(https://kleintools.hms.harvard.edu/paper_websites/wagner_zebrafish_timecourse2018/sprin

gViewer.html?coarse_grained_tree) and the selected candidates were obtained from list of

enriched genes in developing Zebrafish notochords.

References

Adams, D.S. & Keller, R.E., Koehl, M.A.R 1990. Mechanical Development of the Notochord in Xenopus Early Tail-Bud Embryos. In *Biomechanics of Active Movement and Deformation of Cells*. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 471–485.

Briggs, J.A. et al., 2018. The dynamics of gene expression in vertebrate embryogenesis at single-cell resolution. *Science*, 360(6392), pp.eaar5780–17.

Corallo, D., Trapani, V. & Bonaldo, P., 2018. The notochord: structure and functions. *Cellular and Molecular Life Sciences*, pp.1–20.

Dale, R.M. & Topczewski, J., 2011. Identification of an evolutionarily conserved regulatory element of the zebrafish col2a1a gene. *Developmental Biology*, 357(2), pp.518–531.

Delva, E., Tucker, D.K. & Kowalczyk, A.P., 2009. The desmosome. *Cold Spring Harbor Perspectives in Biology*, 1(2), pp.a002543–a002543.

Ellis, K., Bagwell, J. & Bagnat, M., 2013. Notochord vacuoles are lysosome-related organelles that function in axis and spine morphogenesis. *The Journal of Cell Biology*, 200(5), pp.667–679.

Ellis, K., Hoffman, B.D. & Bagnat, M., 2014. The vacuole within. *BioArchitecture*, 3(3), pp.64–68.

Farrell, J.A. et al., 2018. Single-cell reconstruction of developmental trajectories during zebrafish embryogenesis. *Science*, 360(6392), pp.eaar3131–15.

Fleming, A., 2004. A central role for the notochord in vertebral patterning. *Development*, 131(4), pp.873–880.

Fouquet, B. et al., 1997. Vessel Patterning in the Embryo of the Zebrafish: Guidance by Notochord. *Developmental Biology*, 183(1), pp.37–48.

Garcia, J. et al., 2017. Sheath Cell Invasion and Trans-differentiation Repair Mechanical Damage Caused by Loss of Caveolae in the Zebrafish Notochord. *Current Biology*, 27(13), pp.1982–1989.e3.

Garrod, D. & Chidgey, M., 2008a. Desmosome structure, composition and function. *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1778(3), pp.572–587.

Garrod, D. & Chidgey, M., 2008b. Desmosome structure, composition and function. *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1778(3), pp.572–587.

Goonesinghe, A. et al., 2012. Desmosomal cadherins in zebrafish epiboly and gastrulation. *BMC Developmental Biology*, 12(1), p.1.

Gray, R.S. et al., 2014. Loss of col8a1a function during zebrafish embryogenesis results in congenital vertebral malformations. *Developmental Biology*, 386(1), pp.72–85.

Hebrok, M., Kim, S.K. & Melton, D.A., 1998. Notochord repression of endodermal Sonic hedgehog permits pancreas development. *Genes & Development*, 12(11), pp.1705–1713.

Ito, K. et al., 2014. Differential reparative phenotypes between zebrafish and medaka after
cardiac injury. *Developmental Dynamics*, 243(9), pp.1106–1115.

Jiang, D. & Smith, W.C., 2007. Ascidian notochord morphogenesis. *Developmental*
Dynamics, 236(7), pp.1748–1757.

Kaloulis, K. et al., 2004. Reactivation of developmental programs: the cAMP-response
element-binding protein pathway is involved in hydra head regeneration. *Proceedings of*
the National Academy of Sciences, 101(8), pp.2363–2368.

Kang, J. et al., 2016. Modulation of tissue repair by regeneration enhancer elements. *Nature*,
532(7598), pp.201–206.

Katogi, R. et al., 2004. Large-scale analysis of the genes involved in fin regeneration and
blastema formation in the medaka, Oryzias latipes. *Mechanisms of Development*, 121(7-
8), pp.861–872.

Koehl, M.A.R., Quillin, K.J. & Pell, C.A., 2000. Mechanical Design of Fiber-Wound Hydraulic
Skeletons: The Stiffening and Straightening of Embryonic Notochords. *American*
Zoologist, 40(1), pp.28–041.

Krzic, U., Gunther, S., Saunders, T.E., Streichan, S.J. & Hufnagel, L., 2012a. Multiview light-
sheet microscope for rapid in toto imaging. *Nature methods*, 9(7), pp.730–733.

Krzic, U., Gunther, S., Saunders, T.E., Streichan, S.J. & Hufnagel, L., 2012b. Multiview light-
sheet microscope for rapid in toto imaging. *Nature methods*, 9(7), pp.730–733.

Lai, S.-L. et al., 2017. Reciprocal analyses in zebrafish and medaka reveal that harnessing
the immune response promotes cardiac regeneration. *eLife*, 6, p.1382.

Lim, Y.-W. et al., 2017. Caveolae Protect Notochord Cells against Catastrophic Mechanical
Failure during Development. *Current Biology*, 27(13), pp.1968–1981.e7.

Lischik, C.Q., Adelmann, L. & Wittbrodt, J., 2018. Enhanced in vivo-imaging in fish by
optimized anaesthesia, fluorescent protein selection and removal of pigmentation.

Lleras Forero, L. et al., 2018. Segmentation of the zebrafish axial skeleton relies on
notochord sheath cells and not on the segmentation clock. *eLife*, 7, p.15.

LoCascio, S.A., Lapan, S.W. & Reddien, P.W., 2017. Eye Absence Does Not Regulate
Planarian Stem Cells during Eye Regeneration. *Developmental Cell*, 40(4), pp.381–
391.e3.

Lopez-Baez, J.C. et al., 2018. Wilms Tumor 1b defines a wound-specific sheath cell
subpopulation associated with notochord repair. *eLife*, 7, p.239.

Lust, K. & Wittbrodt, J., 2018. Activating the regenerative potential of Müller glia cells in a
regeneration-deficient retina. *eLife*, 7, p.7028.

Melby, A.E., Warga, R.M. & Kimmel, C.B., 1996. Specification of cell fates at the dorsal
margin of the zebrafish gastrula. *Development*, 122(7), pp.2225–2237.

Nixon, S.J. et al., 2007. Caveolin-1 is required for lateral line neuromast and
notochord development. *Journal of Cell Science*, 120(13), pp.2151–2161.
938
Pourquie, O. et al., 1993. Control of dorsoventral patterning of somitic derivatives by
notochord and floor plate. *Proceedings of the National Academy of Sciences*, 90(11),

pp.5242–5246.

Rembold, M. et al., 2006. Transgenesis in fish: efficient selection of transgenic fish by co-
injection with a fluorescent reporter construct. *Nature protocols*, 1(3), pp.1133–1139.

Rodrigo Albors, A. et al., 2015. Planar cell polarity-mediated induction of neural stem cell
expansion during axolotl spinal cord regeneration. *eLife*, 4, p.664.

Satoh, N., Tagawa, K. & Takahashi, H., 2012. How was the notochord born? *Evolution &*
Development, 14(1), pp.56–75.

Seleit, A., Krämer, I., Ambrosio, E., et al., 2017. Sequential organogenesis sets two parallel
sensory lines in medaka. *Development*, 144(4), pp.687–697.

Seleit, A., Krämer, I., Riebesehl, B.F., et al., 2017. Neural stem cells induce the formation of
their physical niche during organogenesis. *eLife*, 6, p.505.

Stemmer, M., Schuhmacher, L.N., et al., 2015. Cavefish eye loss in response to an early
block in retinal differentiation progression. *Development*, 142(4), pp.743–752.

Stemmer, M., Thumberger, T., et al., 2015. CCTop: An Intuitive, Flexible and Reliable
CRISPR/Cas9 Target Prediction Tool. S. Maas, ed. *PloS one*, 10(4), p.e0124633.

Stemple, D.L., 2005. Structure and function of the notochord: an essential organ for chordate
development. *Development*, 132(11), pp.2503–2512.

Stemple, D.L., 2004. The notochord. *Current Biology*, 14(20), pp.R873–R874.

Stemple, D.L. et al., 1996. Mutations affecting development of the notochord in zebrafish.
Development, 123, pp.117–128.

Stoeger, T. et al., 2018. Large-scale investigation of the reasons why potentially important
genes are ignored. T. Freeman, ed. *PLOS Biology*, 16(9), p.e2006643.

Tada, M. & Heisenberg, C.P., 2012. Convergent extension: using collective cell migration
and cell intercalation to shape embryos. *Development*, 139(21), pp.3897–3904.

Talbot, W.S. et al., 1995. A homeobox gene essential for zebrafish notochord development.
Nature, 378(6553), pp.150–157.

Tanaka, E. & Galliot, B., 2009. Triggering the regeneration and tissue repair programs. In
Development (Cambridge, England). pp. 349–353.

Trubiroha, A. et al., 2018. A Rapid CRISPR/Cas-based Mutagenesis Assay in Zebrafish for
Identification of Genes Involved in Thyroid Morphogenesis and Function. *Scientific*
Reports, pp.1–19.

Van Wettere, A.J. et al., 2013. Immunohistochemical characterization of the hepatic
progenitor cell compartment in medaka (Oryzias latipes) following hepatic injury. *Journal*
of comparative pathology, 149(4), pp.434–445.

Wopat, S. et al., 2018. Spine Patterning Is Guided by Segmentation of the Notochord
Sheath. *CellReports*, 22(8), pp.2026–2038.

Wu, R.S. et al., 2018. A Rapid Method for Directed Gene Knockout for Screening in G0
Zebrafish. *Developmental Cell*, 46(1), pp.112–125.e4.

Yamada, T. et al., 1991. Control of cell pattern in the developing nervous system: Polarizing
activity of the floor plate and notochord. *CELL*, 64(3), pp.635–647.

Yamada, T. et al., 1993. Control of cell pattern in the neural tube: Motor neuron induction by
diffusible factors from notochord and floor plate. *CELL*, 73(4), pp.673–686.

Yamamoto, M. et al., 2010. Mib-Jag1-Notch signalling regulates patterning and structural
roles of the notochord by controlling cell-fate decisions. *Development*, 137(15), pp.2527–
2537.

Acknowledgments

We would like to thank Sylvia Urbansky and Steffen Lemke for critical input on the manuscript. J. Wittbrodt for the use of confocal and light sheet microscopes and generous support. We would like to thank C. Funaya, S. Gold and S. Hillmer from the EMCF facility at Heidelberg University for the preparation, imaging and processing of the electron microscopy data. We would like to thank Nihad Softic for injections for spare labeling and Pui Oui Hong for help with the genotyping of Desmogon mutants. We thank E. Leist, M. Majewski and A. Saraceno for fish husbandry.

Funding

Deutsche Forschungsgemeinschaft (SFB 873)
• Ali Seleit
• Karen Gross
• Lazaro Centanin

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Competing interests

No competing interests

Seleit et al, Figure 1

Seleit et al, Figure 2

Seleit et al, Figure 3

Seleit et al, Figure 4

A *oca2* gRNA 1,2 + Cas9

B *desmogon* gRNA 1,2,3 + Cas9

C *desmogon* gRNA 1,2,3+Cas9

D *stable desmogon* mutant line

E *wild-type*

F *desmogon*

1um

1um

G

G'

collapsed **VC**

5 um

G'

0,5 um

H

1um

Seleit et al, Figure 5

A *oca2* gRNA1, 2 + Cas9

B *desmogon* gRNA 1, 2 ,3 +Cas9

C *desmogon* gRNA 1, 2 ,3 +Cas9

D *vgll2b* gRNA 1, 2 +Cas9

E *arrdc3a* gRNA 1, 2 +Cas9

F *desmogon* knockout

gross notochordal phenotypes

local collapse of vacuolated cells

F0 CRISPR/Cas9 screen

desmogon::GFP

screening tool

vgll2b
arrdc3a

notochordal phenotypes

Seleit et al, Figure 6

Gene Name	Full name	proposed roles structure and localization	CRISPR Injected F0 Phenotypes %	Phenotypic description
dsgon	*desmogon*	constituent of desmosomal structures, plasma membrane	40,6	local collapses of vacuolated cells, notochord bending, local notochord collapses, no pleotropic effects, highly specific to notochord, 90% of embryos survive to stage 42
vgll2b	*vestigial-like family member 2b*	predicted transcription factor TF/co-factor, nuclear	55	strong phenotypes in the notochord, bending and twisted notochords, spirals of notochord, notochord tube coiling, local collapses of notochord no big pleotropic effects, 79% of embryos survive to stage 42
arrdc3a	*arrestin domain containing 3a*	regulating signal transduction at G protein-coupled receptors	30	wavy notochords, twisted notochords, no strong pleiotropic effects, 77% of embryos survive to stage 42
kcnk6	*potassium two pore domain channel subfamily K member 6*	ion transport, plasma membrane	44,3	kinks and wavy notochords, shorter axis elongation, general growth retardation 73% survive to stage 42
si:dkey-261h17.1	ENSORLG00000015828	cell adhesion, membrane bound	42,1	notochord spirals and twists, wavy notochord, kinks, perturbed axis elongation, 69% of embryos survive to stage 42
pmp22b	*peripheral myelin protein 22b*	helical transmembarne, integral component of membranes,	31,2	strong pleiotropic effects, growth retardation and developmental delays, heart edemas, smaller eyes, wavy notochords, spiral notochords and twists, short axis/failure of axis extension, patterning defects, 58% survive to stage 42

Supplementary figure 1

Species	Protein name	Protein entry	Sequence identity %
Oryzias latipes	uncharacterized	H2MRM9	100,0
Fundulus heteroclitus	uncharacterized	A0A147B2M0	66,7
Poeciliopsis prolifica	DSG2	A0A0S7JQR1	65,0
Xiphophorus maculatus	uncharacterized	M4AQR9	59,9
Nothobranchius furzeri	uncharacterized	A0A1A7ZRJ5	59,5
Astyanax mexicanus	uncharacterized	W5KFL2	59,5
Poecilia formosa	uncharacterized	A0A087X6J9	57,3
Danio rerio	dsg2.1	H2EQR6	46,6
Xenopus laevis	dsg2	A0A1L8FYU6	41,6
Homo sapiens	DSG2	Q14126	40,2
Gallus gallus	DSG2	F1NJD5	40,0
Mus musculus	Dsg2	O55111	39,2
Rattus norvegicus	Dsg2	F1LYX9	38,7

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

A

Invading sheath cell

5 um

B

1 µm

Supplementary Figure 5

C

Species	Protein name	Protein entry	Sequence identity %
Oryzias latipes	vgll2b	H2MAV4	100
Nothobranchius furzeri	VGLL2B	A0A1A8B070	87,2
Poecilia formosa	uncharacterized	A0A087XF56	78,6
Xiphophorus maculatus	uncharacterized	M4AFN1	77,6
Danio rerio	vgll2b	Q499A6	66
Lepisosteus oculatus	uncharacterized	W5MDI3	57,4
Astyanax mexicanus	uncharacterized	W5KS66	56,5
Gallus gallus	VGLL2	C0KDW3	40,7
Mus musculus	Vgll2	Q8BGW8	38,7
Homo sapiens	VGLL2	Q8N8G2	38

D

Species	Protein name	Protein entry	Sequence identity %
Oryzias latipes	arrdc3a	H2M3K3	100
Oreochromis niloticus	arrdc3	I3K178	96,4
Tetraodon nigroviridis	arrdc3a	H3CX04	95,4
Xiphophorus maculatus	uncharacterized	M4AYA0	93
Poeciliopsis prolifica	ARRD3	A0A0S7HB53	92,7
Astyanax mexicanus	uncharacterized	W5L537	92
Danio rerio	arrdc3a	Q1ECU3	89,9
Gallus gallus	ARRDC3	R4GLF2	85,6
Mus musculus	Arrdc3	Q7TPQ9	85,3
Homo sapiens	ARRDC3	Q96B67	84,8