Altered directed functional connectivity of the right amygdala in depression: high-density

EEG study

Alena Damborská[1,2,3 a *], Eliška Honzírková[2], Richard Barteček[2,3], Jana Hořínková[2,3], Sylvie

Fedorová[2,3], Šimon Ondruš[2,3], Christoph M. Michel[1,4], Maria Rubega[1a,b]

[1]Department of Basic Neurosciences, University of Geneva, Campus Biotech, Geneva,

Switzerland

[2]Department of Psychiatry, Faculty of Medicine, Masaryk University, Brno, Czech Republic

[3]Department of Psychiatry, University Hospital Brno, Brno, Czech Republic

[4]Lemanic Biomedical Imaging Centre (CIBM), Lausanne and Geneva, Switzerland

[a]these authors contributed equally to this work

[b]present address: Department of Neuroscience, University of Padova, Padova, Italy

* Correspondence:

Alena Damborská

adambor@med.muni.cz

Abstract

The cortico-striatal-pallidal-thalamic and limbic circuits are suggested to play a crucial role in

the pathophysiology of depression. Stimulation of deep brain targets might improve

symptoms in treatment-resistant depression. However, a better understanding of connectivity

properties of deep brain structures potentially implicated in deep brain stimulation (DBS)

treatment is needed. Using high-density EEG, we explored the directed functional

connectivity at rest in 25 healthy subjects and 26 patients with moderate to severe depression

within the bipolar affective disorder, depressive episode, and recurrent depressive disorder.

We computed the Partial Directed Coherence on the source EEG signals focusing on the

amygdala, anterior cingulate, putamen, pallidum, caudate, and thalamus. The global

efficiency for the whole brain and the local efficiency, clustering coefficient, outflow, and

strength for the selected structures were calculated. In the right amygdala, all the network

metrics were significantly higher (p<0.001) in patients than in controls. The global efficiency

was significantly higher (p<0.05) in patients than in controls, showed no correlation with

status of depression, but decreased with increasing medication intake ($R^2 = 0.59$ and $p = 1.52e − 05$). The amygdala seems to play an important role in neurobiology of depression.

Practical treatment studies would be necessary to assess the amygdala as a potential future

DBS target for treating depression.

Introduction

Affective disorders belong to the most common and most serious psychiatric disorders [1]. A

crucial role of the cortico-striatal-pallidal-thalamic and limbic circuits in the neurobiology of

depression was repeatedly reported [2][3][4]. Magnetic resonance imaging, functional magnetic

resonance imaging (fMRI), magnetoencephalographic, and electroencephalographic (EEG)

studies have confirmed that depressive patients show structural impairments and functional

disbalances of brain networks that involve structures engaged in a) emotions, i.e. amygdala,

subgenual anterior cingulate, caudate, putamen and pallidum [5][3][6][7][8][9][10][11][12]; b) self-

referential processes, i.e. medial prefrontal cortex, precuneus, and posterior cingulate cortex

[13][14]; c) memory, i.e. hippocampus, parahippocampal cortex [15]; d) visual processing, i.e.

fusiform gyrus, lingual gyrus, and lateral temporal cortex [16]; and e) attention, i.e. dorsolateral

prefrontal cortex, anterior cingulate cortex (ACC), thalamus, and insula [17][10][11][12]. Moreover,

post-mortem morphometric measurements revealed smaller volumes of the hypothalamus,

pallidum, putamen and thalamus in patients with affective disorders [18].

Many depressive patients fail to respond to pharmacological therapy resulting in 1 – 3%

prevalence of treatment-resistant depression (TRD) [19]. One of the newest therapeutic

approaches for TRD is an invasive direct electrical stimulation of relevant deep brain

structures [20]. Both unipolar and bipolar depression patients might benefit from deep brain

stimulation (DBS) treatment [21], although an optimal approach, including selection of an

optimal target structure, has yet to be established. Selection of the brain structures, that are

currently being tested as DBS targets for treating depression [20], is mostly supported with the

evidence from lesional [22][23], animal [24][25][26][27][28][29][30], and neuroimaging [31][32][33][34][35][36][37][38]

studies. The latter approach provides evidence from a network perspective [39][40] showing

dysbalances in the intrinsic functional architecture of the brain. During a resting state,

patients with depression as compared to healthy controls show hyperconnectivity within the

default mode network [13][33][38], hypoconnectivity within the frontoparietal network [41][42],

hyperconnectivity between the default mode and frontoparietal networks [43], and dysbalances

in connectivity within the salience [44][45] and dorsal attention [46] networks. Functional

connectivity anomalies between the hippocampus, cortical and subcortical regions [47] similar

to those observed in humans with depression, were also observed in a genetic rat model of major depression. The pathophysiological basis of depression, however, still remains incompletely understood. Particularly, better understanding of the connectivity properties of deep brain structures potentially implicated in DBS treatment could have an important value.

Neuroimaging techniques, such as fMRI and EEG, allow to investigate the integration of functionally specialized brain regions in a network. Inferring the dynamical interactions among simultaneously recorded brain signals can reveal useful information in abnormal connectivity patterns due to pathologies.

The connectivity studies based on fMRI are usually based on correlation analyses without providing knowledge about the direction of the information flow between the examined regions. Understanding the directionality is, however, crucial when searching for suitable DBS targets for treating TRD, because the antidepressant effect of DBS treatment might be caused by changes in the activity of remote structures that receive inputs from the stimulated region. For example, it has been hypothesized that DBS applied in the nucleus accumbens might influence the activity in the ventral (subgenual ACC, orbitofrontal and insular cortices) and dorsal (dorsal ACC, prefrontal and premotor cortices) subnetworks of the depression neurocircuitry [48]. Causal link between a functional inhibition of the lateral habenula and reduction of the default mode network hyperconnectivity was shown on a rat model of depression [30], which might explain the therapeutic effect of the lateral habenula DBS in TRD patients [49]. In other words, the functional inhibition of a deep brain structure via DBS might cure depression through reduction of the hyperconnectivity in the large-scale brain network. Another example of a particular role of the stimulated structure in the large-scale neural communication is the ACC, whose possible integrative role in cognitive processing [50] [51] might explain the most recently reported high efficacy of DBS to subgenual ACC in treating depression [52].

The growing interest in investigating the dynamical causal interactions that characterize resting-state or task-related brain networks has increased the use of adaptive estimation algorithms during recent years. Particularly, Granger causality based on adaptive filtering algorithms is a well suited procedure to study dynamical networks consisting of highly non-stationary neural signals such as EEG signals [53] [54]. The adaptive filtering enables to deal with time-varying multivariate time-series and test direct causal links among brain regions. A signal x is said to Granger-cause another signal y if the history of x contains information that helps to predict y above and beyond the information contained in the history of y alone [55].

Aberrant functional EEG-based connectivity in depressive patients was reported in studies
where network metrics were computed directly between sensor recordings [56] [57] [58] [59] [60] [61].
Since each EEG channel is a linear mixture of simultaneously active neural and other
electrophysiological sources, whose activities are volume conducted to the scalp electrodes,
the utility of such observations on the sensor level is limited [62] [63]. This limitation is
particularly remarkable in connectivity studies which aim to identify the real active relations
between brain regions. Connectivity analysis performed in the source space enables to
partially overcome this issue [62]. Indeed, Partial Directed Coherence estimators do not take
into account zero-lag interactions that describe the instantaneous propagation of activity,
considering the zero-phase connectivity as noise added to lagged connectivity patterns of
interest. For this reason, directed functional connectivity analysis based on electrical source
imaging proved to be a promising tool to study the dynamics of spontaneous brain activity in
healthy subjects and in various brain disorders [64] [65] [66]. Despite this fact, the electromagnetic
imaging has not been yet used in patients with depression to study the directed connectivity
of resting-state networks.

In the current study, we explored the directed functional connectivity at rest in depression
using high-density EEG. We computed the Partial Directed Coherence on the source EEG
signals focusing on the role of the amygdala, anterior cingulate, putamen, pallidum, caudate,
and thalamus in large-scale brain network activities. We hypothesized that the resting-state
directed functional connectivity in these deep brain structures might be disrupted in patients
with depression compared to healthy controls.

Results

In line with the aim of the study we focused on resting-state electrophysiological activity of
twelve regions of interest (ROIs) of selected deep brain structures. Further details on results
on the ROIs of the whole brain are reported in the Supplementary Information.

Power spectra. We found an overall increase in power in theta and alpha frequency bands in
patients compared to controls at both the *population* and *single-subject* levels. At the
population level, significantly higher power ($p<0.05$) in patients was found in all investigated
subcortical regions in both frequency bands (see Figure 1). At the *single-subject* level, a
significantly higher power ($p<0.05$) in patients than in controls was observed in the [4-12] Hz
frequency range bilaterally in the thalamus, pallidum, putamen, and caudate. Moreover, a

significant left-lateralized power increase (p<0.05) in patients vs controls was observed in the
anterior cingulate and amygdala in this frequency range (see Figure 2b).

We found a significantly decreased power in delta [1-4] Hz and beta [12-18] Hz frequency
bands in patients compared to controls in all investigated ROIs, when evaluating the results at
the *population* level (Figure 1). At the *single-subject* level, delta power was significantly
decreased in patients vs controls in the right caudate, putamen, and pallidum (Figure 2a).
There was no significant difference in beta power between the two groups in any investigated
ROI at the *single-subject* level (see Figure 2c).

Network metrics. The connectivity network measures that we performed in the [4-12] Hz
frequency range, showed increased values in patients compared to controls at both levels. At
the *population* level, the local efficiency measured in patients was higher than in controls in
all examined subcortical ROIs (see Figure 3). At the *single-subject* level, the global
efficiency was significantly higher (p<0.05) in patients (mean $\pm$ standard deviation: 0.0129
$\pm$0.0021) than in controls (mean $\pm$ standard deviation: 0.0126$\pm$0.0019). Considering all
brain regions, the local efficiency tended to be higher in patients compared to controls (see
Supplementary Fig. S2 online) but the significant differences corresponded only to the right
precentral, amygdala and caudate regions (p<0.05). We observed significant correlations
between the local efficiency and power in the [4-12] Hz frequency range in subcortical ROIs
but it was not generalized among all twelve subcortical ROIs (see Supplementary Fig. S3
online). No significant correlations were found between the local efficiency and power in
delta and beta bands. All the network measures computed on the twelve selected ROIs
showed significantly higher values in patients than in controls in the right amygdala. The
strength, local efficiency, and clustering coefficient of the right caudate were significantly
higher in patients than controls, while there was no significant difference between the groups
in the outflow from this ROI. There were no significant differences in any network metric in
the anterior cingulate, thalamus, pallidum, or putamen (see Figure 4).

There were no statistical differences in the network metrics estimated between the left and
right hemisphere in each subject. The laterality indices showed that neither controls, nor
patients had a lateralization in connectivity results of the six investigated deep brain
structures. No significant differences in the laterality indices were observed comparing
controls and patients.

Effect of medication on network impairments. We found no correlation of the connectivity

results with the intake of benzodiazepines, while there was a significant relationship between

the global efficiency as predictor of the intake of AD/AP/MS medication (AD/AP/MS ~ 1 +

GE + GE^2; Root Mean Squared Error: 0.716;   F-statistic vs. constant model:

18.7,   ). The global efficiency decreased with increasing medication score (see

Figure 5). We observed no significant correlation (   ) between the

connectivity results and any of the parameters that describe the status of depression (MADRS

score, CGI score, illness duration, and the number of episodes) or the demographic profile

(age and education level).

Figure 1

171   (A)   (B)

173   Figure 2

181   Figure 3

Average local efficiency in the subcortical regions in [4-12] Hz

Controls

L   R

Left Anterior Cingulum
Left Thalamus
                    Left Caudate
Left Amygdala   Left Putamen
                    Left Pallidum

0.02   0.03

(a)

Patients

L   R

0.02   0.03

(b)

184   Figure 4

Network metrics in the subcortical regions in [4-12] Hz

Local Efficiency

Clustering coefficient

(a)   (b)

Strength

Outflow

(c)   (d)

186   Figure 5

187
Discussion

In this study, we investigated resting-state network alterations using iPDC on source signals

of high-density EEG in patients with depression compared to healthy controls. We explored

the directed functional connectivity of the amygdala, anterior cingulate, putamen, pallidum,

caudate, and thalamus, among them and with all the other brain regions in the time and

frequency domain. We exploited the Kalman filter algorithm [67] assuming that resting state

EEG segments were multiple realizations of the same process. Although we collapsed the

temporal dimension to evaluate the network metrics, we decided to use a time-varying

adaptive algorithm instead of a stationary autoregressive model to take into account the

possible non-stationarity of the EEG signal and to more accurately capture this variability

before collapsing the time with a summary measure, e.g., the median.

To sum up, we demonstrated that in patients with moderate to severe depression: (1) the

directed functional connectivity was significantly increased compared to controls in the right

amygdala and the right caudate; (2) the power in theta and alpha frequency bands was

significantly increased compared to controls in all investigated brain anatomical structures;

(3) higher medication intake was associated with lower overall driving from the investigated

regions.

Increased right amygdala directed functional connectivity in depression. The most robust

finding in our study was an abnormally increased directed functional connectivity in the right

amygdala during resting-state in depressive patients. Even though the left-right asymmetry

was not demonstrated by the laterality indices, a right-lateralized hyper-connectivity, as
revealed with all the computed network metrics, was observed in the amygdala. We observed
an increase in outgoing connections from the right amygdala as reflected with significantly
higher outflow and strength in patients compared to controls. Moreover, we found a hyper-
connectivity in the local networks of the right amygdala as reflected with significantly higher
local efficiency and clustering coefficient in patients compared to controls.

We also found a significantly higher global efficiency in patients compared to healthy
controls. This network feature had the same trend at the population level. Namely, we
observed abnormally increased local efficiency of all examined deep brain structures in
depressive patients. The efficiency measures the ability of a neural network to integrate and
combine information. The deeper regions have a key role as hubs of the large-scale brain
networks, so changes in their local connectivity properties might have also led to connectivity
changes in the whole brain.

The amygdala is involved in processing salient stimuli [68] [69] and has been implicated as one of
the central hubs within the affective salience network [70] [71] [72]. There is converging evidence
from the neuroimaging studies that points to an abnormally increased connectivity and
heightened activation of the amygdala in major depressive disorder (MDD) patients [73] [74],
Reduced connectivity [75] [76] and anomalous subregional functional resting-state connectivity
of the amygdala [77] were also reported. Distinct network dysfunctions in MDD were suggested
to underlie adult-specific amygdala resting-state fMRI connectivity impairment within the
affective network, presumably reflecting an emotional dysregulation in MDD [76].
Hyperconnectivity between the amygdala, default mode network and salience network was
also found to be related to depressive symptoms suggesting to underlie the poststroke
depression in temporal lobe lesions [78]. Unfortunately, the directionality of connections, which
might be of interest when considering a structure as a potential DBS-target for treatment of
TRD, cannot be inferred from these functional studies. There are only rare EEG-based
connectivity studies focusing on depressive symptoms [58] [59] [60] [79] that are, however, conducted
only on a non-clinical population [79] or with connectivity parameters calculated at the sensor
level [57] [58] [59] [60] [61]. Authors of one of these studies [79] suggested an inability of the left
dorsolateral prefrontal cortex to modulate the activation of the left temporal lobe structures to
be a crucial condition for ruminative tendencies. Interestingly, in the current study we
demonstrated an abnormal increase in directed functional connectivity arising from the right
amygdala. This increased connectivity in depressive patients could reflect an abnormal

functioning of the right amygdala. Such dysfunction might represent an impaired bottom-up

signaling for top-down cortical modulation of limbic regions, leading to an abnormal affect

regulation in depressive patients.

The increased functional connectivity in amygdala is likely related to structural changes

observed in depression. Enlarged amygdala volumes was found in first-episode depressive

patients that positively correlated with severity of depression [80]. Higher grey matter volume

was detected in bilateral amygdala of TRD patients compared to non-TRD patients,

irrespective whether the patients presented bipolar or unipolar features and was suggested to

reflect vulnerability to chronicity, revealed by medication resistance [81]. Larger right

amygdala volume was, however, also suggested to be associated with greater chances of

remission in bipolar disorder [82].

In our study we aimed to investigate the directed functional connectivity in amygdala to

provide knowledge on neurobiology of depression that is needed to evaluate this structure as

a possible candidate for DBS treatment in depression. Despite myriad of DBS targets for

treating depression tested in humans [20], the amygdala is not among them. The possible safety

and utility of DBS in the amygdala could only be inferred from studies, in which the

amygdala-DBS was performed for other neuropsychiatric diagnoses, such as epilepsy [83 84 85]

[86], post-traumatic stress disorder [87 88], and autism [89]. In one of these studies transient

stimulation-related positive shift in mood was observed [84]. Particularly, the stimulation of the

right amygdala induced a transient decrease in the negative affective bias, i.e. the tendency to

interpret ambiguous or positive events as relatively negative. In this case study, an epileptic

patient with MDD rated the emotional facial expressions as more positive with stimulation

than without. The stimulation effect might have been associated with a transient

normalization of likely impaired function of the right amygdala in that patient. We can only

speculate, whether this dysfunction was in terms of hyper-connectivity similar to that

observed in our study and whether it was temporally decreased by inhibitory effect of the

stimulation.

Increased right caudate directed functional connectivity in depression. We demonstrated

that during resting state, patients had significantly higher right caudate directed functional

connectivity than healthy controls. Despite no significant difference between groups in the

caudate outflow, we observed an abnormally increased strength of outgoing connections from

the right caudate in patients. Moreover, we found a hyper-connectivity in the local networks

of the right caudate as reflected with significantly higher local efficiency and clustering

coefficient in patients compared to controls. Caudate hyperactivation and increased caudate-
amygdala and caudate-hippocampus fMRI connectivity during stress was previously reported
in remitted individuals with recurrent depression [90]. The here observed EEG-based functional
caudate hyperconnectivity suggests striatal dysfunction even during resting-state in depressed
patients. Our finding is consistent with a compelling evidence directly associating cortico-
basal ganglia functional abnormalities with primary bipolar and unipolar spectrum disorders
[91]. Deficits in resting-state default-mode network connectivity with the bilateral caudate were
suggested to be an early manifestation of MDD [92]. Reduced grey matter volume in the
bilateral caudate [93 94 95 12], diffusion tensor imaging-based hypoconnectivity between the right
caudate and middle frontal gyrus [96], and altered functional connectivity of the right caudate
with the frontal regions [94] was observed in MDD patients. In a post-mortem morphometric
study in late-life depressive subjects, reduction in neuronal density was found in both the
dorsolateral and ventromedial areas of the caudate nucleus [97]. Associations between increased
white matter lesion volumes and a decreased right caudate volume in the late-life depression
was reported [98]. In mild to moderately depressed patients no change in caudate gray matter
volumes were found [99] suggesting inverse correlation between the caudate volume and
severity of depression.

We found no significant differences in any network metric in the putamen, pallidum,
thalamus, and anterior cingulate. It is possible, however, that examining these structures as a
whole might be insensitive to different changes in their relevant subregions. Only the medial
part of the thalamus is expected to play a role in the experience of affect [73 100]. Reduced
activity in the dorsal ACC but increased activity in the subgenual ACC have been found in
acute depression in functional imaging studies [101 102]. Moreover, we must take into account
the limitations of our methodological approach, i.e. the source localization of the EEG
activity in the subcortical regions. Previous work using simulations and source reconstruction
provided indirect evidence for the detectability of subcortical sources in non-invasive EEG
and MEG recordings [103 104 105 106]. Moreover, direct evidence that subcortical activity is
detectable with high density EEG [107] or magnetoencephalography [108] was presented recently.
Nevertheless, we have to keep in mind that the spatial resolution in detecting and
distinguishing neighboring brain regions is about 24 mm [107]. Therefore, our results in the
caudate, putamen and pallidum are probably overlapping due to smearing of the sources.
Keeping in mind these limitations and with respect to the lower robustness of our findings in
the   caudate,   we   can   just   encourage   researchers   to   further   investigate   the

neuropathophysiology of depression associated with the caudate nucleus functioning. More

evidence from neuroimaging studies is needed to provide arguments for the next caudate-

DBS tests in treating TRD. In an early case study, DBS of the ventral caudate nucleus

markedly improved symptoms of depression in a patient with MDD and comorbid obsessive-

compulsive disorder [109]. No change in depressive symptoms, however, was recently observed

during the stimulation of the caudate in a study of three TRD patients [110] and authors

concluded the caudate to be less promising DBS target than the nucleus accumbens.

Increased theta and alpha powers in depression. We found a significantly higher power in

the theta and alpha frequency bands in the depressed compared to the healthy control group

in all the investigated subcortical structures consistently at both the population and single-

subject levels. The power decrease in the beta and delta frequency bands was observed only

in the right striatum at both levels.

Our findings might be in line with previous observations in the sensor space of the scalp

EEG. Abnormally high power in alpha [111] [112] [113] and theta [111] [114] [113] frequency bands in

parietal and occipital regions were found in depressed patients, lower than normal beta and

delta power were also reported [113]. Recent evidence points, however, to opposite power

changes showing that theta and alpha power might decrease, while beta power increases in

depression [115]. Moreover, the same study reported negative association of the posterior alpha

power with the depression severity. While changes in cortical theta and alpha activity were

suggested to be inversely related to the level of cortical activation, enhancement of the

cortical beta power was suggested to reflect higher level of anxiety symptoms in depressed

patients [111]. To the best of our knowledge there is only one study that directly recorded

electrophysiological activity in subcortical structures in depressive patients. In this study, a

larger alpha activity in MDD patients compared with obsessive compulsive disorder was

found in the limbic DBS targets (the anterior cingulate and the bed nucleus of the stria

terminalis) [116]. Moreover, in the same study, the increased alpha power correlated with

severity of depressive symptoms. Nevertheless, in spite of parallels with prior reports, the

current link between the power changes in subcortical structures and depression awaits

replication.

Lower network impairments with more medication. We found an inverse relationship

between the intake of medication and the impairment of the investigated networks.

Particularly, increased intake of antidepressants, antipsychotics, and mood stabilizers was

associated with reduction of the global efficiency. This finding might be related to the

pharmacological effect on the brain activity, i.e. a change towards the normalization of the
hyper-connectivity in the cortico-striatal-pallidal-thalamic and limbic networks. The low
sample size and great variability in medication made it, however, impossible to examine any
potential influence of medication on the network impairments by comparing patients
receiving a specific drug with those not receiving it. To summarize the various medications,
an ordinal variable was used that is only a rough measurement of medication usage.
Moreover, the duration of the illness rather than the duration of the specific drug intake was
considered in our study. Only doses of medication actually taken at the time of experiment
were taken into consideration. The possible accumulated effect of specific drugs on
connectivity results, thus, cannot be assessed. Therefore, the observed relationship between
the global efficiency and medication should be viewed with caution. Interestingly, we have
not found significant correlation between the global efficiency and intake of benzodiazepines.
This negative finding suggests that even though benzodiazepines are known to have an effect
on electrophysiological correlates of brain functions, the network properties might not be
influenced. There were no significant correlations between the connectivity results and
depressive symptom severity or other parameters describing the status of depression within
the patient group. We suppose that heterogeneity of our dataset, in which patients with
different disorders were included, might underlie this observation. We also found no relation
between the connectivity results and education level or age. This finding suggests
independence of the observed impairment on these demographic variables, however, the
current sample size might be insufficient for such investigations.

Conclusions

We found an overall increase in power in theta and alpha frequency bands in depressive
patients compared to healthy controls in the subcortical regions constituting the cortico-
striatal-pallidal-thalamic and limbic circuits. The network measures showed a higher than
normal functional connectivity arising from the right amygdala in depressive patients. The
amygdala seems to play an important role in neurobiology of depression. Resting-state EEG
directed functional connectivity is a useful tool for studying abnormal brain activity in
depression.

Methods

Subjects. Data were collected from 26 depressive patients and 25 healthy controls. The two groups were matched by gender and there were no significant differences in age or education (see Table 1). On a subsample of this dataset we recently showed that the severity of depressive symptoms correlates with resting-state microstate dynamics[117]. The patients were recruited at the Department of Psychiatry, Faculty of Medicine, Masaryk University and University Hospital Brno, Czech Republic. The diagnostic process had two steps and was determined based on the clinical evaluation by two board-certified psychiatrists. First, the diagnosis was made according to the criteria for research of the International Classification of Disorders (ICD-10). Second, the diagnosis was confirmed by the Mini International Neuropsychiatric interview (M.I.N.I.) according to the Diagnostic and Statistical Manual (DSM-V). All patients were examined in the shortest time period after the admission and before the stabilization of treatment, typically during their first week of hospitalization. All patients met the criteria for at least a moderate degree of depression within the following affective disorders: bipolar affective disorder (F31), depressive episode (F32), recurrent depressive disorder (F33). Exclusion criteria for patients were any psychiatric or neurological comorbidity, IQ < 70, organic disorder with influence on the brain function, alcohol dependence or other substance dependence. All patients were in the on-medication state with marked interindividual variability in specific medicaments received. Control subjects were recruited by general practitioners from their database of clients. Control subjects underwent the M.I.N.I. by board-certified psychiatrists, to ensure that they had no previous or current psychiatric disorder according to the DSM-V criteria. The scores on the Montgomery-Åsberg Depression Rating Scale (MADRS), a specific questionnaire validated for patients with mood disorders [118] and the Clinical Global Impression (CGI) [119], a general test validated for mental disorders, were used to evaluate the severity of depressive symptoms in patients. The status of depression was further described with life time count of depressive episodes and illness duration. Medication in 24 hours preceding the EEG examination was also recorded (see Table 2). This study was carried out in accordance with the recommendations of Ethics Committee of University Hospital Brno with written informed consent from all subjects.

EEG - data acquisition and pre-processing steps. Subjects were sitting in a comfortable upright position in an electrically shielded room with dimmed light. They were instructed to stay as calm as possible, to keep their eyes closed and to relax for 15 minutes. They were asked to stay awake. All participants were monitored by the cameras and in the event of signs

of nodding off or EEG signs of drowsiness detected by visual inspection, the recording was
stopped. The EEG was recorded with a high density 128-channel system (EGI System 400;
Electrical Geodesic Inc., OR, USA), f_s = 1kHz, and Cz as acquisition reference.

Five minutes of EEG data were selected and visually assessed. Noisy channels with abundant
artifacts were identified. EEG signal was band-pass filtered between 1 and 40 Hz with a 2nd-
order Butterworth filter avoiding phase-distortion. Subsequently, in order to remove
physiological artifacts, e.g. ballistocardiogram and oculo-motor artifacts, infomax-based
Independent Component Analysis [120] was applied on all but one or two noisy channels. Only
components related to ballistocardiogram, saccadic eye movements, and eye blinking were
removed based on the waveform, topography and time course of the component. Then, the
cleaned EEG recording was down-sampled at f_s = 250 Hz and the previously identified noisy
channels were interpolated using a three-dimensional spherical spline [121], and re-referenced to
the average reference. For the following analyses, thirty 2-s EEG epochs free of artifacts were
selected per subject. All the pre-processing steps were done using the freely available Cartool
Software 3.70, programmed by Denis Brunet [122] and custom functions in MATLAB®
R2018b.

EEG source estimation. We applied the LAURA algorithm implemented in Cartool [122] to
compute the source reconstruction taking into account the patient's age to calibrate the skull
conductivity [123] [124] [125]. The method restricts the solution space to the gray matter of the brain.
Then, the cortex was parcellated into the 90 Automated Anatomical Labeling brain regions
[126]. The dipoles in each ROI were represented with one unique time-series by a singular-value
decomposition [127].

Time-variant multivariate autoregressive modeling. The cortical waveforms computed
after applying the singular-value decomposition, were fitted against a time-variant (tv)
multivariate (MV) autoregressive (AR) model to overcome the problem of non-stationarity of
the EEG data. If the EEG data are available as several trials of the same length, the cortical
waveforms computed from the EEG data generates a collection of realizations of a
multivariate stochastic process which can be combined in a multivariate, multi-trial time
series [127] [128] [67]. The tv-MVAR matrices containing the model coefficients were computed in
the framework of a MATLAB toolbox (code available upon reasonable request to the
authors) that implements the adaptive Kalman filtering and information Partial Directed
Coherence (iPDC) in the source space [67] [129] [130]. The model order of the tv-MAR and the
Kalman filter adaptation constant were chosen applying the method proposed by Rubega and

colleagues [128], i.e., evaluating the partial derivatives of a residual minimization function

obtained varying simultaneously both p (p $\square$ [1, 15]) and c (c $\square$ [0, 0.03]). By means of the

model coefficients, we computed the parametric spectral power density and the iPDC absolute

values for each subject. For each patient, we obtained a 4-dimensional matrix [ROIs x ROIs x

frequency x time] that represented the directed information flow from one ROI to another for

each frequency at each time sample. In this way we performed the analysis on the *single-*

subject level to compare the two groups quantitatively.

Since the features in the power spectra were consistent among subjects in the same

population (patients vs controls), we also performed the analysis on the *population* level. A

population subject was built by estimating the tv-MVAR model, where each trial in the input

was a different subject. One power spectral density matrix and one connectivity matrix [ROIs

x ROIs x frequency x time] were obtained for each group (controls and patients). In other

words, subjects were combined as trials, assuming respectively humans as multiple

realizations of their own brain processes, with the purpose to show that the two approaches,

i.e., *single subject* and *population*, give equivalent results in differentiating patients vs

controls. In the last decade, population-based approaches were successfully exploited in

computer simulations engineered to evaluate the safety and limitations of closed-loop control

treatment algorithms [131] [132]. Population-based approaches for MVAR/PDC modelling are

currently lacking and this might be considered a first attempt justified by the consistent

features estimated in the frequency domain among subjects belonging to the same population

(patients vs controls). Further details on the connectivity estimation are reported in the

Supplementary Information.

Network metrics. In order to study the peculiarities of the brain network in patients vs

controls, the brain was represented as a digraph defined by a collection of nodes and directed

links (directional edges). Nodes in the brain network represent brain regions, i.e., the 90

ROIs, while the directed links represent the values computed by iPDC. Thus, the weight of

such link can vary in the interval [0-1] and it represents the amount of mutual information

flowing between ROIs. We defined twelve ROIs, including the bilateral amygdala, anterior

cingulum, thalamus, putamen, caudate, and pallidum, to examine the directed functional

connectivity between these seeds and the whole brain. Significant differences in power

between patients and controls were observed in the *single-subject* level in alpha and theta

frequency bands in all these six anatomical structures. Therefore, we restricted the network

analysis to this [4-12] Hz frequency range. To evaluate how much the system is fault tolerant

and how much the communication is efficient, the global efficiency for the whole brain and

the local efficiency, clustering coefficient, strength and outflow for each of these twelve

investigated ROIs were computed. To compute all the graph measures, the scripts and

functions implemented on the freely available MATLAB toolbox [133] were customized.

Global efficiency. Global efficiency is defined as the average minimum path length between

two nodes in the network. This measure is inversely related to topological distance between

nodes and is typically interpreted as a measure of the capacity for parallel information

transfer and integrated processing [134].

Local efficiency. Local efficiency is defined as the average efficiency of the local subgraphs

[135], i.e. the global efficiency computed on the neighborhood of the node. It reflects the ability

of a network to transmit information at the local level. This quantity plays a role similar to the

clustering coefficient since it reveals how much the system is fault tolerant, i.e., it shows how

efficient the communication is between the first neighbors of i when i is removed.

483
$$\overrightarrow{E_{loc}} = \frac{1}{2n} \sum_{i \in N} \frac{\sum_{j,h \in N, j \neq i}(a_{ij}+a_{ji})(a_{ih}+a_{hi})([\overrightarrow{d_{jh}}(N_i)]^{-1}+[\overrightarrow{d_{hj}}(N_i)]^{-1})}{(k_i^{out}+k_i^{in})(k_i^{out}+k_i^{in}-1)-2\sum_{j \in N} a_{ij}a_{ji}} \quad (1)$$

where k_i^{out} is the out-degree of node i, k_i^{in} is the in-degree of node i, and a_{ij} is the

connection status between node i and node j, i.e., $a_{ij} = 1$ if the link between i and j exists,

$a_{ij} = 0$ otherwise. N is the set of nodes in the network. n is the number of nodes and

$\overrightarrow{d_{jh}}(N_i)$ is the length of the shortest directed path between j (any node in the network) and h

(any node that neighbors with i).

Clustering coefficient. Clustering coefficient reflects the prevalence of clustered

connectivity around an individual brain region [136]:

492
$$cc_i = \frac{2t_i}{k_i(k_i-1)} \quad (2)$$

493 where t_i are the number of triangles around the node i, and k_i is the degree of node i, i.e., the

494 number of links connected to node i. In our case of a weighted directed network, a weighted

495 directed version of clustering coefficient was used [137]:

496
$$\overrightarrow{cc_i} = \frac{\overrightarrow{t_i}}{\left(k_i^{out}+k_i^{in}\right)\left(k_i^{out}+k_i^{in}-1\right)-2\sum_{j\in N}a_{ij}a_{ji}} \quad (3)$$

where $\overrightarrow{t_i}$ are the number of directed triangles around the node i, k_i^{out} is the out-degree of

node i, k_i^{in} is the in-degree of node i, and a_{ij} is the connection status between the nodes i and

j, i.e., $a_{ij} = 1$ if the link between i and j exists, $a_{ij} = 0$ otherwise. N is the set of nodes in

the network.

Strength and outflow. Finally, the connectivity patterns between the different cortical

regions were summarized by representing the strength that quantifies for each node the sum

of weights of all links connected to the node and the total outflow from a region toward the

others, generated by the sum of all the statistically significant links obtained by application of

the iPDC. The greatest amount of information outflow depicts the ROI as one of the main

sources (drivers) of functional connections to the other ROIs

[138].

Laterality. For all the network metrics explained in the previous paragraph, we also

computed a laterality index, which is defined as $\frac{Left_{metric}-Right_{metric}}{Left_{metric}+Right_{metric}}$ to test if the measures

significantly differentiate between the two hemispheres. Laterality index and all network

metrics were calculated for both groups.

Statistical analysis. To assess whether or not the changes in the network metrics were

statistically significant between patients and controls, paired Student's t-tests were computed

under the hypothesis of normal distribution of samples (Lilliefors test), otherwise Wilcoxon

rank-sign tests were considered. To test whether the age and education level predict the

values of the spectral power distribution and the network metrics in patients, a multiple linear

regression was performed. We also tested the influence of the clinical data on the

connectivity results. A multiple linear regression was performed exploiting correlation of the

connectivity results with four variables describing the status of depression and two variables

describing the medication status in terms of the intake of benzodiazepines (BZP),

antidepressants, antipsychotics, and mood stabilizers (AD/AP/MS). These six clinical

variables are provided for each patient in Table 2. We checked through the following

multiple linear regression models (4) (5), if the response variable Y depends on a number of

predictor variables X_i:

525
$$Y = \beta_0 + \beta_1 X_1 + \cdots \beta_k X_k + \varepsilon \qquad (4)$$

$$526 \qquad Y = \beta_0 + \beta_1 X + \beta_2 X^2 + \varepsilon \qquad\qquad (5)$$

where the ε are the residual terms of the model and β_0, β_1, β_2, ..., β_k are the k regression

coefficients. Both the clinical data and the power and network metrics were used once as

predictors and once as response variables.

Ethics statement. All participants gave their written informed consent prior to the

experiment and the study received the approval of the Ethics Committee of University

Hospital Brno in Brno, Czech Republic. All experiments of this study were performed in

accordance with relevant guidelines and regulations.

References

1.　Andrade, L. *et al.* The epidemiology of major depressive episodes: results from the

International Consortium of Psychiatric Epidemiology (ICPE) surveys. *Int. J. Methods*

Psychiatr. Res. **12**, 3–21 (2003).

2.　Bora, E., Harrison, B. J., Davey, C. G., Yü Cel, M. & Pantelis, C. Meta-analysis of volumetric

abnormalities in cortico-striatal-pallidal-thalamic circuits in major depressive disorder.

doi:10.1017/S0033291711001668

3.　Yang, J. *et al.* Amygdala Atrophy and Its Functional Disconnection with the Cortico-Striatal-

Pallidal-Thalamic Circuit in Major Depressive Disorder in Females. *PLoS One* **12**, e0168239

(2017).

4.　Zhang, B. *et al.* Mapping anhedonia-specific dysfunction in a transdiagnostic approach: an ALE

meta-analysis. *Brain Imaging Behav.* **10**, 920–939 (2016).

5.　Disner, S. G., Beevers, C. G., Haigh, E. A. P. & Beck, A. T. Neural mechanisms of the cognitive

model of depression. *Nat. Rev. Neurosci.* **12**, 467–477 (2011).

6.　Surguladze, S. *et al.* A differential pattern of neural response toward sad versus happy facial

expressions in major depressive disorder. *Biol. Psychiatry* **57**, 201–209 (2005).

7.　Sheline, Y. I. *et al.* Increased amygdala response to masked emotional faces in depressed

subjects resolves with antidepressant treatment: an fMRI study. *Biol. Psychiatry* **50**, 651–658

(2001).

8.   Siegle, G. J., Thompson, W., Carter, C. S., Steinhauer, S. R. & Thase, M. E. Increased Amygdala
and Decreased Dorsolateral Prefrontal BOLD Responses in Unipolar Depression: Related and
Independent Features. *Biol. Psychiatry* **61**, 198–209 (2007).

9.   Nugent, A. C., Robinson, S. E., Coppola, R., Furey, M. L. & Zarate, C. A. Group differences in
MEG-ICA derived resting state networks: Application to major depressive disorder.
Neuroimage **118**, 1–12 (2015).

10. Knyazev, G. G. *et al.* Task-positive and task-negative networks in major depressive disorder: A
combined fMRI and EEG study. *J. Affect. Disord.* **235**, 211–219 (2018).

11. Lu, Y. *et al.* The volumetric and shape changes of the putamen and thalamus in first episode,
untreated major depressive disorder. *NeuroImage. Clin.* **11**, 658–666 (2016).

12. Kim, M. J., Hamilton, J. P. & Gotlib, I. H. Reduced caudate gray matter volume in women with
major depressive disorder. *Psychiatry Res. Neuroimaging* **164**, 114–122 (2008).

13. Sheline, Y. I., Price, J. L., Yan, Z. & Mintun, M. A. Resting-state functional MRI in depression
unmasks increased connectivity between networks via the dorsal nexus. *Proc. Natl. Acad. Sci.*
107, 11020–11025 (2010).

14. Kuhn, S. & Gallinat, J. Resting-State Brain Activity in Schizophrenia and Major Depression: A
Quantitative Meta-Analysis. *Schizophr. Bull.* **39**, 358–365 (2013).

15. Lorenzetti, V., Allen, N. B., Fornito, A. & Yücel, M. Structural brain abnormalities in major
depressive disorder: A selective review of recent MRI studies. *J. Affect. Disord.* **117**, 1–17
(2009).

16. Veer, I. M. Whole brain resting-state analysis reveals decreased functional connectivity in
major depression. *Front. Syst. Neurosci.* **4**, (2010).

17. Hamilton, J. P. *et al.* Functional Neuroimaging of Major Depressive Disorder: A Meta-Analysis
and New Integration of Baseline Activation and Neural Response Data. *Am. J. Psychiatry* **169**,
693–703 (2012).

18. Bielau, H. *et al.* Volume deficits of subcortical nuclei in mood disorders. *Eur. Arch. Psychiatry*
Clin. Neurosci. **255**, 401–412 (2005).

19. Holtzheimer, P. E. & Mayberg, H. S. Stuck in a rut: rethinking depression and its treatment.

Trends Neurosci. **34**, 1–9 (2011).

20.   Drobisz, D. & Damborská, A. Deep brain stimulation targets for treating depression. *Behav.*
584   *Brain Res.* **359**, 266–273 (2019).

585   21.   Holtzheimer, P. E. *et al.* Subcallosal Cingulate Deep Brain Stimulation for Treatment-Resistant
586   Unipolar and Bipolar Depression. *Arch. Gen. Psychiatry* **69**, 150 (2012).

587   22.   KNIGHT, G. STEREOTACTIC TRACTOTOMY IN THE SURGICAL TREATMENT OF MENTAL ILLNESS.
588   *J. Neurol. Neurosurg. Psychiatry* **28**, 304–310 (1965).

589   23.   Dougherty, D. D. *et al.* Cerebral metabolic correlates as potential predictors of response to
590   anterior cingulotomy for treatment of major depression. *J. Neurosurg.* **99**, 1010–1017 (2003).

591   24.   Hamani, C. *et al.* Deep brain stimulation in rats: Different targets induce similar
592   antidepressant-like effects but influence different circuits. *Neurobiol. Dis.* **71**, 205–214 (2014).

593   25.   Hamani, C. & Nóbrega, J. N. Deep brain stimulation in clinical trials and animal models of
594   depression. *Eur. J. Neurosci.* **32**, 1109–1117 (2010).

595   26.   Hamani, C. *et al.* Antidepressant-Like Effects of Medial Prefrontal Cortex Deep Brain
596   Stimulation in Rats. *Biol. Psychiatry* **67**, 117–124 (2010).

597   27.   Moshe, H. *et al.* Prelimbic Stimulation Ameliorates Depressive-Like Behaviors and Increases
598   Regional BDNF Expression in a Novel Drug-Resistant Animal Model of Depression. *Brain*
599   *Stimul.* **9**, 243–250 (2016).

600   28.   Thiele, S., Furlanetti, L., Pfeiffer, L. M., Coenen, V. A. & Döbrössy, M. D. The effects of
601   bilateral, continuous, and chronic Deep Brain Stimulation of the medial forebrain bundle in a
602   rodent model of depression. *Exp. Neurol.* **303**, 153–161 (2018).

603   29.   Rummel, J. *et al.* Testing different paradigms to optimize antidepressant deep brain
604   stimulation in different rat models of depression. *J. Psychiatr. Res.* **81**, 36–45 (2016).

605   30.   Clemm Von Hohenberg, C. *et al.* Lateral habenula perturbation reduces default-mode
606   network connectivity in a rat model of depression. *Transl. Psychiatry* **8**, (2018).

607   31.   Baeken, C., Duprat, R., Wu, G. R., De Raedt, R. & van Heeringen, K. Subgenual Anterior
608   Cingulate–Medial Orbitofrontal Functional Connectivity in Medication-Resistant Major

Depression: A Neurobiological Marker for Accelerated Intermittent Theta Burst Stimulation
Treatment? *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* **2**, 556–565 (2017).

32.   Johansen-Berg, H. *et al.* Anatomical connectivity of the subgenual cingulate region targeted
with deep brain stimulation for treatment-resistant depression. *Cereb. Cortex* **18**, 1374–1383
(2008).

33.   Greicius, M. D. *et al.* Resting-State Functional Connectivity in Major Depression: Abnormally
Increased Contributions from Subgenual Cingulate Cortex and Thalamus. *Biol. Psychiatry* **62**,
429–437 (2007).

34.   Riva-Posse, P. *et al.* Defining critical white matter pathways mediating successful subcallosal
cingulate deep brain stimulation for treatment-resistant depression. *Biol. Psychiatry* **76**, 963–
969 (2014).

35.   Quevedo, K. *et al.* Ventral Striatum Functional Connectivity during Rewards and Losses and
Symptomatology in Depressed Patients. *Biol. Psychol.* **123**, 62–73 (2017).

36.   Gutman, D. A., Holtzheimer, P. E., Behrens, T. E. J., Johansen-Berg, H. & Mayberg, H. S. A
Tractography Analysis of Two Deep Brain Stimulation White Matter Targets for Depression.
Biol. Psychiatry **65**, 276–282 (2009).

37.   Bracht, T. *et al.* White matter microstructure alterations of the medial forebrain bundle in
melancholic depression. *J. Affect. Disord.* **155**, 186–193 (2014).

38.   Kaiser, R. H., Andrews-Hanna, J. R., Wager, T. D. & Pizzagalli, D. A. Large-scale network
dysfunction in major depressive disorder: A meta-analysis of resting-state functional
connectivity. *JAMA Psychiatry* **72**, 603–611 (2015).

39.   Smith, S. M. *et al.* Functional connectomics from resting-state fMRI. *Trends in Cognitive*
Sciences **17**, 666–682 (2013).

40.   Fox, M. D. & Raichle, M. E. Spontaneous fluctuations in brain activity observed with
functional magnetic resonance imaging. *Nature Reviews Neuroscience* **8**, 700–711 (2007).

41.   Hamilton, J. P. *et al.* Default-Mode and Task-Positive Network Activity in Major Depressive
Disorder: Implications for Adaptive and Maladaptive Rumination. *Biol. Psychiatry* **70**, 327–333
(2011).

42.   Lui, S. *et al.* Resting-state functional connectivity in treatment-resistant depression. *Am. J.*
Psychiatry **168**, 642–648 (2011).

43.   Whitton, A. E. *et al.* Electroencephalography Source Functional Connectivity Reveals
Abnormal High-Frequency Communication Among Large-Scale Functional Networks in
Depression. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* **3**, 50–58 (2018).

44.   Sikora, M. *et al.* Salience Network Functional Connectivity Predicts Placebo Effects in Major
Depression. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* **1**, 68–76 (2016).

45.   Gong, J. Y. *et al.* Disrupted functional connectivity within the default mode network and
salience network in unmedicated bipolar II disorder. *Prog. Neuro-Psychopharmacology Biol.*
Psychiatry **88**, 11–18 (2019).

46.   Sacchet, M. D. *et al.* Large-scale hypoconnectivity between resting-state functional networks
in unmedicated adolescent major depressive disorder. *Neuropsychopharmacology* **41**, 2951–
2960 (2016).

47.   Williams, K. A., Mehta, N. S., Redei, E. E., Wang, L. & Procissi, D. Aberrant resting-state
functional connectivity in a genetic rat model of depression. *Psychiatry Res. - Neuroimaging*
222, 111–113 (2014).

48.   Kopell, B. H., Greenberg, B. & Rezai, A. R. Deep Brain Stimulation for Psychiatric Disorders. *J.*
Clin. Neurophysiol. **21**, 51–67 (2004).

49.   Sartorius, A. *et al.* Remission of Major Depression Under Deep Brain Stimulation of the Lateral
Habenula in a Therapy-Refractory Patient. *Biological Psychiatry* **67**, (2010).

50.   Kukleta, M., Bob, P., Brázdil, M., Roman, R. & Rektor, I. The level of frontal-temporal beta-2
band EEG synchronization distinguishes anterior cingulate cortex from other frontal regions.
Conscious. Cogn. **19**, 879–886 (2010).

51.   Brázdil, M. *et al.* Directional functional coupling of cerebral rhythms between anterior
cingulate and dorsolateral prefrontal areas during rare stimuli: A directed transfer function
analysis of human depth EEG signal. *Hum. Brain Mapp.* **30**, 138–146 (2009).

52.   Kibleur, A. *et al.* Stimulation of subgenual cingulate area decreases limbic top-down effect on
ventral visual stream: A DBS-EEG pilot study. *Neuroimage* **146**, 544–553 (2017).

53.   Pereda, E., Quiroga, R. Q. & Bhattacharya, J. Nonlinear multivariate analysis of
666   neurophysiological signals. *Prog. Neurobiol.* **77**, 1–37 (2005).

667   54.   Seth, A. K., Barrett, A. B. & Barnett, L. Granger Causality Analysis in Neuroscience and
668   Neuroimaging. *J. Neurosci.* **35**, 3293–3297 (2015).

669   55.   Granger, C. W. J. Investigating Causal Relations by Econometric Models and Cross-spectral
670   Methods. *Econometrica* **37**, 424–438 (1969).

671   56.   Leistritz, L. *et al.* Connectivity Analysis of Somatosensory Evoked Potentials in Patients with
672   Major Depression. *Methods Inf. Med.* **49**, 484–491 (2010).

673   57.   Yu Sun, Sijung Hu, Chambers, J., Yisheng Zhu & Shanbao Tong. Graphic patterns of cortical
674   functional connectivity of depressed patients on the basis of EEG measurements. in *2011*
675   *Annual International Conference of the IEEE Engineering in Medicine and Biology Society*
676   1419–1422 (IEEE, 2011). doi:10.1109/IEMBS.2011.6090334

677   58.   Tang, Y. *et al.* The altered cortical connectivity during spatial search for facial expressions in
678   major depressive disorder. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* **35**, 1891–1900
679   (2011).

680   59.   Mao, W., Li, Y., Tang, Y., Li, H. & Wang, J. The coherence changes in the depressed patients in
681   response to different facial expressions. in *Lecture Notes in Computer Science (including*
682   *subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)* **6064**
683   **LNCS**, 392–399 (2010).

684   60.   Wang, C. *et al.* The brain network research of poststroke depression based on partial directed
685   coherence (PDC). *Chinese J. Biomed. Eng.* **34**, 385–391 (2015).

686   61.   Sun, Y., Li, Y., Zhu, Y., Chen, X. & Tong, S. Electroencephalographic differences between
687   depressed and control subjects: An aspect of interdependence analysis. *Brain Res. Bull.* **76**,
688   559–564 (2008).

689   62.   Schoffelen, J.-M. & Gross, J. Source connectivity analysis with MEG and EEG. *Hum. Brain*
690   *Mapp.* **30**, 1857–1865 (2009).

691   63.   He, B. *et al.* Electrophysiological Brain Connectivity: Theory and Implementation. *IEEE Trans.*
692   *Biomed. Eng.* **66**, 2115–2137 (2019).

693   64.   Coito, A., Michel, C. M., van Mierlo, P., Vulliemoz, S. & Plomp, G. Directed Functional Brain
694   Connectivity Based on EEG Source Imaging: Methodology and Application to Temporal Lobe
695   Epilepsy. *IEEE Trans. Biomed. Eng.* **63**, 2619–2628 (2016).

696   65.   Sperdin, H. F. *et al.* Early alterations of social brain networks in young children with autism.
697   *Elife* **7**, (2018).

698   66.   Coito, A., Michel, C. M., Vulliemoz, S. & Plomp, G. Directed functional connections underlying
699   spontaneous brain activity. *Hum. Brain Mapp.* **40**, 879–888 (2019).

700   67.   Milde, T. *et al.* A new Kalman filter approach for the estimation of high-dimensional time-
701   variant multivariate AR models and its application in analysis of laser-evoked brain potentials.
702   *Neuroimage* **50**, 960–969 (2010).

703   68.   Pessoa, L. & Adolphs, R. Emotion processing and the amygdala: from a 'low road' to 'many
704   roads' of evaluating biological significance. *Nat. Rev. Neurosci.* **11**, 773–782 (2010).

705   69.   Zheng, J. *et al.* Amygdala-hippocampal dynamics during salient information processing. *Nat.*
706   *Commun.* **8**, 14413 (2017).

707   70.   Freese, Jennifer L.; Amaral, D. G. *Neuroanatomy of the primate amygdala. - PsycNET.*
708   (Guilford Press, 2009).

709   71.   Kober, H. *et al.* Functional grouping and cortical–subcortical interactions in emotion: A meta-
710   analysis of neuroimaging studies. *Neuroimage* **42**, 998–1031 (2008).

711   72.   Thomas Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic
712   functional connectivity. *J. Neurophysiol.* **106**, 1125–1165 (2011).

713   73.   Price, J. L. & Drevets, W. C. Neurocircuitry of Mood Disorders. *Neuropsychopharmacology* **35**,
714   192–216 (2010).

715   74.   Hamilton, J. P., Chen, M. C. & Gotlib, I. H. Neural systems approaches to understanding major
716   depressive disorder: An intrinsic functional organization perspective. *Neurobiol. Dis.* **52**, 4–11
717   (2013).

718   75.   Ramasubbu, R. *et al.* Reduced Intrinsic Connectivity of Amygdala in Adults with Major
719   Depressive Disorder. *Front. Psychiatry* **5**, (2014).

720   76.   Tang, S. *et al.* Abnormal amygdala resting-state functional connectivity in adults and
721   adolescents with major depressive disorder: A comparative meta-analysis. *EBioMedicine* **36**,
722   436–445 (2018).

723   77.   Tang, S. *et al.* Anomalous functional connectivity of amygdala subregional networks in major
724   depressive disorder. *Depress. Anxiety* **36**, 712–722 (2019).

725   78.   Zhang, X. F., He, X., Wu, L., Liu, C. J. & Wu, W. Altered Functional Connectivity of Amygdala
726   with the Fronto-Limbic-Striatal Circuit in Temporal Lobe Lesion as a Proposed Mechanism for
727   Poststroke Depression. *Am. J. Phys. Med. Rehabil.* **98**, 303–310 (2019).

728   79.   Ferdek, M. A., van Rijn, C. M. & Wyczesany, M. Depressive rumination and the emotional
729   control circuit: An EEG localization and effective connectivity study. *Cogn. Affect. Behav.*
730   *Neurosci.* **16**, 1099–1113 (2016).

731   80.   van Eijndhoven, P. *et al.* Amygdala Volume Marks the Acute State in the Early Course of
732   Depression. *Biol. Psychiatry* **65**, 812–818 (2009).

733   81.   Sandu, A.-L. *et al.* Amygdala and regional volumes in treatment-resistant *versus*
734   nontreatment-resistant depression patients. *Depress. Anxiety* **34**, 1065–1071 (2017).

735   82.   Bauer, I. E. *et al.* Amygdala enlargement in unaffected offspring of bipolar parents. *J.*
736   *Psychiatr. Res.* **59**, 200–205 (2014).

737   83.   Inman, C. S. *et al.* Direct electrical stimulation of the amygdala enhances declarative memory
738   in humans. *Proc. Natl. Acad. Sci.* **115**, 98–103 (2018).

739   84.   Bijanki, K. R. *et al.* Case Report: Stimulation of the Right Amygdala Induces Transient Changes
740   in Affective Bias. *Brain Stimul.* **7**, 690–693 (2014).

741   85.   Tyrand, R., Seeck, M., Pollo, C. & Boëx, C. Effects of amygdala–hippocampal stimulation on
742   synchronization. *Epilepsy Res.* **108**, 327–330 (2014).

743   86.   Tyrand, R. *et al.* Effects of amygdala–hippocampal stimulation on interictal epileptic
744   discharges. *Epilepsy Res.* **99**, 87–93 (2012).

745   87.   Langevin, J.-P. *et al.* Deep Brain Stimulation of the Basolateral Amygdala: Targeting Technique
746   and Electrodiagnostic Findings. *Brain Sci.* **6**, 28 (2016).

747   88.   Koek, R. J. *et al.* Deep brain stimulation of the basolateral amygdala for treatment-refractory
748   combat post-traumatic stress disorder (PTSD): study protocol for a pilot randomized
749   controlled trial with blinded, staggered onset of stimulation. *Trials* **15**, 356 (2014).

750   89.   Sturm, V. *et al.* DBS in the basolateral amygdala improves symptoms of autism and related
751   self-injurious behavior: a case report and hypothesis on the pathogenesis of the disorder.
752   *Front. Hum. Neurosci.* **6**, 341 (2013).

753   90.   Admon, R. *et al.* Striatal hypersensitivity during stress in remitted individuals with recurrent
754   depression. *Biol. Psychiatry* **78**, 67–76 (2015).

755   91.   Marchand, W. R. & Yurgelun-Todd, D. Striatal structure and function in mood disorders: a
756   comprehensive review. *Bipolar Disord.* **12**, 764–785 (2010).

757   92.   Bluhm, R. *et al.* Resting state default-mode network connectivity in early depression using a
758   seed region-of-interest analysis: Decreased connectivity with caudate nucleus. *Psychiatry*
759   *Clin. Neurosci.* **63**, 754–761 (2009).

760   93.   Butters, M. A. *et al.* Three-Dimensional Surface Mapping of the Caudate Nucleus in Late-Life
761   Depression. *Am. J. Geriatr. Psychiatry* **17**, 4–12 (2009).

762   94.   Ma, C. *et al.* Resting-State Functional Connectivity Bias of Middle Temporal Gyrus and
763   Caudate with Altered Gray Matter Volume in Major Depression. *PLoS One* **7**, e45263 (2012).

764   95.   Krishnan, K. R. R. Magnetic Resonance Imaging of the Caudate Nuclei in Depression. *Arch.*
765   *Gen. Psychiatry* **49**, 553 (1992).

766   96.   Tymofiyeva, O. *et al.* DTI-based connectome analysis of adolescents with major depressive
767   disorder reveals hypoconnectivity of the right caudate. *J. Affect. Disord.* **207**, 18–25 (2017).

768   97.   Khundakar, A., Morris, C., Oakley, A. & Thomas, A. J. Morphometric Analysis of Neuronal and
769   Glial Cell Pathology in the Caudate Nucleus in Late-Life Depression. *Am. J. Geriatr. Psychiatry*
770   **19**, 132–141 (2011).

771   98.   Hannestad, J. *et al.* White matter lesion volumes and caudate volumes in late-life depression.
772   *Int. J. Geriatr. Psychiatry* **21**, 1193–1198 (2006).

773   99.   Pillay, S. A quantitative magnetic resonance imaging study of caudate and lenticular nucleus
774   gray matter volume in primary unipolar major depression: relationship to treatment response

775  and clinical severity. *Psychiatry Res. Neuroimaging* **84**, 61–74 (1998).

776 100. Price, J. L. & Drevets, W. C. Neural circuits underlying the pathophysiology of mood disorders.
777  *Trends Cogn. Sci.* **16**, 61–71 (2012).

778 101. Limbic-cortical dysregulation: a proposed model of depression. *J. Neuropsychiatry Clin.*
779  *Neurosci.* **9**, 471–481 (1997).

780 102. Pizzagalli, D. A. Frontocingulate Dysfunction in Depression: Toward Biomarkers of Treatment
781  Response. *Neuropsychopharmacology* **36**, 183–206 (2011).

782 103. Mégevand, P. *et al.* Electric source imaging of interictal activity accurately localises the
783  seizure onset zone. *J. Neurol. Neurosurg. Psychiatry* **85**, 38–43 (2014).

784 104. Michel, C. M. *et al.* 128-Channel EEG source imaging in epilepsy: Clinical yield and localization
785  precision. *J. Clin. Neurophysiol.* **21**, 71–83 (2004).

786 105. Attal, Y. & Schwartz, D. Assessment of Subcortical Source Localization Using Deep Brain
787  Activity Imaging Model with Minimum Norm Operators: A MEG Study. *PLoS One* **8**, 59856
788  (2013).

789 106. Krishnaswamy, P. *et al.* Sparsity enables estimation of both subcortical and cortical activity
790  from MEG and EEG. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E10465–E10474 (2017).

791 107. Seeber, M. *et al.* Subcortical electrophysiological activity is detectable with high-density EEG
792  source imaging. *Nat. Commun.* **10**, 753 (2019).

793 108. Pizzo, F. *et al.* Deep brain activities can be detected with magnetoencephalography. *Nat.*
794  *Commun.* **10**, 971 (2019).

795 109. Aouizerate, B. *et al.* Deep brain stimulation of the ventral caudate nucleus in the treatment of
796  obsessive—compulsive disorder and major depression. *J. Neurosurg.* **101**, 682–686 (2004).

797 110. Millet, B. *et al.* Limbic versus cognitive target for deep brain stimulation in treatment-
798  resistant depression: Accumbens more promising than caudate. *Eur. Neuropsychopharmacol.*
799  **24**, 1229–1239 (2014).

800 111. Grin-Yatsenko, V. A., Baas, I., Ponomarev, V. A. & Kropotov, J. D. EEG Power Spectra at Early
801  Stages of Depressive Disorders. *J. Clin. Neurophysiol.* **26**, 401–406 (2009).

802   112.   Pollock, V. E. & Schneider, L. S. Topographic Quantitative EEG in Elderly Subjects with Major
803   Depression. *Psychophysiology* **27**, 438–444 (1990).

804   113.   Roemer, R. A., Shagass, C., Dubin, W., Jaffe, R. & Siegal, L. Quantitative EEG in elderly
805   depressives. *Brain Topogr.* **4**, 285–290 (1992).

806   114.   Kwon, J. S., Youn, T. & Jung, H. Y. Right hemisphere abnormalities in major depression:
807   Quantitative electroencephalographic findings before and after treatment. *J. Affect. Disord.*
808   **40**, 169–173 (1996).

809   115.   Jiang, H. *et al.* Predictability of depression severity based on posterior alpha oscillations. *Clin.*
810   *Neurophysiol.* **127**, 2108–2114 (2016).

811   116.   Neumann, W.-J. *et al.* Different patterns of local field potentials from limbic DBS targets in
812   patients with major depressive and obsessive compulsive disorder. *Mol. Psychiatry* **19**, 1186–
813   1192 (2014).

814   117.   Damborská, A. *et al.* EEG Resting-State Large-Scale Brain Network Dynamics Are Related to
815   Depressive Symptoms. *Front. Psychiatry* **10**, (2019).

816   118.   Williams, J. B. W. & Kobak, K. A. Development and reliability of a structured interview guide
817   for the Montgomery-Åsberg Depression Rating Scale (SIGMA). *Br. J. Psychiatry* **192**, 52–58
818   (2008).

819   119.   Guy, W. *ECDEU assessment manual for psychopharmacology.* (U.S. Dept. of Health Education
820   and Welfare Public Health Service Alcohol Drug Abuse and Mental Health Administration
821   National Institute of Mental Health Psychopharmacology Research Branch, 1976).

822   120.   Jung, T.-P. *et al.* Removal of eye activity artifacts from visual event-related potentials in
823   normal and clinical subjects. *Clin. Neurophysiol.* **111**, 1745–1758 (2000).

824   121.   Perrin, F., Pernier, J., Bertrand, O. & Echallier, J. F. Spherical splines for scalp potential and
825   current density mapping. *Electroencephalogr. Clin. Neurophysiol.* **72**, 184–187 (1989).

826   122.   "the Cartool Community group," [Online]. Available: cartoolcommunity.unige.ch.

827   123.   Grave de Peralta Menendez, R., Murray, M. M., Michel, C. M., Martuzzi, R. & Gonzalez
828   Andino, S. L. Electrical neuroimaging based on biophysical constraints. *Neuroimage* **21**, 527–
829   539 (2004).

830   124.   Michel, C. M. & Brunet, D. EEG Source Imaging: A Practical Review of the Analysis Steps.
831   *Front. Neurol.* **10**, 325 (2019).

832   125.   Spinelli, L., Andino, S. G., Lantz, G., Seeck, M. & Michel, C. M. Electromagnetic Inverse
833   Solutions in Anatomically Constrained Spherical Head Models. *Brain Topogr.* **13**, 115–125
834   (2000).

835   126.   Tzourio-Mazoyer, N. *et al.* Automated Anatomical Labeling of Activations in SPM Using a
836   Macroscopic Anatomical Parcellation of the MNI MRI Single-Subject Brain. *Neuroimage* **15**,
837   273–289 (2002).

838   127.   Rubega, M. *et al.* Estimating EEG Source Dipole Orientation Based on Singular-value
839   Decomposition for Connectivity Analysis. *Brain Topogr.* 1–16 (2018). doi:10.1007/s10548-
840   018-0691-2

841   128.   Rubega M., Pascucci D., Rué Queralt J., Van Mierlo P., Hagmann P., Plomp G., M. C. M. Time-
842   varying effective EEG source connectivity: The optimization of model parameters. 2019, *41st*
843   *Annual International Conference of the IEEE Engineering in Medicine and Biology Society*
844   *(EMBC), SaC05.6.*

845   129.   Takahashi, D. Y., Baccalá, L. A. & Sameshima, K. Information theoretic interpretation of
846   frequency domain connectivity measures. *Biol. Cybern.* **103**, 463–469 (2010).

847   130.   Sameshima, K., Baccala, L. A. & Baccala, L. A. *Methods in Brain Connectivity Inference through*
848   *Multivariate Time Series Analysis.* **20145078**, (CRC Press, 2014).

849   131.   Vettoretti, M., Facchinetti, A., Sparacino, G. & Cobelli, C. Type-1 Diabetes Patient Decision
850   Simulator for In Silico Testing Safety and Effectiveness of Insulin Treatments. *IEEE Trans.*
851   *Biomed. Eng.* **65**, 1281–1290 (2018).

852   132.   Man, C. D. *et al.* The UVA/PADOVA Type 1 Diabetes Simulator. *J. Diabetes Sci. Technol.* **8**, 26–
853   34 (2014).

854   133.   [Online]. Available: http://www.brain-connectivity-toolbox.net.

855   134.   Bullmore, E. & Sporns, O. The economy of brain network organization. *Nat. Rev. Neurosci.* **13**,
856   336–349 (2012).

135.   Latora, V. & Marchiori, M. Efficient Behavior of Small-World Networks. *Phys. Rev. Lett.* **87**,

198701 (2001).

136.   Watts, D. J., Strogatz, S. H. Collective dynamics of 'small-world' networks. *Nature* **393**, 440–
442 (1998).

137.   Fagiolo, G. Clustering in complex directed networks. *Phys. Rev. E* **76**, 026107 (2007).

138.   Babiloni, F. *et al.* Estimation of the cortical functional connectivity with the multimodal
integration of high-resolution EEG and fMRI data by directed transfer function. *Neuroimage*
24, 118–131 (2005).

139.   Bazire Stephen. *Benzodiazepine equivalent doses. Psychotropic Drug Directory.* (Lloyd-
Reinhold Communications, 2014).

Acknowledgments

This study was supported by the European Union Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 739939, by Ministry of Health, Czech Republic - conceptual development of research organization (University Hospital Brno - FNBr, 65269705), by the Swiss National Science Foundation (grant No. 320030_184677), and by the National Centre of Competence in Research (NCCR) "SYNAPSY–The Synaptic Basis of Mental Diseases" (NCCR Synapsy Grant # "51NF40 – 185897). CMM and MR were supported by the Swiss National Science Foundation (Sinergia project CRSII5_170873). The funding sources had no role in the design, collection, analysis, or interpretation of the study. The authors wish to thank Patrik Wahlberg, David Pascucci, and Gijs Plomp for providing useful comments on the manuscript.

Author Contributions

AD – designed the study, performed the preprocessing, and wrote the initial draft; RB and JH – were responsible for patient recruitment and clinical assessment; EH – collected the EEG data; SF and ŠO – were involved in the clinical assessment; CMM – served as an advisor, MR – performed the analysis, wrote the initial draft, and was responsible for the overall oversight of the study. All authors revised the manuscript.

Competing Interests

The authors declare no competing interests.

Table legends

Table 1. ^aEducation was classified into three levels: 1 = no high school, 2 = high school, 3 =
university studies

Table 2. F31.3 - Bipolar affective disorder, current episode mild or moderate depression;
F31.4 - Bipolar affective disorder, current episode severe depression without psychotic
symptoms; F31.5 - Bipolar affective disorder, current episode severe depression with
psychotic symptoms; F32.1 - Moderate depressive episode; F32.2 - Severe depressive
episode without psychotic symptoms; F32.3 - Severe depressive episode with psychotic
symptoms; F33.1 - Recurrent depressive disorder, current episode moderate; F33.2 -
Recurrent depressive disorder, current episode severe without psychotic symptoms; F33.3 -
Recurrent depressive disorder, current episode severe with psychotic symptoms; BZD:
benzodiazepine equivalent dose [139] AD - antidepressants (mirtazapine, citalopram,
venlafaxine, vortioxetine, sertraline, trazodone); AP - antipsychotics (risperidone, olanzapine,
quetiapine, amisulpride, aripiprazole); MS - mood stabilizers (valproate, lamotrigine,
carbamazepine); AD/AP/MS medication scale: 1 – one medication in sub-therapeutic doses, 2
– one medication in therapeutic doses, 3 – combination of medications with one in
therapeutic doses, 4 – combination of medications with more than one in therapeutic doses;
MADRS (Montgomery–Åsberg Depression Rating Scale): score is between 0 and 60, the
higher the score the higher the depressive symptom severity; CGI (Clinical Global
Impression scale): healthy (1) – most extremely ill (7). Four patients were undergoing the
first (patient 3) and second (patient 4 and 9) week of electroconvulsive therapy and the first
week of repetitive transcranial magnetic stimulation (patient 5). No clinical effect of these
neurostimulation treatments was apparent.

Table 1. Demographic data

Characteristic	Patients (n = 26)	Controls (n = 25)	*t*-value	df	p-value
Age: mean ± SD	51.9 ± 9.1	49.5 ± 8.7	0.97	49	0.34
Gender: female, *n*	11	10			

Education[a]: mean ± SD   1.9 ± 0.9   2.3 ± 0.7   -1.70   49   0.10

918   Table 2. Patient characteristics

Patient	ICD-10 diagnose	Number of episodes	Illness duration (years)	MADRS score	CGI score	BZD	ADP/AP/MS	AD/AP/MS medication scale
1	F31.4	3	2	27	4	2	AD, AP, MS	3
2	F32.2	1	0.5	24	5	0	AD	2
3	F32.1	1	1	15	4	2	AD	2
4	F31.5	5	20	39	6	0	AP	2
5	F33.1	3	7	18	4	0	AD	1
6	F33.1	2	8	9	3	1.33	AD	1
7	F32.1	1	1	24	4	1.33	AD, AP	3
8	F31.4	4	27	29	5	2	AP	2
9	F33.3	2	5	36	6	1	AD, AP	4
10	F33.1	3	19	21	4	1	AD	1
11	F33.3	2	2	38	5	6	AD, AP	4
12	F33.2	2	1	39	5	3	AD, AP	4
13	F32.3	1	0.08	21	5	2	AD, AP	4
14	F33.2	5	21	32	5	0	AD, AP	3
15	F33.3	2	2	38	6	3	AD, AP	4
16	F32.3	1	0.08	37	6	2	AD, AP	4
17	F33.1	3	4	18	4	0	AD, AP	4
18	F31.3	2	16	28	4	0	AP, MS	4
19	F31.3	11	24	23	4	1	AP, MS	4
20	F32.2	0	0,17	23	4	1	AD, AP	4
21	F33.1	1	9	34	5	2	AD	2
22	F32.3	0	0,04	37	6	1	AD, AP	4
23	F33.3	1	11	49	6	3	AD, AP	4
24	F33.1	3	20	23	4	0	AD	2
25	F33.1	5	24	26	4	2	AD, AP, MS	4
26	F32.1	0	0,17	23	4	3	AD, AP	3

 Figure legends

Figure 1 Parametric power spectral density (PSD) of the *population subjects* representing
controls (A) vs patients (B) in the subcortical regions of interest. Power significantly
increases within the interval [4-12] Hz (indicated with vertical dashed lines) in theta ([4-8]
Hz) and alpha ([8-12] Hz) bands and decreases in delta ([1-4] Hz) and beta ([12-18] Hz)
bands in patients compared to controls ($p<0.05$) in the subcortical regions of interest.
Continuous and dashed lines indicate the results for structures in the right and left
hemispheres, respectively.

Figure 2 Boxplots to graphical illustrate the distribution of power of controls (green boxes)
and patients (red boxes) in (a) [1-4] Hz, (b) [4-12] Hz and (c) [12-18] Hz. One star (*) stands
for significant statistical difference with $p<0.05$ and two stars (**) for $p<0.001$. Power in [4-
12] Hz significantly increases in patients compared to controls in all examined anatomical
brain structures.

Figure 3 Local efficiency computed in the two *population subjects* representing (a) controls
and (b) patients. Note that all subcortical regions of interest (ROIs) revealed higher values for
patients than controls corresponding to the same tendency observed in all ROIs of the brain at
the *single-subject* level (see Supplementary Fig. S1 – S2 online). The efficiency for each
ROI is represented by a sphere centered on the cortical region, whose radius is linearly
related to the magnitude. Such information is also coded through a color scale.

Figure 4 Boxplots to graphically illustrate the distribution of (a) local efficiency, (b)
clustering coefficient, (c) strength, and (d) outflow in controls (green boxes) and patients (red
boxes). One star (*) stands for significant statistical difference with $p<0.05$ and two stars (**)
for $p<0.001$. All network metrics that refer to the right amygdala significantly differ between
controls and patients ($p<0.001$), applying the Bonferroni correction ($p<0.05/12 \rightarrow p<0.0042$).

Figure 5 Relationship between the intake of antidepressants/antipsychotics/mood stabilizers
(AD/AP/MS) and the global efficiency (GE). Note that higher medication intake is associated
with lower GE. The orange dotted line stands for the predicted value of AD/AP/MS for each
patient using GE as predictor. For values of the AD/AP/MS medication scale the reader is
referred to the legend of Table 2.