Clusters of sub-Saharan African countries based on

sociobehavioural characteristics and associated HIV incidence

Aziza Merzouki[a],*, Janne Estill[a,b], Erol Orel[a], Kali Tal[c], Olivia Keiser[a]

[a]Institute of Global Health, University of Geneva, Geneva, Switzerland

[b]Institute of Mathematical Statistics and Actuarial Science, University of Bern, Bern,

Switzerland

[c]Institute of Primary Health Care (BIHAM), University of Bern, Bern, Switzerland

*Corresponding author: Aziza Merzouki

Institute of Global Health, University of Geneva

Chemin des Mines 9, 1202 Geneva, Switzerland

Tel. +41 78 712 56 46

fatmaaziza.merzouki@unige.ch

Keywords: dimensionality reduction; principal component analysis; hierarchical clustering;

unsupervised machine learning; HIV incidence; sociobehavioural characteristics

Conflicts of interest and source of funding: Authors declare no competing interest. This study

was funded by the Swiss National Science Foundation (SNF professorship grant n° 163878 to

O Keiser).

Abstract

Introduction: HIV incidence varies widely between sub-Saharan African (SSA) countries. This variation coincides with a substantial sociobehavioural heterogeneity, which complicates the design of effective interventions. In this study, we investigated how sociobehavioural heterogeneity in sub-Saharan Africa could account for the variance of HIV incidence between countries.

Methods: We analysed aggregated data, at the national-level, from the most recent Demographic and Health Surveys of 29 SSA countries [2010-2017], which included 594'644 persons (183'310 men and 411'334 women). We preselected 48 demographic, socio-economic, behavioural and HIV-related attributes to describe each country. We used Principal Component Analysis to visualize sociobehavioural similarity between countries, and to identify the variables that accounted for most sociobehavioural variance in SSA. We used hierarchical clustering to identify groups of countries with similar sociobehavioural profiles, and we compared the distribution of HIV incidence (estimates from UNAIDS) and sociobehavioural variables within each cluster.

Results: The most important characteristics, which explained 69% of sociobehavioural variance across SSA among the variables we assessed were: religion; male circumcision; number of sexual partners; literacy; uptake of HIV testing; women's empowerment; accepting attitude toward people living with HIV/AIDS; rurality; ART coverage; and, knowledge about AIDS. Our model revealed three groups of countries, each with characteristic sociobehavioural profiles. HIV incidence was mostly similar within each cluster and different between clusters (median(IQR); 0.5/1000(0.6/1000), 1.8/1000(1.3/1000) and 5.0/1000(4.2/1000)).

Introduction

The burden of HIV in sub-Saharan Africa (SSA) is the heaviest in the world; in 2017, 70% of HIV-infected people lived in this region ("Fact sheet - World AIDS Day 2018"). However, HIV prevalence and incidence vary widely between SSA countries. The region is heterogeneous and sociobehavioural and cultural factors vary widely within and between countries, complicating the design of effective interventions. This heterogeneity ensures that no "one-size-fits-all" approach will stop the epidemic. This is why WHO ("Global Health Sector Strategy on HIV 2016-2021: Towards Ending AIDS") highlights the need to use data and numerical methods to tailor interventions to specific populations and countries based on quantitative evidence.

So far, studies of HIV risk factors or risk factors for the uptake of interventions against HIV have generally been limited to specific sub-populations (Sangowawa & Owoaje, 2012; Kidman & Anglewicz, 2016; Ashaba et al., 2018), sub-national regions (Bailey et al., 2007; Gray et al., 2007; Eaton et al., 2014; Pons-Duran et al., 2016) or single countries (Antelman et al., 2007; Gregson et al., 2010; Tsai & Venkataramani, 2015; Lakew, Benedict & Haile, 2015; Kelly, Weiser & Tsai, 2016; Smith Fawzi et al., 2016; Kim et al., 2016; McGillen et al., 2018; Merzouki et al., 2020). Recent studies included up to 31 SSA countries, but narrowly focused their inquiries to examine, for example, the association between socioeconomic inequalities (Hajizadeh et al., 2014), high-risk sexual behaviour (Kenyon, Buyze & Schwartz, 2018), or HIV-related stigma (Chan & Tsai, 2015; Kelly, Weiser & Tsai, 2016) with HIV testing, treatment uptake, ART (antiretroviral treatment) adherence, or HIV prevalence. Most used standard statistical methods like descriptive statistics (Sangowawa & Owoaje, 2012; Smith Fawzi et al., 2016), linear or logistic regression (Mondal & Shitan, 2013; Delavande, Sampaio & Sood, 2014; Chan & Tsai, 2015; Kidman & Anglewicz, 2016; Ashaba et al., 2018), or concentration indices (Hajizadeh et al., 2014; Pons-Duran et al., 2016; Kim et al., 2016), to assess health inequity and the impact of a small number of variables (5 to 13) on the HIV epidemic. In a previous work, we also investigated the associations and possible causal relationships between sociobehavioural factors at the individual-level that are potentially related to the risk of acquiring HIV in 29 SSA countries using Bayesian Network models (Baranczuk et al., 2019). But, these methods do not inform us on how HIV risk factors vary across SSA and which characteristic sociobehavioural patterns at the country-level are actually associated with different rates of new HIV infections in the region.

Recent studies have shown that unsupervised learning and clustering analysis allows us to find hidden sub-groups of people with varying drivers and potentially different risk levels of having or acquiring HIV(Engl, Smittenaar & Sgaier, 2019; Merzouki et al., 2020). At the country-level, comparing and characterising SSA countries would allow us to test the hypothesis that sociobehavioural heterogeneity might account for spatial variance of HIV epidemic, and inform effective country-specific interventions.

In this study, we used dimensionality reduction and unsupervised machine learning techniques (Principal Component Analysis and hierarchical clustering) to identify the most important factors of 48 national attributes that might account for the variability of HIV incidence across SSA, and identified the sociobehavioural profiles that characterized different levels of HIV incidence, based on Demographic and Health Survey ("The DHS Program - Quality information to plan, monitor and improve population, health, and nutrition programs") data from 29 SSA countries that were aggregated at the national-level.

Methods

Data

We used aggregated data from the most recent Demographic and Health Surveys (DHS) of 29 SSA countries completed between 2010 and 2017, that were available up to July 2018 (**Table S1**). DHS typically gathers nationally representative("DHS Sampling and Household Listing Manual (English)") data on health (including HIV-related data) and population (including social, behavioural, geographic and economic data) every 5 years. These data are publicly accessible at the individual-, district- and country-level. We used data aggregated at the national-level.

We attempted to include in our analysis all variables possibly influencing the risks of HIV acquisition among population aged 15-49 years that were available for the 29 SSA countries through the StatCompiler tool ("STATcompiler"). We only included variables that varied significantly across the region, and did not strongly correlate with other variables. We finally pre-selected the following variables: age (under 25 vs older); rurality (rural vs urban); religion (Christian, Muslim, Folk/Popular religions, unaffiliated, others); marital status (married or in union vs widowed/divorced/other), number of wives for men (1, ≥2) or co-wives for women (0, 1, ≥2); literacy (literate vs illiterate); media access (with access to newspaper, television and radio at least once a week vs without such access); employment (worked in the last 12 months and currently working vs others); wealth (Gini coefficient); age at first sexual intercourse (first sexual intercourse by age 15 vs older); general fertility (number of births to women of reproductive age in the last 3 years); contraception use (using any method of contraception vs not using any); condom use (belief that a woman is justified in asking condom use if she knows her husband has a Sexually Transmitted Infection (STI) vs belief that she is not justified); number of sexual partners in lifetime; unprotected higher risk sex (men who had sex with a non-marital, non-cohabiting partner in the last 12 months and did not use condom during last sexual intercourse vs not); paid sex (men who ever paid for sexual intercourse vs never paid for sex); unprotected paid sex (men who used condom during the last paid sexual intercourse in the last 12 months vs did not use condom); gender-based violence (wife beating justified for at least one specific reason vs not justified for any reason); married women participation to decision making (yes vs no); gender of household head (female vs male); comprehensive correct knowledge about AIDS (yes vs no); HIV testing (ever receiving an HIV test vs never tested); male circumcision (yes vs no); ART coverage (i.e. percentage of people on antiretroviral treatment among those living with HIV); and accepting attitudes toward

people living with HIV/AIDS (would buy fresh vegetables from a shopkeeper with AIDS vs
would not); see **Table 1** for a complete summary of the variables.

We represented each country using 48 dimensions. Each dimension corresponded to an
attribute in **Table 1**, such as the percentage of women married or in union, the mean number
of sexual partners in a lifetime for men, the percentage of Christian populations and the Gini
coefficient in this country. Data were represented as percentages; the mean number of sexual
partners in lifetime was scaled using min-max normalisation. Most of these country-level data
were exported from the DHS with the StatCompiler tool, except for data on religion that we
obtained from Pew-Templeton Global Religious Futures Project ("Religions in Africa | African
Religions | PEW-GRF"), and ART coverage that we obtained from UNAIDS' AIDSinfo
("AIDSinfo | UNAIDS"). We used the latest (2018) UNAIDS estimates of national HIV
incidence for the year 2016 ("AIDSinfo | UNAIDS"; "Estimates Methods 2018").

Ethics approval

133 The study was conducted using aggregated data that are publicly accessible through
134 StatCompiler (for DHS data), Pew-Templeton Global Religious Futures Project, and UNAIDS'
135 AIDSinfo. No ethics approval was needed from our side.

Analysis

We used Principal Component Analysis (PCA) (Trevor Hastie, Robert Tibshirani & Jerome
Friedman; James et al., 2013) to reduce the dimensionality of data from 48 to two dimensions
(2D) so we could visualise sociobehavioural similarity between SSA countries. PCA allows to
extract directions, called principal components (PCs), along which the variation of data is
maximal. The first two PCs, which explain the most variance, represent the axes of the 2D-
space used for visualisation. Countries closest to each other when projected on the 2D-space
correspond to similar countries in terms of demographic, socioeconomic and behavioural
characteristics. PCs consist of a linear combination of the initial 48 dimensions and can
therefore be interpreted in terms of the original sociobehavioural variables ("PCA - Principal
Component Analysis Essentials - Articles - STHDA"). We can then identify the variables
among those we included in the analysis, that explain most of the sociobehavioural
heterogeneity across SSA.

We identified groups of similar SSA countries in terms of sociobehavioural characteristics
using hierarchical clustering. Pairwise country dissimilarity was calculated using the Euclidian

distance (**Equation S1**). These distances were used by the hierarchical clustering algorithm to

create a *dendrogram* with 29 terminal nodes representing the countries to be grouped. Cutting

the dendrogram at a certain height produces clusters of similar countries. The number of

clusters depends on the height at which the tree is cut. We measured the quality of the clustering

results using the Silhouette Index, and selected the optimal number of clusters such that it

maximised this index (**Equation S4**).

We compared the HIV incidence of countries between the identified clusters, and visualised

the distribution of HIV incidence within each cluster using *box plots*. We also identified the

sociobehavioural variables that characterized the resulting clusters, and visualized the

distribution of these variables within each cluster using *density plots*.

We used the open source R language, version 3.5.1 for our analysis. Code and country-level

data are available on GitLab (https://gitlab.com/AzizaM/dhs_ssa_countries_clustering), in two

separate folders "code" and "data_countries", respectively.

Results

We analysed aggregated data from surveys that included 594,644 persons in total,183,310 men and 411,334 women("DHS Sampling and Household Listing Manual (English)") , ranging from 9,552 in Lesotho to 56,307 in Nigeria. Adult HIV incidence ranged from 0.14/1000 in Niger to 19.7/1000 in Lesotho in 2016. HIV prevalence ranged from 0.4% in Niger to 23.9% in Lesotho (**Table S1**). Sociobehavioural characteristics varied widely between SSA countries (**Table 1**).

Visualizing the SSA countries: Geographical and sociobehavioural similarities

Using PCA, we found that the first principal component (PC) explained 49.5% and the second 19.5% of the total sociobehavioural variance across SSA among the 48 variables we considered (**Figure 1**). The original sociobehavioural variables that contributed most to these PCs were religion (12.6% for Muslim and 12.1% for Christian populations), male circumcision (9.4%), number of sexual partners (7.8% for men and 3.4% for women), literacy (6.1% for women and 3.2% for men), HIV testing (5.5% for men and 5.4% for women), women's participation in decision making (3.8%), an accepting attitude towards those living with HIV/AIDS (3.6% for women and 3.2% for men), rurality (3.0% for women and 2.7% for men), ART coverage (2.5%), and women's knowledge about AIDS (2.5%) (**Figure 1B** and **Figure S1**).

Projecting the 29 SSA countries in two dimensions produced a roughly V-shaped scatterplot (**Figure 1A**). **Figure 1B** shows how the 48 original sociobehavioural variables vary over the 2D-space. At the end of the V-shape's left branch, Eastern and Southern African countries, such as Namibia, Zimbabwe, Malawi, Zambia and Uganda, lied next to each other. In these countries, fewer men are circumcised, but the percentage of literate people who had accepting attitudes toward people living with HIV/AIDS (PLWHA) was higher and so was the uptake of HIV testing. Knowledge about AIDS and ART coverage were also high. The end of the right branch, in the upper right quadrant, included countries from the Sahel region, like Senegal, Burkina Faso, Mali, Niger and Chad, where the percentage of Muslims is higher and people have fewer sexual partners. The lower tip of the V-shape included countries from West and Central Africa, like Liberia, Ghana, Côte d'Ivoire, Democratic Republic of the Congo, and Gabon, where people have more sexual partners, more men are circumcised, and the population is less rural.

Clustering the SSA countries and analysis of the associated HIV incidence

The hierarchical clustering of the 29 SSA countries produced a dendrogram (**Figure 2A)**.

Cluster compactness and separation were optimal (maximum silhouette index = 0.3) when we

cut the dendrogram at a height that separated countries into three groups (**Figure 2B**).

The countries of the first cluster, in yellow, had the lowest HIV incidence (median of 0.5/1000

population) (**Figure 3**). This cluster included countries from the Sahel Region, where the

population was mostly rural (median of 71.1% for men) and Muslim (median of 86.2%). On

the one hand, countries of this cluster were characterized by many factors that could account

for low HIV incidence and prevalence. Countries were characterized by high proportions of

circumcised men (median of 95.0%), high percentages of women who were married or lived

in union (median of 70.6%), late sexual initiation for men (median of 1.9% of men who had

their first sexual intercourse by the age of 15), low numbers of sexual partners (median of 3.5

partners for men), low percentages of unprotected higher-risk sex (median of 9.7% for men)

and low percentages of men having ever paid for sex (median of 3.9%). Polygyny (Reniers &

Tfaily, 2012; Eaton et al., 2014), an institutionalized form of sexual concurrency, was also

frequent in this region (median of 22.3%). On the other hand, this cluster was also characterized

by frequent belief that wife beating is justified (median of 61.2% for women), and low levels

of literacy (median of 29.0% for women). Participation of married women in decision making

(median of 18.5%), contraceptive prevalence (median of 13.9%), and knowledge about AIDS

(median of 23.7% for women) was also low. These countries had low percentages of people

ever tested for HIV (median of 19.2% for men; 36.6% for women), low ART coverage (median

of 38.0%) and low levels of acceptance of PLWHA (Median of 47.4% for men); see **Figure 4**

and **Figure 5**.

The countries of the second cluster, coloured in orange, included countries from West and

Central Africa. These countries had a rather low HIV incidence (median of 1.8/1000

population), though Mozambique was a remarkable outlier, with a high HIV incidence

(9.8/1000 population) (**Figure 3**). Like the first cluster, these countries had a high percentage

of circumcised men (median of 97.0%, except in Mozambique where only 48.4% of men were

circumcised). However, these countries were also characterized by the lowest proportions of

rural populations (median of 49.0% for men), the highest numbers of sexual partners (median

of 10.1 for men), early sexual initiation (median of 12.0% of men who had their first sexual

intercourse by the age of 15), and more frequent unprotected high-risk sex (median of 24.3%

for men) and paid sexual intercourse (median of 9.5% for men). HIV testing uptake (median

of 25.8% for men and 48.6% for women), knowledge about AIDS (median of 23.6% for

women), and ART coverage (median of 31.0%) were all low.

The third cluster, in red, included Southern and East African countries. These countries had

high HIV incidence (median of 5.0/1000 population), except two countries that had a lower

HIV incidence: Rwanda (1.1/1000 population) and Burundi (0.5/1000) (**Figure 3**). Countries

belonging to the third cluster were characterized by the lowest percentage of circumcised men

(median of 27.9%). But they were also the ones with the highest uptake of HIV testing (median

of 65.2% for men; 83.3% for women) and ART (median of 61.0%), and the highest percentage

with knowledge about AIDS (median of 54.6% for women) and accepting attitudes towards

PLWHA (median of 84.4% for men). This cluster was also characterized by the highest

percentage of literacy (median of 80.2% for women), high use of contraceptives (median of

42.6%), low percentages of unprotected high-risk sex (median of 9.8% for men) and higher

percentages of married women participating in decision making (median of 67.7%) and

women-headed households (median of 31.0%). Rwanda and Burundi had the lowest HIV

incidence and were characterized by a lower number of sexual partners (Rwanda, 2.6; Burundi,

2.1) vs a median of 6.3 partners for men in the other countries of the third cluster. They also

had larger per capita rural populations (Rwanda, 80.4%; Burundi, 89.4%) vs a median of 61.3%

for women in the other countries of the same cluster.

Discussion

Using hierarchical clustering, we identified the most important characteristics that explained 69% of the sociobehavioural variance among the variables we assessed in SSA. The variables that contributed the most to the sociobehavioural heterogeneity across the 29 SSA countries and among the 48 attributes we included in this analysis are religion, number of sexual partners, literacy, HIV testing, women's participation in decision making, accepting attitude towards those living with HIV/AIDS, rurality, ART coverage, and women's knowledge about AIDS. We found three groups of countries with similar sociobehavioural patterns, and HIV incidence was also similar within each cluster.

In the first cluster, PLWHA were not widely accepted, and the population had an overall low-level knowledge about AIDS. Stigma may be more widespread in this region and explain the lower uptake of interventions among people who are HIV-positive. The relatively low number of people who are living with HIV lowers the general public's exposure to this group and may increase stigma (Chan & Tsai, 2017). Stigma can also result from cultural and religious beliefs that link HIV/AIDS with sexual transgressions, immorality and sin (Campbell et al., 2005; Mbonu, van den Borne & De Vries, 2009).

We think that the apparent contradiction between the presence of many high-risk factors and the low HIV incidence in most countries of the second cluster could be explained by the high proportion of circumcised men. In line with this theory, Mozambique, the only country in this cluster with very high HIV prevalence and incidence, had few circumcised men. Previous observational studies and trials have confirmed the protective effect of male circumcision (Bailey et al., 2007; Gray et al., 2007; Lei et al., 2015; Sharma et al., 2018).

Countries of the third cluster, with the highest HIV incidence, were also the ones with the highest knowledge about AIDS (Chan & Tsai, 2017), ART coverage, uptake of HIV testing, and with the most accepting attitudes toward PLWHA. They also had the lowest percentage of unprotected higher risk sex. These findings are consistent with earlier studies that found broad ART coverage may reduce social distancing towards PLWHA and HIV-related stigma in the general population(Chan & Tsai, 2015; Chan, Tsai & Siedner, 2015). Reduced social distancing and stigma is associated with higher uptake of voluntary HIV counselling and testing (Kalichman, 2003; Kelly, Weiser & Tsai, 2016), and less sexual risk-taking among HIV positive people (Delavande, Sampaio & Sood, 2014).

The high HIV incidence in Mozambique could be caused by any combination of the following
factors: a high number of sexual partners; a low level of male circumcision; and a low level of
literacy and knowledge about AIDS. The latter two factors can also be responsible for low
uptake of HIV testing and ART. However, many West and Central African countries with
population characteristics like Mozambique in terms of sexual practices, literacy, knowledge
about AIDS, HIV testing and ART coverage, had much lower HIV prevalence and incidence,
possibly because males were circumcised at twice the rate. Compared to other West and Central
African countries, an aggravating factor in Mozambique can also be the high prevalence of
female genital schistosomiasis(Hotez et al., 2019; Yegorov et al., 2019). On the other side of
the spectrum, despite a low uptake of male circumcision, it is also possible that the combination
of lower numbers of sexual partners, higher per capita rural populations, more literacy, more
accurate knowledge about AIDS, more HIV testing, and broader ART coverage could account
for lower HIV incidence, like in Rwanda and Burundi.

In contrast to more classical regression techniques that usually estimate the relation between
independent sociobehavioural factors and a single outcome, dimensionality reduction and
clustering techniques allow us to visualize the sociobehavioural heterogeneity across SSA
countries, and identify characteristic patterns of factors that are shared by groups of countries
and are found to be associated with different levels of HIV incidence. These results provide
policy makers with a quick and clearer overview of sociobehavioural factors across SSA, and
help identify targeted interventions that can reduce HIV transmission in a specific country or
group of countries.

The cross-sectional nature of our data makes it impossible to determine precedence and
causality between the sociobehavioural characteristics we measured and HIV prevalence and
incidence. But the associations we identified can open lines of inquiry for researchers. Our
study had the advantage of allowing us to compare countries and regions (clusters of countries),
but ecological studies that use aggregated data are prone to confounding and ecological fallacy
(Levin). Africa is an exceedingly diverse continent with many distinct sub-populations, so a
study based on national population averages cannot explain HIV variation within countries.
Recent studies have shown the complex geographical variation of the HIV epidemic and its
drivers within Eastern and Southern African countries (Cuadros et al., 2017; Bulstra et al.,
2020). Therefore, we intend to repeat our study at a lower level of granularity, using regional-
, district- and individual-level data to capture differences within countries and learn more about

the complex patterns of sociobehavioural factors that affect the sub-populations that are most

at risk. We believe that machine learning is a promising innovative way to further explore and

improve our understanding of these questions, in parallel to more classical techniques.

Our work has some other limitations. We used model estimates for HIV incidence, which may

diverge from reality (Nsanzimana et al., 2017). Sociobehavioural data from the DHS are self-

reported, therefore potentially biased; for instance, literate individuals may be more reluctant

to admit accepting gender-based violence or having higher risk sex. And even though we

included many more variables than is common practice (Hajizadeh et al., 2014; Lakew,

Benedict & Haile, 2015; Kidman & Anglewicz, 2016; Kim et al., 2016; Ashaba et al., 2018;

Kenyon, Buyze & Schwartz, 2018), we still had to exclude many factors that can play a critical

role in HIV transmission, including the prevalence of excessive alcohol consumption, STIs

(Looker et al., 2017; Torrone et al., 2018) (ex. genital herpes, gonorrhea and syphilis), endemic

infections (Hotez et al., 2019; Yegorov et al., 2019)(ex. malaria and helminthiases), ART

adherence and drug resistance, as well as the distribution of different HIV strains and subtypes

(Buonaguro, Tornesello & Buonaguro, 2007; Esbjörnsson et al., 2019; Gartner et al., 2020).

Some of the latter variables were not available from the DHS or were missing for at least one

country included in our analysis.

Conclusions

Our use of PCA allowed us to identify the most important characteristics among the variables

we assessed that explained 69% of the sociobehavioural variance in SSA countries. With

hierarchical clustering, we captured complex patterns of sociobehavioural characteristics

shared by countries with similar HIV incidence, suggesting that the combination of

sociobehavioural factors plays a key role in determining the course of the HIV epidemic. Our

findings can help to design and predict the effect of targeted country-specific interventions to

impede HIV transmission.

Authors' contributions

AM, JE and OK designed the study. AM and OK reviewed and selected the variables to be included in the analysis. AM wrote the code and performed the analysis with support from JE. EO reviewed analysis and code, and checked results reproducibility. AM interpreted the results with support from OK and KT. AM, OK and KT reviewed the literature. AM wrote the manuscript, which was reviewed by OK, JE, and KT. All authors have read and approved the final manuscript.

Acknowledgements

We thank Zofia Baranczuk for helpful discussions.

Conflict of interest

We declare no competing interests.

References

AIDSinfo | UNAIDS. *Available at http://aidsinfo.unaids.org/* (accessed March 6, 2019).

Antelman G, Kaaya S, Wei R, Mbwambo J, Msamanga GI, Fawzi WW, Smith Fawzi MC. 2007. Depressive Symptoms Increase Risk of HIV Disease Progression and Mortality Among Women in Tanzania: *JAIDS Journal of Acquired Immune Deficiency Syndromes* 44:470–477. DOI: 10.1097/QAI.0b013e31802f1318.

Ashaba S, Cooper-Vince C, Maling S, Rukundo GZ, Akena D, Tsai AC. 2018. Internalized HIV stigma, bullying, major depressive disorder, and high-risk suicidality among HIV-positive adolescents in rural Uganda. *Global Mental Health* 5. DOI: 10.1017/gmh.2018.15.

Bailey RC, Moses S, Parker CB, Agot K, Maclean I, Krieger JN, Williams CFM, Campbell RT, Ndinya-Achola JO. 2007. Male circumcision for HIV prevention in young men in Kisumu, Kenya: a randomised controlled trial. 369:14.

Baranczuk Z, Estill J, Blough S, Meier S, Merzouki A, Maathuis MH, Keiser O. 2019. *Socio-behavioural characteristics and HIV: findings from a graphical modelling analysis of 29 sub-Saharan African countries*. Epidemiology. DOI: 10.1101/600510.

Bulstra CA, Hontelez JAC, Giardina F, Steen R, Nagelkerke NJD, Bärnighausen T, de Vlas SJ. 2020. Mapping and characterising areas with high levels of HIV transmission in sub-Saharan Africa: A geospatial analysis of national survey data. *PLOS Medicine* 17:e1003042. DOI: 10.1371/journal.pmed.1003042.

Buonaguro L, Tornesello ML, Buonaguro FM. 2007. Human Immunodeficiency Virus Type 1 Subtype Distribution in the Worldwide Epidemic: Pathogenetic and Therapeutic Implications. *Journal of Virology* 81:10209–10219. DOI: 10.1128/JVI.00872-07.

Campbell C, Foulis CA, Maimane S, Sibiya Z. 2005. "I Have an Evil Child at My House": Stigma and HIV/AIDS Management in a South African Community. *American Journal of Public Health* 95:808–815. DOI: 10.2105/AJPH.2003.037499.

Chan B, Tsai A. 2015. Trends in HIV-Related Stigma in the General Population During the Era of Antiretroviral Treatment Expansion: An Analysis of 31 Sub-Saharan African Countries. *Open Forum Infectious Diseases* 2. DOI: 10.1093/ofid/ofv133.280.

Chan BT, Tsai AC. 2017. Personal contact with HIV-positive persons is associated with reduced HIV-related stigma: cross-sectional analysis of general population surveys from 26 countries in sub-Saharan Africa. *Journal of the International AIDS Society* 20:21395. DOI: 10.7448/IAS.20.1.21395.

Chan BT, Tsai AC, Siedner MJ. 2015. HIV Treatment Scale-Up and HIV-Related Stigma in

Sub-Saharan Africa: A Longitudinal Cross-Country Analysis. *American Journal of Public Health* 105:1581–1587. DOI: 10.2105/AJPH.2015.302716.

Cuadros DF, Li J, Branscum AJ, Akullian A, Jia P, Mziray EN, Tanser F. 2017. Mapping the spatial variability of HIV infection in Sub-Saharan Africa: Effective information for localized HIV prevention and control. *Scientific Reports* 7. DOI: 10.1038/s41598-017-09464-y.

Delavande A, Sampaio M, Sood N. 2014. HIV-related social intolerance and risky sexual behavior in a high HIV prevalence environment. *Social Science & Medicine* 111:84–93. DOI: 10.1016/j.socscimed.2014.04.011.

DHS Sampling and Household Listing Manual (English). *Available at https://dhsprogram.com/publications/publication-dhsm4-dhs-questionnaires-and-manuals.cfm* (accessed October 8, 2020).

Eaton JW, Takavarasha FR, Schumacher CM, Mugurungi O, Garnett GP, Nyamukapa C, Gregson S. 2014. Trends in Concurrency, Polygyny, and Multiple Sex Partnerships During a Decade of Declining HIV Prevalence in Eastern Zimbabwe. *The Journal of Infectious Diseases* 210:S562–S568. DOI: 10.1093/infdis/jiu415.

Engl E, Smittenaar P, Sgaier SK. 2019. Identifying population segments for effective intervention design and targeting using unsupervised machine learning: an end-to-end guide. *Gates Open Research*.

Esbjörnsson J, Månsson F, Kvist A, Silva ZJ da, Andersson S, Fenyö EM, Isberg P-E, Biague AJ, Lindman J, Palm AA, Rowland-Jones SL, Jansson M, Medstrand P, Norrgren H, N'Buna B, Biague AJ, Biai A, Camara C, Esbjörnsson J, Jansson M, Karlson S, Lindman J, Medstrand P, Månsson F, Norrgren H, Palm AA, Sahin GÖ, Silva ZJ da, Wilhelmson S. 2019. Long-term follow-up of HIV-2-related AIDS and mortality in Guinea-Bissau: a prospective open cohort study. *The Lancet HIV* 6:e25–e31. DOI: 10.1016/S2352-3018(18)30254-6.

Estimates Methods 2018. *Available at http://aidsinfo.unaids.org/documents/estimates_methods_2018.pdf* (accessed January 24, 2020).

Fact sheet - World AIDS Day 2018. *Available at https://www.unaids.org/sites/default/files/media_asset/UNAIDS_FactSheet_en.pdf* (accessed January 24, 2020).

Gartner MJ, Roche M, Churchill MJ, Gorry PR, Flynn JK. 2020. Understanding the mechanisms driving the spread of subtype C HIV-1. *EBioMedicine* 53. DOI: 10.1016/j.ebiom.2020.102682.

Global Health Sector Strategy on HIV 2016-2021: Towards Ending AIDS. *Available at*

https://apps.who.int/iris/bitstream/handle/10665/246178/WHO-HIV-2016.05-

eng.pdf?sequence=1 (accessed January 24, 2020).

Gray RH, Kigozi G, Serwadda D, Makumbi F, Watya S, Nalugoda F, Kiwanuka N, Moulton

LH, Chaudhary MA, Chen MZ, Sewankambo NK, Wabwire-Mangen F, Bacon MC, Williams

CFM, Opendi P, Reynolds SJ, Laeyendecker O, Quinn TC, Wawer MJ. 2007. Male

circumcision for HIV prevention in men in Rakai, Uganda: a randomised trial. 369:10.

Gregson S, Gonese E, Hallett TB, Taruberekera N, Hargrove JW, Lopman B, Corbett EL,

Dorrington R, Dube S, Dehne K, Mugurungi O. 2010. HIV decline in Zimbabwe due to

reductions in risky sex? Evidence from a comprehensive epidemiological review. *International*

Journal of Epidemiology 39:1311–1323. DOI: 10.1093/ije/dyq055.

Hajizadeh M, Sia D, Heymann S, Nandi A. 2014. Socioeconomic inequalities in HIV/AIDS

prevalence in sub-Saharan African countries: evidence from the Demographic Health Surveys.

International Journal for Equity in Health 13:18. DOI: 10.1186/1475-9276-13-18.

Hotez PJ, Harrison W, Fenwick A, Bustinduy AL, Ducker C, Sabina Mbabazi P, Engels D,

Floerecke Kjetland E. 2019. Female genital schistosomiasis and HIV/AIDS: Reversing the

neglect of girls and women. *PLOS Neglected Tropical Diseases* 13:e0007025. DOI:

10.1371/journal.pntd.0007025.

James G, Witten D, Hastie T, Tibshirani R (eds.). 2013. *An introduction to statistical learning:*

with applications in R. New York: Springer.

Kalichman SC. 2003. HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and

testing in a black township in Cape Town, South Africa. *Sexually Transmitted Infections*

79:442–447. DOI: 10.1136/sti.79.6.442.

Kelly JD, Weiser SD, Tsai AC. 2016. Proximate Context of HIV Stigma and Its Association

with HIV Testing in Sierra Leone: A Population-Based Study. *AIDS and Behavior* 20:65–70.

DOI: 10.1007/s10461-015-1035-9.

Kenyon C, Buyze J, Schwartz IS. 2018. Strong association between higher-risk sex and HIV

prevalence at the regional level: an ecological study of 27 sub-Saharan African countries.

F1000Research 7:1879. DOI: 10.12688/f1000research.17108.1.

Kidman R, Anglewicz P. 2016. Are adolescent orphans more likely to be HIV-positive? A

pooled data analyses across 19 countries in sub-Saharan Africa. *Journal of Epidemiology and*

Community Health 70:791–797. DOI: 10.1136/jech-2015-206744.

Kim SW, Skordis-Worrall J, Haghparast-Bidgoli H, Pulkki-Brännström A-M. 2016. Socio-

economic inequity in HIV testing in Malawi. *Global Health Action* 9:31730. DOI:

10.3402/gha.v9.31730.

Lakew Y, Benedict S, Haile D. 2015. Social determinants of HIV infection, hotspot areas and
subpopulation groups in Ethiopia: evidence from the National Demographic and Health Survey
in 2011. *BMJ Open* 5:e008669. DOI: 10.1136/bmjopen-2015-008669.

Lei J hao, Liu L ren, Wei Q, Yan S bing, Yang L, Song T run, Yuan H chao, Lv X, Han P.
2015. Circumcision Status and Risk of HIV Acquisition during Heterosexual Intercourse for
Both Males and Females: A Meta-Analysis. *PLOS ONE* 10:e0125436. DOI:
10.1371/journal.pone.0125436.

Levin KA. Study Design VI - Ecological Studies. :1.

Looker KJ, Elmes JAR, Gottlieb SL, Schiffer JT, Vickerman P, Turner KME, Boily M-C.
2017. Effect of HSV-2 infection on subsequent HIV acquisition: an updated systematic review
and meta-analysis. *The Lancet. Infectious Diseases* 17:1303. DOI: 10.1016/S1473-
3099(17)30405-X.

Mbonu NC, van den Borne B, De Vries NK. 2009. Stigma of People with HIV/AIDS in Sub-
Saharan Africa: A Literature Review. *Journal of Tropical Medicine* 2009:1–14. DOI:
10.1155/2009/145891.

McGillen JB, Stover J, Klein DJ, Xaba S, Ncube G, Mhangara M, Chipendo GN, Taramusi I,
Beacroft L, Hallett TB, Odawo P, Manzou R, Korenromp EL. 2018. The emerging health
impact of voluntary medical male circumcision in Zimbabwe: An evaluation using three
epidemiological models. *PLOS ONE* 13:e0199453. DOI: 10.1371/journal.pone.0199453.

Merzouki A, Styles A, Estill J, Orel E, Baranczuk Z, Petrie K, Keiser O. 2020. Identifying
groups of people with similar sociobehavioural characteristics in Malawi to inform HIV
interventions: a latent class analysis. *Journal of the International AIDS Society* 23:e25615.
DOI: 10.1002/jia2.25615.

Mondal M, Shitan M. 2013. Factors affecting the HIV/AIDS epidemic: an ecological analysis
of global data. *African Health Sciences* 13:301–310. DOI: 10.4314/ahs.v13i2.15.

Nsanzimana S, Remera E, Kanters S, Mulindabigwi A, Suthar AB, Uwizihiwe JP,
Mwumvaneza M, Mills EJ, Bucher HC. 2017. Household survey of HIV incidence in Rwanda:
a national observational cohort study. *The Lancet HIV* 4:e457–e464. DOI: 10.1016/S2352-
3018(17)30124-8.

PCA - Principal Component Analysis Essentials - Articles - STHDA. *Available at*
http://www.sthda.com/english/articles/31-principal-component-methods-in-r-practical-
guide/112-pca-principal-component-analysis-essentials/ (accessed January 6, 2020).

Pons-Duran C, González R, Quintó L, Munguambe K, Tallada J, Naniche D, Sacoor C, Sicuri
E. 2016. Association between HIV infection and socio-economic status: evidence from a

semirural area of southern Mozambique. *Tropical Medicine & International Health* 21:1513–

1521. DOI: 10.1111/tmi.12789.

Religions in Africa | African Religions | PEW-GRF. *Available at*

http://www.globalreligiousfutures.org/regions/sub-saharan-

africa#/?region_map_religion=All Religious Groups (accessed January 24, 2019).

Reniers G, Tfaily R. 2012. Polygyny, Partnership Concurrency, and HIV Transmission in Sub-

Saharan Africa. *Demography* 49:1075–1101. DOI: 10.1007/s13524-012-0114-z.

Sangowawa AO, Owoaje ET. 2012. Experiences of discrimination among youth with

HIV/AIDS in Ibadan, Nigeria. *Journal of Public Health in Africa* 3:10. DOI:

10.4081/jphia.2012.e10.

Sharma SC, Raison N, Khan S, Shabbir M, Dasgupta P, Ahmed K. 2018. Male circumcision

for the prevention of human immunodeficiency virus (HIV) acquisition: a meta-analysis. *BJU*

International 121:515–526. DOI: 10.1111/bju.14102.

Smith Fawzi MC, Ng L, Kanyanganzi F, Kirk C, Bizimana J, Cyamatare F, Mushashi C, Kim

T, Kayiteshonga Y, Binagwaho A, Betancourt TS. 2016. Mental Health and Antiretroviral

Adherence Among Youth Living With HIV in Rwanda. *PEDIATRICS* 138:e20153235–

e20153235. DOI: 10.1542/peds.2015-3235.

STATcompiler. *Available at https://www.statcompiler.com/en/* (accessed January 24, 2020).

The DHS Program - Quality information to plan, monitor and improve population, health, and

nutrition programs. *Available at https://dhsprogram.com/* (accessed September 27, 2019).

Torrone EA, Morrison CS, Chen P-L, Kwok C, Francis SC, Hayes RJ, Looker KJ, McCormack

S, McGrath N, van de Wijgert JHHM, Watson-Jones D, Low N, Gottlieb SL, on behalf of the

STIMA Working Group. 2018. Prevalence of sexually transmitted infections and bacterial

vaginosis among women in sub-Saharan Africa: An individual participant data meta-analysis

of 18 HIV prevention studies. *PLOS Medicine* 15:e1002511. DOI:

10.1371/journal.pmed.1002511.

Trevor Hastie, Robert Tibshirani, Jerome Friedman. *The Elements of Statistical Learning:*

Data Mining, Inference, and Prediction. Springer.

Tsai AC, Venkataramani AS. 2015. The causal effect of education on HIV stigma in Uganda:

Evidence from a natural experiment. *Social Science & Medicine* 142:37–46. DOI:

10.1016/j.socscimed.2015.08.009.

Yegorov S, Joag V, Galiwango RM, Good SV, Okech B, Kaul R. 2019. Impact of Endemic

Infections on HIV Susceptibility in Sub-Saharan Africa. *Tropical Diseases, Travel Medicine*

and Vaccines 5:22. DOI: 10.1186/s40794-019-0097-5.

Table 1 - Socio-economic and behavioural variables included in the analysis.

Median values across all 29 countries are shown with the minimum and maximum values.

*ART = antiretroviral therapy.

Attribute	Topic	Variable	Stratification	Categories	Median (min - max)
1		Age under 25	Men		37.6% (28.1%-44.1%)
2			Women		39.9% (34.4%-45.0%)
3		Rurality	Men		56.5% (12.9%-85.1%)
4			Women		59.7% (11.3%-89.4%)
5		Religion		Christian	74.9% (0.8%-97.8%)
6				Muslim	13.9% (0.0%-98.5%)
7				Folk/Popular	1.7% (0.0%-35.7%)
8				Unaffiliated	2.5% (0.0%-18.0%)
9				Others	0.2% (0.0%-2.7%)
10		Married or in union	Men		50.5% (28.8%-65.2%)
11	Demographic		Women		63.5% (34.0%-88.5%)
12		Number of wives (for men) and co-wives (for women)	Men	1[1]	87.5% (72.0%-97.5%)
13				≥2[2]	12.5% (2.5%-28.0%)
14			Women	0[3]	75.5% (57.6%-93.2%)
15				1[4]	17.2% (1.9%-30.4%)
16				≥2[5]	4.3% (0.4%-12.3%)
17		Female headed household			28.0% (9.3%-43.9%)
18		Literacy	Men		79.0% (37.6%-94.2%)
19			Women		58.1% (14.0%-97.0%)
20		Access to media at least once a week	Men		9.9% (1.7%-47.5%)
21			Women		5.6% (0.3%-21.3%)

---

[1] Percentage of currently married or in union men who have one wife
[2] Percentage of currently married or in union men who have two or more wives
[3] Percentage of currently married or in union women whose husband has no other wives
[4] Percentage of currently married or in union women whose husband has one other wife
[5] Percentage of currently married or in union women whose husband has two or more wives

22	Employment	Worked in the last 12 months and is currently working	Men		76.9% (55.9%-92.8%)
23			Women		61.8% (24.5%-77.8%)
24	Wealth	Gini coefficient[6]			30.0% (10.0%-50.0%)
25	Sexual behaviour	First sex by age 15	Men		8.0% (0.8%-25.4%)
26			Women		18.0% (2.6%-28.8%)
27		General fertility rate[7]	Women		17.5 (11.8-26.9)
28		Use of contraception	Women		21.7% (5.4%-50.2%)
29		Woman is justified asking for condom if husband has a sexually transmitted infection (STI)	Men		88.2% (70.3%-98.5%)
30			Women		81.5% (14.3%-97.3%)
31		Mean number of sexual partners in lifetime	Men		6.3 (1.9-15.3)
32			Women		2.2 (1.2-5.1)
33		Unprotected higher risk sex	Men		15.7% (1.6%-43.2%)
34			Women		11.10% (0.3%-30.3%)
35		Ever paid for sexual intercourse	Men		7.7% (1.4%-35.0%)
36		Unprotected paid sexual intercourse	Men		0.8% (0.1%-8.1%)
37	Gender-based violence	Wife beating justified	Men		32.3% (12.5%-59.5%)
38			Women		45.7% (16.2%-76.3%)
39	Women empowerment	Married women participating in decision making[8]			49.9% (9.1%-78.0%)
40		Married women who disagree with all reason justifying wife beating[9]			47.7% (18.7%-80.9%)
41	HIV/AIDS	Comprehensive correct knowledge about AIDS[10]	Men		35.8% (17.4%-68.8%)
42			Women		27.8% (10.9%-66.9%)
43		Ever received an HIV test	Men		30.5% (7.8%-80.8%)

44			Women		49.6% (14.5%-85.5%)
45		Male circumcision			94.0% (14.3%-99.4%)
46		ART* coverage 2015			41.0% (18.0%-76.0%)
47	Accepting attitudes toward PLWHA	Would buy vegetables from shopkeeper with AIDS	Men		57.5% (32.4%-92.1%)
48			Women		53.1% (23.7%-89.2%)

[6] The Gini coefficient indicates the level of wealth concentration in a country.

[7] Average number of children currently being born to women of reproductive age in the three years preceding the survey, expressed per 100 women age 15-44.

[8] Percentage of currently married women age 15-49 who usually make all three specific decisions either alone or jointly with their husband for (1) own health care, (2) large household purchases, and (3) visits to family or relatives.

[9] Percentage of currently married women age 15-49 who disagree with all five specific reasons justifying wife-beating: (1) burning food, (2) arguing with husband, (3) going out without telling him, (4) refusing sexual intercourse with him and (5) neglecting children.

[10] Percentage of men and women who correctly identify the two major ways of preventing the sexual transmission of HIV (using condoms and limiting sex to one faithful, uninfected partner), who reject the two most common local misconceptions about HIV transmission (ex. AIDS cannot be transmitted by mosquito bites, and it cannot be transmitted by supernatural means), and who know that a healthy-looking person can have HIV.

Figure 1 - Visualization of the sociobehavioural similarity between SSA countries using PCA.

(A) Projection of the SSA countries on a 2D-space, based on their socio-economic and behavioural factors. The two dimensions (first two PCs), Dim1 and Dim2, explained 69% of the variance in the data. Countries are coloured based on their HIV incidence per 1000 population (15-49) in 2016. **(B) Correlation plot of the original variables with the first and second dimensions (Dim1, Dim2).** The variable transparency represents its contribution (in %) to the two dimensions. Moving along a variable's vector leads toward a region of the 2D-space where the variable levels tend to be higher, e.g. upper right quadrant contains mainly Muslim countries, while upper left quadrant contains countries with higher levels of HIV testing and knowledge about AIDS.

Figure 2 - Hierarchical clustering of 29 sub-Saharan African countries

(A) Dendrogram. Cutting the tree at the height of the red dashed line results in three

clusters, highlighted in yellow, orange and red. **(B) Average Silhouette width for different**

numbers of clusters. The number of clusters (X axis), from 2 to 10, corresponds to different

heights at which the dendrogram was cut. The maximum average Silhouette width was

obtained for 3 clusters (red circle).

Figure 3 - Analysis of the resulting clusters.

(A) Map of clustered sub-Saharan countries. Countries are coloured based on the cluster to

which they belong. **(B) Box plots of the HIV incidence distribution within each cluster**.

Figure 4 - Analysis of the resulting clusters in terms of their sociobehavioural characteristics. Density plots per cluster of (A) the percentage of Muslim population, (B) the percentage of circumcised men, (C) the mean number of sexual partners in a man's lifetime, (D) the percentage of literate women, (E) the percentage of men who have ever received an HIV test, (F) the percentage of men who say they would buy fresh vegetables from a vendor whom they knew was HIV+, (G) the percentage of women with a comprehensive knowledge about AIDS and (H) the ART coverage in 2015.

Figure 5 - Analysis of the resulting clusters in terms of their HIV-related attributes.
Density plots per cluster of (A) the percentage of men who have ever received an HIV test, (B) the percentage of men who say they would buy fresh vegetables from a vendor whom they knew was HIV+, (C) the percentage of women with a comprehensive knowledge about AIDS and (D) the ART coverage in 2015.