Personalized deep learning of individual immunopeptidomes to identify neoantigens for cancer vaccines

Ngoc Hieu Tran[1,†], Rui Qiao[2,†], Lei Xin[3], Xin Chen[3], Baozhen Shan[3,*], Ming Li[1,*]

[1] David R. Cheriton School of Computer Science, University of Waterloo, Waterloo, Ontario, Canada.

[2] Department of Statistics and Actuarial Science, University of Waterloo, Waterloo, Ontario, Canada.

[3] Bioinformatics Solutions Inc., Waterloo, Ontario, Canada.

[†] Joint-first authors.

[*] Corresponding authors.

Emails: mli@uwaterloo.ca (Ming Li); bshan@bioinfor.com (Baozhen Shan).

Abstract. Tumor-specific neoantigens play the main role for developing personal vaccines in
cancer immunotherapy. We propose, for the first time, a personalized *de novo* sequencing
workflow to identify HLA-I and HLA-II neoantigens directly and solely from mass spectrometry
data. Our workflow trains a personal deep learning model on the immunopeptidome of an
individual patient and then uses it to predict mutated neoantigens of that patient. This
personalized learning and mass spectrometry-based approach enables comprehensive and
accurate identification of neoantigens. We applied the workflow to datasets of five melanoma
patients and substantially improved the accuracy and identification rate of *de novo* HLA
peptides by 14.3% and 38.9%, respectively. This subsequently led to the identification of 10,440
HLA-I and 1,585 HLA-II new peptides that were not presented in existing databases. Most
importantly, our workflow successfully discovered 17 neoantigens of both HLA-I and HLA-II,
including those with validated T cell responses and those novel neoantigens that had not been
reported in previous studies.

INTRODUCTION

Neoantigens are tumor-specific mutated peptides that are brought to the surface of tumor cells by major histocompatibility complex (MHC) proteins and can be recognized by T cells as "foreign" (non-self) to trigger immune response. As neoantigens carry tumor-specific mutations and are not found in normal tissues, they represent ideal targets for the immune system to distinguish cancer cells from non-cancer ones [1-3]. The potential of neoantigens for cancer vaccines is supported by multiple evidences, including the correlation between mutation load and response to immune checkpoint inhibitor therapies [4, 5], neoantigen-specific T cell responses detected even before vaccination (naturally occurring) [6-8]. Indeed, three independent studies have further demonstrated successful clinical trials of personalized neoantigen vaccines for patients with melanoma [6-8]. The vaccination was found to reinforce pre-existing T cell responses and to induce new T cell populations directed at the neoantigens. In addition to developing cancer vaccines, neoantigens may help to identify targets for adoptive T cell therapies, or to improve the prediction of response to immune checkpoint inhibitor therapies.

And thus began the "gold rush" for neoantigen mining [1-3]. The current prevalent approach to identify candidate neoantigens often includes two major phases: (i) exome sequencing of cancer and normal tissues to find somatic mutations and (ii) predicting which mutated peptides are most likely to be presented by MHC proteins for T cell recognition. The first phase is strongly backed by high-throughput sequencing technologies and bioinformatics pipelines that have been well established through several genome sequencing projects during the past decade. The second phase, however, is still facing challenges due to our lack of knowledge of the MHC antigen processing pathway: how mutated proteins are processed into peptides; how those peptides are delivered to the endoplasmic reticulum by the transporter associated with antigen processing; and how they bind to MHC proteins. To make it further complicated, human leukocyte antigens (HLA), those genes that encode MHC proteins, are located among the most genetically variable regions and their alleles basically change from one individual to another. The problem is especially more challenging for HLA class II (HLA-II) peptides than HLA class I (HLA-I), because the former are longer, their motifs have greater variations, and very limited data is available.

Current *in silico* methods focus on predicting which peptides bind to MHC proteins given the HLA alleles of a patient, e.g. NetMHC [9, 10]. However, usually very few, less than a dozen from thousands of predicted candidates are confirmed to be presented on the tumor cell surface and even less are found to trigger T cell responses, not to mention that real neoantigens may not be among top predicted candidates [1, 2]. Several efforts have been made to improve the MHC binding prediction, including using mass spectrometry data in addition to binding affinity data for more accurate prediction of MHC antigen presentation [11, 12]. Recently, proteogenomic approaches have been proposed to combine mass spectrometry and exome sequencing to identify neoantigens directly isolated from MHC proteins, thus overcoming the limitations of MHC binding prediction [13, 14]. In those approaches, exome sequencing was performed to build a customized protein database that included all normal and mutated protein sequences. The database was further used by a search engine to identify endogenous peptides, including neoantigens, that were obtained by immunoprecipitation assays and mass spectrometry.

Existing database search engines, however, are not designed for HLA peptides and may be biased towards tryptic peptides [15, 16]. They may have sensitivity and specificity issues when

dealing with a very large search space created by (i) all mRNA isoforms from exome
sequencing and (ii) unknown digestion rules for HLA peptides. Furthermore, recent
proteogenomic studies reported a weak correlation between proteome- and genome-level
mutations, where the number of identified mutated HLA peptides was three orders of
magnitudes less than the number of somatic mutations that were provided to the database
search engines [13,14]. A large number of genome-level mutations were not presented at the
proteome level, while at the same time, some mutated peptides might be difficult to detect at the
genome level. For instance, Faridi *et al.* found evidence of up to 30% of HLA-I peptides that
were cis- and trans-splicing, which couldn't be detected by exome sequencing nor protein
database search [25]. Thus, an independent approach that does not rely heavily on genome-
level information to identify mutated peptides directly from mass spectrometry data is needed,
and *de novo* sequencing is the key to address this problem.

In this study, we propose, for the first time, a personalized *de novo* sequencing workflow to
identify HLA-I and HLA-II neoantigens directly and solely from mass spectrometry data. *De novo*
sequencing is invented for the purpose of discovering novel peptides and proteins, genetic
variants or mutations. Thus, its application to identify neoantigens is a perfect match. We bring
de novo sequencing to the "personalized" level by training a specific machine learning model for
each individual patient using his/her own data. In particular, we use the collection of normal HLA
peptides, i.e. the immunopeptidome, of a patient to train a model and then use it to predict
mutated HLA peptides of that patient. Learning an individual's immunopeptidome is made
possible by our recent deep learning model, DeepNovo [17, 18], which uses a long-short-term
memory (LSTM) recurrent neural network (RNN) to capture sequence patterns in peptides or
proteins, in a similar way to natural languages [19]. This personalized learning workflow
significantly improves the accuracy of *de novo* sequencing for comprehensive and reliable
identification of neoantigens. Furthermore, our *de novo* sequencing approach predicts peptides
solely from mass spectrometry data and does not depend on genomic information as existing
approaches. We applied the workflow to datasets of five melanoma patients and successfully
identified in average 154 HLA-I and 47 HLA-II candidate neoantigens per patient, including
those with validated T cell responses and those novel neoantigens that had not been reported in
previous studies.

RESULTS

Personalized *De novo* Sequencing of Individual Immunopeptidomes

Figure 1 describes five steps of our personalized *de novo* sequencing workflow to predict HLA
peptides of an individual patient from mass spectrometry data: (1) build the immunopeptidome
of the patient; (2) train personalized machine learning model; (3) personalized *de novo*
sequencing; (4) quality control of *de novo* peptides; and (5) neoantigen selection. The details of
each step on two example datasets, HLA-I and HLA-II of patient Mel-15 from [13], are provided
in Supplementary Table S1 for illustration.

In step 1 of the workflow, to build the immunopeptidome of the patient, we searched the mass
spectrometry data against the standard Swiss-Prot human protein database. As digestion rules
for HLA peptides are unknown, a search engine that supports no-enzyme-specific digestion is
needed (we used PEAKS X [18]). Identified normal HLA peptides and their peptide-spectrum
matches (PSMs) at 1% false discovery rate (FDR) represent the patient's immunopeptidome
and its spectral library. Mutated HLA peptides were not presented in the protein database, so

their spectra remained unlabeled. For example, we identified 341,216 PSMs of 35,551 HLA-I
peptides and 67,021 PSMs of 9,664 HLA-II peptides from Mel-15 datasets. The numbers of
unlabeled spectra were 596,915 and 135,490, respectively (Supplementary Table S1).

In step 2, we used the identified normal HLA peptides and their PSMs to train DeepNovo, a
neural network model for *de novo* peptide sequencing [17, 18]. In addition to capturing fragment
ions in tandem mass spectra, DeepNovo learns sequence patterns of peptides by modelling
them as a special language with an alphabet of 20 amino acid letters. This unique advantage
allowed us to train a personalized model to adapt to a specific immunopeptidome of an
individual patient and achieved much better accuracy than a generic model (results are shown
in a later section). At the same time, it was essential to apply counter-overfitting techniques so
that the model could predict new peptides that it had not seen during training. We partitioned the
PSMs into training, validation, and test sets (ratio 90-5-5, respectively) and restricted them not
to share common peptide sequences. We stopped the training process if there was no
improvement on the validation set and evaluated the model performance on the test set. As a
result, our personalized model was able to both achieve very high accuracy on an individual
immunopeptidome and detect mutated peptides. This approach is particularly useful for
missense mutations (the most common source of neoantigens) because they still preserve most
patterns in the peptide sequences.

In step 3, we used the personalized DeepNovo model to perform *de novo* peptide sequencing
on both labeled spectra (i.e., the PSMs identified in step 1) and unlabeled spectra. Results from
labeled spectra were needed for accuracy evaluation and calibrating prediction confidence
scores. Peptides identified from unlabeled spectra ***and*** not presented in the protein database
were defined as "*de novo* peptides" and would be further analyzed in the next steps to find
candidate neoantigens of interest.

In step 4, a quality control procedure was designed to select high-confidence *de novo* peptides
and to estimate their FDR. We first calculated the accuracy of *de novo* sequencing on the test
set of PSMs by comparing the predicted peptide to the true one for each spectrum. DeepNovo
also provides a confidence score for each predicted peptide, which can be used as a filter for
better accuracy. Since the test set did not share common peptides with the training set, we
expected the distribution of accuracy versus confidence score on the test set to be close to that
of *de novo* peptides which the model had not seen during training. Thus, we calculated a score
threshold at a precision of 95% on the test set and used it to select high-confidence *de novo*
peptides (Figure 2b). Finally, to estimate the FDR of high-confidence *de novo* peptides, we
performed a second-round PEAKS X search of all spectra against a combined list of those
peptides and the database peptides (i.e. normal HLA peptides identified in step 1). Only *de novo*
peptides identified at 1% FDR were retained. Thus, our stringent procedure of quality control
guaranteed that each *de novo* peptide was supported by solid evidences from two independent
tools, DeepNovo and PEAKS X. For instance, we found 16,226 HLA-I and 2,717 HLA-II high-
confidence *de novo* peptides from Mel-15 datasets. Among them, 5,320 HLA-I and 863 HLA-II
de novo peptides passed 1% FDR filter (Supplementary Table S1).

We applied this workflow to train personalized models for another four patients and to predict
their HLA-I and HLA-II *de novo* peptides. The five patients, Mel-5, Mel-8, Mel-12, Mel-15, Mel-
16, were selected because their neoantigens had been identified and validated by a
proteogenomic database search approach in [13]. In total, we identified 10,440 HLA-I and 1,585
HLA-II *de novo* peptides at 1% FDR (Table 1). The number of identified database peptides in

step 1 of the workflow was 97,526 HLA-I and 15,835 HLA-II. Thus, our *de novo* sequencing
results expanded the immunopeptidomes by approximately 10%.

Advantages of Personalized Model over Generic Model

To demonstrate the advantages of our personalized approach, we compared the personalized
model of patient Mel-15's HLA-I to a generic model, which had the same neural network
architecture but was trained on a combined HLA-I dataset of 9 other patients from the same
study [13]. All datasets were derived from the same experiment and instrument, the only
difference is the immunopeptidomes of the patients. The combined dataset has 477,482 PSMs,
which is 39.9% larger than the Mel-15 dataset (477,482 / 341,216 = 1.399). Figure 1a shows the
accuracy of the personalized model versus the generic model on the Mel-15 test set. As
mentioned earlier, this test set did not share common peptides with the Mel-15 training set, so
both models had not seen the test peptides during training. The personalized model achieved
14.3% higher accuracy at the peptide level (0.6939 / 0.6070 = 1.143) and 3.8% higher accuracy
at the amino acid level (0.8668 / 0.8349 = 1.038), despite its smaller training set. The superiority
of the personalized model over the generic one can also be seen from the accuracy-versus-
score distribution in Figure 2b. At the same level of amino acid accuracy, e.g. 95%, the
personalized model required a lower score cutoff, thus allowing more *de novo* peptides to be
identified. Indeed, Figure 1c shows that the personalized model identified 87.8% more high-
confidence *de novo* peptides (16,226 / 8,642 = 1.878) and 38.9% more *de novo* peptides at 1%
FDR (5,320 / 3,829 = 1.389). More importantly, the personalized model was able to capture 6 of
8 target neoantigens of patient Mel-15 (Table 2), while the generic model only recovered 3 of
them. Those results demonstrate that our personalized approach substantially improves the
accuracy and identification rate of *de novo* peptides by adapting to a specific immunopeptidome
of an individual patient.

Analysis of Immune Characteristics of *De novo* HLA peptides

In this section, we studied common immune features of *de novo* HLA peptides and compared
them to normal HLA peptides, i.e. those identified by the database search engine in step 1 of
the workflow. We also compared to previously reported human epitopes from the Immune
Epitope Database (IEDB) [20].

Figure 2d shows the distribution of PEAKS X identification scores of *de novo* PSMs against
those of database and decoy PSMs for HLA-I peptides of patient Mel-15. The distributions
confirm that the *de novo* peptides have strong supporting PSMs as the database peptides and
are clearly distinguishable from the decoy ones. The supporting PSMs of all *de novo* HLA
peptides of five patients are provided in Supplementary Table S2.

Next, we compared *de novo* and database HLA-I peptides of patient Mel-15 to 18,022 IEDB
epitopes, which were retrieved according to the patient's six alleles (HLA-A03:01, HLA-A68:01,
HLA-B27:05, HLA-B35:03, HLA-C02:02, HLA-C04:01). The Venn diagram in Figure 2e shows
that 56 *de novo* peptides have been reported as epitopes in earlier studies. Note that the *de*
novo peptides were specific to an individual patient and were not presented in the protein
database, so the chance to find them in IEDB is rare. Even 81.4% (28,943 / 35,551) of the
database peptides were not found in IEDB. This is due to the large variation of HLA peptides
and further emphasizes the importance of our personalized approach. Figure 2e further shows
that both *de novo* and database peptides have the same characteristic length distribution as
IEDB epitopes. For the other four patients Mel-5, Mel-8, Mel-12, and Mel-16, we also found that

the length distributions of their *de novo* HLA-I peptides are very similar to those of database
peptides (Supplementary Figures S1a-d). However, for HLA-II, the *de novo* peptides tend to be
longer than the database ones (Supplementary Figures S1e, f). We hypothesize that it might be
challenging for the database search engine to identify long HLA-II peptides when the digestion
rule is unknown.

One of the most widely used measures to assess HLA peptides is their binding affinity to MHC
proteins. We used NetMHCpan [10] to predict the binding affinity of the *de novo*, database, and
IEDB peptides for HLA-I alleles of patient Mel-15. Figures 2g shows that the *de novo* peptides
have the same level of binding affinity as database and IEDB peptides (p-value > 0.23 for
Mann-Whitney U test between *de novo* and IEDB peptides). Furthermore, majority of the *de*
novo peptides were predicted as good binders by multiple criteria: 79.3% (4,220 / 5,320) weak-
binding, 51.8% (2,757 / 5,320) strong-binding, and 74.0% (3,938 / 5,320) with binding affinity
less than 500 nM (Supplementary Figure S2). Similar results were observed for *de novo*
peptides of different HLA-I alleles of the other four patients (Supplementary Figure S3). We also
applied GibbsCluster [23], an unsupervised alignment and clustering method to identify binding
motifs without the need of HLA allele information. We found that the *de novo* peptides of patient
Mel-15 were clustered into four groups of which motifs corresponded exactly to four alleles of
the patient (Figure 2h). Note that both *de novo* sequencing and unsupervised clustering
methods do not use any prior knowledge such as protein database or HLA allele information,
yet their combination still revealed the correct binding motifs of the patient. This suggests that
our workflow can be used to identify novel HLA peptides of unknown alleles. Results from the
database peptides also yielded the same binding motifs (Supplementary Figure S4).

Finally, we used an IEDB tool (http://tools.iedb.org/immunogenicity/) [24] to predict the
immunogenicity of *de novo* HLA-I peptides and then compared to database, IEDB, and human
immunogenic peptides that were used in that original study (Figure 2i). We found that 38.8%
(2,065 / 5,320) of the *de novo* peptides had positive predicted immunogenicity (log-likelihood
ratio of immunogenic over non-immunogenic [24]). The *de novo* peptides had lower predicted
immunogenicity than the database and IEDB peptides, which in turn were less immunogenic
than the original peptides (Calis *et al.*). This was expected because the tool had been
developed on a limited set of a few thousands well-studied peptides. The predicted
immunogenicity of *de novo* HLA-I peptides of the other four patients are provided in
Supplementary Figure S5.

Overall, our analysis results confirmed the correctness, and more importantly, the essential
characteristics of *de novo* HLA peptides for immunotherapy. The remaining question is to select
candidate neoantigens from *de novo* HLA peptides based on their characteristics.

Neoantigen Selection and Evaluation

We considered several criteria that had been widely used in previous studies for neoantigen
selection [6-8, 13, 14, 21, 22]. Specifically, we checked whether a *de novo* HLA peptide carried
one amino acid substitution by aligning its sequence to the Swiss-Prot human protein database,
and whether that substitution was caused by one single nucleotide difference in the encoding
codon. In this paper, we refer to those substitutions as "missense-like mutations". For each
mutation, we recorded whether the wild-type peptide was also detected and whether the
mutated amino acid was located at a flanking position. For expression level information of a
peptide, we calculated the number of its PSMs, their total identification score, and their total

abundance. Finally, we used NetMHCpan and IEDB tools [10, 24] to predict the binding affinity
and the immunogenicity of a peptide. The results for 10,440 HLA-I and 1,585 HLA-II *de novo*
peptides of five patients are provided in Supplementary Table S3.

To select candidate neoantigens, we focused on *de novo* HLA peptides that carried one single
missense-like mutation. This criterion reduced the number of peptides considerably, e.g. from
5,320 to 328 HLA-I and from 863 to 154 HLA-II peptides of patient Mel-15. We further filtered
out peptides with only one supporting PSM or with mutations at flanking positions because they
were more error-prone and less stable to be effective neoantigens. In average, we obtained 154
HLA-I and 47 HLA-II candidates per patient. Expression level, binding affinity, and
immunogenicity can be further used to prioritize candidates for experimental validation of
immune response; we avoided using those information as hard filters (Supplementary Table
S3).

We cross-checked our *de novo* HLA peptides against the nucleotide mutations and mRNA
transcripts in the original publication [13]. We identified seven HLA-I and ten HLA-II candidate
neoantigens that matched missense variants detected from exome sequencing (Tables 2 and
3). The first seven were among eleven neoantigens reported by the authors using a
proteogenomic approach that required both exome sequencing and proteomics database
search. Two HLA-I neoantigens, "GRIAF**F**LKY" and "K**L**ILWRGLK", had been experimentally
validated to elicit specific T-cell responses. We indeed observed that those two peptides had
superior immunogenicity, and especially, expression level of up to one order of magnitude
higher than the other neoantigens (Table 2). This observation confirms the critical role of
peptide-level expression for effective immunotherapy, in addition to immunogenicity and binding
affinity.

The ten HLA-II candidate neoantigens were novel and had not been reported in [13]. They were
clustered around a single missense mutation and were a good example to illustrate the
complicated digestion of HLA-II peptides (Table 3). Eight of them were predicted as strong
binders by NetMHCIIpan (rank <= 2%), two as weak binders (rank <= 10%). The peptide
located at the center of the cluster, "TSTRTYSL**S** SALRPS", showed both highest expression
level and binding affinity, thus representing a promising target for further experimental
validation. Interestingly, another peptide, "SL**S**SALRPSTSRSLY", showed up in both HLA-I and
HLA-II datasets with very high expression level (Tables 2 and 3). Using a consensus method of
multiple binding prediction tools from IEDB to double-check, we found that this peptide had a
binding affinity rank of 0.08%, instead of 4.5% as predicted by NetMHCIIpan, and exhibited a
different binding motif from the rest of the cluster. Thus, given its superior binding affinity and
expression level, this peptide would also represent a great candidate for immune response
validation.

We also investigated the four HLA-I neoantigens that had been reported in [13] but were not
detected by our method. Three of them were not supported by good PSMs, and in fact,
DeepNovo and PEAKS X identified alternative peptides that better matched the corresponding
spectra (Supplementary Figure S6). The remaining neoantigen was missed due to a *de novo*
sequencing error. We noticed that all four peptides had been originally identified at 5% FDR
instead of 1%, so their signals were possibly too weak for identification.

DISCUSSION

In this study, we proposed a personalized *de novo* sequencing workflow to identify HLA neoantigens directly and solely from mass spectrometry data. The key advantage of our method is the ability of its deep learning model to adapt to a specific immunopeptidome of an individual patient. This personalized approach greatly improved the performance of *de novo* sequencing and allowed to accurately identify mutated HLA peptides. For instance, we showed that a personalized model achieved up to 14.3% higher accuracy than a generic model, identified 38.9% more *de novo* peptides at 1% FDR, and doubled the number of validated neoantigens. We applied the workflow to five melanoma patients and identified 10,440 HLA-I and 1,585 HLA-II *de novo* peptides at 1% FDR, expanding their immunopeptidomes by approximately 10%. Our analysis also demonstrated that the *de novo* HLA peptides exhibited the same immune characteristics as previously reported human epitopes, including binding affinity, immunogenicity, and expression level, which are essential for effective immunotherapy. We cross-checked our *de novo* HLA peptides against exome sequencing results and found ten novel HLA-II neoantigens that had not been reported earlier. This result demonstrated the capability of our *de novo* sequencing approach to overcome the challenges of unknown digestion rules and binding prediction for HLA-II peptides. Last but not least, our *de novo* sequencing workflow directly predicted neoantigens from mass spectrometry data and did not require genome-level information nor HLA alleles of the patient as in existing approaches. Such an independent approach allows to discover novel mutated peptides that might be difficult to detect at the genome level, e.g. cis- and trans-spliced peptides. Thus, our personalized *de novo* sequencing workflow to predict mutated peptides from the cancer cell surface presents a simple and direct solution to discover neoantigens for cancer immunotherapy. As Newton said, "nature is pleased with simplicity".

Acknowledgements

This work was funded in part by the NSERC OGP0046506 grant and the Canada Research Chair program. N.H.T. was supported by the Mitacs Elevate Fellowship. The authors thank Dr. Kwok Pui Choi for proofreading the manuscript.

Author Contributions

M.L. and B.S. conceived the research idea. N.H.T. designed the neoantigen discovery workflow. N.H.T and R.Q. implemented the software and analyzed the results. X.C. and L.X. contributed to model design, software development, and data analysis. N.H.T, M.L., and R.Q. wrote the manuscript. M.L., B.S., and L.X. supervised the research project.

Competing Interests Statement

The workflow in Figure 1 has been applied for patent in the USPTO Provisional Application by Bioinformatics Solutions Inc., Waterloo, Canada. The authors are named inventors in the patent application. L.X., X.C., and B.S. are employees of Bioinformatics Solutions Inc.

Software Availability

328 DeepNovo and the workflow are implemented in Python. The latest version is provided to the
329 editor and reviewers during the review process and shall be made available on GitHub
330 (https://github.com/nh2tran/DeepNovoAA) upon publication.

References

1. Hu, Z., Ott, P.A., & Wu, C.J. Towards personalized, tumour-specific, therapeutic vaccines
for cancer. *Nat. Rev. Immunol.* **18**, 168-182 (2018).
2. Editorial. The problem with neoantigen prediction. *Nat. Biotechnol.* **35**, 97 (2017).
3. Vitiello, A. & Zanetti M. Neoantigen prediction and the need for validation. *Nat. Biotechnol.*
35, 815-817 (2017).
4. Van Allen, E.M. et al. Genomic correlates of response to CTLA-4 blockade in metastatic
melanoma. *Science* **350**, 207-211 (2015).
5. Rizvi, N.A. *et al.* Mutational landscape determines sensitivity to PD-1 blockade in non–
small cell lung cancer. *Science* **348**, 124-128 (2015).
6. Ott, P.A. *et al.* An immunogenic personal neoantigen vaccine for patients with melanoma.
Nature **547**, 217-221 (2017).
7. Sahin, U. *et al.* Personalized RNA mutanome vaccines mobilize poly-specific therapeutic
immunity against cancer. *Nature* **547**, 222-226 (2017).
8. Carreno, B.M. *et al.* A dendritic cell vaccine increases the breadth and diversity of
melanoma neoantigen-specific T cells. *Science* **348**, 803-808 (2015).
9. Andreatta, M. & Nielsen, M. Gapped sequence alignment using artificial neural networks:
application to the MHC class I system. *Bioinformatics* **32**, 511-517 (2016).
10. Jurtz, V. *et al.* NetMHCpan-4.0: improved peptide-MHC class I interaction predictions
integrating eluted ligand and peptide binding affinity data. *J. Immunol.* **199**, 3360-3368
(2017).
11. Abelin J.G. *et al.* Mass spectrometry profiling of HLA-associated peptidomes in mono-
allelic cells enables more accurate epitope prediction. *Immunity* **46**, 315-326 (2017).
12. Bulik-Sullivan, B. *et al.* Deep learning using tumor HLA peptide mass spectrometry
datasets improves neoantigen identification. *Nat. Biotechnol.* **37**, 55-63 (2019).
13. Bassani-Sternberg M. *et al.* Direct identification of clinically relevant neoepitopes
presented on native human melanoma tissue by mass spectrometry. *Nat. Commun.* **7**,
13404 (2016).
14. Laumont, C.M. *et al.* Noncoding regions are the main source of targetable tumor-specific
antigens. *Sci. Transl. Med.* **10**, eaau5516 (2018).
15. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.*
26, 1367-1372 (2008).
16. Zhang, J. *et al.* PEAKS DB: De novo sequencing assisted database search for sensitive
and accurate peptide identification. *Mol. Cell. Proteomics* **11**, M111.010587 (2012).
17. Tran, N.H, Zhang, X., Xin, L., Shan, B. & Li, M. *De novo* peptide sequencing by deep
learning. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 8247-8252 (2017).
18. Tran, N.H *et al.* Deep learning enables *de novo* peptide sequencing from data-
independent-acquisition mass spectrometry. *Nat. Methods* **16**, 63-66 (2019).

19. Sutskever, I., Vinyals, O. & Le, Q. Sequence to sequence learning with neural networks.
Adv. Neural Inf. Process. Syst. **27**, 3104-3112 (2014).
20. Vita, R. *et al.* The Immune Epitope Database (IEDB): 2018 update. *Nucleic Acids Res.* **47**,
D339-D343 (2018).
21. Bassani-Sternberg, M., Pletscher-Frankild, S., Jensen L.J., & Mann, M. Mass
spectrometry of human leukocyte antigen class I peptidomes reveals strong effects of
protein abundance and turnover on antigen presentation. *Mol. Cell. Proteomics* **14**, 658-
673 (2015).
22. Keskin, D.B. *et al.* Neoantigen vaccine generates intratumoral T cell responses in phase
Ib glioblastoma trial. *Nature* **565**, 234-239 (2019).
23. Andreatta, M., Alvarez, B. & Nielsen, M. GibbsCluster: unsupervised clustering and
alignment of peptide sequences. *Nucleic Acids Res.* **45(W1)**, W458-W463 (2017).
24. Calis, J.J.A. *et al.* Properties of MHC class I presented peptides that enhance
immunogenicity. *PLoS Comput. Biol.* **9**, e1003266 (2013).
25. Faridi, P. et al. A subset of HLA-I peptides are not genomically templated: Evidence for
cis- and trans-spliced peptide ligands. *Sci. Immunol.* **3**, eaar3947 (2018).

Figure Legends

Figure 1. Personalized *de novo* sequencing workflow for neoantigen discovery.
(HLA: Human Leukocyte Antigen; FDR: False Discovery Rate).

Figure 2. Accuracy and immune characteristics of *de novo* HLA-I peptides from patient Mel-15
dataset. (a) Accuracy of *de novo* peptides predicted by personalized and generic models. (b)
Distribution of amino acid accuracy versus DeepNovo confidence score for personalized and
generic models. (c) Number of *de novo* peptides identified at high-confidence threshold and at
1% FDR by personalized and generic models. (d) Distribution of identification scores of *de novo*,
database, and decoy peptide-spectrum matches. The dashed line indicates 1% FDR threshold.
(e) Venn diagram of *de novo*, database, and IEDB peptides. (f) Length distribution of *de novo*,
database and IEDB peptides. (g) Distribution of binding affinity ranks of *de novo*, database, and
IEDB peptides. Lower rank indicates better binding affinity. The two dashed lines correspond to
the ranks of 0.5% and 2%, which indicate strong and weak binding, respectively, by
NetMHCpan. (h) Binding sequence motifs identified from *de novo* peptides by GibbsCluster. (i)
Immunogenicity distribution of *de novo*, database, IEDB, and Calis *et al.*'s peptides [24].
(HLA: Human Leukocyte Antigen; FDR: False Discovery Rate; IEDB: Immune Epitope
Database).

Individual patient's normal & mutated HLA peptides

Mass spectrometry →

Tandem mass spectra

Step 1
Build the immunopeptidome of the patient
- Standard database search engine
- UniProt/Swiss-Prot human database
- No enzyme specificity

Identified normal HLA peptides &
their peptide-spectrum matches

Unlabeled spectra

Step 2
Train personalized machine learning model
- Neural networks to learn peptidome
- Counter overfitting to detect mutations

Step 3
Personalized *de novo* sequencing
- labeled spectra for quality control
- unlabeled spectra for mutated peptides

Identified mutated HLA peptides &
their peptide-spectrum matches

Step 4
Quality control
- *De novo* confidence score
- Second-round database search with FDR

Step 5
Neoantigen selection
- Missense mutation against wild-type
- Peptide expression level
- Binding affinity & immunogenicity

Table 1. Number of *de novo* and database HLA peptides identified at 1% FDR.

Patient ID	HLA-I		HLA-II	
	Database	*De novo*	Database	*De novo*
Mel-5	12,998	1,272	MS data not available	
Mel-8	13,635	1,235	MS data not available	
Mel-12	10,068	1,354	MS data not available	
Mel-15	35,551	5,320	9,664	863
Mel-16	25,274	1,259	6,171	722

FDR: False Discovery Rate

MS: Mass Spectrometry

Table 2. HLA-I candidate neoantigens that matched to RNA-Seq results.

Patient ID	Genename	Transcript ID	Aa change	Wild-type peptide	*De novo* peptide	Peptide expression	Binding affinity rank %	Immunogenicity
Mel-5	GABPA	ENST00000354828	Glu161Lys	ETSEQVTRW	ETSKQVTRW	2.4E+06	0.23	-0.28
Mel-8	NOP16	ENST00000621444	Pro169Leu	SPGPVKLEP	SPGPVKLEL	1.0E+07	0.06	-0.11
Mel-15	SYTL4	ENST00000263033	Ser363Phe	GRIAFSLKY	GRIAFFLKY	1.3E+08	0.22	0.12
Mel-15	RBPMS	ENST00000517860	Pro46Leu	RPFKGYEGSLIK	RLFKGYEGSLIK	5.8E+06	0.03	-0.19
Mel-15	SEC23A	ENST00000307712	Pro52Leu	PPIQYEPVL	LPIQYEPVL	7.2E+06	0.02	-0.01
Mel-15	NCAPG2	ENST00000441982	Pro134Leu	KPILWRGLK	KLILWRGLK	1.3E+07	0.04	0.27
Mel-15	AKAP6	ENST00000280979	Met1482Ile	KLKLPMIMK	KLKLPIIMK	3.6E+06	0.03	-0.21
Mel-15	VIM	ENST00000224237	Gly41Ser	SLGSALRPSTSRSLY	SLSSALRPSTSRSLY	3.4E+07	not available	not available

HLA: Human Leukocyte Antigen

PSM: Peptide-Spectrum Match

Underlined red letters indicate mutated amino acids.

Binding affinity and immunogenicity information are not available for "SLSSALRPSTSRSLY" because this is actually an HLA-II, not HLA-I peptide.

Table 3. HLA-II candidate neoantigens that matched to RNA-Seq results.

... S Y V T T S T R T Y S L G S A L R P S T S R S L Y ...	Peptide expression	Binding affinity rank %
Y V T T S T R T Y S L S S A L R P S T S	4.3E+06	1.8
V T T S T R T Y S L S S A L R P S T S	2.3E+06	1.8
S L S S A L R P S T S R S L Y	1.7E+07	4.5
S Y V T T S T R T Y S L S S A L R P S T S	9.8E+06	1.7
V T T S T R T Y S L S S A L R P S	3.0E+06	1.7
T T S T R T Y S L S S A L R P S	1.1E+07	1.0
Y V T T S T R T Y S L S S A L R P S T	3.7E+05	2.0
T T S T R T Y S L S S A L R P S T S	7.5E+05	1.6
Y V T T S T R T Y S L S S A L R P S	8.5E+05	2.5
T S T R T Y S L S S A L R P S	1.3E+07	0.6

HLA: Human Leukocyte Antigen
The mutated amino acid is underlined and highlighted in red for the reference (top row) and the neoantigens.
This mutation site is located at Gly41Ser, transcript ENST00000224237, gene name VIM, patient Mel-15.