Environmental DNA facilitates accurate, indigenous-led, inexpensive, and multi-year population estimates of millions of Eulachon fish

Meredith Pochardt[1,2,3], Jennifer M. Allen[1], Ted Hart[2], Sophie D. L. Miller[4], Douglas W. Yu[4,5,6], Taal Levi[1]

[1] Department of Fisheries and Wildlife, Oregon State University, Corvallis, OR 97331, USA

[2] Chilkoot Indian Association, Haines, AK 99827 USA

[3] Takshanuk Watershed Council, Haines, AK 99827 USA

[4] State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan, 650223, China

[5] Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences, Kunming Yunnan, 650223 China

[6] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, Norfolk, NR47TJ, UK

Corresponding author E-mail: Taal.Levi@oregonstate.edu,

Keywords: Alaska**,** anadromous, eDNA, eulachon**,** fisheries, *Thaleichthys pacificus*, Tlingit

Running head: indigenous eDNA monitoring of eulachon

Abstract

Although environmental DNA shed from an organism is now widely used for species detection in a wide variety of contexts, mobilizing environmental DNA for management requires estimation of population size and trends rather than simply assessing presence or absence. However, the efficacy of environmental-DNA-based indices of abundance for long-term population monitoring have not yet been assessed. Here we report on the relationship between six years of mark-recapture population estimates for eulachon (*Thaleichthys pacificus*) and 'eDNA rates,' which are calculated from the product of stream flow and DNA concentration. Eulachon are a culturally and biologically important anadromous fish that have significantly declined in the southern part of their range but were historically rendered into oil and traded. Both the peak eDNA rate and the area under the curve of the daily eDNA rate were highly predictive of the mark-recapture population estimate, explaining 84.96% and 92.53% of the deviance respectively. Even in the absence of flow correction, the peak of the daily eDNA concentration explained an astonishing 89.53% while the area under the curve explained 90.74% of the deviance. These results support the use of eDNA to monitor eulachon population trends and represent a >80% cost savings over mark-recapture, which could be further increased with automated water sampling, reduced replication, and focused temporal sampling. Due to its logistical ease and affordability, eDNA sampling can facilitate monitoring a larger number of rivers and in remote locations where mark-recapture is infeasible.

Introduction

While the environmental DNA shed from an organism is now widely used for species

detection in a wide variety of contexts (Barnes & Turner, 2016; Rees, Maddison, Middleditch,

Patmore, & Gough, 2014; Jerde, Mahon, Chadderton, & Lodge, 2011; Laramie, Pilliod, &

Goldberg, 2015; Mächler, Deiner, Steinmann, & Altermatt, 2014; Rees et al., 2014; Takahara,

Minamoto, Yamanaka, Doi, & Kawabata, 2012), mobilizing environmental DNA for

management requires estimation of population size and trends rather than simply assessing

presence or absence. Recent research suggests that eDNA quantified with real-time quantitative

polymerase chain reaction (PCR) or digital-droplet PCR can provide a proxy for actual

abundance in controlled experiments (Rees, Maddison, Middleditch, Patmore, & Gough, 2014),

in ponds (Lacoursière-Roussel, Côté, Leclerc, & Bernatchez, 2016; Takahara et al., 2012) in

streams (Doi et al., 2015; Levi et al., 2019; Lodge et al., 2012; Tillotson et al., 2018; Wilcox et

al., 2016) and in marine bays (Plough et al., 2018) However, the efficacy of environmental DNA

based indices of abundance in natural settings have produced mixed results (Yates, Fraser, &

Derry, 2019) and have not yet been assessed in a management context for long-term population

monitoring.

Anadromous fish enter freshwater systems to spawn, often in large number, representing

the possibility to quantify the size of the spawning population with environmental DNA to

inform management and population trends. While recent research has suggested that daily eDNA

counts correlate well with daily entry by adult salmon or daily outmigration of salmon smolts

(Levi et al., 2019), a more important question is whether total run sizes can be accurately

predicted for long interannual population monitoring programs.

Owing to their short run time and large spawning aggregations, Eulachon (*Thaleichthys*

pacificus), a lipid-rich, anadromous smelt of the family Osmeridae (Mecklenburg et al. 2002),

make an ideal case study to test eDNA for long-term population monitoring of anadromous fish.

Adult eulachon have an average size of 18 to 22 cm (Spangler, 2002). The historic range of

eulachon stretched from southern California to the Bering Sea in southwest Alaska (Hart, 1973).

The majority of eulachon populations have been declining since the 1990s (Hay & Mccarter,

2000). In 2010, the National Marine Fisheries Service (NMFS) listed the southern distinct

population segment in Washington, Oregon, and California as Threatened under the Endangered

Species Act (NOAA, 2010). Because there is no commercial eulachon fishery in northern

Southeast Alaska, there is no harvest regulation or management, agency oversight, or monitoring

of population trends. While some eulachon population declines have been well documented (Hay

& Mccarter, 2000), the status of most eulachon populations is either unknown or anecdotal.

In Southeast Alaska, they are the first anadromous fish to return after the long winter, and

as a result, are a key resource for indigenous people and for wildlife. For the Northwest Coast

native people, eulachon are a culturally significant staple food source that is consumed fresh,

dried, or smoked, and are frequently rendered into oil (Betts, 1994). Historically, eulachon oil

was the most important trade item on a network of 'grease trails' between coastal and interior

peoples, and it is still used and traded (Betts, 1994; Moody & Pitcher, 2010). Eulachon spawn

just prior to the breeding season of many consumers, including marine mammals, thus providing

a high-energy prey resource at an energetically demanding time (Sigler, Womble, &

Vollenweider, 2004). The eulachon spawning aggregation draws enormous congregations of

seabirds, bald eagles (*Haliaeetus leucocephalus*), Steller sea lions (*Eumetopias jubatus*), harbor

seals (*Phoca vitulina*), and humpback whales (*Megaptera novaeangliae*) among many other

smaller predators and scavengers.   A lack of eulachon population information and the

cultural and subsistence value of the species led to the development of an indigenous-led

eulachon monitoring program in northern Southeast Alaska. In 2010 the Chilkoot Indian

Association and the Takshanuk Watershed Council initiated a modified Lincoln-Petersen

(Chapman, 1951, Lincoln, 1930; Petersen, 1896) mark-recapture population estimate on the

Chilkoot River near Haines, Alaska at the northern end of southeast Alaska (Fig. 1). This

program was successful in gathering baseline eulachon population data where none existed

previously; however, monitoring is challenging and expensive (~$20,000 annually), limiting the

feasibility of conducting long-term monitoring and limiting the possible geographic scope of

monitoring. In an effort to develop a more cost-effective monitoring method, in 2014 we began

pairing the mark-recapture program with daily water sampling to evaluate the efficacy of

environmental DNA (eDNA) to produce an index of eulachon abundance.

Here we compare six years (2014-2019) of mark-recapture eulachon abundance estimates

with eulachon eDNA quantification to test whether long-term, affordable, indigenous-led

monitoring of eulachon populations could be effectively achieved with environmental DNA.

This method could facilitate intertribal cooperation for affordable monitoring of a culturally

important subsistence and economic resource on a regional scale. Such regional monitoring is

particularly important for eulachon, which exhibit low site-fidelity and thus regional broad-scale

population structure (Flannery, Spangler, Norcross, Lewis, & Wenburg, 2013) such that a true

population decline can only be verified by monitoring multiple river systems.

Methods

Methods

Study System

The Chilkoot River near Haines, Alaska has long been a culturally and ecologically

important river. The lower Chilkoot River flows 1.5 km from Chilkoot Lake to the ocean at the

terminus of a large fjord. The Chilkoot Tlingit village and fishcamp was historically located

along the banks of the Chilkoot River, which is still utilized for eulachon fishing and processing

today (Betts, 1994; Olds, 2016). Eulachon typically spawn in the lower reaches of the Chilkoot

River (Hay & Mccarter, 2000) where mostly indigenous harvesters capture large quantities for

smoking, frying, and rendering into oil in pits.

The Chilkoot Indian Association initiated a eulachon mark-recapture study to develop the

first population baseline for the Chilkoot River, which is now the longest eulachon population

dataset in Southeast Alaska (Alaska Department of Fish and Game Aquatic Resources Permit:

SF-2014-027, SF2015-066, SF2016-113, SF2017-062, SF2018-072). This effort was initiated

because anecdotal observation suggested that the run size and timing on the Chilkoot River

differed from traditional knowledge, and because the decline of the southern distinct population

segment of eulachon was substantial enough to warrant threatened status under the Endangered

Species Act (NOAA, 2010). The Endangered Species Act listing of eulachon led to concern by

Chilkoot Indian Association tribal members that a decline in northern Southeast Alaska, where a

strong subsistence fishery remained, would go undocumented, and thus un-remediated, without

quantification of the current run size (Olds, 2016).

Mark-Recapture

At the mouth of the Chilkoot River, eulachon were captured using a modified fyke net

trap and dip nets. The initial captured eulachon (M group) were transferred in small groups to

plastic dishpans where they could be easily handled to clip off the adipose fin using retina

scissors and returned to the river. To avoid excessive increases in temperature and to reduce the

possibility of disease transmission, the water in the dishpans was changed between each group

and the dishpans were rinsed with river water. To allow time for the marked fish to mix with the

unmarked fish, the recapture group was captured approximately 0.75 km upstream of the trap

location (C and R group) (Fig 1). Eulachon in the second capture group were collected by field

crews wading through the river with dip nets making sure to sample all portions of the river and

with the help of subsistence harvesters. The captured fish were examined for a clipped adipose

fin before releasing. To avoid repetitive sampling of the same fish, field crews started at a

downstream point and worked their way upstream. Eulachon are thought to be semelparous

(spawning only once), which negates recapturing fish marked in a previous year (Clarke, Lewis,

Telmer, & Shrimpton, 2007). A modified Lincoln-Peterson estimator equation (Chapman, 1951)

was used $N = \frac{(M+1)(C+1)}{R+1} - 1$ where N = total population size, M = marked initially, C = total in

second sample, and R = marked recaptures. The standard error was calculated using the equation

$SE = \sqrt{[(M+1)(C+1)(M-R)(C-R)]/[(R+1)^2(R+2)]}$. The 95% confidence interval

was calculated as $CI = N \pm (1.96)(SE)$. Mark-recapture data were collected from 2010 through

2019, excluding 2013 where a lapse in funding prohibited collection.

Environmental DNA

We collected daily water samples for eulachon eDNA quantification just below the mark-

recapture trap location near the mouth of the Chilkoot River (Fig. 1) from 2014 through 2019.

The samples were taken as close to low tide as was feasible to avoid either DNA intrusion from

the estuary and/or dilution with an influx of tidal flow. Three replicate 1 L water samples were

collected from the same location each sampling day in sterile Whirl Pak bags starting in early to

mid-April and continuing for at least one week beyond the end of the mark-recapture study

duration (Table 2). The exception to this was 5 days in 2019 for which field crews mistakenly

filtered only 750 ml. We multiplied DNA concentrations from these days by 1.33 to account for

the reduced volume. We sampled for 8, 11, 19, 13, 17, and 25 days during each run from 2014

through 2019 respectively.

Each sample was transported from the field to the Takshanuk Watershed Council office

immediately after collection and was filtered through a Nalgene 47mm 0.45 micron cellulose

nitrate filter using either a peristaltic pump (Proactive Alexis peristaltic pump) or vacuum pump

(Gast model DOA-P704-AA) with a three-sample manifold. Filters were stored immediately in

100% ethanol within 2 mL cryovials and refrigerated until shipped to Oregon State University

for extraction. Filters were removed from ethanol and air-dried overnight in sterile, disposable

weight boats in a hepafiltered and UV-irradiated cabinet within a PCR-free laboratory to avoid

contamination. DNA was then extracted using the Qiagen DNeasy Blood and Tissue kit modified

to include a >48 hour soak in ATL buffer, which was found to produce higher and more

consistent yields. DNA was eluted in a total volume of 100 μl.

DNA Quantification

We developed a species-specific quantitative PCR assay for eulachon targeting a 187-bp

region of the *Cytochrome oxidase I* (COI) region of the mitochondrial genome based on

observed sequence divergence among Osmeridae fish species in the Pacific Northwest region of

North America including longfin smelt (*Spirinchus thaleichthys*), capelin (*Mallotus villosus*), and

rainbow smelt (*Osmerus mordax*). Specifically, we ensured at least 2 bp mismatch on the

forward primer and at least 3 bp mismatches on the probe to the other Osmeridae fishes. The

reverse primer contained a 2 bp mismatch to longfin smelt, a 3 bp mismatch to capelin, and a 1bp

mismatch to rainbow smelt. We tested our primers *in vitro* against longfin smelt tissue to ensure

no nonspecific binding, and *in natura* on water samples from a diversity of rivers in southeast

Alaska (Chilkoot, Chilkat, Taiya, Ferebee, Katzehin, Auke, Berners, Lace, Antler, Mendenhall)

and Oregon (Columbia, Cowlitz) outside of the eulachon run to ensure no nonspecific binding to

non-Osmeridae fishes.

The probe was labeled with a 5' FAM fluorescent marker and a minor-groove-binding

non-fluorescent quencher on the 3' end. Primer3 software (Untergasser et al. 2012) was used to

select the following primers: Euc_COI_R (5'- CTCCCTCCTTCCTTCTCCTT-3'), Euc_COI_R

(5'- GGTCTGGTACTGGGAAATGG-3') and the internal probe Euc_COI_I (5'-

6FAM*AGCGGGAGCCGGGACTGGCT*MGBNFQ).

A Bio-Rad QX200 AutoDG Droplet Digital PCR system (Hercules, CA. USA) at the

Oregon State University Center for Genome Research and Biocomputing was used to quantify

DNA concentrations in duplicate. A 22 µl reaction was carried out containing (final

concentrations) 1 x ddPCR Supermix for probes (no dUTP), 900 nM of both forward and reverse

primers, 250 nM internal probe and 4 µl of DNA extract. Droplets were generated using the

QX200 AutoDG system. Cycling consisted of 95 °C for 10 mins, followed by 45 cycles of 94 °C

for 30 secs, and 60 °C for 1 min, ending with 96 °C for 10 mins, allowing for a ramp rate of 2

°C/sec between steps. PCR setup occurred in a hepafiltered and UV-irradiated cabinet within a

PCR-free laboratory to avoid contamination. After the reaction, the droplets were read on a

Droplet Reader and analyzed with QuantaSoft Analysis Pro software (version 1.0.596). We

included extraction blanks every 35 samples, and every ddPCR plate included two no-template

controls (DI water), and positive controls (eulachon tissue extracts). We did not observe false

positives of eulachon in negative controls nor false negatives of eulachon tissue or water samples

when eulachon were observed in the river.

The concentration of eDNA is a function of both the amount of eDNA shed into the river

and dilution of eDNA due to increased stream flow. To calculate the flow-corrected eDNA rate,

we multiplied each day's ddPCR DNA concentration $\left(\frac{copies}{ul}\right)$ against the day's stream flow

$\frac{cubic\ feet}{sec}$. We refer to this as an eDNA rate because once the volume units cancel the result is

proportional to DNA copies/second (Levi et al. 2019). Stream flow measurements were taken

each day that an eDNA sample was collected immediately following the collection of the eDNA

sample. To measure streamflow, we used a rating curve developed by the Alaska Department of

Fish and Game for the Chilkoot River. To validate this rating curve, a stream flow measurement

was taken at the beginning of each field season on the Chilkoot following the USGS velocity-

area method using a type AA current meter (Turnipseed & Sauer, 2010). Following the initial

calibration of the rating curve, the daily river height was measured in feet off of an established

benchmark using surveying equipment, which was then transformed into a river discharge based

on the rating curve (Sowa, 2015).

Analysis

We evaluated the flow-corrected eDNA rate as an index of eulachon abundance based on

two metrics. First, we use the maximum eDNA rate (i.e. size of peak). Second, we used area

under the curve of the eDNA rate throughout the duration of the run. In each case, the daily

eDNA concentration was the average of 6 replicates (2 ddPCR within 3 replicate water samples).

DNA concentration Particularly in cases of multimodal runs, the area under the curve was

expected to provide a more accurate representation of the overall biomass. We computed the area

under the curve with the AUC function in the *DescTools* package version 0.99.27 (Signorell,

2019) in RStudio version 1.1.383 (RStudio Team, 2015). We additionally assessed the need for

flow correction by evaluating the relationship between uncorrected eDNA concentrations using

the same two metrics and mark-recapture population estimates. We used quasipoisson regression

to model the mark-recapture population estimates as a function of the natural log of the two

measures of eDNA rate and the uncorrected eDNA concentration.

Results

Mark-Recapture

The mark-recapture population estimate was initiated in 2010 and continued annually

through 2019, excluding 2013 due to funding constraints. Eulachon exhibited substantial

population fluctuations with a potential 5-6-year cyclic pattern for large returns (Fig. 2). The

average eulachon population estimate for the mark-recapture method from 2010-2019 (excluding

2013) was 8.4 million, with a maximum of 26.7 million in 2019 ($\pm$1,840,573), and a minimum of

319,568 in 2015 ($\pm$158,934) (Table 1). Eulachon arrival in the Chilkoot River was documented

as early as April 20th (2016) and as late as May 6th (2017), with run durations lasting between 4

(2015) and 13 days (2019).

Two notable anomalies occurred during the mark-recapture study period. In 2016 the run

consisted of multiple pulses with what appeared to be a definitive end of the run that was

followed by a final pulse of fish five days later. This final pulse of fish was not recorded as part

of the mark-recapture estimate but was captured in the eDNA data. Additionally, the 2015 run

was a failure, with the smallest number of fish during our study period, but anecdotal reports

suggested that other rivers in Northern Southeast Alaska received unusually large runs.

Environmental DNA

During eulachon runs, all ddPCR replicates of all technical replicates amplified with the

exception of the period that appeared in the field to be prior to any obvious eulachon entry in

which we either observed no amplification or very low copy number amplification of one

replicate but not both. eDNA concentrations varied substantially from near zero to a high of

328000 copies / μL during the peak of the run in 2017. The product of streamflow and eDNA

concentration, which we refer to as 'flow-corrected eDNA rate' (Fig. 3, see also Levi et al.

2019), was highly predictive of the eulachon population estimate generated through the mark-

recapture method. The natural log of the eDNA peak was significantly related to, and explained

84.96% of the deviance in, the mark-recapture population estimate (β=0.533, 95% CI [0.271,

0.898], $p = 0.027$), despite a multimodal eulachon run in 2016 that contained three distinct peaks.

The area under the curve eDNA rate explained 92.53% of the deviance in the mark-recapture

population estimate (β=0.502, 95% CI [0.338, 0.697], $p = 0.005$) (Fig. 4). The peak eDNA

concentration without flow correction explained 89.53% of the deviance in the mark-recapture

population estimate (β=0.503, 95% CI [0.310, 0.742], $p = 0.01$). The area under the curve even

without flow correction still explained 90.74% of the deviance in the mark-recapture population

estimate (β=0.443, 95% CI [0.292, 0.620], $p = 0.006$) (Fig. 4). The quasipoisson regression

models using either the flow-corrected eDNA rate peak (i.e. maximum of flow x DNA

concentration) or the area under the curve as a single predictor produced highly representative

predictions of mark-recapture population estimates (Fig. 5).

267 **Discussion**

The utility of eDNA for the detection of organisms has been widely documented

(Ficetola, Miaud, Pompanon, & Taberlet, 2008; Rees et al., 2014; Wilcox et al., 2016). Recently,

the next generation of eDNA science has evaluated the efficacy of quantifying the abundance of

species using eDNA (Doi et al., 2015; Levi et al., 2019; Takahara et al., 2012; Tillotson et al.,

2018). However, the expansion of eDNA beyond academic settings and into species

management and monitoring is just beginning. eDNA methods may be particularly promising for

the management of neglected species such as eulachon. This is true even if eDNA provides less

accurate or precise results than do traditional methods, because lower quality data from more

streams could result in more robust management decision-making than higher quality data from

just a few streams (Dowling et al., 2008), particularly for a fish that exhibits low site fidelity. In

addition, for many taxa, knowing population trends is just as important as precisely enumerating

abundance. However, our results suggest that this tradeoff is largely inconsequential; both the

flow-corrected and non-flow-corrected eDNA rate was predictive of the eulachon mark-recapture

population estimates at a small fraction of the cost. Further, eDNA was predictive of mark-

recapture population estimates even without flow-correction (Fig. 4D), which suggests the

possibility that eDNA-based quantitation of eulachon could be implemented in systems where

flow measurements cannot be obtained.

Unlike the mark-recapture method, which produced a single population estimate for the

eulachon run, the eulachon eDNA rate captured within-run phenology as eulachon abundance

varied in the Chilkoot River above the sampling location. eDNA was very effective at

quantifying run timing and was particularly effective at demonstrating that the 2016 eulachon

run was multimodal with three distinct pulses of eulachon that were separated by 4-5 days of

inactivity (Fig. 3). The third pulse in 2016 was not represented in the mark-recapture estimate

because field personnel had assumed that the run had terminated, but, due to the minimal labor

required for eDNA sampling, we continued sampling and were able to capture the full 2016 run

with eDNA.

Given the results presented here, eDNA has great potential as an indigenous-led,

efficient, and effective means of monitoring this culturally and ecologically important species.

Because eulachon in many regions, like northern Southeast Alaska, are not a commercially

harvested species, they do not receive agency funding for management. This leaves a financial

gap in monitoring that is often filled through indigenous-led efforts. A primary benefit of using

eDNA is the vastly reduced cost of monitoring under-funded species, such as eulachon. The

mark-recapture study on the Chilkoot River costs approximately \$28,000 per year; largely

because two 5-person crews are needed to properly implement the mark-recapture method. The

use of eDNA at the current Oregon State University rate of \$42/sample for 3 samples/day for

~13 days is \$3,000. Further cost savings would be accrued by reducing sampling prior to and

after the run, when DNA concentrations are low, and instead focusing measurement during the

~1 week peak of the run, because both the peak eDNA concentration and area under the curve

eDNA metrics are invariant to sampling additional days with very low DNA concentrations. The

economic viability of eDNA could be further increased with automated water samplers, currently

in development, or the use of citizen scientists (Biggs et al., 2015; Wilken, 2018).

An additional benefit of eDNA methods is that mark-recapture estimation is not

logistically feasible on all rivers. The Chilkoot poses a unique set of characteristics – single

channel, road accessible, and with a relatively distinct upper limit to spawning activity. Many

rivers in Southeast Alaska where eulachon spawn are glacially-fed, with wide, braided river

mouths that are in remote, road-less areas. A mark-recapture method at these locations would be

logistically challenging, in large part due to a large field-crew requirement. The appeal of eDNA

is the ability to simply collect and filter a water sample to derive an index of abundance, which

can be done by a single person in under one hour. The use of eDNA allows population data to be

gathered on rivers that otherwise would not be possible, which is vital in monitoring a population

that exhibits a regional genetic population structure where a decline in spawning biomass in any

one river system might not necessarily represent a decline in the regional eulachon population

(Flannery et al., 2013). The use of affordable and logistically feasible eDNA methods could

facilitate regional studies of eulachon population size, run timing, and synchrony among rivers,

which would allow for inference on regional population trends, environmental drivers of

population dynamics, and environmental drivers of spawning river selection (Bryant, 2009).

Measurement of eDNA concentrations at a point in space and time represents a simple

sampling process of mtDNA molecule counts per unit reaction volume, which can be modelled

by a Poisson distribution assuming that eDNA is well mixed. The maximum likelihood estimate

of the actual concentration of eDNA, λ, is equal to the sample mean of the N replicate eDNA

concentrations, and the variance around this estimate is equal to the sample mean of the eDNA

concentrations divided by N. Thus, the variance of the estimated 'true' eDNA concentration

declines quickly with the number of replicates from a maximum equal to the mean. In contrast,

mark-recapture analysis is a relatively complex statistical sampling process with a variance

around the population estimate of $([(M+1)(C+1)(M-R)(C-R)]/[(R+1)^2(R+2)])$,

which can be much larger than the mean population estimate (Table 1; SE^2 can be thousands of

times larger than N) if the number of recaptured individuals, R, is small relative to the number of

marked individuals, M, or the number captured in the second session, C (Chapman, 1951). Thus,

although eDNA concentrations are not in the useful units of individual animals, they can be

estimated precisely with limited replication, but the same is not true for mark-recapture

population estimates. Of course, while the observation error of eDNA concentrations in water

samples may be low, the process error linking eDNA concentrations to fish abundance can be

quite high due to the complexity of eDNA transport and degradation, variance in eDNA

production among individuals and through time, the random spatial location of organisms

relative to the sampling site, and more challenging stream morphologies among other

complexities.

Although there is substantial process error linking eDNA concentrations to fish

abundance, mark-recapture presents its own suite of problems. For example, the demographic-

closure assumptions of mark-recapture estimators are difficult to meet with an anadromous fish

that quickly enters and leaves the river (Pollock, 2018). The Chilkoot River mark-recapture study

lasts for the duration of the run (typically 4-8 days), beginning on the first day that fish are

observed in the river (typically late April) and ending once recapture sampling has exhausted all

new fish into the system (i.e. when recaptures are identifying double-marked fish). During this

time, new fish immigrate into the river while subsistence fishing activities actively remove fish,

thus violating closure. However, mark-recapture population estimates can be robust to moderate

violations of closure (Kendall, 1999). In this study, the closed-population assumption is thought

to be reasonably met because (1) initial marking efforts remained relatively constant and

continued until no new fish appear to be entering the system (i.e. approximately all individuals

were potentially subject to marking), (2) there was an equal probability of capture of marked and

unmarked fish by subsistence harvesters, and (3) we secured participation of subsistence

harvesters to search their catch for marked fish.

We have demonstrated that eDNA provides reliable quantification of anadromous

eulachon abundance. eDNA is thus a promising tool that can be mobilized by indigenous people

to affordably monitor noncommercial species that are neglected by agencies but are culturally

and/or economically important to native people. However, substantial uncertainty persists. In

particular, it is unknown whether our model correlating eulachon run size with flow-corrected

eDNA will be transferable to other rivers. This is unlikely to be the case when rivers have

different morphologies, such as braided floodplains with pockets of eulachon spawning

throughout. In such circumstances, a within-river index of abundance might be achieved by

monitoring several braids where eulachon congregate, or perhaps the estuary where mixing of

water might homogenize the sample. Once sampling specifications for rivers of various

morphologies are determined, the final step toward large-scale implementation is regional

cooperation among Alaska natives, First Nations in British Columbia, and tribal governments in

Washington and Oregon to facilitate monitoring of eulachon spawning rivers throughout the

Pacific Northwest.

Acknowledgements

We thank Oregon State University, The National Geographic Society (#9493-14), US Fish &

Wildlife Service Tribal Wildlife grant (AK-U-28-NA-1), North Pacific Landscape Conservation

Cooperative (FW13AP01047), and Bureau of Indian Affairs (A18AP00229) for funding this

work. Additionally, we thank the Chilkoot Indian Association Eulachon Tribal Working Group

for incorporating the cultural significance of eulachon into this research and all the tribal

members that worked as part of the mark-recapture crews. D.W. Yu was supported by the

National Natural Science Foundation of China (41661144002, 31670536, 31400470, 31500305),

the Key Research Program of Frontier Sciences, CAS (QYZDY-SSW-SMC024), the Bureau of

International Cooperation (GJHZ1754), the Strategic Priority Research Program of the Chinese

Academy of Sciences (XDA20050202, XDB31000000), the Ministry of Science and Technology

References

Barnes, M. A., & Turner, C. R. (2016). The ecology of environmental DNA and implications for
conservation genetics. *Conservation Genetics*. https://doi.org/10.1007/s10592-015-0775-4
Betts, M. F. (1994). The subsistence hooligan fishery of the Chilkat and Chilkoot Rivers.
Technical Paper Series, (213), 1–69.
Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C., Dejean, T., Griffiths, R. A., … Dunn, F.
(2015). Using eDNA to develop a national citizen science-based monitoring programme for
the great crested newt (Triturus cristatus). *Biological Conservation*, *183*, 19–28.
https://doi.org/10.1016/J.BIOCON.2014.11.029
Bryant, M. D. (2009). Global climate change and potential effects on Pacific salmonids in
freshwater ecosystems of southeast Alaska. *Climatic Change*, *95*(1–2), 169–193.
https://doi.org/10.1007/s10584-008-9530-x
Clarke, A. D., Lewis, A., Telmer, K. H., & Shrimpton, J. M. (2007). Life history and age at
maturity of an anadromous smelt, the eulachon Thaleichthys pacificus (Richardson).
Journal of Fish Biology, *71*(5), 1479–1493. https://doi.org/10.1111/j.1095-
8649.2007.01618.x
Doi, H., Uchii, K., Takahara, T., Matsuhashi, S., Yamanaka, H., & Minamoto, T. (2015). Use of
Droplet Digital PCR for Estimation of Fish Abundance and Biomass in Environmental
DNA Surveys. *PLOS ONE*, *10*(3), e0122763. https://doi.org/10.1371/journal.pone.0122763
Dowling, N. A., Smith, D. C., Knuckey, I., Smith, A. D. M., Domaschenz, P., Patterson, H. M.,
& Whitelaw, W. (2008). Developing harvest strategies for low-value and data-poor
fisheries: Case studies from three Australian fisheries. *Fisheries Research*, *94*(3), 380–390.
https://doi.org/10.1016/J.FISHRES.2008.09.033
Ficetola, G. F., Miaud, C., Pompanon, F., & Taberlet, P. (2008). Species detection using
environmental DNA from water samples. *Biology Letters*, *4*(4), 423–425.
https://doi.org/10.1098/rsbl.2008.0118
Flannery, B. G., Spangler, R. E., Norcross, B. L., Lewis, C. J., & Wenburg, J. K. (2013).
Microsatellite Analysis of Population Structure in Alaska Eulachon with Application to
Mixed-Stock Analysis. *Transactions of the American Fisheries Society*, *142*(4), 1036–1048.
https://doi.org/10.1080/00028487.2013.790841
Hart, J. (1973). Pacific fishes of Canada. *Bulletin of the Fisheries Research Board of Canada*,
180, 148–150.
Hay, D., & Mccarter, P. B. (2000a). Status of Eulachon Thaleichtheys pacificus in Canada.
Fisheries and Oceans Canada. Canadian Stock Assessment, *2000/145*.
Hay, D., & Mccarter, P. B. (2000b). Status of eulachon Thaleichthys pacificus in Canada.
Fisheries and Oceans Canada. Canadian Stock Assessment, *9R*.
Jerde, C. L., Mahon, A. R., Chadderton, W. L., & Lodge, D. M. (2011). "Sight-unseen" detection
of rare aquatic species using environmental DNA. *Conservation Letters*.
https://doi.org/10.1111/j.1755-263X.2010.00158.x
Kendall, W. (1999). Robustness of Closed Capture – Recapture Methods To Violations of the

Closure Assumption. *Ecology*, *80*(8), 2517–2525.

Lacoursière-Roussel, A., Côté, G., Leclerc, V., & Bernatchez, L. (2016). Quantifying relative

fish abundance with eDNA: a promising tool for fisheries management. *Journal of Applied*

Ecology. https://doi.org/10.1111/1365-2664.12598

Laramie, M. B., Pilliod, D. S., & Goldberg, C. S. (2015). Characterizing the distribution of an

endangered salmonid using environmental DNA analysis. *Biological Conservation*.

https://doi.org/10.1016/j.biocon.2014.11.025

Levi, T., Allen, J. M., Bell, D., Joyce, J., Russell, J. R., Tallmon, D. A., … Yu, D. W. (2019).

Environmental DNA for the enumeration and management of Pacific salmon. *Molecular*

Ecology Resources. https://doi.org/10.1101/394445

Lincoln, F. C. (1930). Calculating waterfowl abundance on the basis of banding returns. *US*

Department of Agriculture, *118*, 1–4.

Lodge, D. M., Turner, C. R., Jerde, C. L., Barnes, M. A., Chadderton, L., Egan, S. P., …

Pfrender Michael E. (2012). Conservation in a cup of water: estimating biodiversity and

population abundance from environmental DNA. *Molecular Ecology*, *21*(11), 2555–2558.

https://doi.org/10.1111/j.1365-294X.2012.05600.x

Mächler, E., Deiner, K., Steinmann, P., & Altermatt, F. (2014). Utility of environmental DNA

for monitoring rare and indicator macroinvertebrate species. *Freshwater Science*.

https://doi.org/10.1086/678128

Moody, M. F., & Pitcher, T. J. (2010). Eulachon (Thaleichthys pacificus): past and present. *The*

Fisheries Centre, University of British Columbia, Canada, *18*(2), 197.

https://doi.org/10.14288/1.0074735

NOAA. (2010). Endangered and Threatened Wildlife and Plants: Threatened Status for Southern

Distinct Population Segment of Eulachon. In *Federal Register* (Vol. 75).

https://doi.org/10.1021/j100299a032

Olds, A. (2016). *Integrating local and traditional knowledge and historical sources to*

characterize run timing and abundance of eulachon in the Chilkat and Chilkoot Rivers.

University of Alaska Fairbanks.

Petersen, C. G. J. (1896). The yearly immigration of young plaice into the Limfjord from the

German Sea. *Report of the Danish Biological Station*, *6*, 1–77.

Plough Id, L. V, Ogburn, M. B., Fitzgerald, C. L., Geranio, R., Marafino, G. A., & Richie, K. D.

(2018). *Environmental DNA analysis of river herring in Chesapeake Bay: A powerful tool*

for monitoring threatened keystone species. https://doi.org/10.1371/journal.pone.0205578

Pollock, K. H. (2018). Modeling Capture, Recapture and Removal Statistics for Estimation of

Demographic Parameters From Fish and Wildlife Populations: Past, Present and Future.

American Statistical Association, *86*(413), 225–238.

Rees, H. C., Maddison, B. C., Middleditch, D. J., Patmore, J. R. M., & Gough, K. C. (2014).

REVIEW: The detection of aquatic animal species using environmental DNA - a review of

eDNA as a survey tool in ecology. *Journal of Applied Ecology*, *51*(5), 1450–1459.

https://doi.org/10.1111/1365-2664.12306

Sigler, M. F., Womble, J. N., & Vollenweider, J. J. (2004). Availability to Steller sea lions (

Eumetopias jubatus) of a seasonal prey resource: a prespawning aggregation of eulachon (

Thaleichthys pacificus). *Canadian Journal of Fisheries and Aquatic Sciences*, *61*(8), 1475–

1484. https://doi.org/10.1139/f04-086

Signorell, A. (2019). *DescTools: Tools for description statistics*.

Sowa, J. (ADFG). (2015). *Hydrologic Investigations in Support of Reservations of Water for the*

Lost River , Alaska by.

Spangler, E. A. K. (2002). *The Ecology of Eulachon (Thaleichthys pacificus) in Twentymile*
River, Alaska. Univeristy of Alaska Fairbanks.

Takahara, T., Minamoto, T., Yamanaka, H., Doi, H., & Kawabata, Z. (2012). Estimation of Fish
Biomass Using Environmental DNA. *PLoS ONE*, *7*(4), e35868.
https://doi.org/10.1371/journal.pone.0035868

Team, Rs. (2015). *RStudio: Integrated Development for R*. Retrieved from www.rstudio.com

Tillotson, M. D., Kelly, R. P., Duda, J. J., Hoy, M., Kralj, J., & Quinn, T. P. (2018).
Concentrations of environmental DNA (eDNA) reflect spawning salmon abundance at fine
spatial and temporal scales. https://doi.org/10.1016/j.biocon.2018.01.030

Turnipseed, D. P., & Sauer, V. B. (2010). *Discharge Measurements at Gaging Stations*.
Retrieved from http://pubs.usgs.gov/tm/tm3-a8/

Wilcox, T. M., McKelvey, K. S., Young, M. K., Sepulveda, A. J., Shepard, B. B., Jane, S. F., …
Schwartz, M. K. (2016). Understanding environmental DNA detection probabilities: A case
study using a stream-dwelling char Salvelinus fontinalis. *Biological Conservation*, *194*,
209–216. https://doi.org/10.1016/j.biocon.2015.12.023

Wilken, U. (2018). Lakes, Labs and Learning: How an Environmental DNA Citizen Science
Project... *K-12 STEM Education*, *4*(4), 391–399. Retrieved from
https://www.learntechlib.org/p/209557/

Yates, M., Fraser, D., & Derry, A. (2019). Meta-analysis supports further refinement of eDNA
for monitoring aquatic species-specific abundance in nature. *Environmental DNA*, *1*, 5–13.
https://doi.org/10.1002/edn3.7

Data Accessibility Statement

498 All data and code are made available in the supporting information.

Author Contributions

501 TH and MP organized the eulachon mark-recapture program. TL initiated the eDNA monitoring
502 program based on discussion with DY and SM. TL, MP, JA, collected eDNA samples that were
503 processed in the lab by JA and MP. JA developed primers. MP and TL analyzed data and wrote
504 the paper with input and edits from other authors.

Tables

Table 1. Annual eulachon population estimates for the Chilkoot River using a modified Lincoln-Peterson equation (excluding 2013). eDNA monitoring began in 2014.

Measurement	2010	2011	2012	2014	2015	2016	2017	2018	2019
M = Marked Initially-adipose clipped	8,017	49,814	27,525	24,084	306	9,384	33,681	30,542	70,127
C = Total in second sample captured above weir	20,210	143,444	48,376	19,886	3,122	8,865	47,654	18,636	80,859
R = Marked recaptures above weir with clip	72	568	186	140	2	45	126	64	210
N1 = Population Estimate	2.2 Million	12.6 Million	7.1 Million	3.4 Million	319,586	1.8 Million	12.6 Million	8.7 Million	26.7 Million
SE^2 = Standard Error	256,415	521,961	516,583	283,226	158,934	262,518	1,113,520	1,074,932	1,840,573
CI^3 = 95% Confidence Interval	1.7 to 2.7 Million	11.5 to 13.6 Million	6.1 to 8.1 Million	2.9 to 3.9 Million	8,074 to 631,098	2.3 to 1.3 Million	10.5 to 14.8 Million	6.6 to 10.9 Million	23.2 to 30.4 Million

Table 2: Annual field effort for the mark-recapture study and number of eDNA sample days, water samples, and ddPCR replicates.

Year	Start Date	End Date	Number of Mark-Recapture field days	Number of eDNA sample days	Number of water samples	Number of ddPCR replicates
2010	4/23/2010	4/27/2010	5	*NA*	*NA*	*NA*
2011	4/27/2011	5/8/2011	12	*NA*	*NA*	*NA*
2012	5/2/2012	5/7/2012	6	*NA*	*NA*	*NA*
2014	5/5/2014	5/9/2014	5	8	24	48
2015	4/26/2015	4/29/2015	4	11	33	66
2016	4/20/2016	4/24/2016	4	19	57	114
2017	4/28/2017	5/5/2017	8	13	39	78
2018	5/6/2018	5/11/2018	6	17	51	102
2019	4/25/2019	5/7/2019	13	25	75	150

Figure 1. The Chilkoot River (A) is located in northern Southeast Alaska. The study area is located in the approximately 1.75 km section of river between the outlet of Chilkoot Lake and where the river meets the estuary (B). Three 1L water samples were collected at low tide just below the initial capture location, where (C) the adipose fin of eulachon captured in the trap was clipped into a "shark fin" for easy identification. The beginning of the recapture reach is located approximately 0.75 km upstream of the initial capture location, although eulachon are not exclusively recaptured within this reach. (D) Crews dip net or cast net to capture eulachon. (E) Depicts the Chilkoot River during a large eulachon run.

533 **Figure 2** Results of mark-recapture population estimate for eulachon on the Chilkoot River using
534 a modified Lincoln-Petersen method. Error bars represent one standard error.
535

Figure 3. Daily results of the Chilkoot River flow-corrected eDNA rate (copies/nl * discharge in cubic-feet/sec.) in 2014-2016 (A), 2017-2019 (B). The boxplots illustrate the variability among the three daily water samples, each quantified in two ddPCR replicates. The lower and upper portions of the box correspond to the 25th and 75th quartile respectively around the median (line). Whiskers extend to the most extreme data points up to 1.5 times the interquartile range. Variability increased with the mean such that boxplots at low flow-corrected eDNA rates are very small and appear as points.

Figure 4. Results of quasipoisson regression models relating log-transformed mark-recapture population estimate to (A) the size of the peak flow-corrected eDNA rate ($p = 0.027$, 84.96% Deviance explained), (B) the area under the curve of the flow-corrected eDNA rate ($p = 0.005$, 92.53% Deviance explained), (C) the size of the peak uncorrected eDNA concentration ($p = 0.01$, 89.53% Deviance explained), and (D) the area under the curve of the uncorrected eDNA concentration ($p = 0.006$, 90.74% Deviance explained). Gray shading denotes 95% confidence interval.

555 **Figure 5**. Mark recapture population estimate of eulachon runs (black dots) from 2014 to 2019
556 and the predicted number of eulachon based on the peak (A) or area under the curve (B) of the
557 flow-corrected eDNA rate and the peak (C) and area under the curve (D) of the non-flow
558 corrected eDNA concentration in the quasipoisson regression model (blue dashed lines). Gray
559 shading denotes the 95% confidence interval.