Evolution of conspicuousness in defended species involved in Müllerian mimicry

Ombeline Sculfort[1,2], Thomas G. Aubier[3,4], Ludovic Maisonneuve[1], Marianne Elias[1],

Violaine Llaurens[1]

[1] Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, CNRS, Sorbonne-Université, EPHE, Université des Antilles, 45 rue Buffon, 75005 Paris, France

[2] Molécules de Communication et Adaptations des Micro-organismes (MCAM), Muséum National d'Histoire Naturelle, CNRS, 63 rue Buffon, 75005 Paris, France

[3] Centre d'Ecologie Fonctionnelle et Evolutive, CNRS, Université de Montpellier, Université Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France

[4] Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

Corresponding author: Ombeline Sculfort, ombelinesculfort@hotmail.fr

Keywords: aposematism, chemical defences, community, mimicry, signalling, toxins, unpalatability

Abstract

Variation in warning pattern resulting in different levels of conspicuousness is observed

within and among chemically-defended prey species. Toxin acquisition and predator

avoidance behaviour have been shown to affect the evolutionary processes underlying such

warning signal diversity. Predators learn to associate warning colour pattern with

unpalatability and this results in a decrease in predation risk for individuals displaying a

locally abundant warning pattern. This positive number-dependent selection generated by

predators causes evolutionary convergence across defended prey species toward a common

warning colour pattern (Müllerian mimicry). Müllerian mimicry has a strong effect on the

local diversity of warning patterns in tight connexion with the local abundance of mimetic

species. Moreover, because chemical defences often stem from diet, the levels of defences

might depend on resource specialisation, therefore impacting species abundance and the

evolution of aposematic colour patterns.

To investigate the effect of Müllerian mimicry on the evolution of the conspicuousness of

warning signal and unpalatability, we built a single-species model where warning signal and

unpalatability can evolve, assuming that (1) predation risk depends on both the number of

individuals sharing a warning signal within the studied species and the local abundance of

mimetic species and (2) the level of unpalatability depends on resource specialisation,

implying that specialists suffer from reduced carrying capacity. We found that Müllerian

mimicry can promote the evolution of more conspicuous colouration in cryptic species.

Nevertheless, by favouring locally abundant signals, Müllerian mimicry stabilizes the

persistence of mildly conspicuous patterns, explaining for instance the evolution of partially

transparent colour patterns in some chemically-defended butterflies. Our model also show

that Müllerian mimicry limits the emergence of new aposematic signals associated with

reduced unpalatability. Altogether, our theoretical approach highlights complex interplays

between resource specialisation, mimicry and predator behaviour on the evolution of

aposematic signals and associated defences.

Impact summary

Aposematism, whereby chemically-defended species display conspicuous colouration acting as warning signal is still a debated question, involves the joint evolution of two independent traits, namely colour pattern and chemical defences. Evolution of new conspicuous warning signal is an evolutionary puzzle, because individuals exhibiting increased conspicuousness suffer from increased detection by predators. Convergence in warning pattern is frequently observed among defended prey (*i.e.* Müllerian mimicry), because sharing a common warning signal decreases individual predation risk. Mimicry among defended prey species may therefore profoundly influence the evolution of warning signals conspicuousness and chemical defences both within and among prey species.

The origin of variations in chemical defences is still largely unknown for a wide range of prey species, but toxic compounds are frequently sequestered from food intake so that specialized species may be more likely to successfully accumulate toxins from their diet. In turn, such specialization may limit species abundance locally, because of restricted range of resource and have a profound importance on the dynamics of aposematic signals. Understanding the evolution of aposematism therefore requires integrating ecological parameters linked to the evolution of defence levels and species abundance.

Here we use a theoretical approach to investigate the evolution of colour pattern conspicuousness in a chemically-defended species, involved in Müllerian mimicry. We explore the impact of ecological parameters, such as composition of local mimetic communities and diet specialisation on the evolution of aposematic signal and chemical

66   defences. We show that (i) mimetic interactions are a key factor enhancing fixation of new

67   patterns with increased conspicuousness (ii) even mildly conspicuous signals can persist in

68   species involved in Müllerian mimicry and that (iii) the evolution of chemical defences in

69   Müllerian mimics tightly depends on the resource acquisition pathway.

INTRODUCTION

The evolution of warning colour patterns in chemically-defended species (*i. e.* aposematism) has received considerable attention. Aposematism evolution relies on the interplay of three main factors, namely levels of defence, warning colouration, and abundance. Predators can learn to associate a warning colour pattern with unpalatability and subsequently avoid any individual displaying a known warning colour pattern. Predator's' learning generates number-dependent selection favouring the most abundant warning pattern (Chouteau *et al.*, 2016). Predators' learning depends on their negative experience associated with attacking a chemically-defended prey, linked to the level of distastefulness of that prey (Alcock, 1970a, 1970b; Goodale & Sneddon, 1977). The origins of unpalatability in defended prey are still largely unknown in many defended species, but in aposematic butterflies, one of the best-known examples of aposematic organisms, toxic compounds are frequently synthesized and/or sequestered from diet (Edgar, 1983; Engler-Chaouat & Gilbert, 2006; Zagrobelny *et al.*, 2014; de Castro *et al.*, 2017), resulting in variations in the level of defences both within species relying on the same diet, and among species relying on drastically different resources. Specialisation on a given source of toxins might enhance toxin sequestering capacity. For instance, specialist butterfly species *Heliconius sara* and *H. sapho* that sequester toxins from their host-plant *Passiflora auriculata* and *P. pittieri*, respectively, were shown to contain drastically smaller amounts of toxins when reared on other *Passiflora* host-plants, highlighting their increased sequestration capacities linked to host-plant specialisation (Engler-Chaouat & Gilbert, 2006). Nevertheless, compared to generalist species, specialist species have a reduced resource range, which may limit their abundance.

By influencing jointly unpalatability and abundance, specialisation on toxin resources may

play a complex role in the evolution of warning colour patterns.

Although higher conspicuous entails higher detectability, it has been shown that

when detected conspicuous colour patterns are be less attacked than cryptic patterns often

displayed by profitable prey because of innate aversion (Sillén-Tullberg, 1985; Lindström *et*

al., 2001), and might also trigger more rapid avoidance-learning due to higher memorability

(Roper & Redston, 1987; Ham *et al.*, 2006; Siddall & Marples, 2008). Unpalatability level can

impact the evolution of warning colour patterns and *vice versa*. In particular, theoretical

models showed that such coevolution should lead to negative or positive correlations

between levels of defence and warning pattern conspicuousness. Some models predict a

positive correlation between signal conspicuousness and unpalatability (Speed & Ruxton,

2005a; Broom *et al.*, 2006, 2008) because warning patterns can act as "honest" signals of

unpalatability, whereby only well-defended individuals display conspicuous pattern

associated with elevated attack risk or other fitness costs. Other models predict instead a

negative correlation, when defences induce immediate predator deterrence even in the

absence of costly conspicuousness (Leimar *et al.*, 1986; Speed & Ruxton, 2005b). Depending

on the availability of toxin resources and level of competition, positive and negative

correlations between signal conspicuousness and unpalatability can also be expected (Speed

& Ruxton, 2007; Blount *et al.*, 2009), so that the directions of such correlation within and

across species might differ (Speed, Ruxton, *et al.*, 2010). In empirical studies, positive

correlations between conspicuousness and chemical defences have been detected across

populations of the Asian ladybird *Harmonia axyridis* (Bezzerides *et al.*, 2007; María Arenas *et*

al., 2015), the poison frog *Dendrobates pumilio* (Maan & Cummings, 2012) and the paper

wasp *Polistes dominula* (Vidal-Cordero *et al.*, 2012) or across species of Dendrobatidae frogs

(Summers & Clough, 2001; Santos & Cannatella, 2011), ladybirds (María Arenas *et al.*, 2015)

and Opisthobranchia (marine gastropods) species (Cortesi & Cheney, 2010). However

empirical investigations across species have also revealed negative correlation between

conspicuousness and unpalatability, across poison frog *Epipedobates* species (Darst &

Cummings, 2006), Nudibranch (marine gastropods) species (Winters *et al.*, 2018) and burnet

moth (Briolat *et al.*, 2019).

In many cases, warning patterns are shared across defended species living in

sympatry (Mallet & Gilbert, 1995): because of the positive number-dependent selection

acting on colour patterns, evolutionary convergence of warning colour pattern among

species within the same locality is favoured, a phenomenon described as Müllerian mimicry

(Müller, 1879; Guilford, 1988; Sherratt, 2008; Joron, 2009). Mimetic interactions with other

defended species can influence the evolution of warning patterns and unpalatability .For

instance, defence levels in *Limenitis archippus* 'viceroy' butterfly tend to increase when the

co-mimetic species *Danaus gilippus* is less abundant (Prudic *et al.*, 2019). In this case, the

observed variations of unpalatability are independent from variations in host-plant toxic

compounds, suggesting direct selection on unpalatability, linked to the decreased

abundance of co-mimetic species. Warning patterns may also evolve in response to

variations in unpalatability displayed by co-mimics: Franks et al. (2009) showed theoretically

that the evolution of aposematic signals can be promoted in a defended species, if the

abundance of a palatable, parasitic mimetic species (*i. e.* Batesian mimics) is high. Different mimicry patterns may vary in their conspicuousness, resulting in contrasted warning colour patterns observed even between closely related species. A striking example is the neotropical butterfly tribe Ithominii, whereby some species exhibit brightly coloured wing patterns while others harbour transparent wings patterns (Beccaloni, 1997; Arias *et al.*, 2019). The evolution of new colour patterns with a different level of conspicuousness is not straightforward, as individuals with new colour patterns may suffer from reduced fitness. In addition to fitness reduction incurred by potential metabolic cost of pigmentation (Hill, 1996; McGraw *et al.*, 2005) and possible increased attack rates stemming from higher detectability for more conspicuous patterns or from higher resemblance to profitable prey for more cryptic patterns (Arias. et. 2019), the initially low abundance of new patterns limits number-dependent advantages linked to predation avoidance learning (Greenwood *et al.*, 1989; Lindstrom *et al.*, 2001). However, when the local community contains multiple mimetic species, including species with different colour patterns as often observed in aposematic butterfly communities (*e. g.*, Willmott et al. 2017), the emergence of new patterns within a species can be facilitated if it resembles a warning pattern already displayed by other defended species. Therefore, mimetic interactions between species need to be taken into account when investigating the evolution of conspicuousness in defended species.

Here we used a theoretical approach to investigate the evolution of conspicuousness and unpalatability as heritable traits, in a population of a defended species involved in Müllerian mimicry. In particular, we investigate whether the presence of a diverse

community of mimetic prey species can facilitate the evolution of new colour pattern, even

in the absence of direct deterrence provided by more conspicuous colour pattern (*i. e.*,

innate aversion against conspicuousness) and unpalatability. Contrary to other studies, in

our model prey do not survive an attack whatever its unpalatability level. We considered

that mutant warning patterns that differ in conspicuousness from the wild-type pattern can

either (i) be recognized as the same signal as the ancestral pattern or (ii) be considered as a

new warning signal, distinguished by predators from the ancestral pattern. In the latter case,

ancestral and emergent patterns could then match local mimicry rings (community of

individuals living in sympatry and sharing the same warning pattern) of various abundances,

and therefore benefit from enhanced protection. We assumed that conspicuousness

enhances not only detectability by predators but also signal memorability, and therefore

influences predator's learning process in associating a given warning colour pattern with

unpalatability. Finally, because unpalatability is linked to resource specialisation in some

prey species (*e. g.*, butterflies), we assume that increase in unpalatability stems from

specialisation towards a limited type of resources, thereby restricting population size. This

hypothesis contrasts with classical assumption assuming a physiological cost of

unpalatability that might limit investment in costly pigments. We also assumed no direct

effect of enhanced unpalatability on predator deterrence, so that the effect of unpalatability

only impacts individual protection because of predator learning.

MATERIAL AND METHODS

General model

Using ordinary differential equations, we model a single population of chemically-defended

species, initially composed of a number of wild-type individuals N_{wt}. Wild-type individuals

have a fixed unpalatability u_{wt} and display a warning colour pattern with a level of

conspicuousness c_{wt} and a basal predation rate p_{wt}, which depends on the abundance of co-

mimetic species (see table 1). We introduce a small number of mutant individuals N_m

harbouring either the same or a different warning colour pattern as the wild-types. Mutant

individuals may have different conspicuousness c_m, basal predation rate p_m or unpalatability

u_m as compared to wild-type (see table 1). We then track the evolution of the number of

wild-types and mutants through time.

The evolution of these numbers of wild-types and mutants through time follows the general

equations [1] and [2], respectively:

$\frac{dN_{wt}(t)}{dt} = \frac{dR_{wt}(t)}{dt} + \frac{dP_{wt}(t)}{dt}$ [1],

and $\frac{dN_m(t)}{dt} = \frac{dR_m(t)}{dt} + \frac{dP_m(t)}{dt}$ [2],

including a reproduction term given by $\frac{dR_{wt}(t)}{dt}$; $\frac{dR_m(t)}{dt}$ and a predation term given by

$\frac{dP_{wt}(t)}{dt}$; $\frac{dP_m(t)}{dt}$ described below.

Reproduction and resource specialisation

Wild-types and mutants belong to the same species. As such, they have equal growth rate

(r), compete for same resources and have the same carrying capacity (K) (see table 1).

We consider no physiological cost associated with unpalatability u, but we assume that

increased unpalatability (u_{wt} and u_m for wild-type and mutant respectively) stems from

higher resource specialisation, so that unpalatability affects the range of available resources

and thus modulates the carrying capacity (see equation [3] and [4]). We implement a

parameter α to specifically investigate the effect of resource dependence on unpalatability

acquisition in the species: when α=0, level of unpalatability is independent from resource

and unpalatability has no effect on carrying capacity; the higher α, the stronger the

dependence between unpalatability and resource specialisation and therefore the impact on

carrying capacity. Changes in numbers of wild-type and mutant individuals due to

reproduction (R_{wt} and R_m respectively) are thus described as follows:

$$\frac{dR_{wt}(t)}{dt} = r * N_{wt}(t)\left[1 - \frac{N_{wt}(t)+N_m(t)}{K\left(\frac{1}{1+u_{wt}*\alpha}\right)}\right] \qquad [3],$$

$$\frac{dR_m(t)}{dt} = r * N_m(t)\left[1 - \frac{N_{wt}(t)+N_m(t)}{K\left(\frac{1}{1+u_m*\alpha}\right)}\right] \qquad [4].$$

Predation and mimicry

We assume that the predation risk decreases as the number of individuals sharing the same

warning colour pattern in the population increases, as classically assumed in models of

Müllerian mimicry (e. g., Joron & Iwasa, 2005). In case of mimicry with other species, the

basal level of predation applied to the wild-type and the mutant (p_{wt} and p_m respectively)

depend on the abundances of their respective co-mimetic species. Predation risk faced by

wild-type and mutant individuals then depends on their warning colour pattern

conspicuousness c, the signal abundance and their unpalatability u.

Conspicuousness has two opposite effects on predator's behaviour, detectability and

memorability. Detectability is described as the direct effect of conspicuousness on predation

rate, assuming more conspicuous warning colour patterns are more likely to be detected and

attacked. Cryptic individuals (c=0) have the lowest level of detectability by predators (but

even those individuals are detectable to some extent). Memorability is then described as an

indirect effect on the number-dependent advantage associated with mimetic signals: we

assume that more conspicuous signals might be easier for predators to remember and

therefore enhance protection brought by mimicry, as in previous models of Sherratt, 2002

and Broom *et al.*, 2006. We introduce a parameter β that modulates the effects of

conspicuousness on detectability compared to memorability. Specifically, for a given

memorability effect, low values of β incur a low detectability, while high values incur a high

detectability. By investigating the effect of parameter β, we investigate the relative effect of

conspicuousness on the balance between detectability and memorability.

First, we investigate the evolution of conspicuousness assuming that variations in

conspicuousness are generalized by predators, so that wild-type and mutant individuals

share the same warning signal ('single signal hypothesis' hereafter), (equations [5] and [6]).

We thus assume that warning colour patterns exhibited by wild-type or mutant individuals

belong to the same mimicry ring (and therefore face the same basal predation rate, $p_{wt}=p_m$)

and benefit from the same number-dependent advantage, but that they can differ in their

conspicuousness level. Under the single signal hypothesis, changes in numbers of wild-type

and mutant individuals due to predation (P_{wt} and P_m respectively) are thus described as

follows:

$$\frac{dP_{wt}(t)}{dt} = \frac{-p_{wt}(1+\beta*c_{wt})N_{wt}(t)}{1+(u_{wt}*c_{wt}*N_{wt}(t)+u_m*c_m*N_m(t))} \qquad [5],$$

$$\frac{dP_m(t)}{dt} = \frac{-p_m(1+\beta*c_m)N_m(t)}{1+(u_{wt}*c_{wt}*N_{wt}(t)+u_m*c_m*N_m(t))}$$ [6].

Alternatively, variations in conspicuousness can result in warning signal recognized as

different by predators and thus do not share the cost of predator's learning process ('two-

signal hypothesis' hereafter) (equations [7] and [8]). Other mimetic species in the

community are not explicitly modelled, but we assume that differences of abundances

between mimicry rings result in different basal predation rates p_{wt} and p_m. Under the two-

signals hypothesis, changes in numbers of wild-type and mutant individuals due to predation

(P_{wt} and P_m respectively) are thus described as follows:

$$\frac{dP_{wt}(t)}{dt} = \frac{-p_{wt}(1+\beta*c_{wt})N_{wt}(t)}{1+(u_{wt}*c_{wt}*N_{wt}(t))}$$ [7],

$$\frac{dP_m(t)}{dt} = \frac{-p_m(1+\beta*c_m)N_m(t)}{1+(u_m*c_m*N_m(t))}$$ [8].

Testing the evolution of conspicuousness and unpalatability using simulations

We investigate the evolution of conspicuousness and unpalatability by running numerical

simulations in Python ("Python Release Python 3.6.5," 2018). We do not investigate co-

evolution pattern between conspicuousness and unpalatability, as always one out the two is

fixed and the other varies. We first model a population of wild-type individuals and let them

reach equilibrium, (i. e., a stable population size). Then, we introduce mutant individuals in

proportion of 0.01% of the wild-type. We assess whether the mutant colour pattern invades

the population or not, after 100,000 time steps. We specifically focus on four main

questions, investigating the evolution of conspicuousness and unpalatability, implementing

the same default values for global carrying capacity K=1000 and growth rate r=2 and

exploring a range of values for the other parameters, depending on the addressed question

(table 1). Notably, for all four questions investigated, we systematically test the effect of our

hypotheses regarding the link between unpalatability and resource specialisation (α) and the

detectability/memorability balance (β) on the evolutionary outcomes.

1- Evolution of conspicuousness within the same warning signal (single-signal)

Assuming the same fixed level of unpalatability in wild-type and mutant ($u_{wt}=u_m$), we study

the evolution of the level of conspicuousness of a given warning colour pattern, *i.e.* under

the single signal hypothesis, and assuming the same basal predation rate ($p_{wt}=p_m$=0.5). Wild-

type and mutant individuals thus have the same warning signal but differ in their

conspicuousness ($c_{wt}{\neq}c_m$).

2- Evolution of conspicuousness among different mimetic warning signals (two-signal)

Assuming again a fixed level of unpalatability ($u_{wt}=u_m$), we explore scenarios where

variations in conspicuousness evolution results in a distinct warning signal in mutants (*i. e.*

two-signal hypothesis). We consider a mutant whose warning colour pattern is recognized as

different from that exhibited by the wild-type, where initially rare mutant colour patterns do

not benefit from protection gained by resemblance with numerous wild-type individuals.

We consider that wild-type and mutant belong to two distinct mimicry rings, that can have

either similar (*i.e.* $p_{wt}=p_m$=0.5), or uneven abundances, with more abundant mimicry ring

resembling wild-type individuals (p_{wt}=0.5 and p_m=0.75) or mutant individuals (p_{wt}=0.5 and

p_m=0.25).

3- Evolution of unpalatability within the same warning signal

Then we investigate the evolution of unpalatability assuming that wild-type and mutant

individuals shared the same warning colour pattern (single signal hypothesis), therefore

following predation equation [5] and [6] and assuming p_{wt}=p_m and c_{wt}=c_m.

In these simulations, we thus assume that wild-type and mutant individuals can differ in

their level of unpalatability (u_{wt}≠u_m).

4- Evolution of unpalatability among different mimetic warning signal

Finally, we explore the evolution of unpalatability (u_{wt}≠u_m) assuming that wild-type and

mutant individuals display different warning colour patterns (two signals hypothesis), but

with similar basal predation rate and level of conspicuousness p_{wt}=p_m and c_{wt}=c_m.

RESULTS

1- Decrease in conspicuousness is favoured in populations displaying a single warning

signal

We explore the evolution of conspicuousness in a population of individuals sharing the same

warning colour pattern (single signal hypothesis). Overall, less conspicuous colour patterns

are advantaged in unpalatable populations (Fig.1A). Predators recognize wild-type and

mutant patterns as the same warning signal, so that mutants and wild-types both benefit

from the same number-dependent protection, brought by the strength in number. However,

less conspicuous individuals also benefit from reduced detection by predators and therefore

benefit from a low predation rate. Because the signal is generalized by predators whatever

the level of conspicuousness, conspicuousness of warning signal tends to decrease. By

modulating the effect of conspicuousness on detectability/memorability (β), we confirmed

that the selective advantage of decreased conspicuousness is due to reduced detection by

predators, limiting predation (Sup. fig. 1.).

For high levels of unpalatability, we observed that less conspicuous mutants are not

fixed after 100,000 generations when wild-type individuals are highly conspicuous (Fig. 1B).

By increasing the number of generations in our simulations, less conspicuous warning colour

patterns still invade the population (Sup. Fig. 2), confirming that the selection coefficient

associated with less conspicuous mutants is still positive. In that case, predation pressure on

wild-type individuals is very low because number-dependent advantage is increased by high

unpalatability; consequently, selection favouring less conspicuous colour patterns is weak

such that more time is needed to reach fixation.

When unpalatability has no cost, our model suggests that decrease in

conspicuousness might be favoured within a single warning colour pattern because more

cryptic individuals benefit from both reduced detectability and protection brought by

mimicry. As previously suggested by theoretical models (*e.g.* Puurtinen & Kaitala, 2006), our

study highlights that the effect of conspicuousness on detectability should be precisely

characterized to understand the evolution conspicuousness within warning signal.

2- Mimicry promotes the persistence of midly conspicuous warning colour pattern

We then investigate invasion probability of mutants harbouring a different colour pattern from that of the wild-type (two-signal hypothesis). In defended population, mutants displaying a new warning signal rarely invade (Fig. 2), even when less conspicuous than the wild-type signal, because wild-type individuals benefit from a stronger number-dependent selection. Nevertheless, more conspicuous signals can invade when wild-type individuals display a very cryptic signal (Fig. 2). Such invasion of a new signal with higher conspicuousness is favoured when the basal predation rate of mutants (p_m) decreases, *i. e.* when the new signal mimics warning patterns displayed by an abundant local community of defended species (Fig. 2). New signals with increased conspicuousness can be favoured by greater memorability; by diminishing the effect of conspicuousness on detectability while maintaining the same level of memorability (low β), we observe an increase in the invasion range of the more conspicuous mutant (Sup. fig. 3). When the effect of conspicuousness on detectability decreases, its positive effect on memorability further facilitates the invasion of an - initially rare - new conspicuous signal.

Our model therefore suggests that a more conspicuous signal matching a local mimetic community can be advantaged in a population of cryptic prey, but mostly fails to invade populations where wild-type individuals already display a mildly to strongly conspicuous signal. This might explain the persistence of signals with mild conspicuousness in Ithomiini butterflies, whereby many species harbour partly transparent mimetic colour patterns (Beccaloni, 1997; Willmott *et al.*, 2017) despite their high levels of chemical defences (Brown, 1984; Trigo *et al.*, 1996; Trigo, 2010). Transparency in Ithomiini results in a

decrease in detectability by avian predators (Arias *et al.*, 2019). But these mildly conspicuous

colour patterns might still be advantaged by their level of conspicuousness (providing both

low detectability but sufficient memorability) and the abundance of their mimicry ring.

3- Less unpalatable mutants are advantaged in population displaying a single warning

signal

We explored the evolution of unpalatability within a population where all individuals display

the same warning colour signal (single-signal hypothesis). When unpalatability stems from

increased resource specialisation ($\alpha \neq 0$), mutants with lower unpalatability invade the

population (Fig. 3A, C) because they share the same warning colour pattern with more

unpalatable wild-type, while bearing a lower cost due to resource specialisation. As a

consequence, when unpalatability becomes independent from resource specialisation ($\alpha = 0$),

mutant with lower unpalatability no longer invade (Fig. 3B, D). Less unpalatable mutants are

favoured as soon as unpalatability is linked to resource specialisation, regardless of

conspicuousness level (Sup. fig. 4). For instance, in aposematic butterflies, the origin of

unpalatability, sequestered and/or synthesized from host-plants, therefore impacts the

evolution of the level of chemical defences, stressing the need to consider not only

variations in chemical defences within mimicry rings but also the ecological pathway

enabling these variations.

4- Müllerian mimicry limits the emergence of new signal associated with reduced defence

Finally, we investigate the evolution of unpalatability when mutants and wild-types display

distinct warning signals (two-signal hypothesis) but similar level of conspicuousness. Here we

assume that level of unpalatability is positively linked to resource specialisation ($\alpha\neq0$), so

that more unpalatable individuals suffer from increased competition for resource. Mutants

with increased unpalatability can invade only in wild-type populations that are fully palatable

(Fig. 4A). Otherwise, only mutants with drastically lower level of unpalatability compared to

wild-type individuals can invade as (Fig. 4A). In that case, individuals displaying a new colour

pattern suffer from increased predation risk as compared to wild-type individuals, which

benefit from number-dependent advantage. Nevertheless, because high unpalatability

results in a reduction in abundance (Fig. 4C), mildly unpalatable mutant individuals can

quickly reach a higher abundance and replace highly unpalatable wild-type individuals (Fig.

4A). When relaxing the hypothesis linking unpalatability to resource specialisation (*i. e.*

assuming $\alpha=0$), mutants with decreased unpalatability no longer invade (Fig. 4B), confirming

the role of resource specialisation in the evolution of new warning signal associated with

decreased unpalatability. This trend is observed whatever the values of conspicuousness

(Sup. fig. 5).

Because resource specialisation is likely to reduce abundance, it has a strong impact

on the evolution of aposematic signal, enabling for instance signal associated with lower

unpalatability to settle in mimetic population with high unpalatability. This might explain

why generalist species, such as the mimetic butterfly *Heliconius erato*, can be found in large

abundances despite displaying a low level of toxins (Arias et al. 2016). This low unpalatability

level contrasts with the high unpalatability levels observed in closely related specialist

species like *H. sara* (Arias *et al.*, 2016), displaying a different signal and often living in

sympatry.

DISCUSSION

Evolution of conspicuousness depends on mimetic interactions

Overall, our model highlights that the evolution of increased conspicuousness is not possible

when assuming no direct effect of unpalatability or colour pattern conspicuousness on

predator deterrence. In this case, we showed that more conspicuous warning colour

patterns cannot evolve unless considering Müllerian mimicry. Within species harbouring a

single warning signal, protection brought by number-dependent selection facilitates the

emergence of less conspicuous warning colouration.

By contrast, our theoretical approach highlights that the emergence of more

conspicuous warning colour patterns can be favoured by mimetic interactions with local

community. When a diverse mimetic environment is considered and the mutant pattern

belongs to another mimicry ring in the community, we showed that more conspicuous

warning colour patterns can be favoured over cryptic patterns, but not over an already

conspicuous one. This suggests that mildly conspicuous signals can still provide efficient

protection against predation. This might explain the diversity of levels of conspicuousness of

warning colour pattern in closely-related species, such as transparent and opaque patterns

in Ithomiini butterflies (Elias *et al.*, 2008; Chazot *et al.*, 2014). Considering mimetic

community opens a new angle to explain the evolution of the wide diversity of colour

patterns: even if number-dependent selection favours the most abundant signals within

population, its effect can be modulated by variation in the conspicuousness of the different

signals.

Nevertheless, some cognitive biases in predators might favour the invasion of more

conspicuous warning colour patterns in unpalatable species, on top of mimicry: innate

memorability, direct deterrence, diet conservatism or neophobia have indeed been shown

to promote the emergence of new warning patterns (Marples & Kelly, 1999; Marples *et al.*,

2005; Aubier & Sherratt, 2015). Moreover, in our model, conspicuousness variations within a

given warning colour pattern do not affect the predator's generalization behaviour.

However, when difference of conspicuousness becomes very high, predators may be less

likely to generalize, leading to mutant individuals losing protection from the strength in

number.

Environmental variations and stochasticity could also favour the settlement of new

warning colour patterns in defended species. For instance, some colour pattern might

benefit from crypsis in a given habitat by matching the local background but habitat change

might then enhances the evolution of conspicuousness (Merilaita & Tullberg, 2005; Speed &

Ruxton, 2005a; Tullberg *et al.*, 2005; Speed, Brockhurst, *et al.*, 2010). Stochastic events,

including colonization of new environments, have been predicted to facilitate the

emergence of new colour patterns (Sherratt et al 2006; Mallet, 2010), and may allow more

conspicuous patterns to arise. Such emergence of conspicuous colour pattern can also be

enhanced by gregariousness, which generate an initial number-dependent advantage

(Leimar *et al.*, 1986).

The balance between detectability and memorability can modify the emergence of new mimicry pattern

Evolution of warning colour patterns relies on the evolution of conspicuousness: defended prey benefit from aposematism only if predators are able to associate the colour pattern with chemical defences. More specifically, our model highlights that memorability of warning signal has an important influence on predator's behaviour, as it reduces predation probability (Gittleman & Harvey, 1980; Ham *et al.*, 2006). Warning patterns might thus have a different evolutionary fate depending on signal memorability. In the mimicry literature, models exploring the emergence and persistence of multiple mimetic signals (Sherratt et al 2006, Mallet 2010) usually consider that all mimetic signals have the same predation risk, and that the attack rate then depends on local number-dependent selection. Here we highlight that the evolution of mimetic signals in defended species results from complex interactions between signal abundance and unpalatability but also on the detectability/memorability balance of the different mimetic colour patterns.

Unpalatability evolution is underpinned by resource specialisation.

We showed that selection favours individuals with lower unpalatability level, because of the link between unpalatability and resource specialisation assumed in our model. Within a single warning colour pattern, less unpalatable mutants strictly invade the population because they benefit from the same number-dependent selection as wild-types. By considering a diverse mimetic community, mutants with a colour pattern belonging to a

different mimicry ring lose this advantage, which inhibits the invasion of less unpalatable

mutants. In our model, as in some prey species such as butterflies, unpalatable metabolites

are either synthesized and/or sequestered from specialised resources (*e. g.*, butterfly larval

hostplants), and unpalatability is considered as an heritable trait linked to capacities to

survive and acquire toxins from defended resources. This assumption has important impact

on the evolution of unpalatability within and among mimetic signals. The origin of

unpalatability variations in conspicuous prey, their dependence to local resources and the

link between resource specialisation and prey carrying capacity thus needs to be empirically

investigated to improve our understanding of the evolution of mimetic warning patterns.

Furthermore, co-mimetic species sometimes share the same resources (Willmott & Mallet,

2004; Elias *et al.*, 2008; but see Alexandrou *et al.*, 2011), such that resource competition

might be more important within mimicry ring. If level of unpalatability is linked to resource

specialisation, the co-evolution between conspicuous mimetic signal and unpalatability

would therefore be shaped by both mutualistic and competitive interactions between co-

mimetic species.

DATA ACCESSIBILITY

AUTHOR'S CONTRIBUTIONS

V.L designed the study, T.G.A and L.M wrote the code, O.S ran simulations. O.S wrote the

manuscript with contributions from all authors. All authors participated in constructive

discussions and approved manuscript final version.

COMPETING INTERESTS

We declare no competing interests.

FUNDINGS

We acknowledge funding from LabexBCDiv Graduate Scholarship to O.S, and Paris city

council grant *Emergence* to VL.

ACKNOWLEDGMENTS

REFERENCES

ALCOCK, J. (1970a) Punishment levels and the response of black-capped chickadees (Parus atricapillus) to three kinds of artificial seeds. *Animal Behaviour*, **18**, 592–599.

ALCOCK, J. (1970b) Punishment levels and the response of white-throated sparrows (Zonotrichia albicollis) to three kinds of artificial models and mimics. *Animal Behaviour*, **18**, 733–739.

ALEXANDROU, M.A., OLIVEIRA, C., MAILLARD, M., MCGILL, R.A.R., NEWTON, J., CREER, S., & TAYLOR, M.I. (2011) Competition and phylogeny determine community structure in Müllerian co-mimics. *Nature*, **469**, 84–88.

ARIAS, M., MAPPES, J., DESBOIS, C., GORDON, S., MCCLURE, M., ELIAS, M., NOKELAINEN, O., & GOMEZ, D. (2019) Transparency reduces predator detection in mimetic clearwing butterflies. *Functional Ecology*, **0**.

ARIAS, M., MEICHANETZOGLOU, A., ELIAS, M., ROSSER, N., DE-SILVA, D.L., NAY, B., & LLAURENS, V. (2016) Variation in cyanogenic compounds concentration within a Heliconius butterfly community: does mimicry explain everything? *BMC Evolutionary Biology*, **16**, 272.

AUBIER, T.G. & SHERRATT, T.N. (2015) Diversity in Müllerian mimicry: The optimal predator sampling strategy explains both local and regional polymorphism in prey. *Evolution*, **69**, 2831–2845.

BECCALONI, G.W. (1997) Ecology, natural history and behaviour of Ithomiine butterflies and their mimics in Ecuador (Lepidoptera: Nymphalidae: Ithomiinae). *Tropical Lepidoptera Research*, **8**, 103–124.

BEZZERIDES, A.L., MCGRAW, K.J., PARKER, R.S., & HUSSEINI, J. (2007) Elytra color as a signal of chemical defense in the Asian ladybird beetle Harmonia axyridis. *Behav Ecol Sociobiol*, **61**, 1401–1408.

BLOUNT, J.D., SPEED, M.P., RUXTON, G.D., & STEPHENS, P.A. (2009) Warning displays may function as honest signals of toxicity. *Proceedings of the Royal Society B: Biological Sciences*, **276**, 871–877.

BRIOLAT, E.S., ZAGROBELNY, M., OLSEN, C.E., BLOUNT, J.D., & STEVENS, M. (2019) No evidence of quantitative signal honesty across species of aposematic burnet moths (Lepidoptera: Zygaenidae). *Journal of Evolutionary Biology*, **32**, 31–48.

BROOM, M., RUXTON, G.D., & SPEED, M.P. (2008) Evolutionarily stable investment in anti-predatory defences and aposematic signalling. *Mathematical modeling of biological systems, Volume II*. Springer, pp. 37–48.

BROOM, M., SPEED, M.P., & RUXTON, G.D. (2006) Evolutionarily stable defence and signalling of that defence. *Journal of Theoretical Biology*, **242**, 32–43.

BROWN, K.S. (1984) Adult-obtained pyrrolizidine alkaloids defend ithomiine butterflies against a spider predator. *Nature*, **309**, 707–709.

CHAZOT, N., WILLMOTT, K.R., SANTACRUZ ENDARA, P.G., TOPOROV, A., HILL, R.I., JIGGINS, C.D., ELIAS, M., SCHMITZ, A.E.O.J., & BRONSTEIN, E.J.L. (2014) Mutualistic Mimicry and Filtering by Altitude Shape the Structure of Andean Butterfly Communities. *The American Naturalist*, **183**, 26–39.

CHOUTEAU, M., ARIAS, M., & JORON, M. (2016) Warning signals are under positive frequency-dependent selection in nature. *PNAS*, **113**, 2164–2169.

CORTESI, F. & CHENEY, K.L. (2010) Conspicuousness is correlated with toxicity in marine opisthobranchs. *Journal of Evolutionary Biology*, **23**, 1509–1518.

DARST, C.R. & CUMMINGS, M.E. (2006) Predator learning favours mimicry of a less-toxic model in poison frogs. *Nature*, **440**, 208–211.

DE CASTRO, É.C.P., ZAGROBELNY, M., CARDOSO, M.Z., & BAK, S. (2017) The arms race between heliconiine
butterflies and *Passiflora* plants - new insights on an ancient subject: New insights on an
ancient subject. *Biological Reviews*, DOI: 10.1111/brv.12357.
EDGAR, J.A. (1983) Pyrrolizidine alkaloids in insect-plant co-evolution. *Toxicon*, **21**, 97–100.
ELIAS, M., GOMPERT, Z., JIGGINS, C., & WILLMOTT, K. (2008) Mutualistic Interactions Drive Ecological
Niche Convergence in a Diverse Butterfly Community. *PLOS Biology*, **6**, e300.
ENGLER-CHAOUAT, H. & GILBERT, L.E. (2006) De novo synthesis vs. sequestration: negatively correlated
metabolic traits and the evolution of host plant specialization in cyanogenic butterflies. *J
Chem Ecol*, **33**, 25–42.
GITTLEMAN, J.L. & HARVEY, P.H. (1980) Why are distasteful prey not cryptic? *Nature*, **286**, 149–150.
GOODALE, M.A. & SNEDDON, I. (1977) The effect of distastefulness of the model on the predation of
artificial batesian mimics. *Animal Behaviour*, **25**, 660–665.
GREENWOOD, J.J.D., COTTON, P.A., & WILSON, D.M. (1989) Frequency-dependent selection on
aposematic prey: some experiments. *Biological Journal of the Linnean Society*, **36**, 213–226.
GUILFORD, T. (1988) The Evolution of Conspicuous Coloration. *The American Naturalist*, **131**, S7–S21.
HAM, A.D., IHALAINEN, E., LINDSTRÖM, L., & MAPPES, J. (2006) Does colour matter? The importance of
colour in avoidance learning, memorability and generalisation. *Behav Ecol Sociobiol*, **60**, 482–
491.
HILL, G.E. (1996) Redness as a measure of the production cost of ornamental coloration. *Ethology
Ecology & Evolution*, **8**, 157–175.
JORON, M. (2009) Chapter 10 - Aposematic Coloration. *Encyclopedia of Insects (Second Edition)* (Resh,
V.H. & Cardé, R.T. eds). San Diego: Academic Press, pp. 33–38.
JORON, M. & IWASA, Y. (2005) The evolution of a Müllerian mimic in a spatially distributed community.
Journal of Theoretical Biology, **237**, 87–103.
JORON, M. & MALLET, J. (1998) Diversity in mimicry: paradox or paradigm? *Trends in Ecology &
Evolution*, **13**, 461–466.
LEIMAR, O., ENQUIST, M., & SILLEN-TULLBERG, B. (1986) Evolutionary Stability of Aposematic Coloration
and Prey Unprofitability: A Theoretical Analysis. *The American Naturalist*, **128**, 469–490.
LINDSTRÖM, L., ALATALO, R.V., LYYTINEN, A., & MAPPES, J. (2001) Predator experience on cryptic prey
affects the survival of conspicuous aposematic prey. *Proceedings of the Royal Society of
London. Series B: Biological Sciences*, **268**, 357–361.
LINDSTROM, L., ALATALO, R.V., LYYTINEN, A., & MAPPES, J. (2001) Strong antiapostatic selection against
novel rare aposematic prey. *Proceedings of the National Academy of Sciences*, **98**, 9181–
9184.
MAAN, M.E. & CUMMINGS, M.E. (2012) Poison Frog Colors Are Honest Signals of Toxicity, Particularly
for Bird Predators. *The American Naturalist*, **179**, E1–E14.
MALLET, J. (2010) Shift happens! Shifting balance and the evolution of diversity in warning colour and
mimicry. *Ecological Entomology*, **35**, 90–104.
MALLET, J. & GILBERT, L.E. (1995) Why are there so many mimicry rings? Correlations between habitat,
behaviour and mimicry in Heliconius butterflies. *Biological Journal of the Linnean Society*, **55**,
159–180.
MARÍA ARENAS, L., WALTER, D., & STEVENS, M. (2015) Signal honesty and predation risk among a closely
related group of aposematic species. *Scientific Reports*, **5**, 11021.
MARPLES, N.M. & KELLY, D.J. (1999) Neophobia and Dietary Conservatism:Two Distinct Processes?
Evolutionary Ecology, **13**, 641–653.
MARPLES, N.M., KELLY, D.J., THOMAS, R.J., & HARRISON, R. (2005) Perspective: the evolution of warning
coloration is not paradoxical. *Evolution*, **59**, 933–940.

MCGRAW, K.J., HILL, G.E., & PARKER, R.S. (2005) The physiological costs of being colourful: nutritional
control of carotenoid utilization in the American goldfinch, Carduelis tristis. *Animal*
Behaviour, **69**, 653–660.
MERILAITA, S. & TULLBERG, B.S. (2005) Constrained Camouflage Facilitates the Evolution of Conspicuous
Warning Coloration. *Evolution*, **59**, 38–45.
MÜLLER, F. (1879) Ituna and Thyridia; a remarkable case of mimicry in butterflies (transl. by Ralph
Meldola from the original German article in Kosmos, May 1879, 100). *Transactions of the*
Entomological Society of London.
PRUDIC, K.L., TIMMERMANN, B.N., PAPAJ, D.R., RITLAND, D.B., & OLIVER, J.C. (2019) Mimicry in viceroy
butterflies is dependent on abundance of the model queen butterfly. *Communications*
Biology, **2**, 68.
PUURTINEN, M. & KAITALA, V. (2006) Conditions for the spread of conspicuous warning signals: a
numerical model with novel insights. *Evolution*, **60**, 2246–2256.
Python   Release   Python   3.6.5   (2018)   *Python.org*.   URL
https://www.python.org/downloads/release/python-365/.
ROPER, T.J. & REDSTON, S. (1987) Conspicuousness of distasteful prey affects the strength and
durability of one-trial avoidance learning. *Animal Behaviour*, **35**, 739–747.
SANTOS, J.C. & CANNATELLA, D.C. (2011) Phenotypic integration emerges from aposematism and scale
in poison frogs. *PNAS*, **108**, 6175–6180.
SHERRATT, T.N. (2002) The coevolution of warning signals. *Proceedings of the Royal Society of London.*
Series B: Biological Sciences, **269**, 741–746.
SHERRATT, T.N. (2008) The evolution of Müllerian mimicry. *Naturwissenschaften*, **95**, 681.
SIDDALL, E.C. & MARPLES, N.M. (2008) Better to be bimodal: the interaction of color and odor on
learning and memory. *Behav Ecol*, **19**, 425–432.
SILLÉN-TULLBERG, B. (1985) Higher survival of an aposematic than of a cryptic form of a distasteful bug.
Oecologia, **67**, 411–415.
SPEED, M.P., BROCKHURST, M.A., & RUXTON, G.D. (2010) The Dual Benefits of Aposematism: Predator
Avoidance and Enhanced Resource Collection. *Evolution*, **64**, 1622–1633.
SPEED, M.P. & RUXTON, G.D. (2005a) Aposematism: what should our starting point be? *Proceedings of*
the Royal Society B: Biological Sciences, **272**, 431–438.
SPEED, M.P. & RUXTON, G.D. (2005b) Warning Displays in Spiny Animals: One (more) Evolutionary
Route to Aposematism. *Evolution*, **59**, 2499–2508.
SPEED, M.P. & RUXTON, G.D. (2007) How bright and how nasty: explaining diversity in warning signal
strength. *Evolution*, **61**, 623–635.
SPEED, M.P., RUXTON, G.D., BLOUNT, J.D., & STEPHENS, P.A. (2010) Diversification of honest signals in a
predator–prey system. *Ecology Letters*, **13**, 744–753.
SUMMERS, K. & CLOUGH, M.E. (2001) The evolution of coloration and toxicity in the poison frog family
(Dendrobatidae). *Proc Natl Acad Sci U S A*, **98**, 6227–6232.
TRIGO, J.R. (2010) Effects of pyrrolizidine alkaloids through different trophic levels. *Phytochem Rev*,
10, 83–98.
TRIGO, J.R., BROWN, K.S., WITTE, L., HARTMANN, T., ERNST, L., & BARATA, L.E.S. (1996) Pyrrolizidine
alkaloids: different acquisition and use patterns in Apocynaceae and Solanaceae feeding
ithomiine butterflies (Lepidoptera: Nymphalidae). *Biological Journal of the Linnean Society*,
58, 99–123.
TULLBERG, B.S., MERILAITA, S., & WIKLUND, C. (2005) Aposematism and crypsis combined as a result of
distance dependence: functional versatility of the colour pattern in the swallowtail butterfly
larva. *Proceedings of the Royal Society B: Biological Sciences*, **272**, 1315–1321.

VIDAL-CORDERO, J.M., MORENO-RUEDA, G., LÓPEZ-ORTA, A., MARFIL-DAZA, C., ROS-SANTAELLA, J.L., & ORTIZ-
SÁNCHEZ, F.J. (2012) Brighter-colored paper wasps (Polistes dominula) have larger poison
glands. *Frontiers in Zoology*, **9**, 20.
WILLMOTT, K.R. & MALLET, J. (2004) Correlations between adult mimicry and larval host plants in
ithomiine butterflies. *Proceedings of the Royal Society of London B: Biological Sciences*, **271**,
S266–S269.
WILLMOTT, K.R., ROBINSON WILLMOTT, J.C., ELIAS, M., & JIGGINS, C.D. (2017) Maintaining mimicry
diversity: optimal warning colour patterns differ among microhabitats in Amazonian
clearwing butterflies. *Proceedings of the Royal Society B: Biological Sciences*, **284**, 20170744.
WINTERS, A.E., WILSON, N.G., BERG, C.P. van den, HOW, M.J., ENDLER, J.A., MARSHALL, N.J., WHITE, A.M.,
GARSON, M.J., & CHENEY, K.L. (2018) Toxicity and taste: unequal chemical defences in a
mimicry ring. *Proc. R. Soc. B*, **285**, 20180457.
ZAGROBELNY, M., OLSEN, C.E., PENTZOLD, S., FÜRSTENBERG-HÄGG, J., JØRGENSEN, K., BAK, S., MØLLER, B.L., &
MOTAWIA, M.S. (2014) Sequestration, tissue distribution and developmental transmission of
cyanogenic glucosides in a specialist insect herbivore. *Insect Biochemistry and Molecular*
Biology, **44**, 44–53.
641

642   **TABLE LIST**

643   **Table 1**: Summary of parameters used in the model, with their biological meaning and
644   explored range values.

Parameter	Range explored	Description
$N_{wt}(t)$, $N_m(t)$	-	Number of individuals sharing wild-type or mutant warning colour pattern at time t
u_{wt}, u_m	[0;1]	Unpalatability associated with wild-type or mutant warning colour pattern
c_{wt}, c_m	[0;1]	Conspicuousness of wild-type or mutant warning colour pattern
p_{wt}, p_m	[0;1]	Basal predation rate exerted on wild-type or mutant warning colour pattern, depending on the abundance of their respective mimetic communities.
α	[0;1]	Effect of unpalatability on resource specialisation
β	[0;1]	Effect of conspicuousness on detectability
K	1000	Global carrying capacity
r	2	Growth rate

Figure 1: **Evolution of conspicuousness within the same mimicry ring; wild-type and mutant have the same warning colour pattern but with a different conspicuousness.**

Frequency of mutant and wild-type individuals after 100,000 generations, assuming different levels of conspicuousness c_{wt} of wild-type individuals (x-axis) and c_m of mutant individuals (y-axis). Colour scale indicates frequency of mutant individuals, with white representing 0% (100% wild-type individual*s*) and black being 100% (0% wild-type individuals). Red dashed line shows diagonal, where $c_{wt}=c_m$. Here the value of resource specialisation, α, do not influence the invasion pattern. Warning colour pattern are detectable, β=1, thus suffer greater predation. Predation functions followed equations [5] and [6], assuming basal predation rates $p_{wt}=p_m=0.5$. Toxicity values are fixed within each simulation, **A)** $u_{wt}=u_m=0.3$, **B)** $u_{wt}=u_m=1$.

Mutant mimicry ring abundance as compared to the wild-type one

Figure 2: **Evolution of conspicuousness among different mimicry rings; wild-type and mutant have a different warning colour pattern under various predation risks.** Frequency of mutant and wild-type individuals after 100,000 generations, assuming different levels of conspicuousness c_{wt} of wild-type individuals (x-axis) and c_m of mutant individuals (y-axis). The x-axis scale is cropped for visual comfort. Colour scale indicates frequency of mutant individuals, with white representing 0% (100% wild-type individual*s*) and black being 100% (0% wild-type individuals). Red dashed line shows diagonal where $c_{wt}=c_m$. Parameter modulating resource specialisation α=1 but variations in α, gives similar results on invasion pattern. Here, warning colour pattern is detectable, β=1. Unpalatability value is fixed within each simulation, here $u_{wt}=u_m=1$ (high unpalatability). Predation functions followed equations [7] and [8]. Various basal predation rates modelling mutant morph belonging to **A**) a rare mimicry ring ($p_{wt}=0.5$, $p_m=0.75$), **B**) a mimicry ring of same abundance as the wild-type morph ($p_{wt}=p_m=0.5$), and **C**) an abundant mimicry ring ($p_{wt}=0.5$, $p_m=0.25$).

Figure 3: Evolution of unpalatability within the same mimicry ring. Only resource specialisation controls invasion pattern and population size. Frequency of mutant and wild-type individuals after 100,000 generations, assuming different levels of unpalatability u_{wt} of wild-type individuals (x-axis) and u_m of mutant individuals (y-axis). Red and white dashed lines show diagonal where $u_{wt}=u_m$. **A-B)** Mutant proportion at equilibrium. Colour scale indicates proportion of mutant individuals fixed, with white representing 0% (100% wild-type individuals) and black representing 100% (0% wild-type individuals). **C-D)** Population size at equilibrium. Colour scale indicates number of individuals ($N_{wt}+N_m$), with dark blue

representing 0 (extinct population but never reached) and red representing 1000 (saturation of the carrying capacity K=1000, but never observed).

Here the effect of conspicuousness on detectable, β, has no effect. Conspicuousness value is fixed within each simulation, here $c_{wt}=c_m$=1, and same results are observed for every tested value. Predation functions following equations [5] and [6], assuming basal predation rates $p_{wt}=p_m$=0.5.

A-C) Individuals rely partly on resources to acquire unpalatable metabolites. Here α=1 but same invasion pattern is observed as long as α≠0. Individual can sequester and synthesize their unpalatable compounds. **B-D)** There is no link between unpalatability and resources, α=0. Individual can only synthesize their unpalatable compounds.

Resource link   **No resource link**

Unpalatability of the mutant

Unpalatability of the wild-type

Figure 4: Evolution of unpalatability between individuals displaying a different warning signal. Only resource specialisation controls invasion pattern and population size. Frequency of mutant and wild-type individuals after 100,000 generations, assuming different levels of unpalatability u_{wt} of wild-type individuals (x-axis) and u_m of mutant individuals (y-axis). Red and white dashed lines show diagonal where $u_{wt}=u_m$. **A-B)** Mutant proportion at equilibrium. Colour scale indicates proportion of mutant individuals fixed, with white representing 0% (100% wild-type individuals) and black representing 100% (0% wild-type individuals). Red arrows point out mutant invasion when the wild-type is palatable ($u_{wt}=0$), and these areas are zoomed for better visualisation. **C-D)** Population size at equilibrium. Colour scale indicates number of individuals ($N_{wt}+N_m$), with dark blue representing 0 (extinct population but never reached) and red representing 1000 (saturation of the carrying capacity $K=1000$, but never observed). Detectability ($\beta=0$ or $\beta=1$) do not influence invasion pattern. Conspicuousness value is fixed within each simulation, here $c_{wt}=c_m=1$, and similar results are observed for every tested value. Predation functions following equations [7] and [8]. We tested various predation risks modelling mutant morph belonging to a rare mimicry ring ($p_{wt}=0.5$, $p_m=0.75$), a mimicry ring of same abundance as the wild-type morph ($p_{wt}=p_m=0.5$), and an abundant mimicry ring ($p_{wt}=0.5$, $p_m=0.25$). Here we present results for $p_{wt}=p_m=0.5$, as other conditions are similar. Red arrows point out mutant invasion when the wild-type is palatable ($u_{wt}=0$).

A-C) Individuals rely partly on resource to acquire unpalatable metabolites, $\alpha=1$. Individual can sequester and synthesize their unpalatable compounds. **B-D)** There is no link between unpalatability and resource, $\alpha=0$. Individual can only synthesize their unpalatable compounds.