1 Time elapsed between Zika and dengue virus type 2 infections alters the magnitude of antibody
2 and T cell responses but not viremia in rhesus macaques
3

Erick X. Pérez-Guzmán[1#] , Petraleigh Pantoja[1,2] , Crisanta Serrano-Collazo[1] , Mariah A. Hassert[3],
Alexandra Ortiz-Rosa[4] , Idia V. Rodríguez[2] , Luis Giavedoni[5] , Vida Hodara[5] , Laura Parodi[5] ,
Lorna Cruz[1,2] , Teresa Arana[1,2] , Laura J. White[6] , Melween I. Martínez[1,2] , Daniela Weiskopf[7],
James D. Brien[3] , Aravinda de Silva[6] , Amelia K. Pinto[3] & Carlos A. Sariol[1,2,8]*

[1] Department of Microbiology and Medical Zoology, University of Puerto Rico-Medical Sciences
Campus, San Juan, Puerto Rico, United States of America. [2] Unit of Comparative Medicine,
Caribbean Primate Research Center and Animal Resources Center, University of Puerto Rico-
Medical Sciences Campus, San Juan, Puerto Rico, United States of America. [3] Department of
Molecular Microbiology & Immunology, Saint Louis University School of Medicine, Saint Louis,
Missouri, United States of America. [4] Department of Biology, University of Puerto Rico-Río
Piedras Campus, San Juan, Puerto Rico, United States of America. [5] Texas Biomedical Research
Institute, San Antonio, Texas, United States of America. [6] Departments of Microbiology &
Immunology, University of North Carolina-Chapel Hill, North Carolina, United States of America.
[7] Division of Vaccine Discovery, La Jolla Institute for Immunology, La Jolla, California, United
States of America. [8] Department of Internal Medicine, University of Puerto Rico-Medical Sciences
Campus, San Juan 00936, Puerto Rico, United States of America. [#] Current Address: Takeda
Vaccines Inc, Cambridge, Massachusetts, United States of America.

Short Title: Longevity of Zika immunity modulates dengue infection
Keywords: Zika, dengue, immunity, macaques

*Corresponding author: Carlos A. Sariol
carlos.sariol@upr.edu

Abstract

The role of a previous Zika virus (ZIKV) immunity on subsequent dengue virus (DENV) infections is poorly understood. This is relevant to anticipate the dynamics of forthcoming DENV epidemics in areas with previous ZIKV exposure. It is still uncertain if the immunity conferred by the recent ZIKV epidemic may contribute to protection or worsening DENV cases severity. Accordingly, we have studied the effect of ZIKV infection with various strains on subsequent DENV immune response after 10 and 2 months of ZIKV infection. Our results in non-human primates showed that a subsequent DENV infection in animals with early- and middle-convalescent periods to ZIKV do not promote an increase in DENV viremia nor pro-inflammatory status. We found that previous ZIKV exposure increases the magnitude of the antibody and cell-mediated immune responses against DENV and that the different time intervals between infections alter the magnitude and durability of such responses—more after longer ZIKV pre-exposure. Furthermore, our data suggest that the elicited immune modulation between both ZIKV-immune groups after DENV infection are more influenced by the time elapsed between ZIKV and DENV infections and the maturation of the cross-reactive immune memory, rather than a possible effect due to ZIKV strain variation. Collectively, we found no evidence of a detrimental effect of ZIKV immunity in a subsequent DENV infection regardless the period of time between infections tested on this work. This supports the implementation of ZIKV vaccines that could also boost immunity against future DENV epidemics.

Zika virus (ZIKV) is a re-emerging mosquito-borne *Flavivirus* that has captivated the
attention of the scientific community by its explosive spread in The Americas[1], and severe
neurological sequelae following infection[2-5]. ZIKV established itself in tropical and sub-tropical
regions that are endemic to other closely-related flaviviruses such as dengue virus (DENV). Both
viruses belong to the Flaviviridae family and are transmitted by *Aedes spp.* mosquitoes. DENV is
a global public health threat, having two-thirds of world's population at risk of infection, causing
~390 million infections annually[6,7]. DENV exists as four genetically similar but antigenically
different serotypes (DENV1-4)[8]. Exposure to one DENV serotype confers long-lived immunity
against a homotypic secondary infection. However, secondary infection with a heterologous
serotype of DENV is the major risk factor to induce severe DENV disease [9-11].

Due to the established history regarding the influence of cross-reactive immune
interactions in dictating disease outcomes during heterologous infection and the genetic and
consequently antigenic similarities between DENV and ZIKV, concerns have been raised
regarding the impact of DENV-ZIKV cross-reactive immunity on the development of severe clinical
manifestations[12,13]. In the last few years, multiple studies have aimed to understand the role of a
prior DENV exposure in the outcome of ZIKV infection. It has been demonstrated that DENV-
immune sera from humans can enhance ZIKV infection *in vitro*[14,15], and *in vivo* in immune-deficient
mouse models[16]. However, recent results from our group and others have shown that previous
flavivirus exposure—including DENV—may have no detrimental impact on ZIKV infection *in vivo*
in non-human primates (NHP)[17,18] and humans[19]. Moreover, these studies and others suggest
that previous DENV immunity may play a protective role during ZIKV infection involving humoral
and cellular responses[20-24]. On the other hand, little is known about the opposite scenario, the role
of a previous ZIKV exposure on a subsequent DENV infection, which is relevant to anticipate the
dynamics of forthcoming DENV epidemics.

The recent ZIKV epidemic in the Americas resulted in the development of a herd immunity
that may have an impact in subsequent infections with other actively circulating flaviviruses such
as DENV. Thus, human sub-populations such as newborns, international travelers from non-
flavivirus endemic areas or DENV-naïve subjects could be exposed to a ZIKV infection prior to
DENV—since DENV declined in the Americas during ZIKV epidemic[25]. After the epidemic, herd
immunity reduced ZIKV transmission and DENV will re-emerge and potentially infect these DENV-
naïve ZIKV-immune sub-populations in The Americas or potentially in other geographic areas
newly at risk[26,27]. An epidemiological study based on active DENV surveillance in Salvador, Brazil,
suggests that the reduction of DENV cases after the ZIKV epidemic is due to protection from
cross-reactive immune responses between these viruses[28]. Prospective experimental studies are

needed to confirm this hypothesis. For this purpose, we propose the use of NHPs as a suitable

model. NHPs provide advantages such as an immune response comparable to humans, and the

normalization of age, sex, injection route, viral inoculum and timing of infection[29]. Although clinical

manifestations by flaviviral infections are limited in NHPs[30], they have been widely used as an

advanced animal model for the study of DENV and ZIKV immune response, pathogenesis, and

vaccine development[17,18,29,31-34].

103   ZIKV antibodies (Abs) are capable of enhancing DENV infection *in vitro*[35].

Characterization of the specificity of DENV and ZIKV cross-reactive response revealed that ZIKV

monoclonal Abs and maternally acquired ZIKV Abs can increase DENV severity and viral burden

in immune-deficient mouse models[36,37]. However, little evidence is available concerning this

phenomenon occurring *in vivo* in immuno-competent large animal models such as NHPs. *George*

et al., showed that an early convalescence to ZIKV induced a significant higher peak of DENV

viremia and a pro-inflammatory profile compared to ZIKV-naïve status in rhesus macaques[38].

Further characterization of ZIKV early convalescent sera from these macaques indicated a

delayed induction of the cross-reactive Ab response against DENV, supporting no cross-

protection against the outcome of DENV infection[39]. A recent NHPs study showed that clinical

and laboratory parameters of ZIKV-immune animals were not associated with an enhancement

of DENV-2 infection. However, a higher peak of DENV-2 plasma RNAemia in ZIKV-immune

animals was observed compared to DENV-2 serum RNAemia loads in control animals, but the

use of different sample types may account for these differences[40]. Despite these findings, further

studies are needed to dissect the complementary role of the innate, humoral and cellular immune

response to mechanistically explain these findings. Particularly, there is no evidence of the

modulation and functionality of the T cell immune response in the ZIKV-DENV scenario. Available

studies rely upon pathogenesis and antibody studies, but there is no documented evidence as to

whether cell-mediated immunity (CMI)—specifically the functional response of T cells—is

modulated in a subsequent DENV infection by the presence of ZIKV immune memory.

123   The time interval between primary and secondary DENV infections have been shown to

be an important predictor for the development of severe clinical outcomes in humans[10]. Shorter

time interval between DENV infections result in a subclinical secondary infection, while

symptomatic secondary infections and severe DENV cases have been related with longer periods

between infections[41-44]. These findings suggest that high titers of cross-reactive Abs play a time-

dependent protective role between heterotypic DENV infections. Despite this evidence from

DENV sequential infections, it remains poorly understood if the same applies to the time interval

between ZIKV-DENV sequential infections. Specifically, how do longer periods of convalescence

after ZIKV infection impact the outcome of DENV infection. This scenario will more closely
resemble the epidemiological setting and time intervals elapsed between the current circulation
of related flaviviruses in the Americas. So currently, the role of multiple convalescent periods to
ZIKV in the outcome of DENV and other flavivirus infections is in the forefront of discussions
based on the limited studies available in experimental models and a lack of characterized human
prospective cohorts of this scenario yet[28,45-47].

To address these knowledge gaps, the objective of our study is to investigate the immune
modulatory role of an early- and middle-convalescence after ZIKV infection on the outcome of a
subsequent DENV infection in a NHP model. To test this, NHP cohorts who were ZIKV immune
for 10 months (mid-convalescence), 2 months (early-convalescence) or naïve for ZIKV were
exposed to DENV. The 2 months cohort was selected for direct comparison with previous work in
NHP[38], while the 10 months cohort was selected based on availability and to test a longer period
of convalescence to ZIKV. In each of these groups we assessed DENV pathogenesis, the elicited
Ab response, and characterized the CMI. Based on our knowledge, this is the first characterization
of CMI with this scenario in NHPs—taking into account the synergistic effect between the Ab and
cell-mediated responses. This study provides evidence that the presence of ZIKV immune
memory contributes to improve the immune response—more efficient after longer ZIKV pre-
exposure—against a DENV infection, without promoting enhancement of DENV viremia nor
inducing higher levels of pro-inflammatory cytokines.

Results

DENV challenge and clinical status of rhesus macaques. The experimental design includes
three cohorts of rhesus macaques (*Macaca mulatta*), within the age range considered as young
adults (Supplementary Fig. 1k), that were challenged with DENV-2 (NGC-44 strain), monitored
and bled for three months (Fig. 1). Two cohorts were previously exposed to ZIKV: cohort 1
(ZIKVPF-10mo) was comprised of 4 animals that had been exposed to ZIKV H/PF/2013 strain 10
months before DENV-2 challenge (mid-convalescence), and cohort 2 (ZIKVPR-2mo) comprised
of 6 animals that had been exposed to ZIKV PRVABC59 strain two months before DENV-2
challenge (early-convalescence). Both ZIKV strains used for previous exposure of these groups
are >99.99% comparable in amino acid identity (Supplementary Table 1). An additional cohort 3
(Naïve) included four animals naïve to ZIKV/DENV as a control group. After DENV challenge all
macaques were extensively monitored and sample collection was performed at various timepoints
up to 90 days post infection (dpi) for serum and PBMCs isolation.

The clinical status was monitored to determine if the presence of ZIKV immunity affected

the clinical outcome of DENV infection. Vital signs such as weight (kg), and temperature (°C) were

monitored. Also, complete blood cell counts (CBC), and comprehensive metabolic panel (CMP)

were performed before (baseline: day 0) and after DENV infection at multiple timepoints (CBC: 0,

7, 15 dpi; CMP: 0, 7, 15, 30 dpi). Neither symptomatic manifestations nor significant differences

in weight or temperature were observed in any of the animals after DENV infection up to 90 dpi

(Supplementary Fig. 1a-b). Likewise, no significant differences between groups were detected in

CBC parameters: white blood cells (WBC), lymphocytes (LYM), neutrophils (NEU), monocytes

(MON), and platelets (PLT) after DENV infection compared to basal levels of each group

(Supplementary Fig. 1c-g). CMP levels showed no differences in alkaline phosphatase and

aspartate transaminase (AST) (Supplementary Fig. 1h-i). Although within the normal range, levels

of alanine transaminase (ALT) were significantly higher in the ZIKVPR-2mo group compared to

its baseline at 7 dpi (p=0.0379, Two-way Anova Dunnett test), but at 15 and 30 dpi values returned

to baseline levels (Supplementary Fig. 1j). Overall, except for the isolated increase of ALT at 7

dpi in ZIKVPR-2mo, the clinical profile suggests that the presence of ZIKV-immunity did not

significantly influence the clinical outcome of DENV infection.

DENV RNAemia is not enhanced by previous ZIKV immunity. RNAemia levels in NHPs serum

were quantified by qRT-PCR at baseline, 1 to 10, and 15 dpi to determine if the presence of early-

(ZIKVPR-2mo) or mid-convalescence (ZIKVPF-10mo) to ZIKV alters DENV RNAemia kinetics.

No significant differences between groups were observed in detected levels of DENV genome

copies per ml of serum overtime (Fig. 2a). We noted that in the ZIKVPF-10mo group 3 out of 4

animals were able to keep the RNAemia level below 10^3 genome copies the next day after DENV

infection. This group started an early clearance of the RNAemia at 7 dpi, with only 1 out of 4

animals having detectable levels by days 8 and 9 pi. For ZIKVPR-2mo and naïve animals, the

clearance of detectable RNAemia started at 8 dpi, in 4 out of 6 and 1 out of 4 of the animals,

respectively. Naïve animals had the most delayed clearance of RNAemia with at least half of the

animals with detectable levels of viral RNA until day 9 pi. RNAemia was completely resolved in

all animals by 10 dpi. In summary, ZIKVPF-10mo had 7.25, ZIKVPR-2mo 7.5, and naïve animals

8 mean days of detectable RNAemia after DENV infection (Fig. 2b). In addition, the area under

the curve (AUC) was calculated but no statistically significance differences were observed in the

RNAemia peak among groups (Supplementary Fig. 2). However, the AUC trend to be lower in

both ZIKV-immune groups. In terms of the kinetics, a delay in the peak RNAemia set point was

observed in both ZIKV-immune groups (switch from day 2 to days 5 and 6) followed by higher,

but non-significant, levels compared to the naïve group, and a subsequent early RNAemia
clearance in both ZIKV-immune groups. Together these results show that, although no statistically
significant differences among groups were observed, previous immunity to ZIKV is not associated
with an increase in DENV RNAemia; even more, a mid-convalescence to ZIKV tended to develop
a shorter viremic period.

Pro-inflammatory cytokines are not exacerbated by the presence of ZIKV immunity. To
determine if the characterized cytokine profile of an acute DENV infection was modulated by ZIKV
immunity we assessed the serum concentration (pg/ml) of 8 cytokines/chemokines by Luminex
multiplex at baseline, 1, 2, 3, 5, 10, 15 and 30 dpi. The naïve group showed significant higher
levels of Type I interferon alpha (IFN-α) and pro-inflammatory cytokines such as Interleukin-6 (IL-
6), and monokine induced by IFN-gamma (MIG/CXCL9) (Fig. 3a-c). IFN-α was highest at 5 dpi
(Fig. 3a: $p<0.0001$ vs ZIKVPF-10mo and $p=0.0003$ vs ZIKVPR-2mo, Two-way Anova Tukey test).
IFN-α has been demonstrated to be involved in the innate anti-viral immunity and elevated levels
are associated with higher viral load and antigen availability. IL-6, a multifunctional cytokine
involved in immune response regulation and many inflammatory reactions showed the highest
levels at 1 dpi in naïve animals (Fig. 3b: $p=0.0115$ vs ZIKVPF-10mo and $p=0.0185$ vs ZIKVPR-
2mo, Two-way ANOVA Tukey test). Finally, MIG/CXCL9, which is a potent chemoattractant
involved in leucocyte trafficking demonstrated the highest levels at 10 dpi in naïve animals (Fig.
3c: $p=0.0004$ vs ZIKVPR-2mo, Two-way Anova Tukey test). On the other hand, the mid-
convalescent ZIKVPF-10mo group showed higher levels of CXCL10 (IP-10) (Fig. 3g) at day 1
($p=0.0198$ vs ZIKVPR-2mo, Two-way Anova Tukey test), 5 ($p=0.0487$ vs Naïve, Two-way Anova
Tukey test) and 10 pi ($p=0.0009$ vs ZIKVPR-2mo, Two-way Anova Tukey test). CXCL10 is a T
cell-activating chemokine and chemoattractant for many other immune cells. Also, this group
showed higher levels of perforin (Fig. 3h) at day 10 ($p=0.0024$ vs Naïve and $p=0.0190$ vs ZIKVPR-
2mo, Two-way Anova Tukey test) and 15 pi ($p=0.0178$ vs Naïve, Two-way Anova Tukey test).
Perforin is an effector cytolytic protein released by activated cytotoxic CD8+ T cells and natural
killer (NK) cells. No significant differences between groups were observed for other pro-
inflammatory citokines such as monocyte chemoattractant protein 1 (MCP-1), macrophage
inflammatory protein 1-beta (MIP-1β) and IL-1 receptor antagonist (IL-1RA) (Fig. 3d-f).
Collectively, these results demonstrate that the presence of ZIKV immunity does not exacerbate
pro-inflammatory status after DENV infection while mid-convalescence immunity to ZIKV
stimulated levels of mediators mainly involved in the activation of cell-mediated immune response.
231

DENV and ZIKV cross-reactive antibody response is boosted by ZIKV immunity and is influenced by the time span of the previous ZIKV infection. An ELISA-based serological profile was performed to determine the contribution of ZIKV immunity in the cross-reactive Ab response before and after DENV infection. We assessed the levels of DENV IgM and IgG, and cross-reactivity with ZIKV (IgM, IgG, NS1-IgG and EDIII-IgG) at multiple timepoints (Supplementary Fig. 3). Naïve cohort showed a significant higher peak of IgM (Supplementary Fig. 3a) characteristic of a primary DENV infection at 15 and 30 dpi ($p<0.0001$ vs ZIKVPF-10mo and $p=0.0004$ vs ZIKVPR-2mo, $p=0.0044$ vs ZIKVPF-10mo and $p=0.0179$ vs ZIKVPR-2mo, respectively, Two-way Anova Tukey test). This indicates the productive and acute DENV infection, while ZIKV immune groups showed lower levels of IgM resembling a heterotypic secondary infection. Total DENV IgG levels (Supplementary Fig. 3b) of both ZIKV-immune groups were significantly higher compared to naïve since baseline (cross-reactive ZIKV-IgG Abs) and 7, 15, 30, 60 and 90 (the latter for ZIKVPF-10mo only) (ZIKVPF-10mo vs Naïve: $p=0.0010$, $p<0.0001$, $p<0.0001$, $p<0.0001$, $p<0.0001$, $p=0.0016$; ZIKVPF-2mo vs Naïve: $p=0.0029$, $p=0.0002$, $p<0.0001$, $p<0.0001$, $p=0.0006$; Two-way Anova Tukey test). The ZIKVPF-10mo group showed significant higher levels than ZIKVPR-2mo group at 30 and 90 dpi ($p=0.0242$ and $p=0.0348$, Two-way Anova Tukey test). Overall, ZIKVPF-10mo developed higher and long-lasting levels of DENV IgG.

In contrast, ZIKV IgM levels were under or near the limit of detection in all groups over time after DENV infection despite several significant differences between groups (Supplementary Fig. 3c). ZIKV IgG levels (Supplementary Fig. 3d) were high in both ZIKV-immune groups at baseline and 7 dpi compared to naive ($p<0.0001$ vs naïve, Two-way Anova Tukey test), suggesting that although different pre-infecting ZIKV strains, the previous elicited IgG response against both ZIKV strains is comparable. After DENV infection, an increase of ZIKV IgG was shown and remain constantly high at 15, 30, 60 and 90 dpi in both ZIKV-immune groups ($p<0.0001$ vs naïve for all timepoints, Two-way Anova Tukey test), suggesting that DENV has the potential to stimulate ZIKV-binding Ab-producing plasmablasts. In addition, to elucidate the composition of similar ZIKV IgG levels in ZIKV-immune groups, we measured ZIKV-specific NS1 IgG (Supplementary Fig. 3e) and ZIKV-specific EDIII IgG (Supplementary Fig. 3f) levels. Although ZIKVPR-2mo showed significant differences compared to naïve at 30, 60 and 90 dpi ($p<0.0001$, $p=0.0001$, $p=0.0159$; Two-way Anova Tukey test), we observed a significantly higher expansion and long-lasting response of ZIKV NS1-specific Abs in the ZIKVPF-10mo group compared to the ZIKVPR-2mo group at baseline, 60 and 90 dpi ($p=0.0036$, $p=0.0071$, $p=0.0294$; Two-way Anova Tukey test) and also compared to naïve animals at all timepoints ($p<0.0001$, Two-way Anova

Tukey test). Moreover, higher magnitude of ZIKV-specific EDIII-IgG levels in the ZIKVPF-10mo

group than in the ZIKVPR-2mo group was observed compared to naïve at baseline (ZIKVPF-

10mo only), 15, 30 and 60 (ZIKVPF-10mo vs Naïve: $p=0.0092$, $p<0.0001$, $p<0.0001$, $p=0.0034$;

ZIKVPR-2mo vs Naïve: $p=0.0003$, $p=0.0014$, $p=0.0055$; Two-way Anova Tukey test), suggesting

that a ZIKV mid-convalescence promotes an expansion of higher magnitude of ZIKV EDIII-IgG

Abs from ZIKV memory B cells (MBC). However, those higher cross-reacting levels decrease

overtime as expected. In summary, a boost of DENV and ZIKV Abs is triggered by the presence

of ZIKV immunity and the expansion of specific- and cross-reactive Abs is higher on magnitude

and durability when a mid-convalescence immunity to ZIKV is present.

Neutralizing antibody response against DENV-2 and heterologous serotypes is higher in

magnitude and durability in presence of mid-convalescence to ZIKV. Neutralizing antibodies

(NAbs) are essential to combat DENV and ZIKV infection. The maturation and potency of this

response is known to define to a great extent the infection outcome[12,48]. Accordingly, we tested

the neutralization capacity of NAbs in serum from ZIKV-immune and naïve animals before and

after DENV infection, to determine whether an early- or mid-convalescence to ZIKV affected the

NAb response. Plaque Reduction Neutralization Test (PRNT) was performed to elucidate the NAb

titers of all groups against all DENV serotypes and both ZIKV pre-infecting strains. Before

infection with DENV the naïve groups had no detectable NAb levels (<1:20 PRNT60 titers) against

all DENV serotypes, while ZIKV-immune groups showed low cross-NAb titers against DENV-2

and DENV-4 (Fig. 4a). These cross-reactive levels were higher in the ZIKVPF-10mo group than

in the ZIKVPR-2mo group for both viruses. The peak of high NAb titers occurred at 30 days after

DENV infection for all groups (ZIKVPF-10mo>ZIKVPR-2mo>Naïve) against all DENV serotypes

(DENV-2>DENV-4>DENV-3>DENV-1) (Fig. 4b). The ZIKVPF-10mo group neutralized all DENV

serotypes with significant higher potency than naïve animals ($p<0.0001$, $p=0.0337$, $p<0.0001$,

$p<0.0001$ for DENV1-4; Two-way Anova Tukey test) and the ZIKVPR-2mo group, except for

DENV-2, that both ZIKV-immune groups have comparable neutralization magnitude at 30 dpi

($p=0.0002$, $p=0.7636$, $p=0.0016$, $p=0.0004$ for DENV1-4; Two-way Anova Tukey test). However,

the neutralization kinetics by sigmoidal response curves suggest higher percent of neutralization

against DENV-2 overtime in the group with mid-convalescence to ZIKV (Supplementary Fig. 4).

On the other hand, the ZIKVPR-2mo group showed significantly higher potency of the NAb

response only against DENV-1 compared to naive animals ($p=0.0146$; Two-way Anova Tukey

test) (Fig. 4b).

In addition, we tested whether the NAb titers that peak at 30 dpi for all groups remain
constant over time (up to 90 dpi) against all DENV serotypes (Fig. 4c-f). In general, the
neutralizing response of the ZIKVPF-10mo maintained higher NAb titers up to 90 dpi compared
to ZIKVPR-2mo and naïve groups. Significant differences between ZIKVPF-10mo and ZIKVPR-
2mo groups were observed against DENV-1,-3 and -4 at day 30 pi (p=0.0002, p=0.0016,
p=0.0004; Two-way Anova Tukey test) and at day 60 pi against DENV-2 and DENV-3 (p=0.0179,
p=0.0047; Two-way Anova Tukey test). The neutralizing Ab response of the ZIKVPF-10mo group
was even more significantly higher compared to the naïve group at day 15 (only performed for
the infecting serotype to monitor early neutralizing activity), day 30, 60 and 90 pi against DENV-
2 (p=0.0022, p=0.0337, p=0.0146, p=0.0337; Two-way Anova Tukey test); at day 30 pi against
DENV-1 (p<0.0001, Two-way Anova Tukey test); at day 30 and 60 pi against DENV-3 (p<0.0001,
Two-way Anova Tukey test); and at day 30 pi against DENV-4 (p<0.0001, Two-way Anova Tukey
test). In contrast, the ZIKVPR-2mo group showed a neutralizing Ab response with a magnitude
and long-lasting levels comparable to the naïve group, except at day 15 and 30 pi against DENV-
2 and DENV-1, respectively (p=0.0067, p=0.0146; Two-way Anova Tukey test). The neutralizing
response was long-lasting in the ZIKVPF-10mo group compared to other groups as supported by
the data from days 30 and 60 p.i. At day 90 pi, although no significant differences were observed
between all groups, the ZIKVPF-10mo group showed a consistent trend to maintain higher NAb
titers against all DENV serotypes indicating a higher and long-lasting breadth of cross-
neutralization within DENV serocomplex.

Collectively, these results demonstrate that a mid-convalescence to ZIKV provokes a
boost of the magnitude and durability of the neutralizing response against all DENV serotypes
more effectively than in animals with an early-convalescence to ZIKV, but also higher compared
to a *de novo* DENV-specific NAb response of the naïve animals.

ZIKV cross-neutralizing antibody response is strain-independent and higher in magnitude
and durability in the presence of mid-convalescence to ZIKV. Previous exposure to ZIKV
strains in ZIKV-immune groups developed high levels of cross-reactive, non-neutralizing, and
neutralizing Abs before DENV infection (baseline). To determine if this memory Ab response is
strain-specific and if the difference in convalescence period to ZIKV alters the efficacy and
modulation after DENV infection, we assessed the NAb levels in ZIKV-immune (ZIKVPF-10mo
and ZIKVPR-2mo) and ZIKV-naïve serum with both pre-infecting contemporary Asian-lineage
H/PF/2013 and PRVABC59 ZIKV strains at multiple timepoints after DENV infection. At baseline,
both ZIKV-immune groups showed high NAb titers against H/PF/2013 strain, which suggest that

irrespective of pre-exposure to different ZIKV strains and different convalescent periods the Ab

response remains similarly effective (Fig. 5a). As early as day 15 after DENV infection, a potent

boost of NAb titers in both ZIKV-immune groups was developed. However, elevated NAb titers

were significantly higher in the ZIKVPF-10mo group compared to the ZIKVPR-2mo and naïve

groups at day 15 pi (p=0.0005, p<0.0001; Two-way Anova Tukey test) and day 30 pi (p=0.0067,

p=0.0012; Two-way Anova Tukey test). As expected, this elevated ZIKV cross-reactive NAb

levels decreased gradually overtime after 15 dpi in both ZIKV-immune groups. Nevertheless, the

ZIKVPF-10mo group retained higher NAb titers until 90 dpi while the titers of the ZIKVPR-2mo

group returned to baseline levels. Of note, the NAb titers of the naïve group were considered as

negative in all timepoints and failed to neutralize ZIKV throughout DENV infection even at

concentrated levels of the antibodies (Fig. 5a). These results are confirmed by the behavior of

neutralization kinetics by sigmoidal response curves where the ZIKVPF-10mo group retained

elevated magnitude of ZIKV neutralization overtime (Supplementary Fig. 5).

346   To determine if the immune memory induced by different ZIKV strains play a role in the

modulation of the cross-NAb response triggered by a subsequent DENV infection, NAb titers were

measured against both ZIKV strains before and 30 days after DENV infection. The ZIKVPF-10mo

group showed significant higher NAb titers against both ZIKV strains compared to the ZIKVPR-

2mo group before DENV infection (p=0.0093, p=0.0141; Two-way Anova Tukey test) (Fig. 5b).

Subsequently, DENV infection promote an equally 8-fold increase of NAb titers against both

strains in the ZIKVPF-10mo group, significantly higher than the 4-fold increase in the ZIKVPR-

2mo group (p=0.0025, p=0.0011; Two-way Anova Tukey test) (Fig. 5c). To rule out that difference

in fitness between both ZIKV strains would bias the magnitude of the NAbs after DENV infection

we compared in parallel the NAb titers at 30 and 60 days after ZIKV infection (day 60 correspond

to the baseline of the ZIKVPR-2mo group). No significant differences were observed between

ZIKV-immune groups in the NAb titers induced by both strains at the same timepoints after ZIKV

infection (Supplementary Fig. 6). Altogether, these results demonstrate that DENV infection

results in a significant increase in the magnitude and durability of the cross-neutralizing Ab

response against ZIKV in animals with a mid-convalescent period from ZIKV infection. The elicited

changes in neutralization capacity were likely driven more by the longevity of the immune memory

maturation and the associated memory recall of the ZIKV immunity than by a strict dependency

of the specific pre-exposed ZIKV strain.

Frequency, early activation and proliferation of immune cell subsets are modulated by

ZIKV immunity. We performed immunophenotyping by flow cytometry to assess the frequency,

early activation and proliferation of multiple immune cell subsets and how these parameters are

affected by the presence of pre-existing immunity to ZIKV on a subsequent DENV infection

(Supplementary Fig. 7, 8, and 9 for gating strategy; Supplementary Table 2 for Ab panel). As part

of the innate immune response, the frequency of dendritic cells (DCs) and natural killer (NK) cells

subpopulations were measured. Plasmacytoid DCs (pDCs: Lin⁻HLA-DR⁺CD123⁺) are known to

respond to viral infection by production of IFN-α, while myeloid DCs (mDCs: Lin⁻HLA-DR⁺CD11c⁺)

interacts with T cells. The frequency of pDCs was not significantly altered by DENV infection in

any group compared to baseline levels (Supplementary Fig. 10a). At day 2 pi we detected a

significant increase of mDCs in the ZIKVPF-10mo group (p=0.0082; Two-way Anova Dunnett test)

(Supplementary Fig. 10b). Furthermore, we determined the frequency of NK subpopulations

including: NKCD8, NKCD56, NKp30 and NKp46 (Supplementary Fig. 11). In general, no

differences were detected between baseline and after DENV infection in all groups for all NK

subpopulations and receptor expression with the exception of the ZIKVPR-2mo group that

showed a significant increases in the following subpopulations: NKG2A⁺NKp30 and

NKp30⁺NKp46⁺ at 7 dpi (p=0.0495, p=0.0006; Two-way Anova Dunnett test) and NKp46⁺NKp30⁺

at 7 and 10 dpi (p=0.0005, p=0.0001; Two-way Anova Dunnett test) (Supplementary Fig. 11j, o,

s).

We next investigated cell subsets that are part of the bi-phasic (humoral/cellular) adaptive

immune response such as B (CD20+CD3-) and T (CD3+CD20-) cells. No differences were detected

in total B cells between groups following DENV infection compared to baseline levels (Supplementary

Fig. 12a), but ZIKV-immune groups had elevated levels of activated B cells (CD20+CD3-CD69+)

since baseline and a trend to increase these levels more than the naïve group overtime

(Supplementary Fig. 12b). We detected a significant decrease of proliferating B cells (CD20+CD3-

Ki67+) in naïve animals at 7 and 10 dpi (p=0.0031, p=0.0345; Two-way Anova Dunnett test), while

ZIKV-immune groups retained their proliferating levels (Supplementary Fig. 12c). Interestingly, the

ZIKVPF-10mo group showed a significant increase of B cells that were proliferating and activated

simultaneously (CD20+CD3-CD69+Ki67+) as early as in day 1 pi (p=0.0240; Two-way Anova Dunnett

test) and maintained higher levels up to 10 dpi (Supplementary Fig. 12d). Together, these

phenotyping results of B cells are consistent with the early and boosted production of binding and

neutralizing Abs in the ZIKVPF-10mo group compared to naïve animals. The frequency of total T cells

(CD3⁺CD20⁻) and CD4⁺/CD8⁺ T cells subsets, was comparable at all timepoints before and after

DENV infection in all groups of animals (Supplementary Fig. 13a-c).

Previous studies have demonstrated that DENV and ZIKV specific CD4⁺ and CD8⁺ T cells

are enriched in certain memory subsets[24,49]. Thus, we measured whether the early activation of

T cell subpopulations, such as effector memory (CD3$^+$CD4$^+$CD28$^-$CD95$^+$) and central memory (CD3$^+$CD4$^+$CD28$^+$CD95$^+$) T cells (T-EM and T-CM), within each T cell compartment was modulated following DENV infection in presence or absence of convalescence to ZIKV (Fig. 6, and Supplementary Fig. 7 for gating strategy). The ZIKVPF-10mo group showed significant higher frequency of activated CD4$^+$ and CD8$^+$ T-EM (CD3$^+$CD4$^+$CD28$^-$CD95$^+$CD69$^+$ and CD3$^+$CD8$^+$CD28$^-$CD95$^+$CD69$^+$) following DENV infection compared to basal levels (CD4$^+$ T-EM at 7 and 10 dpi: $p=0.0001$, $p=0.0072$; CD8$^+$ T-EM at 2 and 7 dpi: $p=0.0291$, $p=0.0001$; Two-way Anova Dunnett test) (Fig. 6a, d). Interestingly, the ZIKVPR-2mo group showed a very limited frequency and activation of the CD4$^+$ and CD8$^+$ T-EM compared to the ZIKVPF-10mo and naïve groups. However, this group with an early convalescent period to ZIKV, contrary to the other two groups, showed a very limited but significant activation of CD8$^+$ T-CM (CD3$^+$CD8$^+$CD28$^+$CD95$^+$CD69$^+$) at day 7 and 10 pi ($p=0.0007$, $p=0.0147$; Two-way Anova Dunnett test) (Fig. 6e). In contrast, naïve animals did not show any significant activation of these memory cell subsets after DENV infection. Collectively, these results suggest that following DENV infection: (i) animals with a mid-convalescence ZIKV immunity have a more dynamic B cell response and are able to rapidly produce more activated effector memory T cells from both T cell compartments; (ii) animals with an early-convalescence to ZIKV induced activation of central memory T cells in the CD8$^+$ compartment with a very limited T-EM frequency and activation profile compatible with a contraction phase of the T cells compartments; (iii) and animals without previous exposure to ZIKV exhibited a limited B cell response and minimal modulation of memory T cell subpopulations at early timepoints as the ZIKV-immune groups.

T cell functional effector response against DENV and ZIKV is shaped by the longevity of ZIKV-immunity. To further characterize the cross-reactive T cell response, we investigated if different convalescent periods of ZIKV immunity impacted the outcome of the effector role of CD4$^+$ and CD8$^+$ T cells following DENV infection. PBMCs were isolated and stimulated with peptide pools from DENV and ZIKV envelope (E) proteins and from ZIKV non-structural proteins (ZIKV-NS) (Supplementary Table 4 for peptide sequences). Then, intracellular cytokine staining using flow cytometry analysis (Supplementary Fig. 14 for gating strategy; Supplementary Table 3 for Ab panel) was performed to quantify the production of effector immune markers such as the cytotoxic marker CD107a, IFN-γ, and TNF-α by CD4$^+$ and CD8$^+$ T cells at baseline, 30, 60, and 90 days after DENV infection (Fig. 7).

To assess the ZIKV-primed specific- or cross-reactive effector T cell response we studied the response against ZIKV or DENV stimuli before DENV infection. In general, before DENV

infection, we found that the ZIKV-primed effector T cell response was higher in CD8$^+$ (Fig. 7m, q, u) than in CD4$^+$ (Fig 7a, e, i) T cells. Of note, significant higher levels of CD107a, INF-γ and TNF-α producing CD8$^+$ T cells were found only in the ZIKVPF-10mo group before DENV infection (ZIKVPF-10mo vs ZIKVPR-2mo for CD107a: *p*=0.0002; ZIKVPF-10mo vs Naïve for CD107a: *p*=0.0401; ZIKVPF-10mo vs ZIKVPR-2mo for INF-γ: *p*=0.0020; ZIKVPF-10mo vs ZIKVPR-2mo for TNF-α: *p*=0.0033; ZIKVPF-10mo vs Naïve for TNF-α: *p*=0.0354; Two-way Anova Tukey test) (Fig. 7m, q, u). This basal effector response of CD8$^+$ T cells in the ZIKVPF-10mo group is predominated by cross-reactive CD8$^+$ T cells against DENV E protein. Very low effector T cell response against ZIKV NS proteins was detected for all groups (ZIKVPF-10mo>ZIKVPR-2mo>Naïve). In summary, results of T cell functional response before DENV infection suggest that a mid-convalescence to ZIKV provoke a higher CD8$^+$ T cell effector response capable to cross-react efficiently with DENV E protein.

After DENV infection, we were able to determine the modulation of the ZIKV-primed effector CD4$^+$ and CD8$^+$ T cell responses of ZIKV-immune groups and the *de novo* response of ZIKV-naïve animals. The ZIKVPF-10mo and naïve groups significantly boosted their CD107a expression in both T cell compartments stimulated mainly by DENV E protein at 30 and up to 90 days after DENV infection (CD4$^+$ T cells: ZIKVPF-10mo vs ZIKVPR-2mo: *p*<0.0001 at 30 dpi, *p*<0.0001 at 60 dpi; Naïve vs ZIKVPR-2mo: *p*<0.0001 at 30 dpi, *p*=0.0018 at 60 dpi; ZIKVPF-10mo vs Naïve: *p*=0.0204 at 30 dpi. CD8$^+$ T cells: ZIKVPF-10mo vs ZIKVPR-2mo: *p*<0.0001 at 30 dpi, *p*<0.0001 at 60 dpi, *p*=0.0008 at 90 dpi; Naïve vs ZIKVPR-2mo: *p*=0.0039 at 30 dpi, *p*<0.0001 at 60 dpi; *p*=0.0081 at 90 dpi; ZIKVPF-10mo vs Naïve: *p*=0.0194 at 30 dpi; Two-way Anova Tukey test) (Fig. 7b, c, n, o, p). Also, these groups boosted the CD107a cytotoxic signature reacting against ZIKV E and NS proteins by cross-reactive CD4$^+$ T cells 30 days after DENV infection (ZIKVPF-10mo vs ZIKVPR-2mo: *p*=0.0025 for ZIKV E, *p*<0.0001 for ZIKV NS; Naïve vs ZIKVPR-2mo: *p*=0.0025 for ZIKV E, *p*=0.0002 for ZIKV NS; Two-way Anova Tukey test) (Fig. 7b).

The ZIKVPF-10mo group showed a remarkable significant increase of the IFN-γ-producing CD4$^+$ T cells against DENV E protein since 60 dpi and is maintained up to 90 dpi compared to other groups (ZIKVPF-10mo vs ZIKVPR-2mo at 60 and 90 dpi: *p*<0.0001, *p*=0.0024; ZIKVPF-10mo vs Naïve at 60 and 90 dpi: *p*<0.0001, *p*=0.0037; Two-way Anova Tukey test) (Fig. 7g, h), and was the only group with significant increase in the IFN-γ producing CD8$^+$ T cell compartment at 60 dpi (ZIKVPF-10mo vs ZIKVPR-2mo: *p*=0.0253; Two-way Anova Tukey test) (Fig. 7s). On the other hand, the ZIKVPR-2mo group exhibited a significant increase of IFN-γ producing CD4$^+$ T cells earlier than other groups at 30 dpi (ZIKVPR-2mo vs ZIKVPF-10mo: *p*<0.0001; ZIKVPR-2mo vs Naïve: Two-way Anova Tukey test) (Fig. 7f). Interestingly, the naïve

group showed an increase of cross-reactive TNF-α producing CD4$^+$ T cells against ZIKV NS proteins 30 days after DENV infection (Naïve vs ZIKVPR-2mo: p=0.0359; Two-way Anova Tukey test) (Fig. 7j). The ZIKVPF-10mo group developed a significant effector T cell response by TNF-α producing CD4$^+$ T cells against DENV and ZIKV E proteins at 60 days after DENV infection (ZIKVPF-10mo vs ZIKVPR-2mo against DENV/ZIKV E protein: p=0.0163, p=0.0172; Two-way Anova Tukey test) (Fig. 7k). Although all groups showed a boosted TNF-α effector response in the CD8$^+$ T cell compartment up to 90 days after DENV infection, no significant differences between groups were observed.

Collectively, these results after DENV infection suggest that a mid-convalescence to ZIKV translate in a more complete functional T cell response characterized by: (i) a cytotoxic CD107a$^+$ phenotype directed to DENV E protein for both T cell compartments comparable to the DENV-specific *de novo* response of the naïve group, (ii) developed CD107a, IFN-γ and TNF-α producing CD8$^+$ T cell effector response that cross-react efficiently with DENV E protein since baseline and is boosted after DENV infection, (iii) and promoted the higher T cell effector response against ZIKV NS proteins. An early-convalescence to ZIKV results in (iv) a very limited cytotoxic activity (limited expression of CD107a marker) which is in line with a very limited activation of the T-EM, and with failed capability to react efficiently against E or NS proteins. The ZIKV-naïve group response was characterized by: (v) production of a DENV-specific *de novo* functional T cell response with similar magnitude between both T cell compartments, (vi) capable to cross-react against ZIKV E and NS proteins, (vii) and able to mount a DENV-specific cytotoxic CD107a$^+$ phenotype.

Discussion

We found that previous infection to ZIKV modulates the immune response against subsequent DENV infection without an enhancement of DENV viremia nor pro-inflammatory status, and that this modulation is influenced by the longevity of ZIKV convalescence—more after longer ZIKV pre-exposure. The aftermath of the recent ZIKV epidemic has been related to a remarkable decrease in DENV cases in Brazil [28], and also in most of Latin American and Caribbean countries (http://www.paho.org/data/index.php/es/temas/ indicadores-dengue/dengue-nacional/9-dengue-pais-ano.html?start=2)[25]. Yet, little is known about the role of previous ZIKV immunity in the outcome of a subsequent DENV infection in human populations, and if ZIKV immunity is supporting this epidemiological phenomenon observed post-ZIKV epidemic[28]. To evaluate the hypothesis of a potential ZIKV-DENV cross-protection in humans characterizing the immunological history of prospective cohorts[47] will be necessary, but human

samples for this purpose are scarce yet. Because of this, NHPs are key to provide knowledge

and anticipate different immunological scenarios when DENV epidemics re-emerge in human

populations with previous immunity to ZIKV.

Animals with pre-existing ZIKV immunity do not show an enhancement of DENV-induced

RNAemia, regardless of the period of convalescence from previous ZIKV infection (10 or 2

months) and different pre-infecting ZIKV strains. Previous ZIKV immunity is associated with a

trend of less RNAemia days during subsequent DENV infection. This effect is more evident in

animals with a ZIKV convalescence period of 10 months. Previous work reported that a period of

early-convalescence (56 days) to ZIKV (PRVABC59 strain) in rhesus macaques was associated

with a significant increase of DENV-2 RNAemia at day 5 after DENV infection and a pro-

inflammatory cytokine profile. However, very similar to our results, it was noteworthy a delay at

early timepoints and an early clearance in late timepoints of the DENV-2 RNAemia in ZIKV-

immune macaques in comparison to the naïve ones[38]. The lack of significant DENV RNAemia

enhancement found in the group with the early-convalescence period in our work, compared to

previous results[38], may be attributable to the different sample types collected (plasma vs serum),

or different DENV-2 strains used for the challenge [New Guinea/1944 strain vs

Thailand/16681/1964 strain, from Asian II and Asian I Genotype, respectively]. This fact is of

relevance because it suggests that the effect of previous ZIKV immunity on a subsequent DENV

infection may differ between DENV serotypes or even within genotypes. Another possible

explanation is the genetic heterogeneity of rhesus macaques used in these two studies as they

are derived from different breeders. The importance of selecting genetic well-characterized

macaques have been discussed previously[50].

Due to limited availability of ZIKV-immune cohorts we used animals infected with two

different ZIKV strains for our subsequent challenge with DENV-2. However, extensive revision of

the literature up to date reveals a broad consensus that these two contemporary ZIKV strains

behave very similar from an antigenic point of view[12,51-53]. Our results are confirmatory of those

results showing that both ZIKV strains were neutralized with same efficacy by serum within each

ZIKV-convalescent group, explained by the broadly neutralization activity against multiple ZIKV

strains irrespective of the infecting strain[52]. However, the magnitude of the neutralization of both

strains was statistically higher in animals exposed to DENV 10 months (mid-convalescence) after

ZIKV infection compared to the animals with a shorter ZIKV convalescence (2 months). These

results suggest that the differences in the neutralization profile between the two ZIKV-immune

groups are associated to the longevity of ZIKV convalescence which may be attributable to the

maturation of the cross-reactive immune memory elicited by the heterologous DENV infection and

no to the antigenic differences or the different replication capabilities in rhesus macaques of those
two pre-infecting ZIKV strains[17,54].

The period of convalescence further had an impact in the maintenance of the
neutralization magnitude against ZIKV and DENV overtime. We observed a higher activation of
the memory immune response characterized by transiently higher peak levels of serum NAbs
against DENVs and ZIKV in ZIKVPF-10mo immune animals compared to ZIKVPR-2mo immune
animals challenged with DENV-2. However, unlike heterologous infections with different DENV
serotypes, by 90 days after DENV-2 infection, the naïve and ZIKV-immune animals had similar
levels of DENV-2 NAbs. Moreover, in the ZIKV-immune animals, the ZIKV NAbs returned to
steady-state levels similarly observed before DENV-2 challenge. Overall, these results
demonstrate that pre-existing ZIKV immunity leads to a transient increase in neutralizing Ab
responses in animals challenged with DENV-2 compared to naïve animals. This is in contrast with
previous findings were ZIKV-convalescent macaques show a lack of an early and delayed
anamnestic response overtime with limited induction of DENV NAbs compared to ZIKV-naïve
animals after DENV infection[39]. However, our results show the ability of DENV-2 to activate MBCs
stimulated by the previous ZIKV infection, but this activation is modest and short-lived compared
to the robust and sustained activation of MBCs on secondary DENV infections [11,31,55,56]. Is still
uncertain why the ZIKVPF-10mo animals have a slightly higher peak of Ab response compared
to the ZIKVPR-2mo animals. We speculate this may be caused by modification of MBCs overtime,
so that by 10 months the cells are able to better respond to antigen compared to cells at two
months. After ZIKV infection in human DENV-naïve subjects, the ZIKV/DENV cross-reactive MBC
response increased in magnitude (39% of total MBC proportion) after longer periods of ZIKV
convalescence (~8 months post-ZIKV infection)[57], similar to the 10 months in the ZIKV mid-
convalescent group that exhibited higher DENV cross-neutralization. Based upon studies of
human monoclonal Abs, plasmablasts response during secondary DENV infection is mainly of
MBC origin, resulting in a mature response characterized by cross-neutralizing Abs *in vitro*[58].
However, there are very limited studies on how the affinity maturation develops during the initial
viral encounter and whether the affinity of MBCs is modified during a secondary heterologous
infection or as in this work, during a secondary DENV infection following a primary ZIKV infection.
These are seminal contributions to forecast and understand the cross-neutralization capacity of
further heterologous DENV epidemics in the context of previous ZIKV-DENV immunity.
Interestingly, ZIKV-convalescent animals showed some degree of cross-neutralization
against DENV-2 and DENV-4 before DENV infection. This is consistent with our previous results
showing that DENV-naïve ZIKV-infected animals also preferentially neutralized DENV-4 followed

by DENV-2 after ZIKV infection[17]. Longitudinal data of cross-neutralization of DENV serotypes in DENV-naïve ZIKV-infected human subjects showed low cross-neutralization against all DENV serotypes, but DENV-4 followed by DENV-2 were neutralized more efficiently up to 6 months after ZIKV infection with comparable basal titers reported here[59]. There is no data yet that delineates shared cross-neutralizing epitopes between ZIKV and DENV-2/-4, but it is known that DENV-4 genotypic diversity impact the capacity of its neutralization[60].

One factor that plays a critical role in the induction of enhancement and disease severity is the time elapsed between sequential heterologous DENV infections[10]. At this time, it is unknown whether this factor plays a role when DENV accounts for a secondary infection following ZIKV. Based on our previous works[17,54], it is possible to argue that the sequence of DENV-ZIKV infections induce a different immunological response—in terms of the neutralization magnitude, cytokines profile and functionality of the cellular immune response—compared to the ZIKV-DENV scenario shown here. However, in both scenarios, the role of the time interval between infections seems to play a critical role in the quality and quantity of the immune response.

Early studies of T cells associate their contribution towards immunopathogenesis in DENV secondary infections explained by the original antigenic sin[61], but increasing evidence suggest their protective role during primary and secondary DENV infections[62]. Recently, with the introduction of ZIKV into The Americas, T cells from DENV immunity are being implicated in mediating cross-protection against ZIKV[22-24]. We found that animals with a mid-convalescence to ZIKV developed an early activation of CD4$^+$ and CD8$^+$ effector memory T cells after DENV infection. This early activation has been observed for the opposite scenario in DENV-immune ZIKV-infected patients[24]. Interestingly, the ZIKV early-convalescent group displays a modest activation (T-CM>T-EM) early after DENV infection. Since this group was infected with ZIKV only two months before DENV it is possible that after viral clearance and development of ZIKV-specific T cell response, the T cell compartments were still under the contraction phase at the time of the DENV challenge. Yellow fever virus (YFV) and vaccinia virus vaccinations in humans demonstrate that T cell contraction start as early as approximately one-month post-vaccination and at least for almost three months is still ongoing[63]. Also, a study shows that re-stimulation using alphavirus replicons during T cell response contraction does not have significant impact modulating the pre-existing T cell response[64].

The profile of ZIKV-specific CD8$^+$ T cells in humans with convalescence to ZIKV is characterized by the production of IFN-γ, and expression of activation and cytotoxic markers[65]. Presence of sustained levels of IFN-γ prior and early after DENV challenge in vaccinees has been associated with protection against viremia and/or severe disease[66,67]. We observed a similar

phenotype of the functional response of CD8$^+$ T cells prior DENV infection in animals with longer
convalescence to ZIKV. Strikingly, this response recognizes more efficiently peptides from DENV
E protein than from ZIKV E protein. However, ZIKV-specific CD8$^+$ T cells direct 57% of their
response against structural proteins, which may suggest these cells can recognize conserved
epitopes between ZIKV and DENV structural proteins. Cross-reactivity of T cells between
heterologous flavivirus infections is explained by selective immune recall of memory T cells that
recognize conserved epitopes between DENV and ZIKV[24], which also has previously been
demonstrated during secondary heterotypic DENV infections[68,69]. In addition, an increased
cytotoxic profile as demonstrated by the higher frequency of CD107a-expressing CD4$^+$ and CD8$^+$
T cells in the ZIKV mid-convalescent group correlates with the synchronously early activation of
CD4$^+$ and CD8$^+$ effector memory T cells and elevated levels of perforin release.

Higher proportion of IFN-γ and TNF-α producing T cells before a secondary heterologous
DENV infection has been associated to a subsequent subclinical outcome[70]. Herein, we observed
that the ZIKV mid-convalescent group had elevated levels of IFN-γ and TNF-α producing T cells
since baseline. In this group, DENV infection stimulated a higher frequency of these cells, but
remarkably, also increased highly cross-reactive IFN-γ-producing CD4$^+$ T cells directed to DENV
E, and ZIKV E/NS proteins. A study showed that cross-reactive ZIKV-primed CD4$^+$ T cells
recognized conserved homologous sequences of other related flaviviruses such as West Nile
virus (WNV), YFV, and of relevance for our study, cross-react with E protein epitopes of all DENV
serotypes[71]. Moreover, IFN-γ-producing CD4$^+$ T cells have a role in providing help to B cells in
DENV antigens presentation to CD4$^+$ T cells. This interaction produce IFN-γ and other immune
mediators that induced B cell activation and subsequent efficient Ab production[72]. Memory CD4$^+$
T cells are also required to generate an effective humoral response against ZIKV[73]. Based on
this, the higher proportion of DENV-E-reactive IFN-γ-producing CD4$^+$ T cells may play a role in
the induction of the robust Ab response in the ZIKV mid-convalescent group against ZIKV and all
DENV serotypes. On the other hand, we showed that naïve animals with DENV *de novo* response
did not cross-neutralized ZIKV at all, which state that although similar, antigenic differences are
sufficient to mount predominantly type-specific rather than cross-reactive responses during a
primary infection[51,57].

A lack of ZIKV immunity promoted a more pro-inflammatory profile characterized by
significant elevated levels of IL-6 and MIG/CXCL9. IL-6 has been detected in high levels during
secondary DENV infections in children[74], and the day patients suffer from shock (DSS)[75] or died
from DHF[76]. MIG/CXCL9 is known to be a risk factor for DENV severity involved in vascular
permeability[77]. Its detection varies between primary and secondary (higher levels) DENV

infections[78], which may explain non-significant peaks within ZIKV-immune groups during a
secondary DENV infection. Interestingly, higher levels of IFN-α were observed in the ZIKV-naïve
animals. This antiviral cytokine is known to be actively produced during acute DENV infection *in*
vitro and *in vivo*[79]. Elevated levels have been correlated with severity in DHF patients, and to act
as a marker for elevated DENV replication[80,81]. On the other hand, the presence of a longer ZIKV
convalescence is associated with increased levels of CXCL10 and perforin. CXCL10 is an
immune mediator for T cells proliferation, recruitment of CD4$^+$ and CD8$^+$ activated T cells and IFN-
γ-producing CD8$^+$ T cells, required to control DENV infection *in vivo*[82,83]. This correlates with
higher proportion and activation of both T cell compartments and subsequent functional T cell
response against DENV-E-specific peptides in the group with longer convalescence to ZIKV.
Perforin is involved in the cytotoxic degranulation process against virus-infected cells. In DENV
infection, perforin is part of the anti-DENV cytotoxic phenotype of CD8$^+$ and CD4$^+$ T cells[49,84].
Perforin levels were significantly elevated only in the ZIKV mid-convalescent group after DENV
infection. Accordingly, this coincides with a significant activation of CD8$^+$ and CD4$^+$ effector
memory T cells, and degranulation functional response of both T cell compartments, suggesting
an enhanced perforin-producing cytotoxic role of T cells in presence of longer convalescence to
ZIKV. Contrary to our findings, a previously published work found that an approximately two
month ZIKV immunity period resulted in an increase of pro-inflammatory cytokines[38]. However, a
differential effect due to the use of different sample types (plasma vs serum) between both studies
cannot be ruled out.

One limitation of our study is the utilization of low numbers of animals per group. Additional
studies with a larger number of animals are warranted. However, fundamental and seminal
contributions on ZIKV and ZIKV/DENV interactions have been obtained by using similar limited
number of animals per group[17,18,38,54,85-89]. Another limitation is that our study monitored the
immune response up to 90 days after DENV infection. Additional longitudinal studies are needed
to test the immune response over longer periods of time including subsequent DENV heterotypic
challenges to evaluate the efficacy of the memory recall in cross-protection between serotypes.
Finally, we cannot comment about the likelihood to increase or decrease susceptibility to develop
DHF/DSS in the context of ZIKV immunity since DENV clinical manifestations in NHP models are
limited and are characterized to be subclinical infections [29].

In summary, dissecting our main findings per previous ZIKV-immune status we found that
a ZIKV middle-convalescence: (i) results in shorter DENV viremic period, (ii) lowest pro-
inflammatory status with upregulation of cellular immune response mediators, (iii) robust
neutralizing antibody response higher in magnitude and durability against ZIKV strains and DENV

serotypes, (iv) elevated activated and proliferating B cells, (v) early activation of cross-reactive
CD4$^+$ and CD8$^+$ effector memory T cells, (v) and a major breadth of functional T cell response.
For ZIKV early-convalescence we demonstrated: (i) average DENV viremic period and no
exacerbation of pro-inflammatory status, (ii) neutralizing antibody response with high magnitude
but less durability against ZIKV strains and DENV serotypes compared to the ZIKV middle-
convalescent group, (iii) early activation of central memory CD8$^+$ T cells, (iv) and very limited
activation of effector memory T cells. For the ZIKV-naïve group we demonstrated: (i) longer DENV
viremic period and pro-inflammatory status, (ii) a more delayed *de novo* neutralizing antibody
response against DENV serotypes and inability to neutralize ZIKV strains, (iii) a limited B cell
response, (iv) and an overall *de novo* T cell response lower in magnitude and cross-reactivity
compared to ZIKV-immune groups.

This study reinforces the usefulness of NHPs as a suitable model to characterize the
immune response elicited by heterologous and consecutive flavivirus infections and to identify
differential modulation of the immune response influenced by the time interval between infections.
This proof-of-concept and other prospective studies of ZIKV/DENV pathogenesis and cross-
immune relationships are urgently needed even as the peak of the ZIKV epidemic has passed as
there is a high probability for ZIKV to establish a sylvatic transmission cycle using neotropical
primates and mosquitoes in the Americas that will sustain ZIKV circulation and potential re-
emergence[90,91]. Our findings of highly cross-reactive response against DENV in presence of
previous ZIKV immunity with no exacerbation of DENV pathogenesis may contribute to explain
the decrease of detected DENV cases after ZIKV epidemic in the Americas. This scenario has
been suggested recently using a fewer number of animals[40]. Furthermore, our data show a
positive scenario that supports the implementation of ZIKV vaccine programs, since it suggests
that a vaccine-acquired ZIKV-immunity will not worsen DENV pathogenesis and may ameliorate
immune response against a subsequent infection with DENV. Similarly, the implementation of
DENV vaccines is also supported in the context of previous ZIKV immunity, since ZIKV
convalescence may boost the vaccine-acquired anamnestic immune response to DENV without
predisposing to an enhanced pathogenesis. However, the selection of the vaccine schedule may
be critical to induce the optimal immune response when more than one doses are planned.
702
703 **Methods**
704
Cell lines. Aedes *albopictus* cells, clone C6/36 (ATCC CRL-1660), whole mosquito larva cells,
were maintained in Dulbecco Minimum Essential Medium (DMEM) (GIBCO, Life Technologies)

supplemented with 10% fetal bovine serum (FBS) (Gibco) and 1% Penicillin/Streptomycin (P/S)

(Gibco). C6/36 were used to produce previous ZIKV and DENV viral stocks with high titers in 150-

175 cm^2 cell culture flasks (Eppendorf), and incubated at 33°C and 5% CO_2. Vero cells, clone 81

(ATCC CCL-81), African green monkey kidney epithelial cells, were maintained with DMEM

supplemented with 10% FBS and 1% of P/S, HEPES, L-glutamine and non-essential amino acids

(NEAA) in 75 cm^2 cell culture flasks, and incubated at 37°C and 5% CO_2. Vero-81 cells were used

for the cells monolayer in viral titrations by plaque assays and plaque reduction neutralization test

(PRNT) in flat-bottom 24-well plates (Eppendorf).

Viral stocks. The DENV-2 New Guinea 44 (NGC) strain (kindly provided by Steve Whitehead,

NIH/NIAID, Bethesda, Maryland, USA), known to replicate well in rhesus macaques, was used

for the challenge in order to obtain comparative results with previous published studies from our

group on DENV and ZIKV challenge studies[17,34]. We have standardized the assays to quantify

this virus by Plaque assay, as described in our previous work[17]. The titer of DENV-2 for the

challenge was 5x10^7 pfu/ml. In addition, ZIKV H/PF/2013 strain (kindly provided by CDC-Dengue

Branch, San Juan, Puerto Rico), ZIKV PRVABC59 (ATCC VR-1843), DENV-1 Western Pacific

74, DENV-3 Sleman 73, and DENV-4 Dominique strains (kindly provided by Steve Whitehead

from NIH/NIAID, Bethesda, Maryland, USA) were propagated in C6/36 cells, titrated and used for

Plaque Reduction Neutralization Test (PRNT) assays.

Viral titration plaque assay. DENV titrations by plaque assay were performed seeding Vero-81

(~8.5x10^4 cells /well) in flat bottom 24-well cell culture well plates (Eppendorf) in supplemented

DMEM the day before. Viral dilutions (10-fold) were made in diluent media [Opti-MEM (Invitrogen)

with 2% FBS (Gibco) and 1% P/S (Gibco)]. Prior to inoculation, growth medium was removed and

cells were inoculated with 100 ul/well of each dilution in triplicates. Plates were incubated for 1

hr, 37°C, 5% CO_2 and rocking. After incubation, virus dilutions were overlaid with 1 ml of Opti-

MEM [1% Carboxymethylcellulose (Sigma), 2% FBS, 1% of NEAA (Gibco) and P/S (Gibco)]. After

3 to 5 days of incubation (days vary between DENV serotypes), overlay was removed and cells

were washed twice with phosphate buffered saline (PBS), fixed in 80% methanol (Sigma) in PBS,

and incubated at room temperature (RT) for 15 minutes. Plates were blocked with 5% Non-fat dry

milk (Denia) in PBS for 10 minutes. Blocking buffer was discarded and 200 ul/well of primary

antibodies mix [anti-E protein monoclonal antibody (mAb) 4G2 and anti-prM protein mAb 2H2

(kindly provided by Aravinda de Silva and Ralph Baric, University of North Carolina Chapel Hill,

North Carolina, USA), both diluted 1:250 in blocking buffer] were added and incubated for 1 hr,

$37°C$, 5% CO_2 and rocking. Plates were washed twice with PBS and incubated in same conditions

with horseradish peroxidase (HRP)-conjugated goat anti-mouse secondary antibody (Sigma),

diluted 1:1000 in blocking buffer. Plates were washed twice with PBS and 150 ul/well of TrueBlue

HRP substrate (KPL) were added and plates were incubated from 1-10 minutes at RT until plaque-

forming units (pfu) were produced and visible. Then 200 ul/well of distilled water were added to

stop the substrate reaction, plates get dry and pfu were counted to calculate viral titers.

Macaques and viral challenge. From 2008 to 2015, the Caribbean Primate Research Center

(CPRC) funded a large DENV research program. Multiple studies made available several cohorts

of rhesus macaques (*Macaca mulatta*) infected with different DENV serotypes in distinct timelines

and also naïve cohorts were available as well. After our laboratories prioritized ZIKV research

since 2016, DENV pre-exposed and naïve cohorts were infected with ZIKV and pre-exposed

animals became available for this study. All animals were housed within the Animal Resources

Center facilities at the University of Puerto Rico-Medical Sciences Campus (UPR-MSC), San

Juan, Puerto Rico. All the procedures were performed under the approval of the Institutional

Animal Care and Use Committee (IACUC) of UPR-MSC and in a facility accredited by the

Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC file # 000593;

Animal Welfare Assurance number A3421; protocol number, 7890116). Procedures involving

animals were conducted in accordance with USDA Animal Welfare Regulations, *the Guide for the*

Care and use of Laboratory Animals and institutional policies to ensure minimal suffering of

animals during procedures. All invasive procedures were conducted using anesthesia by

intramuscular injection of ketamine at 10-20 mg kg^{-1} of body weight. Rhesus macaques from the

CPRC are very well genetically characterized from a common stock introduced in 1938 at Cayo

Santiago, an islet located in the southeast of Puerto Rico. These macaques with Indian genetic

background are part of the purest colony used in the United States for comparative medicine and

biomedical research[50]

The experimental design was based on 14 young adult male rhesus macaques divided in

three cohorts. Cohort 1 (ZIKVPF-10mo): composed of four animals (5K6, CB52, 2K2, and 6N1)

that were inoculated with $1x10^6$ pfu/500 ul of the ZIKV H/PF/2013 strain subcutaneously[17] 10

months before DENV-2 challenge. Cohort 2 (ZIKVPR-2mo): composed of 6 animals (MA067,

MA068, BZ34, MA141, MA143, and MA085) that were inoculated with $1x10^6$ pfu/500 ul of the

ZIKV PRVABC59 strain two months before DENV-2 challenge. Both ZIKV strains used for

previous exposure of these groups are >99.99% comparable in amino acid identity

(Supplementary Table 1). Cohort 3 (Naïve): composed of four ZIKV/DENV naïve animals (MA123,

MA023, MA029, and MA062) as a control group. Cohort 1 and 3 were challenged on the same day while cohort 2 was challenged 3 months later with the same stock of DENV-2. However, all samples were frozen and analyzed together, except for the immunophenotyping analysis.

The ages of all animals are within the age range for young adults rhesus macaques https://www.nc3rs.org.uk/macaques/macaques/life-history-and-diet/ (ZIKVPF-10mo: 6.8, 6.8, 5.8, and 5.9; ZIKVPR-2mo: 6.4, 6.5, 5.2, 4.3, 5.6, and 5.5; Naïve: 4.8, 6.6, 6.8, and 5.7). Prior to DENV-2 challenge all animals were subjected to quarantine period. All cohorts were bled for baseline and challenged subcutaneously (deltoid area) with $5x10^5$ pfu/500 ul of DENV-2 New Guinea 44 strain. After DENV-2 challenge all animals were extensively monitored by trained and certified veterinary staff for evidence of disease and clinical status: external temperature (°C) with an infrared device (EXTECH Instruments, Waltham, MA), weight (Kg), CBC and CMP. All animals were bled once daily from day 1 to day 10 and after that on days 15, 30, 60 and 90 post-infection (dpi). In all timepoints the blood samples were used for serum separation (Baseline, 7, 30, 60, 90 dpi only). PBMCs were collected at same time points using CPT tubes (BD-Biosciences, San Jose, CA) containing citrate. Additional heparin samples were obtained for immunophenotyping by flow cytometry using fresh whole blood. Fig. 1 shows the experimental design and samples collection timeline.

DENV RNAemia. DENV viral RNA extraction was performed from acute serum samples (Baseline, 1-10, and 15 dpi) using QIAamp Viral RNA mini kit (Qiagen Inc, Valencia, CA, USA) according to the manufacturer's instructions. RNAemia levels were measured by a One-Step qRT-PCR detection kit (Oasig, Primerdesign Ltd., UK) and using DENV RT primer/probe Mix kit (Genesig, Primerdesign Ltd., UK) according to the manufacturer's protocol. Assays were performed in an iCycler IQ5 Real-Time Detection System with Optical System Software version 2.1 (Bio-Rad, Hercules, CA, USA). Limit of detection (LOD) was 20 copies per ml. Furthermore, in order to correlate RNAemia levels with DENV pathogenesis we monitored the clinical status for injury and/or clinical manifestations. Complete Blood Counts (CBC) were performed for all animals in several timepoints (Baseline, 7, and 15 dpi) to determine the absolute number (10^6 cells/ml) and percent (%) of lymphocytes (LYM), monocytes (MON), white blood cells (WBC), neutrophils (NEU) and platelets (PLT). Also, Comprehensive Metabolic Panel (CMP) were evaluated in several timepoints (Baseline, 7, 15 and 30 dpi) to measure concentration (U/L) of alkaline phosphatase and liver enzymes alanine aminotransferase (ALT) and aspartate aminotransferase (AST).

ELISA. Seroreactivity to DENV and cross-reactivity to ZIKV was measured at different timepoints before and after DENV-2 challenge. DENV-IgM (Focus Diagnostics, Cypress, CA, USA) was quantified at baseline, 5, 10, 15 and 30 dpi. DENV-IgG was quantified at baseline, 7, 15, 30, 60 and 90 dpi (Focus Diagnostics, Cypress, CA, USA). To determine the modulation of serological profile against ZIKV we assessed: levels of anti-ZIKV IgM (InBios, Seattle, WA, USA) at baseline, 5, 10, 15 and 30 dpi; anti-ZIKV IgG (XPressBio, Frederick, MD, USA) at baseline, 7, 15, 30, 60 and 90 dpi; anti-ZIKV NS1-IgG (Alpha Diagnostics, San Antonio, TX, USA) at baseline, 30, 60 and 90 dpi (including additional timepoints prior baseline for both ZIKV-immune groups); and anti-ZIKV EDIII-IgG (Alpha Diagnostics International, San Antonio, TX, USA). All ELISA-based assays were performed following the manufacturers' instructions. This serological characterization allows us to assess the dynamics of DENV and ZIKV cross-reactivity but without discerning between cross-reactive binding Abs and cross- or type-specific neutralizing Abs.

Plaque Reduction Neutralization Test (PRNT). Selected serum samples (baseline, 30, 60 and 90 dpi) were challenged to neutralized ZIKV (H/PF/2013, PRVABC59), DENV-1 Western Pacific 74, DENV-2 NGC 44, DENV-3 Sleman 73, and DENV-4 Dominique strains. For the infecting serotype (DENV-2) and ZIKV the NAbs were measured in early timepoints as well (7 and 15 dpi). For the PRNT, serum samples were inactivated, diluted (2-fold), mixed with a constant inoculum of virus (volume necessary to produce ~35 pfu/well) and then incubated for 1 hr at 37°C and 5% CO_2. After incubation, virus-serum mix dilutions were added to Vero-81 cells monolayer in flat bottom 24-well plates seeded the day before for 1 hr at 37°C and 5% CO_2, finally overlay medium was added and incubated by several days (serotype dependent). Results were reported as PRNT60 titers, NAb titer capable of reduce 60% or more of DENV serotypes or ZIKV strains pfu compared with the mock (control of virus without serum). A PRNT60 1:20 titer was considered a positive Neut titer, and <1:20 as a negative Neut titer. Non-neutralizing titers (<1:20) were assigned with one-half of the limit of detection for graphs visualization.

Multiplex cytokine profile. A total of 8 cytokines/chemokines were measured (pg /ml^{-1}) by Luminex at baseline, 1, 2, 3, 5, 10, 15 and 30 dpi, including: interferon alpha (IFN-α), interleukin-6 (IL-6), monokine induced by IFN-gamma (MIG/CXCL9), monocyte chemoattractant protein 1 (MCP-1/CCL2), macrophage inflammatory protein 1-beta (MIP-1β/CCL4), IL-1 receptor antagonist (IL-1RA), C-X-C motif chemokine 10 (CXCL10/IP-10) and perforin. The multiplex assay was conducted as previously described[17,92].

Immunophenotyping. Flow cytometry (MACSQuant Analyzer 10, Miltenyi Biotec) analysis was

performed to determine the frequency, activation and proliferation of cell populations of the innate

and adaptive immune response based on the phenotyping strategy of a previous study[17]

(Supplementary Fig. 7, 8, and 9 for gating strategy; Supplementary Table 2 for Ab panel).

Phenotypic characterization of macaque PBMCs from fresh whole blood samples was performed

by 8-multicolor flow cytometry using fluorochrome conjugated Abs at several timepoints (Baseline,

1, 2, 3, 7, 10 dpi; and 15 and 30 dpi for B/T cell panel only). Single cells (singlets) were selected

by their FSC area (FSC-A) and height (FSC-H) patterns. Lymphocytes (LYM) were gated based

on their characteristic forward and side scatter pattern (FSC, SSC). T cells were selected gating

on the $CD3^+$ population. $CD4^+$ and $CD8^+$ T cells were defined as $CD3^+CD4^+$ and $CD3^+CD8^+$,

respectively. Naive (N; $CD28^+CD95^-$), effector memory (EM; $CD28^-CD95^+$) and central memory

(CM; $CD28^+CD95^+$) T cell subpopulations were determined within $CD4^+$ and $CD8^+$ T cells. B cells

were defined as $CD20^+CD3^-$. The activation of B and T cell memory subpopulations (EM and CM)

was assessed by the presence of the early activation marker CD69. Proliferation of total and

activated B cells was quantified by the expression of the intracellular marker Ki67. Natural killer

(NK) cells were defined as $CD3^-CD20^-CD14^-$ and analyzed by the double positive expression of

the following NK cell markers: CD8, CD56, NKG2A, NKp30, and NKp46 (Supplementary Fig. 9

for gating strategy). Dendritic cells (DC) were separated in two populations within the Lineage-

DR+ ($HLA-DR^+$ $CD3^-$ $CD14^-$ $CD16^-$ $CD20^-$ $CD8^-$ $NKG2A^-$) by the expression of CD123

(plasmacytoid, pDC) or CD11c (myeloid, mDCs) (Supplementary Fig. 8 for gating strategy). Then,

DC percentages were calculated from total PBMCs (total events of the DC subpopulation divided

by total PBMCs and multiplied by 100). The phenotyping assays were optimized and performed

as previously published[17,34,93].

T cell functional response. Intracellular cytokine staining of macaques PBMCs was performed

by multicolor flow cytometry using methods previously described (Supplementary Fig. 14 for

gating strategy; Supplementary Table 3 for Ab panel)[17,93]. Functional effector response of $CD4^+$

and $CD8^+$ T cells was measured before and after DENV infection. Antigen-specific $CD4^+$ and

$CD8^+$ T cell effector responses were measured at baseline to determine basal levels in presence

(ZIKVPF-10mo, ZIKVPR-2mo) or absence (Naïve) of previous immunity to ZIKV. Also, 30, 60 and

90 dpi were assessed to determine how this pre-existing functional response is modulated after

DENV infection and if is maintained over time. For peptide pools stimulation, PBMCs were

stimulated for 6 hr at 37°C and 5% CO_2. The peptides used for DENV-E, ZIKV-E and ZIKV-NS

were 15-mers overlapped by 10 amino acids at 1.25 ug/ml^{-1}, 2.5 ug/ml^{-1}, 475 ng/ml^{-1} per peptide,

respectively (Supplementary Table 4 for peptide sequences). The stimulation with peptides was
performed in presence of brefeldin A at 10 ug/ml^{-1}. After stimulation, the cells were stained for the
following markers: CD3, CD4, CD8, CD20 (excluded), CD107a (functional cytotoxicity). Levels of
IFN-γ and TNF-α also were measured in gated lymphocytes cell populations. Samples were
measured and data was collected on a LSRII (BD).

Statistical analysis. Statistical analyses were performed using GraphPad Prism 7.0 software
(GraphPad Software, San Diego, CA, USA). The statistical significance between the means of all
groups were determined using Two-way ANOVA Multiple Comparison Tukey Test, and to
compare each mean against the baseline mean within same group Two-way ANOVA Multiple
Comparison Dunnett Test was performed. Total number of families and comparisons per family
used for adjustments are depicted in each figure legend. Significant multiplicity adjusted p values
(* <0.05, ** <0.01, *** <0.001, **** <0.0001) show statistically significant difference between
groups (Tukey Test) or timepoints within a group (Dunnett Test).

Data availability. All relevant data is in main figures and supplementary information, any
additional details are available from authors upon request.

Acknowledgments. We thank all the staff of the Caribbean Primate Research Center and Animal
Resources Center for their continuous support with the sample collection, schedule, and
monitoring of the animals. Authors recognize the support provided by Dr. Elmer Rodriguez
reviewing the statistics. This work was supported by the following grants: 2 P40 OD012217 and
2U42OD021458-15 to C.A.S. and M.I.M., K22AI104794 to J.D.B., P51OD011133 (L.G.),
HHSN272201400045C to D.W., and R25GM061838 to E.X.P.-G.

Author contributions. C.A.S. and E.X.P.-G. developed the experimental design. I.V.R.
supervised and performed sample collection and animals monitoring. E.X.P.-G., P.P., C.S.-C.,
M.A.H., A.O.-R., V.H., L.P., L.C., and T.A. performed the experiments. E.X.P.-G., C.A.S., V.H.,
M.A.H., L.J.W., A.d.S., and D.W. analyzed the data. E.X.P.-G. and C.A.S. drafted the manuscript.
C.A.S., E.X.P.-G., D.W., A.K.P., J.D.B., M.A.H., L.G., L.J.W., and A.d.S. revised the manuscript.

Additional Information

Supplementary Information available at:

Competing interests: The authors declare no competing financial interests.

913
References

915
916

1   Metsky, H. C. *et al.* Zika virus evolution and spread in the Americas. *Nature* **546**, 411-415, doi:10.1038/nature22402 (2017).
2   Garcez, P. P. *et al.* Zika virus impairs growth in human neurospheres and brain organoids. *Science (New York, N.Y.)* **352**, 816-818, doi:10.1126/science.aaf6116 (2016).
3   Li, C. *et al.* Zika Virus Disrupts Neural Progenitor Development and Leads to Microcephaly in Mice. *Cell stem cell* **19**, 672, doi:10.1016/j.stem.2016.10.017 (2016).
4   Mlakar, J. *et al.* Zika Virus Associated with Microcephaly. *The New England journal of medicine* **374**, 951-958, doi:10.1056/NEJMoa1600651 (2016).
5   de Oliveira, W. K. *et al.* Zika Virus Infection and Associated Neurologic Disorders in Brazil. *The New England journal of medicine* **376**, 1591-1593, doi:10.1056/NEJMc1608612 (2017).
6   Guo, C. *et al.* Global Epidemiology of Dengue Outbreaks in 1990-2015: A Systematic Review and Meta-Analysis. *Frontiers in cellular and infection microbiology* **7**, 317, doi:10.3389/fcimb.2017.00317 (2017).
7   Bhatt, S. *et al.* The global distribution and burden of dengue. *Nature* **496**, 504-507, doi:10.1038/nature12060 (2013).
8   Katzelnick, L. C. *et al.* Dengue viruses cluster antigenically but not as discrete serotypes. *Science (New York, N.Y.)* **349**, 1338-1343, doi:10.1126/science.aac5017 (2015).
9   Khurram, M. *et al.* Dengue hemorrhagic fever: comparison of patients with primary and secondary infections. *Journal of infection and public health* **7**, 489-495, doi:10.1016/j.jiph.2014.05.005 (2014).
10   Guzman, M. G. *et al.* Enhanced severity of secondary dengue-2 infections: death rates in 1981 and 1997 Cuban outbreaks. *Rev Panam Salud Publica* **11**, 223-227, doi:S1020-49892002000400003 [pii] (2002).
11   Guzman, M. G., Alvarez, M. & Halstead, S. B. Secondary infection as a risk factor for dengue hemorrhagic fever/dengue shock syndrome: an historical perspective and role of antibody-dependent enhancement of infection. *Archives of virology* **158**, 1445-1459, doi:10.1007/s00705-013-1645-3 (2013).
12   Barba-Spaeth, G. *et al.* Structural basis of potent Zika-dengue virus antibody cross-neutralization. *Nature* **536**, 48-53, doi:10.1038/nature18938 (2016).
13   Butler, D. Brazil asks whether Zika acts alone to cause birth defects. *Nature* **535**, 475-476, doi:10.1038/nature.2016.20309 (2016).
14   Dejnirattisai, W. *et al.* Dengue virus sero-cross-reactivity drives antibody-dependent enhancement of infection with zika virus. *Nature immunology*, doi:10.1038/ni.3515 (2016).
15   Priyamvada, L. *et al.* Human antibody responses after dengue virus infection are highly cross-reactive to Zika virus. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 7852-7857, doi:10.1073/pnas.1607931113 (2016).
16   Bardina, S. V. *et al.* Enhancement of Zika virus pathogenesis by preexisting antiflavivirus immunity. *Science (New York, N.Y.)* **DOI 10.1126/science.aal4365**, doi:10.1126/science.aal4365 (2017).
17   Pantoja, P. *et al.* Zika virus pathogenesis in rhesus macaques is unaffected by pre-existing immunity to dengue virus. *Nature communications* **8**, 15674, doi:10.1038/ncomms15674 (2017).

McCracken, M. K. *et al.* Impact of prior flavivirus immunity on Zika virus infection in rhesus macaques. *PLoS pathogens* **13**, e1006487, doi:10.1371/journal.ppat.1006487 (2017).

Terzian, A. C. B. *et al.* Viral Load and Cytokine Response Profile Does Not Support Antibody-Dependent Enhancement in Dengue-Primed Zika Virus-Infected Patients. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **65**, 1260-1265, doi:10.1093/cid/cix558 (2017).

Gordon, A. *et al.* Prior dengue virus infection and risk of Zika: A pediatric cohort in Nicaragua. *PLoS medicine* **16**, e1002726, doi:10.1371/journal.pmed.1002726 (2019).

Kam, Y. W. *et al.* Cross-reactive dengue human monoclonal antibody prevents severe pathologies and death from Zika virus infections. *JCI insight* **2**, doi:10.1172/jci.insight.92428 (2017).

Wen, J. *et al.* Dengue virus-reactive CD8(+) T cells mediate cross-protection against subsequent Zika virus challenge. *Nature communications* **8**, 1459, doi:10.1038/s41467-017-01669-z (2017).

Wen, J. *et al.* Identification of Zika virus epitopes reveals immunodominant and protective roles for dengue virus cross-reactive CD8(+) T cells. *Nature microbiology* **2**, 17036, doi:10.1038/nmicrobiol.2017.36 (2017).

Grifoni, A. *et al.* Prior Dengue virus exposure shapes T cell immunity to Zika virus in humans. *Journal of virology* **DOI 10.1128/jvi.01469-17**, doi:10.1128/jvi.01469-17 (2017).

Perez, F. *et al.* The decline of dengue in the Americas in 2017: discussion of multiple hypotheses. *Tropical medicine & international health : TM & IH*, doi:10.1111/tmi.13200 (2019).

Solimini, A. G., Manica, M., Rosa, R., Della Torre, A. & Caputo, B. Estimating the risk of Dengue, Chikungunya and Zika outbreaks in a large European city. *Scientific reports* **8**, 16435, doi:10.1038/s41598-018-34664-5 (2018).

Gutierrez-Lopez, R. *et al.* Vector Competence of Aedes caspius and Ae. albopictus Mosquitoes for Zika Virus, Spain. *Emerging infectious diseases* **25**, 346-348, doi:10.3201/eid2502.171123 (2019).

Ribeiro, G. S. *et al.* Does immunity after Zika virus infection cross-protect against dengue? *The Lancet. Global health* **6**, e140-e141, doi:10.1016/s2214-109x(17)30496-5 (2018).

Sariol, C. A. & White, L. J. Utility, limitations, and future of non-human primates for dengue research and vaccine development. *Frontiers in immunology* **5**, 452, doi:10.3389/fimmu.2014.00452 (2014).

Clark, K. B., Onlamoon, N., Hsiao, H. M., Perng, G. C. & Villinger, F. Can non-human primates serve as models for investigating dengue disease pathogenesis? *Frontiers in microbiology* **4**, 305, doi:10.3389/fmicb.2013.00305 (2013).

Koraka, P., Benton, S., van Amerongen, G., Stittelaar, K. J. & Osterhaus, A. D. Characterization of humoral and cellular immune responses in cynomolgus macaques upon primary and subsequent heterologous infections with dengue viruses. *Microbes and infection* **9**, 940-946, doi:10.1016/j.micinf.2007.03.012 (2007).

Borges, M. B. *et al.* Characterization of recent and minimally passaged Brazilian dengue viruses inducing robust infection in rhesus macaques. *PloS one* **13**, e0196311, doi:10.1371/journal.pone.0196311 (2018).

White, L. J. *et al.* An alphavirus vector-based tetravalent dengue vaccine induces a rapid and protective immune response in macaques that differs qualitatively from immunity induced by live virus infection. *Journal of virology* **87**, 3409-3424, doi:10.1128/JVI.02298-12 (2013).

Sariol, C. A. *et al.* Decreased Dengue Replication and an Increased Anti-viral Humoral Response with the use of Combined Toll-Like Receptor 3 and 7/8 Agonists in Macaques. *PloS one* **6**, 9 (2011).

35 Kawiecki, A. B. & Christofferson, R. C. Zika Virus-Induced Antibody Response Enhances
Dengue Virus Serotype 2 Replication In Vitro. *The Journal of infectious diseases* **214**,
1357-1360, doi:10.1093/infdis/jiw377 (2016).
36 Stettler, K. *et al.* Specificity, cross-reactivity, and function of antibodies elicited by Zika
virus infection. *Science (New York, N.Y.)* **353**, 823-826, doi:10.1126/science.aaf8505
(2016).
37 Fowler, A. M. *et al.* Maternally Acquired Zika Antibodies Enhance Dengue Disease
Severity in Mice. *Cell host & microbe* **24**, 743-750.e745, doi:10.1016/j.chom.2018.09.015
(2018).
38 George, J. *et al.* Prior Exposure to Zika Virus Significantly Enhances Peak Dengue-2
Viremia in Rhesus Macaques. *Scientific reports* **7**, 10498, doi:10.1038/s41598-017-
10901-1 (2017).
39 Valiant, W. G. *et al.* Zika convalescent macaques display delayed induction of anamnestic
cross-neutralizing antibody responses after dengue infection. *Emerging microbes &*
infections **7**, 130, doi:10.1038/s41426-018-0132-z (2018).
40 Breitbach, M. E. *et al.* Primary infection with dengue or Zika virus does not affect the
severity of heterologous secondary infection in macaques. *PLoS pathogens* **15**,
e1007766, doi:10.1371/journal.ppat.1007766 (2019).
41 Anderson, K. B. *et al.* A shorter time interval between first and second dengue infections
is associated with protection from clinical illness in a school-based cohort in Thailand. *The*
Journal of infectious diseases **209**, 360-368, doi:10.1093/infdis/jit436 (2014).
42 Montoya, M. *et al.* Symptomatic versus inapparent outcome in repeat dengue virus
infections is influenced by the time interval between infections and study year. *PLoS*
neglected tropical diseases **7**, e2357, doi:10.1371/journal.pntd.0002357 (2013).
43 Kliks, S. C., Nimmanitya, S., Nisalak, A. & Burke, D. S. Evidence that maternal dengue
antibodies are important in the development of dengue hemorrhagic fever in infants. *The*
American journal of tropical medicine and hygiene **38**, 411-419 (1988).
44 Bhoomiboonchoo, P. *et al.* Sequential dengue virus infections detected in active and
passive surveillance programs in Thailand, 1994-2010. *BMC public health* **15**, 250,
doi:10.1186/s12889-015-1590-z (2015).
45 Arien, K. K., Michiels, J., Foque, N., Heyndrickx, L. & Van Esbroeck, M. Can Zika virus
antibodies cross-protect against dengue virus? *The Lancet. Global health* **6**, e494,
doi:10.1016/s2214-109x(18)30122-0 (2018).
46 Ribeiro, G. S. *et al.* Can Zika virus antibodies cross-protect against dengue virus? -
Authors' reply. *The Lancet. Global health* **6**, e495, doi:10.1016/s2214-109x(18)30123-2
(2018).
47 Katzelnick, L. C. & Harris, E. The use of longitudinal cohorts for studies of dengue viral
pathogenesis and protection. *Current opinion in virology* **29**, 51-61,
doi:10.1016/j.coviro.2018.03.004 (2018).
48 Katzelnick, L. C., Montoya, M., Gresh, L., Balmaseda, A. & Harris, E. Neutralizing antibody
titers against dengue virus correlate with protection from symptomatic infection in a
longitudinal cohort. *Proceedings of the National Academy of Sciences of the United States*
of America **113**, 728-733, doi:10.1073/pnas.1522136113 (2016).
49 Weiskopf, D. *et al.* Dengue virus infection elicits highly polarized CX3CR1+ cytotoxic
CD4+ T cells associated with protective immunity. *Proceedings of the National Academy*
of Sciences of the United States of America **112**, E4256-4263,
doi:10.1073/pnas.1505956112 (2015).
50 Kanthaswamy, S. *et al.* The Population Genetic Composition of Conventional and SPF
Colonies of Rhesus Macaques (Macaca mulatta) at the Caribbean Primate Research
Center. *Journal of the American Association for Laboratory Animal Science : JAALAS* **55**,
147-151 (2016).

Collins, M. H. *et al.* Lack of Durable Cross-Neutralizing Antibodies Against Zika Virus from Dengue Virus Infection. *Emerging infectious diseases* **23**, 773-781, doi:10.3201/eid2305.161630 (2017).

Dowd, K. A. *et al.* Broadly Neutralizing Activity of Zika Virus-Immune Sera Identifies a Single Viral Serotype. *Cell reports* **16**, 1485-1491, doi:10.1016/j.celrep.2016.07.049 (2016).

Swanstrom, J. A. *et al.* Dengue Virus Envelope Dimer Epitope Monoclonal Antibodies Isolated from Dengue Patients Are Protective against Zika Virus. *mBio* **7**, DOI 10.1128/mBio.01123-01116, doi:10.1128/mBio.01123-16 (2016).

Serrano-Collazo, C. *et al.* Significant control of Zika infection in macaques depends on the elapsing time after dengue exposure. *bioRxiv*, 625293, doi:10.1101/625293 (2019).

de Alwis, R. *et al.* In-depth analysis of the antibody response of individuals exposed to primary dengue virus infection. *PLoS Negl Trop Dis* **5**, e1188, doi:10.1371/journal.pntd.0001188 (2011).

Wahala, W. M. & Silva, A. M. The human antibody response to dengue virus infection. *Viruses* **3**, 2374-2395, doi:10.3390/v3122374 (2011).

Andrade, P. *et al.* Impact of pre-existing dengue immunity on human antibody and memory B cell responses to Zika. *Nature communications* **10**, 938, doi:10.1038/s41467-019-08845-3 (2019).

Priyamvada, L. *et al.* B Cell Responses during Secondary Dengue Virus Infection Are Dominated by Highly Cross-Reactive, Memory-Derived Plasmablasts. *Journal of virology* **90**, 5574-5585, doi:10.1128/jvi.03203-15 (2016).

Montoya, M. *et al.* Longitudinal Analysis of Antibody Cross-neutralization Following Zika Virus and Dengue Virus Infection in Asia and the Americas. *The Journal of infectious diseases* **218**, 536-545, doi:10.1093/infdis/jiy164 (2018).

Gallichotte, E. N. *et al.* Genetic Variation between Dengue Virus Type 4 Strains Impacts Human Antibody Binding and Neutralization. *Cell reports* **25**, 1214-1224, doi:10.1016/j.celrep.2018.10.006 (2018).

Mathew, A., Townsley, E. & Ennis, F. A. Elucidating the role of T cells in protection against and pathogenesis of dengue virus infections. *Future microbiology* **9**, 411-425, doi:10.2217/fmb.13.171 (2014).

St John, A. L. & Rathore, A. P. S. Adaptive immune responses to primary and secondary dengue virus infections. *Nature reviews. Immunology*, doi:10.1038/s41577-019-0123-x (2019).

Miller, J. D. *et al.* Human effector and memory CD8+ T cell responses to smallpox and yellow fever vaccines. *Immunity* **28**, 710-722, doi:10.1016/j.immuni.2008.02.020 (2008).

Knudsen, M. L. *et al.* Kinetic and phenotypic analysis of CD8+ T cell responses after priming with alphavirus replicons and homologous or heterologous booster immunizations. *Journal of virology* **88**, 12438-12451, doi:10.1128/jvi.02223-14 (2014).

Grifoni, A. *et al.* Cutting Edge: Transcriptional Profiling Reveals Multifunctional and Cytotoxic Antiviral Responses of Zika Virus-Specific CD8(+) T Cells. *Journal of immunology (Baltimore, Md. : 1950)* **201**, 3487-3491, doi:10.4049/jimmunol.1801090 (2018).

Gunther, V. J. *et al.* A human challenge model for dengue infection reveals a possible protective role for sustained interferon gamma levels during the acute phase of illness. *Vaccine* **29**, 3895-3904, doi:10.1016/j.vaccine.2011.03.038 (2011).

Wijeratne, D. T. *et al.* Quantification of dengue virus specific T cell responses and correlation with viral load and clinical disease severity in acute dengue infection. *PLoS neglected tropical diseases* **12**, e0006540, doi:10.1371/journal.pntd.0006540 (2018).

Weiskopf, D. *et al.* Comprehensive analysis of dengue virus-specific responses supports an HLA-linked protective role for CD8+ T cells. *Proceedings of the National Academy of*

Sciences of the United States of America **110**, E2046-2053, doi:10.1073/pnas.1305227110 (2013).

69 Mathew, A. *et al.* Predominance of HLA-restricted cytotoxic T-lymphocyte responses to serotype-cross-reactive epitopes on nonstructural proteins following natural secondary dengue virus infection. *Journal of virology* **72**, 3999-4004 (1998).

70 Hatch, S. *et al.* Intracellular cytokine production by dengue virus-specific T cells correlates with subclinical secondary infection. *The Journal of infectious diseases* **203**, 1282-1291, doi:10.1093/infdis/jir012 (2011).

71 Reynolds, C. J. *et al.* T cell immunity to Zika virus targets immunodominant epitopes that show cross-reactivity with other Flaviviruses. *Scientific reports* **8**, 672, doi:10.1038/s41598-017-18781-1 (2018).

72 Rivino, L. *et al.* Differential targeting of viral components by CD4+ versus CD8+ T lymphocytes in dengue virus infection. *Journal of virology* **87**, 2693-2706, doi:10.1128/jvi.02675-12 (2013).

73 Elong Ngono, A. *et al.* CD4+ T cells promote humoral immunity and viral control during Zika virus infection. *PLoS pathogens* **15**, e1007474, doi:10.1371/journal.ppat.1007474 (2019).

74 Restrepo, B. N. *et al.* Serum levels of interleukin-6, tumor necrosis factor-alpha and interferon-gamma in infants with and without dengue. *Revista da Sociedade Brasileira de Medicina Tropical* **41**, 6-10 (2008).

75 Hober, D. *et al.* Serum levels of tumor necrosis factor-alpha (TNF-alpha), interleukin-6 (IL-6), and interleukin-1 beta (IL-1 beta) in dengue-infected patients. *The American journal of tropical medicine and hygiene* **48**, 324-331 (1993).

76 Chen, L. C. *et al.* Correlation of serum levels of macrophage migration inhibitory factor with disease severity and clinical outcome in dengue patients. *The American journal of tropical medicine and hygiene* **74**, 142-147 (2006).

77 Dalrymple, N. A. & Mackow, E. R. Endothelial cells elicit immune-enhancing responses to dengue virus infection. *Journal of virology* **86**, 6408-6415, doi:10.1128/jvi.00213-12 (2012).

78 Huang, J. *et al.* Serum Cytokine Profiles in Patients with Dengue Fever at the Acute Infection Phase. *Disease markers* **2018**, 8403937, doi:10.1155/2018/8403937 (2018).

79 Gandini, M. *et al.* Dengue virus activates membrane TRAIL relocalization and IFN-alpha production by human plasmacytoid dendritic cells in vitro and in vivo. *PLoS neglected tropical diseases* **7**, e2257, doi:10.1371/journal.pntd.0002257 (2013).

80 Kurane, I. *et al.* High levels of interferon alpha in the sera of children with dengue virus infection. *The American journal of tropical medicine and hygiene* **48**, 222-229 (1993).

81 Singla, M. *et al.* Immune Response to Dengue Virus Infection in Pediatric Patients in New Delhi, India--Association of Viremia, Inflammatory Mediators and Monocytes with Disease Severity. *PLoS neglected tropical diseases* **10**, e0004497, doi:10.1371/journal.pntd.0004497 (2016).

82 Dufour, J. H. *et al.* IFN-gamma-inducible protein 10 (IP-10; CXCL10)-deficient mice reveal a role for IP-10 in effector T cell generation and trafficking. *Journal of immunology (Baltimore, Md. : 1950)* **168**, 3195-3204 (2002).

83 Hsieh, M. F. *et al.* Both CXCR3 and CXCL10/IFN-inducible protein 10 are required for resistance to primary infection by dengue virus. *Journal of immunology (Baltimore, Md. : 1950)* **177**, 1855-1863 (2006).

84 Gagnon, S. J., Ennis, F. A. & Rothman, A. L. Bystander target cell lysis and cytokine production by dengue virus-specific human CD4(+) cytotoxic T-lymphocyte clones. *Journal of virology* **73**, 3623-3629 (1999).

Dudley, D. M. *et al.* Infection via mosquito bite alters Zika virus tissue tropism and replication kinetics in rhesus macaques. *Nature communications* **8**, 2096, doi:10.1038/s41467-017-02222-8 (2017).

Magnani, D. M. *et al.* Neutralizing human monoclonal antibodies prevent Zika virus infection in macaques. *Science translational medicine* **9**, doi:10.1126/scitranslmed.aan8184 (2017).

Masel, J. *et al.* Does prior dengue virus exposure worsen clinical outcomes of Zika virus infection? A systematic review, pooled analysis and lessons learned. *PLoS neglected tropical diseases* **13**, e0007060, doi:10.1371/journal.pntd.0007060 (2019).

Osuna, C. E. *et al.* Zika viral dynamics and shedding in rhesus and cynomolgus macaques. *Nature medicine* **DOI 10.1038/nm.4206**, doi:10.1038/nm.4206 (2016).

Silveira, E. L. V. *et al.* Immune Cell Dynamics in Rhesus Macaques Infected with a Brazilian Strain of Zika Virus. *Journal of immunology (Baltimore, Md. : 1950)* **199**, 1003-1011, doi:10.4049/jimmunol.1700256 (2017).

Althouse, B. M. *et al.* Potential for Zika Virus to Establish a Sylvatic Transmission Cycle in the Americas. *PLoS neglected tropical diseases* **10**, e0005055, doi:10.1371/journal.pntd.0005055 (2016).

Favoretto, S. R. *et al.* Zika Virus in Peridomestic Neotropical Primates, Northeast Brazil. *EcoHealth*, doi:10.1007/s10393-019-01394-7 (2019).

Giavedoni, L. D. Simultaneous detection of multiple cytokines and chemokines from nonhuman primates using luminex technology. *Journal of immunological methods* **301**, 89-101, doi:10.1016/j.jim.2005.03.015 (2005).

Meyer, C., Haberthur, K., Asquith, M. & Messaoudi, I. Flow Cytometry-Based Methods to Characterize Immune Senescence in Nonhuman Primates. *Methods in molecular biology* **1343**, 65-80, doi:10.1007/978-1-4939-2963-4_6 (2015).

Main Figures

Figure 1 | Experimental design for DENV-2 challenge of ZIKV-immune and naïve rhesus macaques. 14 young adult male rhesus macaques (*Macaca mulatta*), matched in age and weight, were divided in three cohorts. ZIKVPF-10mo (n=4): composed of four animals (5K6, CB52, 2K2, and 6N1) that were inoculated with 1×10^6 pfu/500 ul of the ZIKV H/PF/2013 strain subcutaneously 10 months before (middle convalescence) DENV-2 challenge. ZIKVPR-2mo (n=6): composed of six animals (MA067, MA068, BZ34, MA141, MA143, and MA085) that were inoculated with 1×10^6 pfu/500 ul of the contemporary ZIKV PRVABC59 strain two months before (early convalescence) DENV-2 challenge. Both ZIKV strains used for previous exposure of these groups are >99.99% comparable in amino acid identity (Supplementary Table 1). Naïve (n=4): composed of four ZIKV/DENV naïve animals (MA123, MA023, MA029, and MA062) as a control group. Prior to DENV-2 challenge all animals were subjected to quarantine period. All cohorts challenged subcutaneously (deltoid area) with 5×10^5 pfu/500 ul of DENV-2 New Guinea 44 strain (NGC44). After DENV-2 challenge all animals were extensively monitored for evidence of disease and clinical status by vital signs such as external temperature (°C), weight (Kg), CBC and CMP panels at the Caribbean Primate Research Center (CPRC). Blood samples were collected at baseline, 1 to 10, 15, 30, 60 and 90 days after DENV infection. In all timepoints the blood samples were used for serum separation (yellow). PBMCs isolation (red) was performed in different tubes with citrate as anticoagulant at baseline, 1, 2, 3, 7, 10, 15, 30, 60, and 90 days after DENV infection.

Figure 2 | Previous ZIKV immunity does not contribute to an increase of DENV RNAemia. (**a**) DENV-2 RNA kinetics in ZIKV-immune and naïve animals at baseline, day 1 to day 10, and day 15 after DENV infection. RNA genome copies (Log10) per ml of serum were measured by qRT-PCR. Symbols represent individual animals per cohort: blue squares (ZIKVPF-10mo), orange squares (ZIKVPR-2mo) and black circles (Naïve). Box and whiskers show the distribution of log-transformed values per group per timepoint. Boxes include the mean value per group while whiskers depict the minimum and maximum values for each group. Cutted line mark the limit of detection (20 genomes copies). Statistically significant differences between groups were determined using Two-Way Anova adjusted for Tukey's multiple comparisons test including 12 families, and 3 comparisons per family. (**b**) Total days that DENV-2 RNAemia was detected for each animal within cohorts. Bars represent mean days per cohort.

Figure 3 | ZIKV immunity does not exacerbate levels of pro-inflammatory cytokines. Cytokines and chemokines expression levels were determined in serum (pg/ml) by multiplex bead assay (Luminex) at baseline, 1, 2, 3, 5, 10, 15 and 30 days after DENV infection. The panel includes: (**a**) interferon alpha (IFN-α), (**b**) interleukin-6 (IL-6), (**c**) monokine induced by IFN-gamma (MIG/CXCL9), (**d**) monocyte chemoattractant protein 1 (MCP-1/CCL2), (**e**) macrophage inflammatory protein 1-beta (MIP-1β/CCL4), (**f**) IL-1 receptor antagonist (IL-1RA), (**g**) C-X-C motif chemokine 10 (CXCL10/IP-10) and (**h**) perforin. Symbols connected with lines represent mean expression levels detected of each cytokine/chemokine per cohort over time: blue squares (ZIKVPF-10mo), orange squares (ZIKVPR-2mo) and black circles (Naïve). Error bars indicate the standard error of the mean (SEM) for each cohort per timepoint. Cutted line mark the limit of

detection for each individual cytokine/chemokine. Statistically significant differences between groups were calculated using Two-Way Anova adjusted for Tukey's multiple comparisons test including 8 families, and 3 comparisons per family. Significant multiplicity adjusted p values (* <0.05, ** <0.01, *** <0.001, **** <0.0001) are shown colored representing the cohort against that particular point where is a statistically significant difference between groups.

Figure 4 | Neutralization of DENV-2 and heterologous serotypes is higher in magnitude and durability by animals with mid-convalescence to ZIKV. The magnitude of the neutralizing antibody (NAb) response was determined (**a**) before and (**b**) 30 days after DENV infection by Plaque Reduction Neutralization Test (PRNT) against all DENV serotypes. (**c-f**) The durability of the neutralizing response was assessed measuring NAb titers up to 90 dpi against all DENV serotypes. Symbols connected with full lines indicate mean levels of NAb titers detected per cohort over time: blue squares (ZIKVPF-10mo), orange squares (ZIKVPR-2mo) and black circles (Naïve). Error bars represent the standard error of the mean (SEM). PRNT60: NAb titer capable of reduce 60% or more of DENV serotypes plaque-forming units (pfu) compared with the mock (control of virus without serum). A PRNT60 1:20 titer was considered positive, and <1:20 as a negative Neut titer. Dotted line mark <1:20 for negative results. Non-neutralizing titers (<1:20) were assigned with one-half of the limit of detection for graphs visualization (1:10). Statistically significant differences between groups were calculated using Two-Way Anova adjusted for Tukey's multiple comparisons test including 4 and 6 families for heterologous serotypes and DENV-2, respectively, and 3 comparisons per family. Significant multiplicity adjusted *p* values (* <0.05, ** <0.01, *** <0.001, **** <0.0001) are shown. Blue and orange asterisks represent significant difference between the corresponded ZIKV immune groups and naive group, and gray asterisks indicate a significant difference between ZIKV immune groups.

Figure 5 | ZIKV neutralization is more potent and durable in animals with mid-convalescence to ZIKV and is independent of the pre-infecting ZIKV strain. (**a**) NAb titers against ZIKV H/PF/2013 were determined by PRNT60 at baseline, 7, 15, 30, 60 and 90 days after DENV infection. Comparison of NAb titers between pre-infecting ZIKV strains was performed (**b**) before and (**c**) after DENV infection. Symbols connected with full lines indicate mean levels of NAb titers detected per cohort over time: blue squares (ZIKVPF-10mo), orange squares (ZIKVPR-2mo) and black circles (Naïve). Error bars represent the standard error of the mean (SEM). PRNT60: NAb titer capable of reduce 60% or more of ZIKV strains plaque-forming units (pfu) compared with the mock (control of virus without serum). A PRNT60 1:20 titer was considered positive, and <1:20 as a negative Neut titer. Dotted line mark <1:20 for negative results. Non-neutralizing titers (<1:20) were assigned with one-half of the limit of detection for graphs visualization (1:10). Statistically significant differences between groups were calculated using Two-Way Anova adjusted for Tukey's multiple comparisons test including 6 and 2 families for panel a and b-c, respectively, and 3 comparisons per family. Significant multiplicity adjusted p values (* <0.05, ** <0.01, *** <0.001, **** <0.0001) are shown. Blue and orange asterisks represent significant difference between the corresponded ZIKV-immune groups and naive group, and gray asterisks indicate a significant difference between ZIKV-immune groups.

Figure 6 | Effector and central memory T cells within CD4⁺ and CD8⁺ T cell compartments are activated after DENV infection. Activation (CD69$^+$) of effector memory (T-EM: CD3$^+$CD4$^+$CD28$^-$CD95$^+$) and central memory (T-CM: CD3$^+$CD4$^+$CD28$^+$CD95$^+$) T cells within (**a-c**) CD4$^+$ and (**d-f**) CD8$^+$ T cell compartments before and after DENV infection. Percent of cells were determined by immunophenotyping using flow cytometry (Supplementary Fig. 7 for gating strategy). Blue, orange and black squares represent T-EM for ZIKVPF-10mo, ZIKVPR-2mo and Naïve, respectively. Gray squares represent T-CM for each group. Short black lines mark mean value for each group per timepoint. Cutted line divide % of T-EM and T-CM cells quantified before and after DENV infection. Statistically significant differences within groups were determined using Two-Way Anova adjusted for Dunnett's multiple comparisons test (comparison of each group response at each timepoint versus baseline of the same group) including 2 families, and 7 comparisons per family. Significant differences are reported as multiplicity adjusted *p* values (* <0.05, ** <0.01, *** <0.001, **** <0.0001). Asterisks represent significant difference between the corresponded timepoint and baseline within the same group.

Figure 7 | Longevity of ZIKV immunity shapes the functional response of CD4⁺ and CD8⁺ T cells. T cell functional effector response was determined by the quantification (%) of (**a-d; m-p**) CD107a-expressing and (**e-h; q-t**) IFN-γ or (**i-l; u-x**) TNF-α producing CD4⁺ and CD8⁺ T cells before (0) and 30, 60 and 90 days after DENV infection. Responses to several peptide pools that encode for DENV and ZIKV envelope (E) proteins or ZIKV non-structural (NS) protein were quantified. After antigenic stimulation intracellular cytokine staining was performed using flow cytometry analysis (Supplementary Fig. 14 for gating strategy). Individual symbols represent each animal per antigenic stimulation over time: blue squares (ZIKVPF-10mo), orange squares (ZIKVPR-2mo) and black circles (Naïve). Short gray lines mark mean value for each group.

Statistically significant differences between groups were calculated using Two-Way Anova adjusted for Tukey's multiple comparisons test including 3 families, and 3 comparisons per family. Significant multiplicity adjusted p values (* <0.05, ** <0.01, *** <0.001, **** <0.0001) are shown. Asterisks represent significant difference between indicated groups.

Supplementary Information

Supplementary Figure 1 | Clinical status and vital signs kinetics in ZIKV-immune and naïve macaques. (**a**) Weight (kg) was measured at baseline, 1, 3, 5, 7, 9, 15, 30, 60 and 90 dpi. (**b**)

Temperature (°C) was monitored with an infrared device at baseline, 1-10, 15, 30, 60 and 90 dpi. Complete blood cell counts (CBC) parameters (thou/ul and/or % of total WBC) such as (**c**) white blood cells (WBC), (**d**) lymphocytes (LYM), (**e**) neutrophils (NEU), (**f**) monocytes (MON), and (**g**) platelets (PLT) were screened at baseline, 7, and 15 dpi. Comprehensive metabolic panel (CMP) was performed to assess levels (U/L) of (**h**) alkaline phosphatase (ALK PHOSPHATASE) and liver enzymes (**i**) aspartate transaminase (AST), and (**j**) alanine transaminase (ALT) at baseline, 7, 15 and 30 dpi. Normal range of AST and ALT are depicted for reference. (**k**) Age of rhesus macaques are depicted including the range of young adults for reference. Symbols represent mean level detected for each parameter per cohort per timepoint: blue squares (ZIKVPF-10mo), orange squares (ZIKVPR-2mo) and black circles (Naïve). Lines connect mean values detected over time. Error bars indicate the standard error of the mean (SEM) for each cohort per timepoint. Statistically significant differences between groups were determined using Two-Way Anova adjusted for Tukey's multiple comparisons test including 10, 15, 3, 4, and 3 families for panel a, b, c-g, h-j, and k, respectively, and 3 comparisons per family. For differences in ALT levels Two-Way Anova Dunnett's multiple comparisons test (comparison of each group response at each timepoint versus baseline of the same group) was performed including 3 families, and 3 comparisons per family due to divergence of non-specific levels between cohorts at baseline. Statistically differences are reported as multiplicity adjusted p values (* <0.05).

Supplementary Figure 2 | Previous ZIKV immunity modulates DENV RNAemia kinetics and is associated with a lower area under the curve. The area under the curve (AUC) was calculated using log-transformed values of DENV-2 RNAemia in ZIKV-immune and naïve animals. The total area by group is depicted on the graph as light blue, light orange and gray for ZIKVPF-10mo, ZIKVPR-2mo, and Naïve, respectively. Lines mark the mean value of genome copies per group per timepoint. A value of 1 was assigned to all samples below the LOD in order to calculate the means.

Supplementary Figure 3 | Serological cross-reactivity is boosted by ZIKV immunity. Levels of DENV (**a**) IgM and (**b**) IgG, and ZIKV (**c**) IgM, (**d**) IgG, (**e**) NS1-IgG and (**f**) EDIII-IgG were measured by ELISA at multiple timepoints before and after DENV infection. Symbols connected with full lines represent mean levels of Abs detected per cohort over time: blue squares (ZIKVPF-10mo), orange squares (ZIKVPR-2mo) and black circles (Naïve). Panel e includes additional timepoints before DENV infection for ZIKV-immune groups: 30, 60, 90 and 180 days after ZIKV (H/PF/2013) infection for the ZIKVPF-10mo group, and 30 days after ZIKV (PRVABC59) infection for the ZIKVPR-2mo group. Error bars indicate the standard error of the mean (SEM) and dotted line mark the limit of detection for each individual ELISA. Results were read at OD 450, 405 or using ISR (Immune Status Ratio) following manufacturer's instructions. Statistically significant differences between groups were calculated using Two-Way Anova adjusted for Tukey's multiple comparisons test including 5, 6, 9, and 4 families, and 3 comparisons per family. Significant multiplicity adjusted p values (* <0.05, ** <0.01, *** <0.001, **** <0.0001) are shown. Blue and orange asterisks represent significant difference between the corresponded ZIKV immune groups and naive group, and gray asterisks indicate a significant difference between ZIKV immune groups.

Supplementary Figure 4 | Neutralization kinetics against DENV-2 of ZIKV middle- and early-convalescent, and naïve animals. Percentage of DENV-2 neutralization of each animal per

group calculated by the transformation of PRNT60 Neut 2-fold titers into Log10 (1/serum dilution), and sigmoidal-dose response curves were generated. Each column of panels represent the % of DENV-2 neutralization for each group (ZIKVPF-10mo: blue squares/curves; ZIKVPR-2mo: orange squares/curves; Naïve: black circles/curves) and each row of panels represent a timepoint before and after DENV infection (baseline, 7, 15, 30, 60, 90 dpi).

Supplementary Figure 5 | Neutralization kinetics against ZIKV of ZIKV middle- and early-convalescent animals, and naïve animals. Percentage of ZIKV (H/PF/2013) neutralization of

each animal per group calculated by the transformation of PRNT60 Neut 2-fold titers into Log10 (1/serum dilution), and sigmoidal-dose response curves were generated. Each column of panels represent the % of ZIKV neutralization for each group (ZIKVPF-10mo: blue squares/curves; ZIKVPR-2mo: orange squares/curves; Naïve: black circles/curves) and each row of panels represent a timepoint before and after DENV infection (baseline, 7, 15, 30, 60, 90 dpi).

Supplementary Figure 6 | Similar neutralizing titers induced by two different ZIKV strains in rhesus macaques. NAb titers against H/PF/2013 and PRVABC59 ZIKV strains for ZIKVPF-10mo and ZIKVPR-2mo groups, respectively, were determined by PRNT60 at 30 and 60 after ZIKV infection. Symbols indicate levels of NAb titers detected per animal: blue squares (ZIKVPF-10mo), and orange squares (ZIKVPR-2mo). Error bars represent the standard error of the mean (SEM). PRNT60: NAb titer capable of reduce 60% or more of ZIKV strains plaque-forming units (pfu) compared with the mock (control of virus without serum). A PRNT60 1:20 titer was considered positive, and <1:20 as a negative Neut titer. Dotted line mark <1:20 for negative results. Statistically significant differences (ns: not significant) between two groups were calculated using Two-Way Anova corrected for Sidak's multiple comparisons test including 1 family, and 2 comparisons within the family.

Supplementary Figure 7 | Gating strategy for immunophenotyping and activation of B cells, and memory T cell subpopulations.

Supplementary Figure 8 | Gating strategy for immunophenotyping of plasmacytoid and myeloid dendritic cells.

Supplementary Figure 9 | Gating strategy for Natural killer cell subpopulations.

Supplementary Figure 10 | Dendritic cells subsets modulation depends on the presence or absence of ZIKV immunity. The frequency (% of total PBMCs) of dendritic cells (DCs) subsets including (**a**) plasmacytoid (pDCs: Lin-HLA-DR$^+$CD123$^+$) and (**b**) myeloid (mDCs: Lin-HLA-DR$^+$CD11c$^+$) was assessed before and up to 10 days after DENV infection by immunophenotyping using flow cytometry analysis. Symbols represent individual animals per group for each timepoint: blue squares (ZIKVPF-10mo), orange squares (ZIKVPR-2mo) and black circles (Naïve). Short gray lines depict mean value for each group detected overtime. Cutted line divide % of DCs quantified before and after DENV infection. Statistically significant differences within groups were determined using Two-Way Anova Dunnett's multiple comparisons test (comparison of each group response at each timepoint versus baseline of the same group) including 3 families, and 5 comparisons per family. Significant differences are reported as multiplicity adjusted *p* values (* <0.05, ** <0.01, *** <0.001, **** <0.0001). Asterisks represent significant difference between the corresponded timepoint and baseline within the same group.

Supplementary Figure 11 | Natural killer cell subpopulations and their differential receptors expression in presence or absence of ZIKV immunity. Natural killer (NK) cell subpopulations and the relative expression of multiple NK receptors within each subpopulation: (**a-d**) NKCD8, (**e-h**) NKCD56, (**i-l**) NKG2A, (**m-p**) NKp30 and (**q-t**) NKp46 were quantified by immunophenotyping using flow cytometry analysis before and up to 10 days after DENV infection. Individual symbols represent each animal per group over time: blue squares (ZIKVPF-10mo), orange squares

(ZIKVPR-2mo) and black circles (Naïve). Short gray lines mark mean value for each group. Cutted line divide % of NK cells quantified before and after DENV infection. Statistically significant differences within groups were determined using Two-Way Anova Dunnett's multiple comparisons test (comparison of each group response at each timepoint versus baseline of the same group) including 3 families, and 5 comparisons per family. Significant differences are reported as multiplicity adjusted p values (* <0.05, ** <0.01, *** <0.001, **** <0.0001). Asterisks represent significant difference between the corresponded timepoint and baseline within the same group. ND (Not Done) in panels *8o* and *8s* refers that for ZIKVPF-10mo and Naïve groups the NKp30$^+$NKp46$^+$ and NKp46$^+$NKp30$^+$ subpopulations were not measured.

Supplementary Figure 12 | B cells proliferation and activation levels are higher in ZIKV middle-convalescent macaques. The (**a**) total (% of total Lymphocytes), (**b**) activated, (**c**) proliferating, and (**d**) proliferating/activated B cells (% of total B cells) were determined at baseline and following DENV infection by immunophenotyping using flow cytometry analysis. B cells proliferation and activation were monitored since baseline up to 10 and 30 dpi, respectively. Symbols represent individual animals per group for each timepoint: blue squares (ZIKVPF-10mo), orange squares (ZIKVPR-2mo) and black circles (Naïve). Short gray lines depict mean value of B cells percent in each group of animals per timepoint. Cutted line divide % of B cells quantified before and after DENV infection. Statistically significant differences within groups were determined using Two-Way Anova Dunnett's multiple comparisons test (comparison of each group response at each timepoint versus baseline of the same group) including 3 families, and 7 and 5 comparisons per family in panels a-b and c-d, respectively. Significant differences are reported as multiplicity adjusted *p* values (* <0.05, ** <0.01). Asterisks represent significant difference between the corresponded timepoint and baseline within the same group.

Supplementary Figure 13 | T cells frequency before and after DENV infection. The (**a**) total T cells (% of total Lymphocytes), (**b**) CD4[+] and (**c**) CD8[+] T cell compartments (% of total T cells) frequencies were quantified at baseline and following DENV infection up to 30 dpi by immunophenotyping using flow cytometry. Symbols represent individual animals per group for each timepoint: blue squares (ZIKVPF-10mo), orange squares (ZIKVPR-2mo) and black circles (Naïve). Short gray lines mark mean value of T cells percent in each cohort per timepoint. Cutted line divide % of T cells quantified before and after DENV infection. Statistically significant differences within groups were determined using Two-Way Anova Dunnett's multiple comparisons test (comparison of each group response at each timepoint versus baseline of the same group) including 3 families, and 7 comparisons per family.

Supplementary Figure 14 | Gating strategy for CD4+ and CD8+ T cell functional response.

Supplementary Table 1 | Sequence alignment and amino acid identity of ZIKV strains PRVABC59 and H/PF/2013.

Pairwise alignment of both ZIKV strains sequences. (Sequences downloaded from ViPR database and global alignment was performed using Blosum62 in Genious Software).	>99.99% amino acid identity
Envelope (E) protein region of both ZIKV strains.	Identical
Amino acids residues changes between both ZIKV strains. Marked in red within sequences. From ZIKV-PR → ZIKV-PF	$T_{80}\rightarrow I$ (Capsid) $G_{892}\rightarrow W$ (NS1) $V_{2611}\rightarrow A$ (NS5) $V_{2634}\rightarrow M$ (NS5)
ZIKV-PRVABC59 Accession number: KX377337	MKNPKKKSGGFRIVNMLKRGVARVSPFGGLKRLPAGLLLGHGPIRMVLAI LAFLRFTAIKPSLGLINRWGSVGKKEAMETIKKFKKDLAAMLRIINARKE KKRRGADTSVGIVGLLLTTAMAAEVTRRGSAYYMYLDRNDAGEAISFPTT LGMNKCYIQIMDLGHMCDATMSYECPMLDEGVEPDDVDCWCNTTSTWVVY GTCHHKKGEARRSRRAVTLPSHSTRKLQTRSQTWLESREYTKHLIRVENW IFRNPGFALAAAAIAWLLGSSTSQKVIYLVMILLIAPAYSIRCIGVSNRD FVEGMSGGTWVDVVLEHGGCVTVMAQDKPTVDIELVTTTVSNMAEVRSYC YEASISDMASDSRCPTQGEAYLDKQSDTQYVCKRTLVDRGWGNGCGLFGK GSLVTCAKFACSKKMTGKSIQPENLEYRIMLSVHGSQHSGMIVNDTGHET DENRAKVEITPNSPRAEATLGGFGSLGLDCEPRTGLDFSDLYYLTMNNKH WLVHKEWFHDIPLPWHAGADTGTPHWNNKEALVEFKDAHAKRQTVVVLGS QEGAVHTALAGALEAEMDGAKGRLSSGHLKCRLKMDKLRLKGVSYSLCTA AFTFTKIPAETLHGTVTVEVQYAGTDGPCKVPAQMAVDMQTLTPVGRLIT ANPVITESTENSKMMLELDPPFGDSYIVIGVGEKKITHHWHRSGSTIGKA FEATVRGAKRMAVLGDTAWDFGSVGGALNSLGKGIHQIFGAAFKSLFGGM SWFSQILIGTLLMWLGLNTKNGSISLMCLALGGVLIFLSTAVSADVGCSV DFSKKETRCGTGVFVYNDVEAWRDRYKYHPDSPRRLAAAVKQAWEDGICG ISSVSRMENIMWRSVEGELNAILEENGVQLTVVVGSVKNPMCRGPQRLPV PVNELPHGWKAWGKSYFVRAAKTNNSFVVDGDTLKECPLKHRAWNSFLVE DHGFGVFHTSVWLKVREDYSLECDPAVIGTAVKGKEAVHSDLGYWIESEK NDTWRLKRAHLIEMKTCEWPKSHTLWTDGIEESDLIIPKSLAGPLSHHNT REGYRTQMKGPWHSEELEIRFEECPGTKVHVEETCGTRGPSLRSTTASGR VIEEWCCRECTMPPLSFRAKDGCWYGMEIRPRKEPESNLVRSMVTAGSTD HMDHFSLGVLVILLMVQEGLKKRMTTKIIISTSMAVLVAMILGGFSMSDL AKLAILMGATFAEMNTGGDVAHLALIAAFKVRPALLVSFIFRANWTPRES MLLALASCLLQTAISALEGDLMVLINGFALAWLAIRAMVVPRTDNITLAI LAALTPLARGTLLVAWRAGLATCGGFMLLSLKGKGSVKKNLPFVMALGLT AVRLVDPINVVGLLLLTRSGKRSWPPSEVLTAVGLICALAGGFAKADIEM AGPMAAVGLLIVSYVVSGKSVDMYIERAGDITWEKDAEVTGNSPRLDVAL

	DESGDFSLVEDDGPPMREIILKVVLMTICGMNPIAIPFAAGAWYVYVKTG KRSGALWDVPAPKEVKKGETTDGVYRVMTRRLLGSTQVGVGVMQEGVFHT MWHVTKGSALRSGEGRLDPYWGDVKQDLVSYCGPWKLDAAWDGHSEVQLL AVPPGERARNIQTLPGIFKTKDGDIGAVALDYPAGTSGSPILDKCGRVIG LYGNGVVIKNGSYVSAITQGRREEETPVECFEPSMLKKKQLTVLDLHPGA GKTRRVLPEIVREAIKTRLRTVILAPTRVVAAEMEEALRGLPVRYMTTAV NVTHSGTEIVDLMCHATFTSRLLQPIRVPNYNLYIMDEAHFTDPSSIAAR GYISTRVEMGEAAAIFMTATPPGTRDAFPDSNSPIMDTEVEVPERAWSSG FDWVTDHSGKTVWFVPSVRNGNEIAACLTKAGKRVIQLSRKTFETEFQKT KHQEWDFVVTTDISEMGANFKADRVIDSRRCLKPVILDGERVILAGPMPV THASAAQRRGRIGRNPNKPGDEYLYGGGCAETDEDHAHWLEARMLLDNIY LQDGLIASLYRPEADKVAAIEGEFKLRTEQRKTFVELMKRGDLPVWLAYQ VASAGITYTDRRWCFDGTTNNTIMEDSVPAEVWTRHGEKRVLKPRWMDAR VCSDHAALKSFKEFAAGKRGAAFGVMEALGTLPGHMTERFQEAIDNLAVL MRAETGSRPYKAAAAQLPETLETIMLLGLLGTVSLGIFFVLMRNKGIGKM GFGMVTLGASAWLMWLSEIEPARIACVLIVVFLLLVVLIPEPEKQRSPQD NQMAIIIMVAVGLLGLITANELGWLERTKSDLSHLMGRREEGATIGFSMD IDLRPASAWAIYAALTTFITPAVQHAVTTSYNNYSLMAMATQAGVLFGMG KGMPFYAWDFGVPLLMIGCYSQLTPLTLIVAIILLVAHYMYLIPGLQAAA ARAAQKRTAAGIMKNPVVDGIVVTDIDTMTIDPQVEKKMGQVLLIAVAVS SAILSRTAWGWGEAGALITAATSTLWEGSPNKYWNSSTATSLCNIFRGSY LAGASLIYTVTRNAGLVKRRGGGTGETLGEKWKARLNQMSALEFYSYKKS GITEVCREEARRALKDGVATGGHAVSRGSAKLRWLVERGYLQPYGKVIDL GCGRGGWSYYV<mark>A</mark>TIRKVQEVKGYTKGGPGHEEP<mark>V</mark>LVQSYGWNIVRLKSGV DVFHMAAEPCDTLLCDIGESSSSPEVEEARTLRVLSMVGDWLEKRPGAFC IKVLCPYTSTMMETLERLQRRYGGGLVRVPLSRNSTHEMYWVSGAKSNTI KSVSTTSQLLLGRMDGPRRPVKYEEDVNLGSGTRAVVSCAEAPNMKIIGN RIERIRSEHAETWFFDENHPYRTWAYHGSYEAPTQGSASSLINGVVRLLS KPWDVVTGVTGIAMTDTTPYGQQRVFKEKVDTRVPDPQEGTRQVMSMVSS WLWKELGKHKRPRVCTKEEFINKVRSNAALGAIFEEEKEWKTAVEAVNDP RFWALVDKEREHHLRGECQSCVYNMMGKREKKQGEFGKAKGSRAIWYMWL GARFLEFEALGFLNEDHWMGRENSGGGVEGLGLQRLGYVLEEMSRIPGGR MYADDTAGWDTRISRFDLENEALITNQMEKGHRALALAIIKYTYQNKVVK VLRPAEKGKTVMDIISRQDQRGSGQVVTYALNTFTNLVVQLIRNMEAEEV LEMQDLWLLRRSEKVTNWLQSNGWDRLKRMAVSGDDCVVKPIDDRFAHAL RFLNDMGKVRKDTQEWKPSTGWDNWEEVPFCSHHFNKLHLKDGRSIVVPC RHQDELIGRARVSPGAGWSIRETACLAKSYAQMWQLLYFHRRDLRLMANA ICSSVPVDWVPTGRTTWSIHGKGEWMTTEDMLVVWNRVWIEENDHMEDKT PVTKWTDIPYLGKREDLWCGSLIGHRPRTTWAENIKNTVNMVRRIIGDEE KYMDYLSTQVRYLGEEGSTPGVL
ZIKV H/PF2013 Accession number: KJ776791	MKNPKKKSGGFRIVNMLKRGVARVSPFGGLKRLPAGLLLGHGPIRMVLAI LAFLRFTAIKPSLGLINRWGSVGKKEAME<mark>T</mark>IKKFKKDLAAMLRIINARKE KKRRGADTSVGIVGLLLTTAMAAEVTRRGSAYYMYLDRNDAGEAISFPTT LGMNKCYIQIMDLGHMCDATMSYECPMLDEGVEPDDVDCWCNTTSTWVVY GTCHHKKGEARRSRRAVTLPSHSTRKLQTRSQTWLESREYTKHLIRVENW IFRNPGFALAAAAIAWLLGSSTSQKVIYLVMILLIAPAYSIRCIGVSNRD FVEGMSGGTWVDVVLEHGGCVTVMAQDKPTVDIELVTTTVSNMAEVRSYC YEASISDMASDSRCPTQGEAYLDKQSDTQYVCKRTLVDRGWGNGCGLFGK GSLVTCAKFACSKKMTGKSIQPENLEYRIMLSVHGSQHSGMIVNDTGHET DENRAKVEITPNSPRAEATLGGFGSLGLDCEPRTGLDFSDLYYLTMNNKH WLVHKEWFHDIPLPWHAGADTGTPHWNNKEALVEFKDAHAKRQTVVVLGS QEGAVHTALAGALEAEMDGAKGRLSSGHLKCRLKMDKLRLKGVSYSLCTA AFTFTKIPAETLHGTVTVEVQYAGTDGPCKVPAQMAVDMQTLTPVGRLIT ANPVITESTENSKMMLELDPPFGDSYIVIGVGEKKITHHWHRSGSTIGKA FEATVRGAKRMAVLGDTAWDFGSVGGALNSLGKGIHQIFGAAFKSLFGGM SWFSQILIGTLLMWLGLNTKNGSISLMCLALGGVLIFLSTAVSADVGCSV DFSKKETRCGTGVFVYNDVEAWRDRYKYHPDSPRRLAAAVKQAWEDGICG

ISSVSRMENIMWRSVEGELNAILEENGVQLTVVVGSVKNPMWRGPQRLPV
PVNELPHGWKAWGKSYFVRAAKTNNSFVVDGDTLKECPLKHRAWNSFLVE
DHGFGVFHTSVWLKVREDYSLECDPAVIGTAVKGKEAVHSDLGYWIESEK
NDTWRLKRAHLIEMKTCEWPKSHTLWTDGIEESDLIIPKSLAGPLSHHNT
REGYRTQMKGPWHSEELEIRFEECPGTKVHVEETCGTRGPSLRSTTASGR
VIEEWCCRECTMPPLSFRAKDGCWYGMEIRPRKEPESNLVRSMVTAGSTD
HMDHFSLGVLVILLMVQEGLKKRMTTKIIISTSMAVLVAMILGGFSMSDL
AKLAILMGATFAEMNTGGDVAHLALIAAFKVRPALLVSFIFRANWTPRES
MLLALASCLLQTAISALEGDLMVLINGFALAWLAIRAMVVPRTDNITLAI
LAALTPLARGTLLVAWRAGLATCGGFMLLSLKGKGSVKKNLPFVMALGLT
AVRLVDPINVVGLLLLTRSGKRSWPPSEVLTAVGLICALAGGFAKADIEM
AGPMAAVGLLIVSYVVSGKSVDMYIERAGDITWEKDAEVTGNSPRLDVAL
DESGDFSLVEDDGPPMREIILKVVLMTICGMNPIAIPFAAGAWYVYVKTG
KRSGALWDVPAPKEVKKGETTDGVYRVMTRRLLGSTQVGVGVMQEGVFHT
MWHVTKGSALRSGEGRLDPYWGDVKQDLVSYCGPWKLDAAWDGHSEVQLL
AVPPGERARNIQTLPGIFKTKDGDIGAVALDYPAGTSGSPILDKCGRVIG
LYGNGVVIKNGSYVSAITQGRREEETPVECFEPSMLKKKQLTVLDLHPGA
GKTRRVLPEIVREAIKTRLRTVILAPTRVVAAEMEEALRGLPVRYMTTAV
NVTHSGTEIVDLMCHATFTSRLLQPIRVPNYNLYIMDEAHFTDPSSIAAR
GYISTRVEMGEAAAIFMTATPPGTRDAFPDSNSPIMDTEVEVPERAWSSG
FDWVTDHSGKTVWFVPSVRNGNEIAACLTKAGKRVIQLSRKTFETEFQKT
KHQEWDFVVTTDISEMGANFKADRVIDSRRCLKPVILDGERVILAGPMPV
THASAAQRRGRIGRNPNKPGDEYLYGGGCAETDEDHAHWLEARMLLDNIY
LQDGLIASLYRPEADKVAAIEGEFKLRTEQRKTFVELMKRGDLPVWLAYQ
VASAGITYTDRRWCFDGTTNNTIMEDSVPAEVWTRHGEKRVLKPRWMDAR
VCSDHAALKSFKEFAAGKRGAAFGVMEALGTLPGHMTERFQEAIDNLAVL
MRAETGSRPYKAAAAQLPETLETIMLLGLLGTVSLGIFFVLMRNKGIGKM
GFGMVTLGASAWLMWLSEIEPARIACVLIVVFLLLVVLIPEPEKQRSPQD
NQMAIIIMVAVGLLGLITANELGWLERTKSDLSHLMGRREEGATIGFSMD
IDLRPASAWAIYAALTTFITPAVQHAVTTSYNNYSLMAMATQAGVLFGMG
KGMPFYAWDFGVPLLMIGCYSQLTPLTLIVAIILLVAHYMYLIPGLQAAA
ARAAQKRTAAGIMKNPVVDGIVVTDIDTMTIDPQVEKKMGQVLLIAVAVS
SAILSRTAWGWGEAGALITAATSTLWEGSPNKYWNSSTATSLCNIFRGSY
LAGASLIYTVTRNAGLVKRRGGGTGETLGEKWKARLNQMSALEFYSYKKS
GITEVCREEARRALKDGVATGGHAVSRGSAKLRWLVERGYLQPYGKVIDL
GCGRGGWSYYAATIRKVQEVKGYTKGGPGHEEPMLVQSYGWNIVRLKSGV
DVFHMAAEPCDTLLCDIGESSSSPEVEEARTLRVLSMVGDWLEKRPGAFC
IKVLCPYTSTMMETLERLQRRYGGGLVRVPLSRNSTHEMYWVSGAKSNTI
KSVSTTSQLLLGRMDGPRRPVKYEEDVNLGSGTRAVVSCAEAPNMKIIGN
RIERIRSEHAETWFFDENHPYRTWAYHGSYEAPTQGSASSLINGVVRLLS
KPWDVVTGVTGIAMTDTTPYGQQRVFKEKVDTRVPDPQEGTRQVMSMVSS
WLWKELGKHKRPRVCTKEEFINKVRSNAALGAIFEEEKEWKTAVEAVNDP
RFWALVDKEREHHLRGECQSCVYNMMGKREKKQGEFGKAKGSRAIWYMWL
GARFLEFEALGFLNEDHWMGRENSGGGVEGLGLQRLGYVLEEMSRIPGGR
MYADDTAGWDTRISRFDLENEALITNQMEKGHRALALAIIKYTYQNKVVK
VLRPAEKGKTVMDIISRQDQRGSGQVVTYALNTFTNLVVQLIRNMEAEEV
LEMQDLWLLRRSEKVTNWLQSNGWDRLKRMAVSGDDCVVKPIDDRFAHAL
RFLNDMGKVRKDTQEWKPSTGWDNWEEVPFCSHHFNKLHLKDGRSIVVPC
RHQDELIGRARVSPGAGWSIRETACLAKSYAQMWQLLYFHRRDLRLMANA
ICSSVPVDWVPTGRTTWSIHGKGEWMTTEDMLVVWNRVWIEENDHMEDKT
PVTKWTDIPYLGKREDLWCGSLIGHRPRTTWAENIKNTVNMVRRIIGDEE
KYMDYLSTQVRYLGEEGSTPGVL

Supplementary Table 2 | Antibody panel for Immunophenotyping.

Cell Subset	Ab	Clone	Dye	Company	Cat. #
B / T cells	CD20	2H7	PacificBlue	BioLegend	302328
	CD3	10D12	PE-Vio770	Miltenyi	130-104-202
	CD4	M-T466	PerCP	Miltenyi	130-101-147
	CD8	BW135/80	VioGreen	Miltenyi	130-096-902
	CD28	15E8	APC-Vio770	Miltenyi	130-104-278
	CD69	FN50	PE	BD	557050
	CD95	DX2	APC	Miltenyi	130-092-417
	Ki67	B56	Alexa 488	BD	558616
NK	CD3	10D12	APC	Miltenyi	130-091-998
	CD16	VEP13	APC-Vio770	Miltenyi	130-096-655
	CD56	AF12-7H3	PE	Miltenyi	130-090-755
	CD14	M5E2	V500	BD	561391
	CD8	SK1	BV421	BioLegend	344748
	NKp30	AF29-4D12	PE-Vio770	Miltenyi	130-104-116
	NKp46	BAB281	PC5	Beck-Coulter	A66902
	NK2GA	REA110	FITC	Miltenyi	130-098-818
DC	CD20	2H7	FITC	BD	555622
	CD3	SP34		BD	556611
	CD14	M5E2		BD	555397
	CD16	3G8		BD	555406
	NKG2A	REA110		Miltenyi	130-098-818
	CD8	SK1		BioLegend	344704
	HLA DR	REA 805	VioGreen	Miltenyi	130-111-795
	CD123	7G3	APC	BD	560087
	CD11c	3.9	PE/Cy7	BioLegend	301608

Supplementary Table 3 | Antibody panel for T cell functional response assessment.

Marker	Stain	Clone	Catalog Number	Vendor	Dilution
CD4	PerCP-Cy-5.5	SK3	566316	BD Biosciences	1:25
CD8β	PE	ECD	6607123	Beckman-Coulter	1:20
CD3	Pacific Blue	SP34-2	558124	BD Biosciences	1:30
CD20	BV605	2H7	563783	BD Biosciences	1:30
CD107a	FITC	H4A3	555800	BD Biosciences	1:10
CD28	PE-Cy-5	CD28.2	555730	BD Biosciences	1:10
CD95	BV510	DX2	305640	Biolegend	1:30
IFN-γ	APC	B27	554702	BD Biosciences	1:30
TNF-α	PE-Cy-7	Mab11	557647	BD Biosciences	1:30

Supplementary Table 4 | Peptide sequences for stimulation of T cell functional response.

Dengue Virus Type 2 Peptides

Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence
1	MRCIGISNRDFVEGV	29	AWLVHRQWFLDLPLPWL	57	MRGAKRMAILGDTAWDF
2	ISNRDFVEGVSGGSWVDI	30	WFLDLPLPWLPGADTQGSNW	58	AILGDTAWDFGSLGGVF
3	GVSGGSWVDIVLEHGSCV	31	PGADTQGSNWIQKETLV	59	WDFGSLGGVFTSIGKALH
4	DIVLEHGSCVTTMAKNK	32	SNWIQKETLVTFKNPHAK	60	VFTSIGKALHQVFGAIY
5	SCVTTMAKNKPTLDFELI	33	LVTFKNPHAKKQDVVVL	61	ALHQVFGAIYGAAFSGV
6	NKPTLDFELIETEAKQPA	34	HAKKQDVVVLGSQEGAMH	62	AIYGAAFSGVSWIMKILI
7	LIETEAKQPATLRKYCI	35	VLGSQEGAMHTALTGA	63	GVSWIMKILIGVIITWI
8	KQPATLRKYCIEAKL	36	GAMHTALTGATEIQM	64	ILIGVIITWIGMNSR
9	LRKYCIEAKLTNTTTDSR	37	ALTGATEIQMSSGNLLF	65	IITWIGMNSRSTSLSVSL
10	KLTNTTTDSRCPTQGEPSL	38	IQMSSGNLLFTGHLKCRL	66	SRSTSLSVSLVLVGVVTL
11	RCPTQGEPSLNEEQDKRF	39	LFTGHLKCRLRMDKLQLK	67	SLVLVGVVTLYLGVMVQA
12	SLNEEQDKRFVCKHSMV	40	RLRMDKLQLKGMSYSM		
13	KRFVCKHSMVDRGWGNGCGL	41	LQLKGMSYSMCTGKFKVV		
14	DRGWGNGCGLFGKGGIV	42	SMCTGKFKVVKEIAETQH		
15	CGLFGKGGIVTCAMFTCK	43	VVKEIAETQHGTIVIRV		
16	IVTCAMFTCKKNMKGKVV	44	TQHGTIVIRVQYEGDGSPCK		
17	CKKNMKGKVVQPENLEY	45	VQYEGDGSPCKIPFEIM		
18	KVVQPENLEYTIVITPH	46	SPCKIPFEIMDLEKRHVL		
19	LEYTIVITPHSGEEHAV	47	IMDLEKRHVLGRLITV		
20	TPHSGEEHAVGNDTGKH	48	RHVLGRLITVNPIVTEK		
21	HAVGNDTGKHGKEIKI	49	ITVNPIVTEKDSPVNIEA		
22	TGKHGKEIKITPQSSI	50	EKDSPVNIEAEPPFGDSY		
23	EIKITPQSSITEAELTGY	51	EAEPPFGDSYIIIGV		
24	SITEAELTGYGTVTM	52	FGDSYIIIGVEPGQLKL		
25	ELTGYGTVTMECSPRTGL	53	IGVEPGQLKLNWFKK		
26	TMECSPRTGLDFNEMVLL	54	GQLKLNWFKKGSSIGQMI		
27	GLDFNEMVLLQMENKAWL	55	KKGSSIGQMIETTMRGAK		
28	LLQMENKAWLVHRQWFL	56	MIETTMRGAKRMAIL		

Supplementary Table 4 | Continuation

Zika Virus Envelope Peptides

Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence
ZIKV59	IRCIGVSNRDFVEGM	ZIKV87	LSVHGSQHSGMIVND	ZIKV115	KGRLSSGHLKCRLKM
ZIKV60	VSNRDFVEGMSGGTW	ZIKV88	SQHSGMIVNDTGHET	ZIKV116	SGHLKCRLKMDKLRL
ZIKV61	FVEGMSGGTWVDVVL	ZIKV89	MIVNDTGHETDENRA	ZIKV117	CRLKMDKLRLKGVSY
ZIKV62	SGGTWVDVVLEHGGC	ZIKV90	TGHETDENRAKVEIT	ZIKV118	DKLRLKGVSYSLCTA
ZIKV63	VDVVLEHGGCVTVMA	ZIKV91	DENRAKVEITPNSPR	ZIKV119	KGVSYSLCTAAFTFT
ZIKV64	EHGGCVTVMAQDKPT	ZIKV92	KVEITPNSPRAEATL	ZIKV120	SLCTAAFTFTKIPAE
ZIKV65	VTVMAQDKPTVDIEL	ZIKV93	PNSPRAEATLGGFGS	ZIKV121	AFTFTKIPAETLHGT
ZIKV66	QDKPTVDIELVTTTV	ZIKV94	AEATLGGFGSLGLDC	ZIKV122	KIPAETLHGTVTVEV
ZIKV67	VDIELVTTTVSNMAE	ZIKV95	GGFGSLGLDCEPRTG	ZIKV123	TLHGTVTVEVQYAGT
ZIKV68	VTTTVSNMAEVRSYC	ZIKV96	LGLDCEPRTGLDFSD	ZIKV124	VTVEVQYAGTDGPCK
ZIKV69	SNMAEVRSYCYEASI	ZIKV97	EPRTGLDFSDLYYLT	ZIKV125	QYAGTDGPCKVPAQM
ZIKV70	VRSYCYEASISDMAS	ZIKV98	LDFSDLYYLTMNNKH	ZIKV126	DGPCKVPAQMAVDMQ
ZIKV71	YEASISDMASDSRCP	ZIKV99	LYYLTMNNKHWLVHK	ZIKV127	VPAQMAVDMQTLTPV
ZIKV72	SDMASDSRCPTQGEA	ZIKV100	MNNKHWLVHKEWFHD	ZIKV128	AVDMQTLTPVGRLIT
ZIKV73	DSRCPTQGEAYLDKQ	ZIKV101	WLVHKEWFHDIPLPW	ZIKV129	TLTPVGRLITANPVI
ZIKV74	TQGEAYLDKQSDTQY	ZIKV102	EWFHDIPLPWHAGAD	ZIKV130	GRLITANPVITESTE
ZIKV75	YLDKQSDTQYVCKRT	ZIKV103	IPLPWHAGADTGTPH	ZIKV131	ANPVITESTENSKMM
ZIKV76	SDTQYVCKRTLVDRG	ZIKV104	HAGADTGTPHWNNKE	ZIKV132	TESTENSKMMLELDP
ZIKV77	VCKRTLVDRGWGNGC	ZIKV105	TGTPHWNNKEALVEF	ZIKV133	NSKMMLELDPPFGDS
ZIKV78	LVDRGWGNGCGLFGK	ZIKV106	WNNKEALVEFKDAHA	ZIKV134	LELDPPFGDSYIVIG
ZIKV79	WGNGCGLFGKGSLVT	ZIKV107	ALVEFKDAHAKRQTV	ZIKV135	PFGDSYIVIGVGEKK
ZIKV80	GLFGKGSLVTCAKFA	ZIKV108	KDAHAKRQTVVVLGS	ZIKV136	YIVIGVGEKKITHHW
ZIKV81	GSLVTCAKFACSKKM	ZIKV109	KRQTVVVLGSQEGAV	ZIKV137	VGEKKITHHWHRSGS
ZIKV82	CAKFACSKKMTGKSI	ZIKV110	VVLGSQEGAVHTALA	ZIKV138	ITHHWHRSGSTIGKA
ZIKV83	CSKKMTGKSIQPENL	ZIKV111	QEGAVHTALAGALEA	ZIKV139	HRSGSTIGKAFEATV
ZIKV84	TGKSIQPENLEYRIM	ZIKV112	HTALAGALEAEMDGA	ZIKV140	TIGKAFEATVRGAKR
ZIKV85	QPENLEYRIMLSVHG	ZIKV113	GALEAEMDGAKGRLS	ZIKV141	FEATVRGAKRMAVLG
ZIKV86	EYRIMLSVHGSQHSG	ZIKV114	EMDGAKGRLSSGHLK	ZIKV142	RGAKRMAVLGDTAWD

Supplementary Table 4 | Continuation

Peptide	Amino Acid Sequence
ZIKV143	MAVLGDTAWDFGSVG
ZIKV144	DTAWDFGSVGGALNS
ZIKV145	FGSVGGALNSLGKGI
ZIKV146	GALNSLGKGIHQIFG
ZIKV147	LGKGIHQIFGAAFKS
ZIKV148	HQIFGAAFKSLFGGM
ZIKV149	AAFKSLFGGMSWFSQ
ZIKV150	LFGGMSWFSQILIGT
ZIKV151	SWFSQILIGTLLMWL
ZIKV152	ILIGTLLMWLGLNTK
ZIKV153	LLMWLGLNTKNGSIS
ZIKV154	GLNTKNGSISLMCLA
ZIKV155	NGSISLMCLALGGVL
ZIKV156	LMCLALGGVLIFLST
ZIKV157	LGGVLIFLSTAVSAD
ZIKV158	IFLSTAVSADVGCSV
ZIKV159	AVSADVGCSVDFSKK

Supplementary Table 4 | Continuation

Zika Virus Non-Structural Peptides

Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence
ZIKV160	VGCSVDFSKKETRCG	ZIKV188	ECPLKHRAWNSFLVE	ZIKV216	GTKVHVEETCGTRGP
ZIKV161	DFSKKETRCGTGVFV	ZIKV189	HRAWNSFLVEDHGFG	ZIKV217	VEETCGTRGPSLRST
ZIKV162	ETRCGTGVFVYNDVE	ZIKV190	SFLVEDHGFGVFHTS	ZIKV218	GTRGPSLRSTTASGR
ZIKV163	TGVFVYNDVEAWRDR	ZIKV191	DHGFGVFHTSVWLKV	ZIKV219	SLRSTTASGRVIEEW
ZIKV164	YNDVEAWRDRYKYHP	ZIKV192	VFHTSVWLKVREDYS	ZIKV220	TASGRVIEEWCCREC
ZIKV165	AWRDRYKYHPDSPRR	ZIKV193	VWLKVREDYSLECDP	ZIKV221	VIEEWCCRECTMPPL
ZIKV166	YKYHPDSPRRLAAAV	ZIKV194	REDYSLECDPAVIGT	ZIKV222	CCRECTMPPLSFRAK
ZIKV167	DSPRRLAAAVKQAWE	ZIKV195	LECDPAVIGTAVKGK	ZIKV223	TMPPLSFRAKDGCWY
ZIKV168	LAAAVKQAWEDGICG	ZIKV196	AVIGTAVKGKEAVHS	ZIKV224	SFRAKDGCWYGMEIR
ZIKV169	KQAWEDGICGISSVS	ZIKV197	AVKGKEAVHSDLGYW	ZIKV225	DGCWYGMEIRPRKEP
ZIKV170	DGICGISSVSRMENI	ZIKV198	EAVHSDLGYWIESEK	ZIKV226	GMEIRPRKEPESNLV
ZIKV171	ISSVSRMENIMWRSV	ZIKV199	DLGYWIESEKNDTWR	ZIKV227	PRKEPESNLVRSMVT
ZIKV172	RMENIMWRSVEGELN	ZIKV200	IESEKNDTWRLKRAH	ZIKV228	ESNLVRSMVTAGSTD
ZIKV173	MWRSVEGELNAILEE	ZIKV201	NDTWRLKRAHLIEMK	ZIKV229	RSMVTAGSTDHMDHF
ZIKV174	EGELNAILEENGVQL	ZIKV202	LKRAHLIEMKTCEWP	ZIKV230	AGSTDHMDHFSLGVL
ZIKV175	AILEENGVQLTVVVG	ZIKV203	LIEMKTCEWPKSHTL	ZIKV231	HMDHFSLGVLVILLM
ZIKV176	NGVQLTVVVGSVKNP	ZIKV204	TCEWPKSHTLWTDGI	ZIKV232	SLGVLVILLMVQEGL
ZIKV177	TVVVGSVKNPMWRGP	ZIKV205	KSHTLWTDGIEESDL	ZIKV233	VILLMVQEGLKKRMT
ZIKV178	SVKNPMWRGPQRLPV	ZIKV206	WTDGIEESDLIIPKS	ZIKV234	VQEGLKKRMTTKIII
ZIKV179	MWRGPQRLPVPVNEL	ZIKV207	EESDLIIPKSLAGPL	ZIKV235	KKRMTTKIIISTSMA
ZIKV180	QRLPVPVNELPHGWK	ZIKV208	IIPKSLAGPLSHHNT	ZIKV236	TKIIISTSMAVLVAM
ZIKV181	PVNELPHGWKAWGKS	ZIKV209	LAGPLSHHNTREGYR	ZIKV237	STSMAVLVAMILGGF
ZIKV182	PHGWKAWGKSYFVRA	ZIKV210	SHHNTREGYRTQMKG	ZIKV238	VLVAMILGGFSMSDL
ZIKV183	AWGKSYFVRAAKTNN	ZIKV211	REGYRTQMKGPWHSE	ZIKV239	ILGGFSMSDLAKLAI
ZIKV184	YFVRAAKTNNSFVVD	ZIKV212	TQMKGPWHSEELEIR	ZIKV240	SMSDLAKLAILMGAT
ZIKV185	AKTNNSFVVDGDTLK	ZIKV213	PWHSEELEIRFEECP	ZIKV241	AKLAILMGATFAEMN
ZIKV186	SFVVDGDTLKECPLK	ZIKV214	ELEIRFEECPGTKVH	ZIKV242	LMGATFAEMNTGGDV
ZIKV187	GDTLKECPLKHRAWN	ZIKV215	FEECPGTKVHVEETC	ZIKV243	FAEMNTGGDVAHLAL

Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence
ZIKV244	TGGDVAHLALIAAFK	ZIKV272	DPINVVGLLLLTRSG	ZIKV300	YVKTGKRSGALWDVP
ZIKV245	AHLALIAAFKVRPAL	ZIKV273	VGLLLLTRSGKRSWP	ZIKV301	KRSGALWDVPAPKEV
ZIKV246	IAAFKVRPALLVSFI	ZIKV274	LTRSGKRSWPPSEVL	ZIKV302	LWDVPAPKEVKKGET
ZIKV247	VRPALLVSFIFRANW	ZIKV275	KRSWPPSEVLTAVGL	ZIKV303	APKEVKKGETTDGVY
ZIKV248	LVSFIFRANWTPRES	ZIKV276	PSEVLTAVGLICALA	ZIKV304	KKGETTDGVYRVMTR
ZIKV249	FRANWTPRESMLLAL	ZIKV277	TAVGLICALAGGFAK	ZIKV305	TDGVYRVMTRRLLGS
ZIKV250	TPRESMLLALASCLL	ZIKV278	ICALAGGFAKADIEM	ZIKV306	RVMTRRLLGSTQVGV
ZIKV251	MLLALASCLLQTAIS	ZIKV279	GGFAKADIEMAGPMA	ZIKV307	RLLGSTQVGVGVMQE
ZIKV252	ASCLLQTAISALEGD	ZIKV280	ADIEMAGPMAAVGLL	ZIKV308	TQVGVGVMQEGVFHT
ZIKV253	QTAISALEGDLMVLI	ZIKV281	AGPMAAVGLLIVSYV	ZIKV309	GVMQEGVFHTMWHVT
ZIKV254	ALEGDLMVLINGFAL	ZIKV282	AVGLLIVSYVVSGKS	ZIKV310	GVFHTMWHVTKGSAL
ZIKV255	LMVLINGFALAWLAI	ZIKV283	IVSYVVSGKSVDMYI	ZIKV311	MWHVTKGSALRSGEG
ZIKV256	NGFALAWLAIRAMVV	ZIKV284	VSGKSVDMYIERAGD	ZIKV312	KGSALRSGEGRLDPY
ZIKV257	AWLAIRAMVVPRTDN	ZIKV285	VDMYIERAGDITWEK	ZIKV313	RSGEGRLDPYWGDVK
ZIKV258	RAMVVPRTDNITLAI	ZIKV286	ERAGDITWEKDAEVT	ZIKV314	RLDPYWGDVKQDLVS
ZIKV259	PRTDNITLAILAALT	ZIKV287	ITWEKDAEVTGNSPR	ZIKV315	WGDVKQDLVSYCGPW
ZIKV260	ITLAILAALTPLARG	ZIKV288	DAEVTGNSPRLDVAL	ZIKV316	QDLVSYCGPWKLDAA
ZIKV261	LAALTPLARGTLLVA	ZIKV289	GNSPRLDVALDESGD	ZIKV317	YCGPWKLDAAWDGHS
ZIKV262	PLARGTLLVAWRAGL	ZIKV290	LDVALDESGDFSLVE	ZIKV318	KLDAAWDGHSEVQLL
ZIKV263	TLLVAWRAGLATCGG	ZIKV291	DESGDFSLVEDDGPP	ZIKV319	WDGHSEVQLLAVPPG
ZIKV264	WRAGLATCGGFMLLS	ZIKV292	FSLVEDDGPPMREII	ZIKV320	EVQLLAVPPGERARN
ZIKV265	ATCGGFMLLSLKGKG	ZIKV293	DDGPPMREIILKVVL	ZIKV321	AVPPGERARNIQTLP
ZIKV266	FMLLSLKGKGSVKKN	ZIKV294	MREIILKVVLMTICG	ZIKV322	ERARNIQTLPGIFKT
ZIKV267	LKGKGSVKKNLPFVM	ZIKV295	LKVVLMTICGMNPIA	ZIKV323	IQTLPGIFKTKDGDI
ZIKV268	SVKKNLPFVMALGLT	ZIKV296	MTICGMNPIAIPFAA	ZIKV324	GIFKTKDGDIGAVAL
ZIKV269	LPFVMALGLTAVRLV	ZIKV297	MNPIAIPFAAGAWYV	ZIKV325	KDGDIGAVALDYPAG
ZIKV270	ALGLTAVRLVDPINV	ZIKV298	IPFAAGAWYVYVKTG	ZIKV326	GAVALDYPAGTSGSP
ZIKV271	AVRLVDPINVVGLLL	ZIKV299	GAWYVYVKTGKRSGA	ZIKV327	DYPAGTSGSPILDKC

Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence
ZIKV328	TSGSPILDKCGRVIG	ZIKV356	IRVPNYNLYIMDEAH	ZIKV384	MGANFKADRVIDSRR
ZIKV329	ILDKCGRVIGLYGNG	ZIKV357	YNLYIMDEAHFTDPS	ZIKV385	KADRVIDSRRCLKPV
ZIKV330	GRVIGLYGNGVVIKN	ZIKV358	MDEAHFTDPSSIAAR	ZIKV386	IDSRRCLKPVILDGE
ZIKV331	LYGNGVVIKNGSYVS	ZIKV359	FTDPSSIAARGYIST	ZIKV387	CLKPVILDGERVILA
ZIKV332	VVIKNGSYVSAITQG	ZIKV360	SIAARGYISTRVEMG	ZIKV388	ILDGERVILAGPMPV
ZIKV333	GSYVSAITQGRREEE	ZIKV361	GYISTRVEMGEAAAI	ZIKV389	RVILAGPMPVTHASA
ZIKV334	AITQGRREEETPVEC	ZIKV362	RVEMGEAAAIFMTAT	ZIKV390	GPMPVTHASAAQRRG
ZIKV335	RREEETPVECFEPSM	ZIKV363	EAAAIFMTATPPGTR	ZIKV391	THASAAQRRGRIGRN
ZIKV336	TPVECFEPSMLKKKQ	ZIKV364	FMTATPPGTRDAFPD	ZIKV392	AQRRGRIGRNPNKPG
ZIKV337	FEPSMLKKKQLTVLD	ZIKV365	PPGTRDAFPDSNSPI	ZIKV393	RIGRNPNKPGDEYLY
ZIKV338	LKKKQLTVLDLHPGA	ZIKV366	DAFPDSNSPIMDTEV	ZIKV394	PNKPGDEYLYGGGCA
ZIKV339	LTVLDLHPGAGKTRR	ZIKV367	SNSPIMDTEVEVPER	ZIKV395	DEYLYGGGCAETDED
ZIKV340	LHPGAGKTRRVLPEI	ZIKV368	MDTEVEVPERAWSSG	ZIKV396	GGGCAETDEDHAHWL
ZIKV341	GKTRRVLPEIVREAI	ZIKV369	EVPERAWSSGFDWVT	ZIKV397	ETDEDHAHWLEARML
ZIKV342	VLPEIVREAIKTRLR	ZIKV370	AWSSGFDWVTDHSGK	ZIKV398	HAHWLEARMLLDNIY
ZIKV343	VREAIKTRLRTVILA	ZIKV371	FDWVTDHSGKTVWFV	ZIKV399	EARMLLDNIYLQDGL
ZIKV344	KTRLRTVILAPTRVV	ZIKV372	DHSGKTVWFVPSVRN	ZIKV400	LDNIYLQDGLIASLY
ZIKV345	TVILAPTRVVAAEME	ZIKV373	TVWFVPSVRNGNEIA	ZIKV401	LQDGLIASLYRPEAD
ZIKV346	PTRVVAAEMEEALRG	ZIKV374	PSVRNGNEIAACLTK	ZIKV402	IASLYRPEADKVAAI
ZIKV347	AAEMEEALRGLPVRY	ZIKV375	GNEIAACLTKAGKRV	ZIKV403	RPEADKVAAIEGEFK
ZIKV348	EALRGLPVRYMTTAV	ZIKV376	ACLTKAGKRVIQLSR	ZIKV404	KVAAIEGEFKLRTEQ
ZIKV349	LPVRYMTTAVNVTHS	ZIKV377	AGKRVIQLSRKTFET	ZIKV405	EGEFKLRTEQRKTFV
ZIKV350	MTTAVNVTHSGTEIV	ZIKV378	IQLSRKTFETEFQKT	ZIKV406	LRTEQRKTFVELMKR
ZIKV351	NVTHSGTEIVDLMCH	ZIKV379	KTFETEFQKTKHQEW	ZIKV407	RKTFVELMKRGDLPV
ZIKV352	GTEIVDLMCHATFTS	ZIKV380	EFQKTKHQEWDFVVT	ZIKV408	ELMKRGDLPVWLAYQ
ZIKV353	DLMCHATFTSRLLQP	ZIKV381	KHQEWDFVVTTDISE	ZIKV409	GDLPVWLAYQVASAG
ZIKV354	ATFTSRLLQPIRVPN	ZIKV382	DFVVTTDISEMGANF	ZIKV410	WLAYQVASAGITYTD
ZIKV355	RLLQPIRVPNYNLYI	ZIKV383	TDISEMGANFKADRV	ZIKV411	VASAGITYTDRRWCF

Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence
ZIKV412	ITYTDRRWCFDGTTN	ZIKV440	GIGKMGFGMVTLGAS	ZIKV468	LMAMATQAGVLFGMG
ZIKV413	RRWCFDGTTNNTIME	ZIKV441	GFGMVTLGASAWLMW	ZIKV469	TQAGVLFGMGKGMPF
ZIKV414	DGTTNNTIMEDSVPA	ZIKV442	TLGASAWLMWLSEIE	ZIKV470	LFGMGKGMPFYAWDF
ZIKV415	NTIMEDSVPAEVWTR	ZIKV443	AWLMWLSEIEPARIA	ZIKV471	KGMPFYAWDFGVPLL
ZIKV416	DSVPAEVWTRHGEKR	ZIKV444	LSEIEPARIACVLIV	ZIKV472	YAWDFGVPLLMIGCY
ZIKV417	EVWTRHGEKRVLKPR	ZIKV445	PARIACVLIVVFLLL	ZIKV473	GVPLLMIGCYSQLTP
ZIKV418	HGEKRVLKPRWMDAR	ZIKV446	CVLIVVFLLLVVLIP	ZIKV474	MIGCYSQLTPLTLIV
ZIKV419	VLKPRWMDARVCSDH	ZIKV447	VFLLLVVLIPEPEKQ	ZIKV475	SQLTPLTLIVAIILL
ZIKV420	WMDARVCSDHAALKS	ZIKV448	VVLIPEPEKQRSPQD	ZIKV476	LTLIVAIILLVAHYM
ZIKV421	VCSDHAALKSFKEFA	ZIKV449	EPEKQRSPQDNQMAI	ZIKV477	AIILLVAHYMYLIPG
ZIKV422	AALKSFKEFAAGKRG	ZIKV450	RSPQDNQMAIIIMVA	ZIKV478	VAHYMYLIPGLQAAA
ZIKV423	FKEFAAGKRGAAFGV	ZIKV451	NQMAIIIMVAVGLLG	ZIKV479	YLIPGLQAAAARAAQ
ZIKV424	AGKRGAAFGVMEALG	ZIKV452	IIMVAVGLLGLITAN	ZIKV480	LQAAAARAAQKRTAA
ZIKV425	AAFGVMEALGTLPGH	ZIKV453	VGLLGLITANELGWL	ZIKV481	ARAAQKRTAAGIMKN
ZIKV426	MEALGTLPGHMTERF	ZIKV454	LITANELGWLERTKS	ZIKV482	KRTAAGIMKNPVVDG
ZIKV427	TLPGHMTERFQEAID	ZIKV455	ELGWLERTKSDLSHL	ZIKV483	GIMKNPVVDGIVVTD
ZIKV428	MTERFQEAIDNLAVL	ZIKV456	ERTKSDLSHLMGRRE	ZIKV484	PVVDGIVVTDIDTMT
ZIKV429	QEAIDNLAVLMRAET	ZIKV457	DLSHLMGRREEGATI	ZIKV485	IVVTDIDTMTIDPQV
ZIKV430	NLAVLMRAETGSRPY	ZIKV458	MGRREEGATIGFSMD	ZIKV486	IDTMTIDPQVEKKMG
ZIKV431	MRAETGSRPYKAAAA	ZIKV459	EGATIGFSMDIDLRP	ZIKV487	IDPQVEKKMGQVLLI
ZIKV432	GSRPYKAAAAQLPET	ZIKV460	GFSMDIDLRPASAWA	ZIKV488	EKKMGQVLLIAVAVS
ZIKV433	KAAAAQLPETLETIM	ZIKV461	IDLRPASAWAIYAAL	ZIKV489	QVLLIAVAVSSAILS
ZIKV434	QLPETLETIMLLGLL	ZIKV462	ASAWAIYAALTTFIT	ZIKV490	AVAVSSAILSRTAWG
ZIKV435	LETIMLLGLLGTVSL	ZIKV463	IYAALTTFITPAVQH	ZIKV491	SAILSRTAWGWGEAG
ZIKV436	LLGLLGTVSLGIFFV	ZIKV464	TTFITPAVQHAVTTS	ZIKV492	RTAWGWGEAGALITA
ZIKV437	GTVSLGIFFVLMRNK	ZIKV465	PAVQHAVTTSYNNYS	ZIKV493	WGEAGALITAATSTL
ZIKV438	GIFFVLMRNKGIGKM	ZIKV466	AVTTSYNNYSLMAMA	ZIKV494	ALITAATSTLWEGSP
ZIKV439	LMRNKGIGKMGFGMV	ZIKV467	YNNYSLMAMATQAGV	ZIKV495	ATSTLWEGSPNKYWN

Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence
ZIKV496	WEGSPNKYWNSSTAT	ZIKV524	KVQEVKGYTKGGPGH	ZIKV552	TSQLLLGRMDGPRRP
ZIKV497	NKYWNSSTATSLCNI	ZIKV525	KGYTKGGPGHEEPVL	ZIKV553	LGRMDGPRRPVKYEE
ZIKV498	SSTATSLCNIFRGSY	ZIKV526	GGPGHEEPVLVQSYG	ZIKV554	GPRRPVKYEEDVNLG
ZIKV499	SLCNIFRGSYLAGAS	ZIKV527	EEPVLVQSYGWNIVR	ZIKV555	VKYEEDVNLGSGTRA
ZIKV500	FRGSYLAGASLIYTV	ZIKV528	VQSYGWNIVRLKSGV	ZIKV556	DVNLGSGTRAVVSCA
ZIKV501	LAGASLIYTVTRNAG	ZIKV529	WNIVRLKSGVDVFHM	ZIKV557	SGTRAVVSCAEAPNM
ZIKV502	LIYTVTRNAGLVKRR	ZIKV530	LKSGVDVFHMAAEPC	ZIKV558	VVSCAEAPNMKIIGN
ZIKV503	TRNAGLVKRRGGGTG	ZIKV531	DVFHMAAEPCDTLLC	ZIKV559	EAPNMKIIGNRIERI
ZIKV504	LVKRRGGGTGETLGE	ZIKV532	AAEPCDTLLCDIGES	ZIKV560	KIIGNRIERIRSEHA
ZIKV505	GGGTGETLGEKWKAR	ZIKV533	DTLLCDIGESSSSPE	ZIKV561	RIERIRSEHAETWFF
ZIKV506	ETLGEKWKARLNQMS	ZIKV534	DIGESSSSPEVEEAR	ZIKV562	RSEHAETWFFDENHP
ZIKV507	KWKARLNQMSALEFY	ZIKV535	SSSPEVEEARTLRVL	ZIKV563	ETWFFDENHPYRTWA
ZIKV508	LNQMSALEFYSYKKS	ZIKV536	VEEARTLRVLSMVGD	ZIKV564	DENHPYRTWAYHGSY
ZIKV509	ALEFYSYKKSGITEV	ZIKV537	TLRVLSMVGDWLEKR	ZIKV565	YRTWAYHGSYEAPTQ
ZIKV510	SYKKSGITEVCREEA	ZIKV538	SMVGDWLEKRPGAFC	ZIKV566	YHGSYEAPTQGSASS
ZIKV511	GITEVCREEARRALK	ZIKV539	WLEKRPGAFCIKVLC	ZIKV567	EAPTQGSASSLINGV
ZIKV512	CREEARRALKDGVAT	ZIKV540	PGAFCIKVLCPYTST	ZIKV568	GSASSLINGVVRLLS
ZIKV513	RRALKDGVATGGHAV	ZIKV541	IKVLCPYTSTMMETL	ZIKV569	LINGVVRLLSKPWDV
ZIKV514	DGVATGGHAVSRGSA	ZIKV542	PYTSTMMETLERLQR	ZIKV570	VRLLSKPWDVVTGVT
ZIKV515	GGHAVSRGSAKLRWL	ZIKV543	MMETLERLQRRYGGG	ZIKV571	KPWDVVTGVTGIAMT
ZIKV516	SRGSAKLRWLVERGY	ZIKV544	ERLQRRYGGGLVRVP	ZIKV572	VTGVTGIAMTDTTPY
ZIKV517	KLRWLVERGYLQPYG	ZIKV545	RYGGGLVRVPLSRNS	ZIKV573	GIAMTDTTPYGQQRV
ZIKV518	VERGYLQPYGKVIDL	ZIKV546	LVRVPLSRNSTHEMY	ZIKV574	DTTPYGQQRVFKEKV
ZIKV519	LQPYGKVIDLGCGRG	ZIKV547	LSRNSTHEMYWVSGA	ZIKV575	GQQRVFKEKVDTRVP
ZIKV520	KVIDLGCGRGGWSYY	ZIKV548	THEMYWVSGAKSNTI	ZIKV576	FKEKVDTRVPDPQEG
ZIKV521	GCGRGGWSYYVATIR	ZIKV549	WVSGAKSNTIKSVST	ZIKV577	DTRVPDPQEGTRQVM
ZIKV522	GWSYYVATIRKVQEV	ZIKV550	KSNTIKSVSTTSQLL	ZIKV578	DPQEGTRQVMSMVSS
ZIKV523	VATIRKVQEVKGYTK	ZIKV551	KSVSTTSQLLLGRMD	ZIKV579	TRQVMSMVSSWLWKE

Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence	Peptide	Amino Acid Sequence
ZIKV580	SMVSSWLWKELGKHK	ZIKV608	LGYVLEEMSRIPGGR	ZIKV636	RLKRMAVSGDDCVVK
ZIKV581	WLWKELGKHKRPRVC	ZIKV609	EEMSRIPGGRMYADD	ZIKV637	AVSGDDCVVKPIDDR
ZIKV582	LGKHKRPRVCTKEEF	ZIKV610	IPGGRMYADDTAGWD	ZIKV638	DCVVKPIDDRFAHAL
ZIKV583	RPRVCTKEEFINKVR	ZIKV611	MYADDTAGWDTRISR	ZIKV639	PIDDRFAHALRFLND
ZIKV584	TKEEFINKVRSNAAL	ZIKV612	TAGWDTRISRFDLEN	ZIKV640	FAHALRFLNDMGKVR
ZIKV585	INKVRSNAALGAIFE	ZIKV613	TRISRFDLENEALIT	ZIKV641	RFLNDMGKVRKDTQE
ZIKV586	SNAALGAIFEEEKEW	ZIKV614	FDLENEALITNQMEK	ZIKV642	MGKVRKDTQEWKPST
ZIKV587	GAIFEEEKEWKTAVE	ZIKV615	EALITNQMEKGHRAL	ZIKV643	KDTQEWKPSTGWDNW
ZIKV588	EEKEWKTAVEAVNDP	ZIKV616	NQMEKGHRALALAII	ZIKV644	WKPSTGWDNWEEVPF
ZIKV589	KTAVEAVNDPRFWAL	ZIKV617	GHRALALAIIKYTYQ	ZIKV645	GWDNWEEVPFCSHHF
ZIKV590	AVNDPRFWALVDKER	ZIKV618	ALAIIKYTYQNKVVK	ZIKV646	EEVPFCSHHFNKLHL
ZIKV591	RFWALVDKEREHHLR	ZIKV619	KYTYQNKVVKVLRPA	ZIKV647	CSHHFNKLHLKDGRS
ZIKV592	VDKEREHHLRGECQS	ZIKV620	NKVVKVLRPAEKGKT	ZIKV648	NKLHLKDGRSIVVPC
ZIKV593	EHHLRGECQSCVYNM	ZIKV621	VLRPAEKGKTVMDII	ZIKV649	KDGRSIVVPCRHQDE
ZIKV594	GECQSCVYNMMGKRE	ZIKV622	EKGKTVMDIISRQDQ	ZIKV650	IVVPCRHQDELIGRA
ZIKV595	CVYNMMGKREKKQGE	ZIKV623	VMDIISRQDQRGSGQ	ZIKV651	RHQDELIGRARVSPG
ZIKV596	MGKREKKQGEFGKAK	ZIKV624	SRQDQRGSGQVVTYA	ZIKV652	LIGRARVSPGAGWSI
ZIKV597	KKQGEFGKAKGSRAI	ZIKV625	RGSGQVVTYALNTFT	ZIKV653	RVSPGAGWSIRETAC
ZIKV598	FGKAKGSRAIWYMWL	ZIKV626	VVTYALNTFTNLVVQ	ZIKV654	AGWSIRETACLAKSY
ZIKV599	GSRAIWYMWLGARFL	ZIKV627	LNTFTNLVVQLIRNM	ZIKV655	RETACLAKSYAQMWQ
ZIKV600	WYMWLGARFLEFEAL	ZIKV628	NLVVQLIRNMEAEEV	ZIKV656	LAKSYAQMWQLLYFH
ZIKV601	GARFLEFEALGFLNE	ZIKV629	LIRNMEAEEVLEMQD	ZIKV657	AQMWQLLYFHRRDLR
ZIKV602	EFEALGFLNEDHWMG	ZIKV630	EAEEVLEMQDLWLLR	ZIKV658	LLYFHRRDLRLMANA
ZIKV603	GFLNEDHWMGRENSG	ZIKV631	LEMQDLWLLRRSEKV	ZIKV659	RRDLRLMANAICSSV
ZIKV604	DHWMGRENSGGGVEG	ZIKV632	LWLLRRSEKVTNWLQ	ZIKV660	LMANAICSSVPVDWV
ZIKV605	RENSGGGVEGLGLQR	ZIKV633	RSEKVTNWLQSNGWD	ZIKV661	ICSSVPVDWVPTGRT
ZIKV606	GGVEGLGLQRLGYVL	ZIKV634	TNWLQSNGWDRLKRM	ZIKV662	PVDWVPTGRTTWSIH
ZIKV607	LGLQRLGYVLEEMSR	ZIKV635	SNGWDRLKRMAVSGD	ZIKV663	PTGRTTWSIHGKGEW

Peptide	Amino Acid Sequence
ZIKV664	TWSIHGKGEWMTTED
ZIKV665	GKGEWMTTEDMLVVW
ZIKV666	MTTEDMLVVWNRVWI
ZIKV667	MLVVWNRVWIEENDH
ZIKV668	NRVWIEENDHMEDKT
ZIKV669	EENDHMEDKTPVTKW
ZIKV670	MEDKTPVTKWTDIPY
ZIKV671	PVTKWTDIPYLGKRE
ZIKV672	TDIPYLGKREDLWCG
ZIKV673	LGKREDLWCGSLIGH
ZIKV674	DLWCGSLIGHRPRTT
ZIKV675	SLIGHRPRTTWAENI
ZIKV676	RPRTTWAENIKNTVN
ZIKV677	WAENIKNTVNMVRRI
ZIKV678	KNTVNMVRRIIGDEE
ZIKV679	MVRRIIGDEEKYMDY
ZIKV680	IGDEEKYMDYLSTQV
ZIKV681	KYMDYLSTQVRYLGE
ZIKV682	LSTQVRYLGEEGSTP
ZIKV683	RYLGEEGSTPGVL