Distribution and density of oxpeckers on giraffes in Hwange National Park, Zimbabwe

Roxanne Gagnon[1], Cheryl T. Mabika[2], and Christophe Bonenfant[3,4]

[1]Université de Sherbrooke, Département de biologie, 2500 Boulevard de l'Université, Sherbrooke, Québec J1K 2R1, Canada.

[2]Zimbabwe Parks and Wildlife Management Authority, Hwange Cluster, Main Camp, Zimbabwe.

[3]Université Claude Bernard – Lyon 1; UMR CNRS 5558, Laboratoire de Biométrie et Biologie Évolutive, F-69 622, Villeurbanne CEDEX, France.

ORCID: 0000-0002-9924-419X

[4]LTSER France, Zone Atelier "Hwange", Hwange National Park, Bag 62, Dete, Zimbabwe – CNRS HERD (Hwange Environmental Research Development) program

Correspondence

C. Bonenfant

Email: christophe.bonenfant@univ-lyon1.fr

Abstract   Oxpeckers (*Buphagus sp.*) are two sub-Saharan bird species closely associated to large mammals, which they use as hosts for foraging. The comparison of opxecker abundance among large mammalian species has been repeatedly investigated in many areas of Southern Africa, and giraffe (*Giraffa camelopardalis*) is a strongly favoured hosts. The foraging behaviour of oxpeckers at the individual and the body levels of giraffes remains little studied, however. By comparing the distribution of oxpeckers among and within individual giraffes, one may infer on the foraging behaviour of birds. Here, we counted every oxpeckers on giraffe's body from pictures and assessed the detection probability of birds on their host. We first tested whether oxpeckers distributed themselves at random or aggregated on individual giraffes. Then we explored the distribution of oxpeckers on pre-defined zones on the giraffe body from which tested the prediction that birds would be preferentially found on the host's body parts providing birds with the greatest amount of ticks. Oxpeckers displayed a strong aggregation behaviour with few hosts carrying many birds while many carried a limited number or no birds, a pattern that differed between sexes. Oxpeckers were not uniformly distributed on the giraffe's body and were mostly found on the mane and back, where the density of ticks is presumably the highest. The high aggregation level of birds we report is typical of parasitic species and would hence suggest oxpeckers behave more like parasites rather than a mutualistic species with giraffes. Alternatively this particular oxpecker distribution may mirror the distribution of ticks among hosts. Abundance of ticks appears indeed as a major driver of the oxpecker foraging behaviour because, as we predicted, birds were most numerous on the giraffes' body part where ectoparasite density is expected to be the greatest (mane and back up to the tail insertion). From an ecological point of view, the oxpecker–large herbivores system proves to be highly relevant and useful for the study of host-parasite dynamics.

1 | INTRODUCTION

The distribution of animals in the environment results from a complex sequence of behavioural decisions aiming at satisfying the energy requirements of individuals while minimizing costs of movements, competition with con-specifics or other species, in balance with the perceived risks of predation for prey species (Krebs, 1972). From an evolutionary perspective, animals should select habitats with the highest suitability (sensu Fretwell & Lucas, 1969), *i.e.* those habitats in which the fitness of individuals will be the greatest. Variation in habitat suitability, in space and time, is of prime importance to the ecology of species with consequences on its distribution range (MacArthur, 1972), its mating tactic (Emlen & Oring, 1977), or its population dynamics (Lack, 1966; Ostfeld & Keesing, 2000).

At the population level, when and where resources are available is a strong predictor of the abundance of animals. Assuming an homogeneous distribution of discrete resources patches in the landscape and a random walk of foraging animal trajectories, the expected number of foragers per patch is given by a Poisson distribution (Hutchinson & Waser, 2007). This simple model of encounters between motile entities underpins most multi-species interaction models, including the Lotka-Volterra (Lotka, 1956; Hutchinson & Waser, 2007) or Nicholson-Bailey models (May, 1978) for, respectively, predators–prey and hosts–parasitoid dynamics. Any deviation from the Poisson distribution is usually interpreted as a sign of aggregation (over-dispersion) or avoidance (under-dispersion) of individuals (Pielou (1969, p. 96) but see Taylor et al. (1979) or Sjöberg et al. (2000) for a discussion and alternatives). Several indices have been proposed to quantify aggregation levels among populations or species from count data (see Kretzschmar and Adler 1993 for a review), the most widely used being the aggregation index k (Shaw et al., 1998). Aggregation may arise from the animal behaviour such as social interactions (Wittenberger, 1981), constrain on mobility among patches (Gueron & Levin, 1995), or if animals perform area-restricted search of food patches (Morales et al., 2010) or do copy what the other con-specifics do when using public information (Clark & Mangel, 1986).

In multi-species interactions such as in bird vs. large mammals associations, the

distribution of the birds is first guided by the one of the mammalian hosts, conceptually

2 equivalent to a resource patch. This scenario fits the association between oxpeckers

(*Buphagus sp.*) and large herbivores, described in the early 20th century (Moreau, 1933).

4 Oxpeckers are two bird species of sub-Saharan Africa associated to savanna ecosystems

(Hustler, 1987; Plantan, 2009; Palmer & Packer, 2018). They live and feed almost

6 exclusively on the body of large herbivores such as African buffaloes (*Syncerus caffer*

caffer), giraffes (*Giraffa camelopardalis*), black rhinoceros (*Diceros bicornis*), white

8 rhinoceros (*Ceratotherium simum*), impalas (*Aepyceros melampus*), greater kudus

(*Tragelaphus strepsiceros*), common elands (*Taurotragus oryx*) and sable antelopes

10 (*Hippotragus niger*) (Hustler, 1987; Stutterheim et al., 1988; Palmer & Packer, 2018).

Oxpeckers mainly prey upon ectoparasites of their large mammalian hosts although they

12 sometimes can snatch tissues from their host (Bezuidenhout & Stutterheim, 1980). In

terms of resource selection, the foraging behavior of oxpeckers can be decomposed into

14 two main steps. The first step for birds is to localize large mammals in the landscape which,

for oxpeckers represent motile and widely dispersed resources patches of varying size. The

16 second event takes place on the host's body where oxpeckers will search for the most

suitable body part in terms of ectoparasites. These two sequential steps, taking place at

18 different spatio-temporal scales (see Johnson, 1980), will drive the distribution of oxpeckers

among hosts and on their body.

20 At the largest spatial scale, the distribution of oxpeckers among the different species of

mammalian hosts has been documented for decades (Moreau, 1933; Grobler, 1980;

22 Hustler, 1987; Ndlovu & Combrink, 2015). Clear preferences were demonstrated and, with

a large body and a high tolerance to the presence of birds, the giraffe appears to be one of

24 the key host for two oxpecker species (Grobler, 1980; Veríssimo et al., 2017). Much less is

known about how oxpeckers are distributed on the different individuals of a given host

26 species, with the exception of maximum records (*e.g.* 51 birds on one side of a single

giraffe, Veríssimo et al. (2017)). At the host level of resource selection, oxpeckers seems to

28 favor some body parts of their hosts but what body parts is most attractive seems to be

different according to the host species (Palmer & Packer, 2018). For instance, Ndlovu &

Combrink (2015) reported that red-billed oxpeckers (*Buphagus erythrorhynchus*) were most

2 frequently perch on the back and the head of buffaloes and white rhinoceros and that the

neck was preferred on giraffes.

4   In this paper we investigated the among-host and within-host distribution of oxpeckers

on giraffes at Hwange National Park, Zimbabwe. We extracted the number and location of

6 yellow-billed (*Buphagus africanus*) and red-billed oxpeckers on $n = 683$ giraffes from 500

pictures collected in 2012, 2015 and 2016. We aimed at testing the random encounter

8 model between foragers and hosts, and the ideal free distribution (IFD, Fretwell & Lucas,

1969) with oxpeckers as foragers and giraffes as their primary large mammalian hosts. We

10 tested the three following predictions:

1. *Distribution of oxpeckers among giraffes*: By comparing the observed distribution of

12   the number of oxpeckers per giraffes with the expected prediction from theoretical

    models, one may make inference on the underlying behaviour of resource selection

14   by and movements of birds. If oxpeckers search for giraffes at random in the

    landscape the random encounter model predicts a Poisson distribution of birds

16   among hosts (Hutchinson & Waser, 2007). Alternatively, if oxpeckers aggregate

    preferentially on some particular hosts because of marked differences in parasite

18   loads among giraffes or because of copying behaviour, the model predicts a negative

    binomial distribution of birds among hosts (Pielou, 1969);

20 2. *Distribution of oxpeckers on giraffe body*: According to the IFD (Fretwell & Lucas,

    1969), oxpeckers should be distributed on the giraffe body parts proportionally to the

22   local ectoparasite load. Therefore, if the ectoparasite density is homogeneous over

    the whole giraffe body, the IFD predicts an homogeneous number of oxpeckers per

24   area unit. Alternatively, if ectoparasites concentrate on some specific body parts, the

    IFD predicts an heterogeneous distribution of birds, with higher densities of oxpeckers

26   on those giraffe body parts with the higher ectoparasite load (Horak et al., 1983;

    Mysterud et al., 2014). From previous observations (Plantan, 2009; Ndlovu &

28   Combrink, 2015), giraffe body parts with the highest oxpecker number should be, in

decreasing order, the mane, the neck, the scapula and the back;

3. *Sex-differences in bird load of giraffes*: Many studies evidenced that the ectoparasite load is proportional to the body mass and skin surface of the host (Horak et al., 1983, 1987; Koenig, 1997). Consequently a bigger host should carry more ectoparasites and hence, more birds than a smaller one. Sexual size dimorphism is observed among many species and is particularly observable between male and female giraffes, reaching a 43% difference for fully grown individuals (Shorrocks, 2016). We will therefore test the prediction that more birds are present on male than on female giraffes;

2 | MATERIAL AND METHODS

2.1 | Study site

This study was undertaken around Main Camp in the northeast of Hwange National Park , the main biological reserve of Zimbabwe (HNP; 19°00'S, 26°30'E, extending from Main Camp to Giraffe's spring and Ngweshla pans; Fig. 1). This park covers $14\,650$ km^2 and supports a population of approx. $2\,800$ giraffes (Shorrocks, 2016). The Bulawayo-Victoria Falls railway line defines the western boundary of HNP while border with Botswana draws the eastern boundary. The long-term mean annual rainfall is ca. 600 mm (CV = 25%) and generally falls between October and April to form seasonal wetlands. Because of this relatively low annual rainfall, a xerophile vegetation covers most of HNP. The woodland vegetation consists primarily of African teak (*Baikiaea plurijuga*) intersected with patches of camel thorn (*Acacia erioloba*) or leadwood (*Combretum imberbe*). Bushland savanna with patches of grassland makes x% of the area, mainly around the many artificially maintained waterholes. HNP hosts many large and mega-herbivore species attractive to oxpeckers, including giraffes, plain zebras (*Equus quagga*), African buffaloes, wildbeest (*Connochaetes taurinus*), greater kudus, waterbuck (*Kobus ellipsiprymnus*) and impala, and the less abundant sable and roan antelopes (*H. equinus*).

2.2 | Oxpecker biology

Red-billed oxpecker (RBO) and yellow-billed oxpecker (YBO) are two sympatric species, strictly african, that can be mostly found in Botswana, Namibia, South Africa and Zimbabwe (Hustler, 1987; Plantan, 2009; Palmer & Packer, 2018). Although those birds feed mainly on ticks, many studies have reported that oxpeckers can feed on wounded tissue, mucus, saliva, ear wax, hair and blood (Bezuidenhout & Stutterheim, 1980; Weeks, 1999; Plantan, 2009). Some authors definitely think that the interaction between oxpekers and mammalian hosts is parasitism, while others support mutualism even though they admit oxpeckers can cause injuries to hosts (van Someren, 1951; Samish & Rehacek, 1999). In fact the relationship between oxpeckers and hosts could be context-dependent, where birds can be opportunists under particular biotic and abiotic conditions (Moreau, 1933; Nunn et al., 2011; Plantan, 2009). For instance, a mutualistic relationship may develop when the ectoparasites load is high on the host, but oxpeckers may become parasitic when hosts carry few ticks with too many birds, and leads to numerous open wounds.

2.3 | Data collection

Oxpecker data derived from the study of giraffe ecology carried out at HNP. We have been monitoring the giraffe population opportunistically in 2012 and 2013, and on a regular basis since 2014, aiming at the photo-identification of individuals. Each year we drove the HNP road network daily (Fig. 1) for at least three weeks in a row and shot pictures of every encountered giraffes. For all encounters, we recorded the date, the location and the time of observation along with group size and composition the individuals belonged to. We sexed giraffes based on the presence of bare skin at the ossicone tip for males, and on shape of the skull or visible male genitals when possible. We classified giraffes into four age-classes (calf, juvenile, sub-adult and adult) assessed in the field from their size and coat color since giraffes darken with age (Dagg, 2014). In this study we used all pictures taken between 2013 and 2015, yielding a total of 500 photographs and a sample size of 683 giraffes. Although we avoided to analyze a sequence of continuous pictures, re-observations of the

same individual in time might have occurred, either during the same field session or from

2 one year to another. Doing so, we assumed very little consequences of pseudo-replication

on the results because of the rapid exchanges and movements of oxpeckers among hosts.

4 We then counted oxpeckers on every giraffes seen on pictures and assigned each detected

bird to one of the 14 pre-defined parts of the giraffe body (see Fig. 3A) – giraffe body parts

6 were chosen so that they could easily be identified from landmark points whatever the point

of view –. We excluded approx. 50% of the pictures because giraffes were too small to

8 reliably spot oxpeckers which roughly corresponded to a giraffe relative size smaller than

$1/3$ of the picture height, or because of a poor picture quality. For a subset of pictures we

10 repeated oxpecker counts twice with two different observers (RG and CB) to estimate bird

detection probability.

12 **2.4 │ Data analyses**

We first estimated the detection probability of individual oxpeckers from pictures by setting

14 a double-observer experiment (Nichols et al., 2000). The two observers reported the total

number of detected birds they found (noted x_{11} and x_{22}) from which we calculated the

16 number of birds seen by observer 1 and missed by 2 (x_{12}) and conversely (x_{21}). The

double-observer method returns the detection probability of observer 1 and 2 (respectively

18 p_1 and p_2), as well as the average detection probability of birds p. The estimated number of

oxpeckers per giraffe is then given by $\hat{N} = (x_{11} + x_{22} + x_{12} + x_{21})/\hat{p}$. We fit the

20 double-observer model to our data with the *unmarked* library (Fiske & Chandler, 2011) and

tested for the effect of host sex on the detection probabilities.

22 We derived the aggregation index k from a particular parameterization of the negative

binomial distribution (where the variance V is related to the mean by $V = \lambda + \lambda^2/k$) that

24 takes values close to zero with increasing levels of aggregation. To compare with previous

studies, we also estimated oxpecker's preference for a given hosts with preference index

26 (PI), calculated as the number of hosts counted divided by the number of oxpeckers

counted. A PI of 5 means one bird is seen every fifth counted hosts on average. We

28 computed confidence limits of point estimate of PI with a non-parametric bootstrap.

We estimated the relative distribution of oxpeckers on giraffe body parts with a
2 multinomial logistic regression. This particular type of GLM estimates the 14 probabilities
(noted π_i) of birds to be located on each body part of the giraffe holding the constrain that
4 $\sum_{i=1}^{14} \pi_i = 1$. To control for the imbalanced number of males and female giraffes among
years, we entered year as a categorical variable before testing for the effect the host sex (a
6 2-levels categorical variable) on the relative density of oxpeckers on body parts with
likelihood ratio tests using the *nnet* library (Venables & Ripley, 2002).
8 Finally, we modeled the number of oxpeckers per giraffe with generalized linear models
(GLMs), testing for effect of host sex (a 2-levels categorical variable), accounting for the
10 effects of time (year as a categorical variable), with likelihood ratio tests using the *MASS*
library (Venables & Ripley, 2002). We ran all analyses in the **R** 3.4 statistical software (R
12 Core Team, 2018). Unless otherwise stated, we reported all estimated parameters as
mean $\pm$ sd and predicted probabilities as the mean with its associated 95% confidence
14 interval in brackets.

3 │ RESULTS

16 From a subsample of $n = 117$ giraffes, the overall detection probability of oxpeckers was
$\hat{p} = 0.99\ (0.97, 1.00)$ but differed substantially between the two observers (RG:
18 $\hat{p_1} = 0.97\ (0.95, 0.98)$; CB: $\hat{p_2} = 0.85\ (0.81, 0.89)$). Although oxperckers have 1.3 more
chances to be seen on a male than on female giraffe, detection probability did not differ
20 significantly between the host sex (estimated difference between male and female hosts on
the logit scale: $\beta = 0.26 \pm 0.34$, $P = 0.45$). On average we estimated oxpecker density to be
22 $D_O = 2.91\ (2.63, 3.23)$ birds per giraffe once we accounted for imperfect detections. In the
following analyses, RG did analyze all pictures.
24 When using $n = 683$ giraffes, mean oxpecker density was 2.16 ± 3.01 birds per host
without accounting for detection probability. The overall preference index (PI) read
26 0.46 ± 0.10 with a maximum number of oxpeckers counted on a single host of 17 (Fig. 2).
In support of the aggregation hypothesis, the estimated aggregation coefficient
28 $\hat{k} = 0.68\ (0.58, 0.79)$ suggested a strong aggregation of oxpeckers on individual giraffes.

Because the estimated aggregation coefficient k approaches zero, the negative binomial distribution converges to the logarithmic series distribution, with a strong skew toward giraffe carrying no bird (Fig. 2). Overall, our results lend support to the hypothesis of a non-random association between oxpeckers and giraffes at HNP (H_1).

The relative distribution of oxpeckers on giraffe's body deviated strongly from uniformity with some body parts being much more used than others (Fig. 3B). Supporting our hypothesis H_2, birds gathered principally on the neck ($\pi = 0.18\,(0.15, 0.20)$) and mane ($\pi = 0.25\,(0.21, 0.27)$) of giraffes, but were rarely seen on the lower limbs ($\pi < 0.01$) or on the tail ($\pi < 0.01$). Oxpeckers did not use the giraffe's body differently according to the host sex (likelihood ratio test: $\chi^2 = 28.68$, df = 13, $P = 0.07$; Fig. 3B), although relatively more birds used the ano-genital ($\beta = 0.40 \pm 56$, $P = 0.47$) and the scapula areas ($\beta = 0.45 \pm 0.53$, $P = 39$) of females compared to males. Conversely relatively more birds were seen on the head ($\beta = 1.53 \pm 0.57$, $P < 0.001$) and croup ($\beta = 1.17 \pm 0.54$, $P = 0.03$) of male than of female giraffes. Our results hence confirm the marked heterogeneous distribution of oxpeckers on the body of large mammalian hosts.

Although GLM with a Poisson distribution and a logarithmic link are usually recommended for count data (Agresti 2002), a preliminary goodness-of-fit (GOF) test suggested an over-dispersion of the data compared to a Poisson distribution ($\chi^2 = 2877.21$, df = 682, $P < 0.001$). A GLM a with negative binomial distribution did fit the data better than with a Poisson distribution (GOF test: $\chi^2 = 609.10$, df = 682, $P = 0.98$). As expected from our last hypothesis H_3, the number of oxpeckers was larger on the giraffes exposing the largest body area to the birds (Fig 4). Accordingly, we found that oxpeckers were 20% more numerous on males than on females (density of 2.60 ± 0.25 and 2.07 ± 0.14 birds per giraffe respectively: $\beta = 0.23 \pm 0.12$, $\chi^2 = 3.75$, df = 1, $P = 0.05$). The sex-specific aggregation coefficient read $k = 0.85 \pm 0.09$ and $k = 0.62 \pm 0.07$ for females and males respectively, and was significantly smaller for male giraffes (bootstrap test: $\beta = 0.23 \pm 0.13$, $P = 0.02$).

4 | DISCUSSION

The foraging behaviour and type of interaction between oxpeckers and their large mammalian hosts is poorly understood and still debated (Weeks, 2000; Nunn et al., 2011). A closer look at the distribution of birds among and within giraffes at HNP clearly shows how heterogeneous it is at the host level with many carrying no bird while a few has > 10 birds on them. We also provide empirical evidences for non-random choice of host body part by oxpeckers, a behaviour likely driven by the amount of ticks birds can find and share with conspecifics. Overall the observed oxpecker distribution among giraffes at HNP matches with distributions generally observed in parasitic organisms, although we reckon it may only mirror the tick load of individual hosts.

The distribution of oxpeckers on individual giraffes results from the host choice from the landscape to the giraffe body part, to prey ultimately upon ectoparasites. At the largest spatio-temporal scale, oxpeckers have to chose among large herbivore species, which is reflected by the host's preference index (PI). The observed PI $= 0.46 \pm 0.10$ for giraffes at HNP is similar to previously reported values in Kruger National Park (KNP), South Africa (0.90 for Grobler (1980); 0.54 for Ndlovu & Combrink (2015)). Surprisingly, Hustler (1987) found PI of 5.39 and 6.71 for giraffes in two separate areas of HNP. Host availability should indeed influence oxpeckers' choice because the decrease in abundance of a key host such as the giraffe may force birds to switch to another less preferred but more numerous host with little fitness costs (Hustler, 1987; Pyke, 1984). Here, host size plays a major role in host detection in flight and giraffes – like other large mammalian hosts – are easier to detect compared with smaller species (Grobler, 1980; Koenig, 1997). This is the main reason why the key host of oxpeckers alternates between buffaloes (Hustler, 1987), white rhinoceros (Ndlovu & Combrink, 2015) and giraffes (Grobler, 1980; Ndlovu & Combrink, 2015). The past high PI of giraffes at HNP suggests that birds must have exploited others hosts in the 80ies such as black rhinoceros, white rhinoceros, roans and sables (see Hustler, 1987, for details). This interpretation is supported by change in the composition of HNP's community of large herbivores over the last decades with, notably, the recent loss of

the white rhinoceros (Valeix et al., 2008), which is consistent with the low PI values we

report here for giraffes.

Focusing on the choice of individual giraffes by oxpeckers, we found a marked

asymmetric distribution of birds (exponential distribution) whereby many carried no bird and

a few ones were seen with up to 17 birds simultaneously (Fig. 2). This non-random

distribution of oxpeckers among individual giraffes usually likely results from an aggregation

behaviours (Palmer & Packer, 2018). The aggregation coefficient k we found for the

oxpecker distribution at HNP, close to 0, is typical of parasitic infections where only a few

individuals are massively infested (Shaw et al., 1998). That oxpeckers similarly aggregate

on a few giraffes would suggest they behave like parasites with their host in agreement with

previous studies (Plantan, 2009; Nunn et al., 2011). Birds could make use of public

information like conspecific density to chose a giraffe in a group (Doligez et al., 2004).

Because oxpeckers mostly feed on ectoparasites, the marked aggregation of birds could

indirectly mirror the distribution of ticks among giraffes. In mammals, infestation is indeed

highly variable among hosts (*e.g.* Shaw et al., 1998; Brunner & Ostfeld, 2008). For

instance, in roe deer (*Capreolus capreolus*), bank vole (*Clethrionomys glareolus*) and

mountain hare (*Lepus timidus*), most host individuals bear few ticks and only a few

individuals bear many (Horak et al., 1983; Talleklint & Jaenson, 1997; Mysterud et al.,

2014). From this hypothesis one could make indirect inference on tick burden of individual

giraffes from the number of hosted oxpeckers, given the birds distribute themselves

according to the ideal free distribution.

Within giraffes, we clearly found preferences for some body parts by oxpeckers. At HNP,

oxpeckers were mostly found on the neck and the back of giraffes followed by the head, the

abdomen, the lower limbs and the tail (see also Plantan, 2009; Ndlovu & Combrink, 2015,

for similar results). The mane seems the most preferred giraffes' body part of oxpeckers

(Koenig, 1997). This row of hairs seems to be a favourable habitat for ectoparasites by

providing shelter from predators (Ndlovu & Combrink, 2015) although oxpeckers use a

scissoring behaviour to easily pick parasites from the hairs Koenig (1997). It has been

noted that oxpeckers gather at the bigger host's mane to hide from predators or when they

are alarmed, which may contribute to increase their number in this area. However, because

giraffes rarely feed directly on the ground (Seeber et al., 2012) we expected a relatively low density of ticks on the giraffe's head. Our results support this assertion but oxpeckers could

sometimes forage the head seeking for other food resources such as saliva, mucus, earwax (Ndlovu & Combrink, 2015) or wounds. This also could be the case for female genitalia

where oxpeckers can also feed on mucus and secretion of their hosts (Weeks, 1999; Plantan, 2009). The abdomen, groins, thighs and tail present the lowest density of by

oxpeckers. Unlike mane, these areas are parts that can be easily groomed by giraffes, depleting tick quickly and making this area less preferred for oxpeckers (Koenig, 1997;

Ndlovu & Combrink, 2015). Assuming that abundance or presence of ticks is highest in the mane running on the neck and back of giraffes, our results would concur with the

distribution of oxpeckers as predicted by the optimal foraging theory (Pyke, 1984).

Density of oxpeckers differed substantially according to the sex of the giraffe, with male

hosts carrying 20% more birds than females (Fig. 4). This higher abundance of oxpeckers on male than on female giraffes likely proceed from their higher load in ectoparasites and

hence, larger food resources for the birds. The simplest explanation is that male giraffes being larger than females, more birds can feed on a male holding a constant *per capita*

food rate. Alternative explanations may also account for the preference of male hosts by oxpeckers, ranging from male-specific parasitic load, to the aggressive behaviour. Previous

studies who investigated the effect of host sex on the ectoparasite load are equivocal in mammals. For instance, Horak et al. (1987) reported more ticks (*Amblyomma hebraeum*) in

male than in female kudus, which could make the former more attractive to oxpeckers. Conversely, another study carried out in Scandinavia found that tick load was similar

whatever the sex and age of red deer (Mysterud et al., 2014). A proximate mechanism for why male giraffes would carry more ticks than females is intra-sexual fights for

reproduction. Neck fight, opponent chasing and female mounting indeed result in males having many injuries and open wounds all over their body (Nunn et al., 2011). Oxpeckers

being opportunist feeders they could benefit from the higher wound- and tick-feeding opportunities on male giraffes (Plantan, 2009). Alternatively, the handicap principle of the

sexual selection (Zahavi, 1975) proposes that, by surviving with many parasites or extravagant sexually selected traits, males would display honestly their intrinsic quality to females. Hence, oxpeckers could play the role of in indirect signal of ectoparasite load of male giraffes.

To evaluate the reliability of the oxpecker detection and location on giraffes from pictures, we carried out a double-observer experiment on a sub-sample of our images. We found the overall detection of the birds from photos to be very high (99%) but ideally all should be analyzed by 2 observers. Although not perfect, one person only (RG) scrutinized the 500 photographs henceforth keeping the detection and condition of observation similar for the complete data set. A major advantage of oxpecker counts from pictures is to make the counts and analyses repeatable and, contrary to what one might think, the use of one side of giraffes to locate and to count birds is an advantage because it avoids the issue of double counting. That said, we acknowledge that the major bias of our study was the location where the giraffe pictures were taken *i.e.* mainly around the many artificially maintained waterholes where large herbivores come to drink. When a large herbivore stand on the shore of waterholes, oxpeckers often use it as platform to rest, to sunbath, and to reach water to drink as well (Stutterheim, 1976). Consequently, the maximum number of oxpeckers per giraffe might read higher than it is elsewhere in HNP. To avoid this bias some studies tend to limit counting within 500 meters of water points (Grobler, 1980) but because HNP is densely covered with trees, observations of giraffes away from waterholes remained very difficult.

5 | CONCLUSION

Our study puts forward that the distribution and abundance of oxpeckers were surprisingly heterogeneous among and within giraffes. Some host body parts are clearly preferred for foraging by birds such as the neck and the mane because those areas could be suitable habitats for ticks. Gregarious hosts (female giraffes, buffaloes) travel and forage as a group thereby increasing local abundance and transmission of ticks (Koenig, 1997) to which oxpeckers could be excellent control agents on wild large herbivores and on domestic ones

too (Ndlovu & Combrink, 2015). From an ecological point of view, the oxpecker–large herbivores system proves to be highly relevant and useful for the study of host-parasite dynamics.

Acknowledgements

We would like to thank Simon Chamaillé-James and Marion Valeix for their critical reading of the manuscript. Thanks are extended to Martin Muzamba and Trust Dube for their invaluable help in data collection at Hwange, and to the park authorities of Zimbabwe for their support.

References

Bezuidenhout, J. D., & Stutterheim, C. (1980). A critical evaluation of the role played by the red-billed oxpecker *Buphagus erythrorhynchus* in the biological control of ticks. *The Onderstepoort Journal of Veterinary Research*, *47*, 51–75.

Brunner, J. L., & Ostfeld, R. S. (2008). Multiple causes of variable tick burdens on small-mammal hosts. *Ecology*, *89*, 2259–2272.

Clark, C. W., & Mangel, M. (1986). The evolutionary advantages of group foraging. *Theoretical population biology*, *30*, 45–75.

Dagg, A. I. (2014). *Giraffe: biology, behaviour and conservation*. Cambridge University Press.

Doligez, B., Pärt, T., Danchin, E., Clobert, J., & Gustafsson, L. (2004). Availability and use of public information and conspecific density for settlement decisions in the collared flycatcher. *Journal of Animal Ecology*, *73*, 75–87.

Emlen, S. T., & Oring, L. W. (1977). Ecology, sexual selection, and the evolution of mating systems. *Science*, *197*, 215–223.

Fiske, I., & Chandler, R. (2011). unmarked: An R package for fitting hierarchical models of wildlife occurrence and abundance. *Journal of Statistical Software*, *43*, 1–23.

Fretwell, S. D., & Lucas, H. L. (1969). On territorial behavior and other factors influencing habitat distribution in birds. *Acta Biotheoretica*, *19*, 16–36. doi:10.1007/bf01601953

Grobler, J. H. (1980). Host selection and species preference of the red-billed oxpecker *Buphagus erythrorhynchus* in the Kruger National Park. *Koedoe*, *23*. doi:10.4102/koedoe.v23i1.637

Gueron, S., & Levin, S. A. (1995). The dynamics of group formation. *Mathematical Biosciences*, *128*, 243–264.

Horak, I., MacIvor, K., Petney, T., & De, V. V. (1987). Some avian and mammalian hosts of *Amblyomma hebraeum* and *Amblyomma marmoreum* (Acari: Ixodidae). *The Onderstepoort Journal of Veterinary Research*, *54*, 397–403.

Horak, I., Potgieter, F., Walker, J. B., De Vos, V., & Boomker, J. (1983). The ixodid tick burdens of various large ruminant species in South African nature reserves. *Onderstepoort Journal of Veterinary Research*, *50*, 221–228.

Hustler, K. (1987). Host preference of oxpeckers in the Hwange National Park, Zimbabwe. *African Journal of Ecology*, *25*, 241–245. doi:10.1111/j.1365-2028.1987.tb01115.x

Hutchinson, J. M., & Waser, P. M. (2007). Use, misuse and extensions of "ideal gas" models of animal encounter. *Biological Reviews*, *82*, 335–359.

Johnson, D. H. (1980). The comparison of usage and availability measurements for evaluating resource preference. *Ecology*, *61*, 65–71.

Koenig, W. D. (1997). Host preferences and behaviour of oxpeckers: co-existence of similar species in a fragmented landscape. *Evolutionary Ecology*, *11*, 91–104. doi:10.1023/a:1018439614008

Krebs, C. J. (1972). The experimental analysis of distribution and abundance. *New York: Harper and Row*.

Lack, D. L. (1966). *Population studies of birds*. Clarendon P.

Lotka, A. J. (1956). Elements of mathematical biology.

MacArthur, R. H. (1972). *Geographical ecology: patterns in the distribution of species*. Princeton University Press.

May, R. M. (1978). Host-parasitoid systems in patchy environments: a phenomenological model. *Journal of Animal Ecology*, *47*, 833–843.

Morales, J. M., Moorcroft, P. R., Matthiopoulos, J., Frair, J. L., Kie, J. G., Powell, R. A., Merrill, E. H., & Haydon, D. T. (2010). Building the bridge between animal movement and population dynamics. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*, 2289–2301.

Moreau, R. E. (1933). The food of the red-billed oxpecker, *Buphagus erythrorhynchus* (Stanley). *Bulletin of Entomological Research*, *24*, 325. doi:10.1017/s000748530003162x

Mysterud, A., Hatlegjerde, I. L., & Sørensen, O. J. (2014). Attachment site selection of life stages of *Ixodes ricinus* ticks on a main large host in Europe, the red deer (*Cervus elaphus*). *Parasites & Vectors*, *7*. doi:10.1186/s13071-014-0510-x

Ndlovu, M., & Combrink, L. (2015). Feeding preferences of oxpeckers in Kruger National Park, South Africa. *Koedoe*, *57*. doi:10.4102/koedoe.v57i1.1316

Nichols, J. D., Hines, J. E., Sauer, J. R., Fallon, F. W., Fallon, J. E., & Heglund, P. J. (2000). A double-observer approach for estimating detection probability and abundance from point counts. *The Auk*, *117*, 393–408.

Nunn, C. L., Ezenwa, V. O., Arnold, C., & Koenig, W. D. (2011). Mutualism or parasitism? Using a phylogenetic approach to characterize the oxpecker-ungulate relationship. *Evolution*, *65*, 1297–1304. doi:10.1111/j.1558-5646.2010.01212.x

Ostfeld, R. S., & Keesing, F. (2000). Pulsed resources and community dynamics of consumers in terrestrial ecosystems. *Trends in Ecology & Evolution*, *15*, 232–237.

Palmer, M. S., & Packer, C. (2018). Giraffe bed and breakfast: Camera traps reveal Tanzanian yellow-billed oxpeckers roosting on their large mammalian hosts. *African Journal of Ecology*. doi:10.1111/aje.12505

Pielou, E. (1969). An introduction to mathematical ecology. *An introduction to mathematical ecology.*

Plantan, T. B. (2009). *Feeding behaviour of wild and captive oxpeckers (*Buphagus spp.*): a case of conditional mutualism.* Ph.D. thesis, University of Miami.

Pyke, G. H. (1984). Optimal foraging theory: A critical review. *Annual Review of Ecology and Systematics*, *15*, 523–575. doi:10.1146/annurev.es.15.110184.002515

R Core Team (2018). *R: A Language and Environment for Statistical Computing.* R Foundation for Statistical Computing, Vienna, Austria.

Samish, M., & Rehacek, J. (1999). Pathogens and predators of ticks and their potential in biological control. *Annual Review of Entomology*, *44*, 159–182. doi:10.1146/annurev.ento.44.1.159

Seeber, P. A., Ndlovu, H. T., Duncan, P., & Ganswindt, A. (2012). Grazing behaviour of the giraffe in Hwange National Park, Zimbabwe. *African Journal of Ecology*, *50*, 247.

Shaw, D., Grenfell, B., & Dobson, A. (1998). Patterns of macroparasite aggregation in wildlife host populations. *Parasitology*, *117*, 597–610.

Shorrocks, B. (2016). *The giraffe: biology, ecology, evolution and behaviour*. John Wiley & Sons.

Sjöberg, M., Albrectsen, B., & Hjältén, J. (2000). Truncated power laws: a tool for understanding aggregation patterns in animals? *Ecology letters*, *3*, 90–94.

Stutterheim, C. (1976). *The biology of the red-billed oxpecker,* Buphagus erythrorhynchus *(Stanley 1814) in the Kruger National Park*. Master's thesis, University of Pretoria, South African.

Stutterheim, I. M., Bezuidenhout, J. D., & Elliot, E. (1988). Comparative feeding behaviour and food preferences of oxpeckers (*Buphagus erythrorhynchus* and *B. africanus*) in captivity. *The Onderstepoort Journal of Veterinary Research*.

Talleklint, L., & Jaenson, T. G. (1997). Infestation of mammals by *Ixodes ricinus* ticks (Acari: Ixodidae) in south-central Sweden. *Experimental & applied acarology*, *21*, 755–771.

Taylor, L., Woiwod, I., & Perry, J. (1979). The negative binomial as a dynamic ecological model for aggregation, and the density dependence of *k*. *The Journal of Animal Ecology*, (pp. 289–304).

Valeix, M., Fritz, H., Chamaillé-Jammes, S., Bourgarel, M., & Murindagomo, F. (2008). Fluctuations in abundance of large herbivore populations: insights into the influence of dry season rainfall and elephant numbers from long-term data. *Animal Conservation*, *11*, 391–400.

van Someren, V. D. (1951). The red-billed oxpecker and its relation to stock in Kenya. *The East African Agricultural Journal*, *17*, 1–11.

Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S*. New York: Springer, fourth edition. ISBN 0-387-95457-0.

Veríssimo, D., Cugnière, J.-C., Cugnière, S., Cugnière, J., Cugnière, G., & Cugnière, L. (2017). Record number of yellow-billed oxpeckers *Buphagus africanus* Linnaeus, 1766 (Aves: Passeriformes: Buphagidae) foraging on a single host. *Journal of Threatened Taxa*, *9*, 9768. doi:10.11609/jott.3055.9.1.9768-9770

Weeks, P. (1999). Interactions between red-billed oxpeckers, *Buphagus erythrorhynchus*, and domestic cattle, *Bos taurus*, in Zimbabwe. *Animal Behaviour*, *58*, 1253–1259. doi:10.1006/anbe.1999.1265

Weeks, P. (2000). Red-billed oxpeckers: vampires or tickbirds? *Behavioral Ecology*, *11*, 154–160. doi:10.1093/beheco/11.2.154

Wittenberger, J. F. (1981). *Animal social behavior*. Duxbury Press.

Zahavi, A. (1975). Mate selection – a selection for a handicap. *Journal of theoretical Biology*, *53*, 205–214.

Fig. 1 Hwange National Park map, Zimbabwe. The giraffe study from which the oxpecker data derived from is located in the Main Camp area, covering the north east of the park from Ngweshla / Somalisa to Giraffe Springs passing through Jambile.

Fig. 2 Distribution of oxpecker (*Buphagus sp.*) number on individual giraffes (*Giraffa cameelopardalis*) at Hwange National Park, Zimbabwe. In black is the observed distribution and in grey the expected distribution according to a Poisson distribution law taking the observed mean as parameter ($\lambda = 2.16$). Note the marked over-representation of giraffes carrying no bird and the long distribution tail of giraffes with numerous birds on their body in the observed data.

(A)

(B)

Fig. 3 (A) Defined zonation and (B) proportion of oxpeckers (*Buphagus sp.*) counted on the 14 different body parts of giraffes (*Giraffa camelopardalis*) observed at Hwange National Park, Zimbabwe. The 14 different body parts we defined were: AB: abdomen, AG: ano-genital, BA: back, BU: buttock, GR: groin, HE: head, LL: lower leg, MA: mane, NE: neck, SC: scapula, SH: shoulder, TA: tail, TH: thigh, UL: upper leg. Although the general patterns were similar the use of giraffe body by oxpeckers differed significantly (*n* = 683 giraffes).

Fig. 4 Distribution of oxpecker (*Buphagus sp.*) number on male and female giraffes (*Giraffa camelopardalis*) at Hwange National Park, Zimbabwe. Curves are the predicted frequencies as given by a negative binomial distribution model which parameter have been estimated separately for the two sexes. Vertical dashed lines represent the mean number of oxpeckers carried by individual giraffes. Note that males with no oxpecker are less frequent than females, and the largest aggregations of oxpeckers have been found on male giraffes.