Fungi form interkingdom microbial communities in the primordial human gut that develop with gestational age

Kent A. Willis[1], John H. Purvis[1], Erin D. Myers[2], Michael M. Aziz[4], Ibrahim Karabayir[5], Charles K. Gomes[3], Brian M. Peters[6], Oguz Akbilgic[5], Ajay J. Talati[1,4], Joseph F. Pierre[3,8]*

[1]Department of Pediatrics, Division of Neonatology, College of Medicine, The University of Tennessee Health Science Center (UTHSC), Memphis 38103 TN, USA
[2]College of Medicine, UTHSC
[3]Deparment of Pediatrics, Obesity, College of Medicine, UTHSC
[4]Department of Obstetrics and Gynecology, College of Medicine, UTHSC.
[5]Department of Health Informatics and Data Science, Stritch School of Medicine, Loyola University Chicago, Chicago 60660 IL, USA
[6]Department of Clinical Pharmacy and Translational Science, College of Pharmacy, UTHSC
[7]Center for Health Outcome and Informatics Research, Loyola University Chicago, Chicago, IL, USA
[8]Deparment of Microbiology, Immunology and Biochemistry, College of Medicine, UTHSC

*Correspondence and request for materials should be addressed to J.F.P. (email: jpierre1@uthsc.edu)

Keywords Meconium; Microbiome; Mycobiome; Premature birth; Gut ecology; Dectin-1

Abstract

Fungal and bacterial commensal organisms play a complex role in the health of the human host. Expansion of commensal ecology after birth is a critical period in human immune development. However, the initial fungal colonization of the primordial gut remains undescribed. To investigate primordial fungal ecology, we performed amplicon sequencing and culture-based techniques of first-pass meconium, which forms in the fetal intestine prior to birth, from a prospective observational cohort of term-born and preterm newborns. Here, we describe fungal ecologies in the primordial gut that develop complexity with advancing gestational age at birth. Our findings suggest homeostasis of fungal commensals may represent an important aspect of human biology present even before birth. Unlike bacterial communities which gradually develop complexity, the domination of the fungal communities of some preterm infants by Saccromycetes, specifically *Candida*, may suggest a pathologic association with preterm birth.

Introduction

Human health is intimately entwined with the commensal microorganisms that constitutionally colonize our barrier surfaces, such as the intestinal tract. The human gut microbiome is host to a complex interkingdom ecology of bacteria, fungi, archaea, protozoa and viruses. While recent research has led to a drastic expansion in our understanding of bacterial communities in the gut[1-4], little is known about the role of fungal organisms, collectively the mycobiome, particularly in relation to their impact on the population dynamics of early-life colonization.

The tightly choreographed maturation of commensal microbial communities is crucial to mammalian immune development[5]. Fungal organisms can form a crucial nidus for initial colonization[6] and may play a vital role in facilitating early colonization of the neonatal gut. Disruption of this process could produce pathogenic fungal overgrowth or invasive disease. Indeed, this mechanism may play a role in the development of asthma and other metabolic and immunological diseases[7]. The intestinal mycobiota likely confers unique physiologic effects to the host, in part by processing nutrients and vitamins either separately or in conjunction with the microbiome. However, characterization of the functional impact of early mycobiome is impaired by limited knowledge of the normal development of these communities. To understand the functional impact of the early-life mycobiome on human health, it is necessary to determine if the human primordial gut hosts fungal communities, if these communities are established before or after birth and if the presence of fungi in the early gut impacts neonatal physiology.

Advances in next-generation sequencing techniques have pushed back the timing of initial commensal colonization[3] which was initially thought to occur at birth[8,9]. Despite these advances, interkingdom interactions within microbial communities and the influence of environmental factors on the dynamics of microbial ecosystems remains poorly understood. Due to delayed progress in sequencing and characterizing fungal communities[10], the appreciation of commensal fungal ecology

and biomedical impact of fungi lags behind the bacterial microbiome. It appears that during the first

several years of life, early fungal communities, like their bacterial counterparts, arise from maternal

communities, but unlike their bacterial counterparts, remain at a relatively low and consistent

diversity[11,12]. The relative consistency of early mycobiome makes understanding initial colonization

important to unravel both the long-term biomedical impact of fungal communities and to explain their

divergence from bacterial communities. While we are beginning to understand the development of the

mycobiome during early-life, the timing of initial fungal colonization remains undescribed.

To characterize the interkingdom ecologies of the fetal microbiome and understand when fungal

colonization first occurs, we performed a molecular and culture-based survey of first-pass meconium

collected from an observational cohort of preterm and term-born human newborns. In this study, we

characterize fungal communities in first-pass meconium within hours of birth and explore the influence

of perinatal factors on the foundational ecology of the human gut, laying the groundwork for

understanding the functional impact of fungal communities on the host. We support our observations

by examining the fetal and neonatal mouse mycobiome. We describe complex bacterial and fungal

commensal communities that develop in abundance and diversity with length of gestation. Our results

suggest that the human fetus is naturally exposed an increasing variety of microbial DNA as gestation

progresses. However, the increased prevalence of fungal order Saccharomyces, especially the genus

Candida, in preterm infants may also suggest a pathologic association with preterm birth.

86
87 **Results**
88
89
90 **Study design and cohort characteristics**
91
92 To characterize the fungal ecology of the primordial human gut, we collected the first-pass meconium

93 from a cohort of very low birthweight preterm (<1500 g, gestational age 23-32 weeks, n = 54) and

94 term-born (gestational age 37-41 weeks, n = 36) newborns and characterized the microbial

95 communities using both culture-independent and culture-dependent techniques (Fig. 1). Of these 90

infants prospectively enrolled from a Level IIIB Neonatal Intensive Care Unit and Well Baby Nursery at single academic medical center, 19 infants (31.5% of preterm and 5.6% of term-born infants, χ^2 p =0.0033) did not produce a meconium stool that could be collected during the first 48 h of life (Fig. 1a). To differentiate the effects of perinatal factors on microbial ecology, we prospectively collected clinical data describing perinatal exposures (Fig. 1b). In addition to lower gestational age, infants born preterm were more likely than term-born infants to have a lower birthweight, a lower Apgar and higher Critical Risk Index for Babies II (CRIB-II) scores, receive postnatal antibiotics and be delivered by cesarean section (Table 1).

Figure 1

Fig. 1. Curation of clinical and microbial data. **a** Schematic of patient allocation. VLBW, very low birthweight (preterm infants < 1500 g). **b** Schematic of data acquisition and analysis. Low biomass samples of meconium, an intralumenal material formed in the intestine prior to birth, were analyzed by culture-based techniques or Illumina MiSeq and quantitative PCR after high-fidelity DNA extraction. Sequence reads were subjected to exacting data processing to remove potential contaminants. Clinical demographics were used to perform ecological modeling of microbial communities and develop sequential rarified random forest classifier machine learning models. A mouse model of normal pregnancy was utilized to quantify intrauterine fungal biomass. 16S rRNA, 16S ribosomal RNA (bacteria and archaea). ITS rDNA, internal transcribed spacer ribosomal DNA (fungi).

121 **Table 1. Cohort Characteristics**

	Preterm	Term	*P* Value
Stooled < 48 h	37	34	
Male Sex	49%	56%	0.5580
Black/African American	78%	91%	0.1361
Maternal Age (yr)	26.94 ± 6.00	25.15 ± 4.64	0.1731
Gestational Age (wk)	27.51 ± 2.71	39 ± 1.15	<0.0001
Birthweight (g)	874.1 ± 266.81	3115 ± 360.06	<0.0001
CRIB-II Score	10 (8:13)	0 (0:0)	<0.0001
Apgar Score (5 min)	6 (5:9)	9 (9:9)	<0.0001
Cesarean Delivery	68%	35%	0.0058
Prenatal Antibiotics	32%	38%	0.5987
Prenatal Antibiotic Doses* (n)	1 (1:1.5)	2 (1:2)	0.2231
Prenatal Antibiotic Doses (n)	0 (0:1)	0 (0:1)	0.7943
Postnatal Antibiotics	92%	0%	<0.0001
16S rRNA Positive	92%	94%	0.5397
ITS rDNA Positive	88%	65%	0.0224

*Number of doses received prenatally in those who received antibiotics. n (quartile 25%: quartile 75%) or n ± SD.

Fungal community composition differs by gestational age

The bacterial community composition of the microbiome differs by gestational age at birth[13]. This may

be related to differences in initial meconium colonization[3]. To explore if fungal colonization of the

microbiome was influenced by gestational age, we performed a binary comparison of infants born at

<33 weeks' (n = 37) and infants > 33 weeks' gestation (n = 34). Meconium samples from preterm

infants were more likely to contain fungal DNA than samples from term-born infants using internal

transcribed spacer region 2 ribosomal DNA (ITS2 rDNA) amplicon sequencing at a read depth of 300x

(88 versus 65%, χ^2 p = 0.0224, Table 1). The alpha diversity of these fungal sequences clearly

increased with increasing gestational age (Shannon diversity index, ANOVA f = 4.5 p =0.04. Principal

coordinates analysis (PCoA) of Bray-Curtis dissimilarity matrices demonstrated fungal community

structure was different by permutational multivariate analysis of variance (PERMANOVA, R^2 =0.308 p

=0.01, Fig. 2c). Discriminant analysis of principal components suggests that the principal variables at

the genus level are the prevalence of *Elmerina*, *Cladosporium, Candida* and *Stereum*. Similarly,

canonical correspondence analysis (CCA, χ^2 distance = 0.71 f = 1.14 p =0.004), and redundancy

analysis (RDA, variance =3.83 f =1.22 p =0.001) were both consistent with a clear differentiation in

fungal community structure by advancing gestational age (Fig. 3d).

Network analysis supports the sparseness of fungal communities, particularly in preterm infants,

which is likely related to the low fungal diversity (Fig 2d). However, as quantified by positive

Spearman's correlations, fungal networks do appear to develop complexity with increasing gestational

age.

The relative composition of the fungal communities varied with gestational age (Fig. 2 and 3). At the

order level, preterm infants were primarily colonized with Saccharomycetales (73.4%), with other

fungal taxa accounting for less than a quarter of assigned reads. Term-born infants possessed

mycobiomes that were less dominated by a singles species than preterm infants. Saccharomycetales

and Polyporales accounted still accounted for about half of the fungal burden in infants with a later

gestation (22.1% and 27.7%, respectfully), but other minor taxa were more markedly prevalent with

increasing gestational age. In addition to Polyporales (5.2% < 33 weeks' and 22.1% > 33 weeks'

gestation), Russulales showed the most pronounced expansion with increasing gestational age (0.8%

< 33 weeks' and 14.5% > 33 weeks' gestation, Fig. 2f, 3a). The most significant expansions in

abundance at the genus level were in *Stereum*, *Malassezia*, *Elmerina* and *Armillaria*. In contrast to

this trend, the domination of the mycobiome by *Candida* in some preterm infants lead to this particular

genus being uniquely more abundant in preterm infants overall. Finally, core microbiome analysis also

supports the expansion of fungal communities with increasing gestational age, with the addition of 2

unique core phyla and 3 unique core OTUs in term-born as compared to preterm infants (Fig. 3e).

To further investigate the association of specific fungal taxa with gestational age, we performed linear

discriminant analysis of effect size (LEfSe) to identify fungal taxa that could perform as high-

dimensional biomarkers for prematurity. The prevalence of the fungal order Saccharomyces and

specifically the genus *Candida* reliably identified preterm samples. In addition, the prevalence of

fungal class Elmerina and the genera *Stereum* and *Malassezia* reliably identified samples from full-

term newborns (Supplemental Fig. 3i).

Figure 2

Fig. 2. Fungal ecologies in human meconium vary by the gestational age at birth.
a Sequencing rarefaction analysis of species richness.
b Alpha diversity quantified by the Shannon index, ANOVA $f = 4.5$ $p = 0.04$.
c Principal coordinates analysis of Bray-Curtis dissimilarity matrices, PERMANOVA $R^2 = 0.308$ $p = 0.01$. The subset displays a discriminant analysis of principal components.
d Network analysis developed using Pearson's correlations. Positive correlations with false discovery rate-adjusted, p-values <0.05 are presented as an edge.
e Heat map of differential distribution of taxa at the order level, ranked by gestation.
f Relative abundance of taxa at the phylum level. Sqrt(TSS), square root total sum normalization (Hellinger transformation).
For all analyses n=71. Preterm samples are displayed in red and term-born samples in blue.

Fig. 3. Fungal communities increase in complexity with advancing gestational age at birth.
a Relative abundance at the phylum level. Sqrt(TSS), square root total sum normalization (Hellinger transformation).
b Distribution of key taxa at the genus level. ANOVA, Bonferroni, * $p < 0.05$, ** $p < 0.01$.
c Twenty most abundant interkingdom taxa at the order level.
d Canonical correspondence analysis (CCA), $\chi^2 = 0.71$ $f = 1.14$ $p = 0.004$, and redundancy analysis (RDA), variance $= 3.83$ $f = 1.22$ $p = 0.001$.
e Core interkingdom OTUs and unique phyla between preterm and term-born infants. OTU, operational taxonomic unit.
Data are median ± IQR (n=71). Preterm samples are displayed in red and term-born samples in blue.

In childhood, fungal colonization has been shown to be influenced by host sex[14]. We examined fungal

community composition in our cohort to see if fungal populations in meconium were influence by the

sex of the host. Alpha diversity of was unaltered by sex (Supplemental Fig. 4g). Relative abundance

differences were most pronounced for Polyporales (24.1% in females and only 6.9% in males), with

other orders having less dramatic differences (Supplemental Fig. 4h). Meconium fungal community

structure was not influenced by host sex (PCoA, PERMANOVA R^2 =0.0209 p =0.529, Fig. 4a).

Similarly, redundancy analysis (RDA, variance =3.19 f =1.02 p =0.348) was also not significant.

Antibiotics can have a profound effect on the compositional structure of the microbiome[15], and lead to

changes in the fungal colonization as well [16]. To characterize the potential effect of maternal antibiotic

exposure on the initial structure of the mycobiome, we examined how any antibiotic exposure altered

fungal composition of the meconium. In our samples, perinatal antibiotic exposure did not produce

significant differences in alpha diversity (Supplemental Fig. 4d). The relative abundance of

Saccharomycetales was increased by maternal antibiotic exposure (60.1% after antibiotic exposure,

and 32.5% without antibiotic exposure, Supplemental Fig. 4e). The distribution of other major taxa

was similar. Fungal community structure was not significantly altered by prenatal antibiotic exposure

(PERMANOVA R^2 =0.202 p =0.598 and RDA, variance =3.51 f =1.12 p =0.083, Fig. 4b).

Whether mode of delivery influences microbial colonization remains controversial[2,17]. To explore

influence of the delivery mode on fungal colonization we examined fungal community structure using

cesarean or vaginal delivery as a binary variable. Alpha diversity was not significantly altered by mode

of delivery, although cesarean born infants did demonstrate more variability (Supplemental Fig. 4a).

Relative composition differences showed some variation by mode of delivery (Supplemental Fig. 4b)

Similar to preterm infants, samples from infants born via caesarean were predominantly colonized by

Saccharomycetales (61.7% after cesarean and 39.7% after vaginal). Both groups expressed similar

levels of Polyporales (15.3% after cesarean and 13.8% after), but Russulales was more prevalent in

vaginally-born infants (13.2% after vaginal and 0.62% after cesarean). While preterm infants were

more likely to be delivered by cesarean in our cohort (68% versus 35% χ^2 p =0.0058, Table 1), as

characterized by PCoA, fungal community structure did not significantly differ after either vaginal or

cesarean delivery (PERMANOVA R^2 =0.202 p =0.598 and RDA, variance =3.51 f =1.12 p =0.083, Fig.

4c), which is consistent with the formation of meconium prior to birth.

Increasing illness severity closely associates with decreasing gestational age and is therefore a

covariate and potential confounder in many diseases related to preterm birth[18]. To gauge a potential

confounding influence of illness severity, we examined fungal community composition in relation to a

critical score of 11 on the CRIB-II. No differences were appreciated in alpha diversity as quantified by

the Chao1 index (Supplemental Fig. 4j). The relative abundance differences were also less markedly

altered than for other perinatal factors (Supplemental Fig. 4k). The abundance of Saccharomycetales

was unaltered at the class level, but the abundance of Polyporales was reduced (15.5% with score

<11 and 6.9% with a score ≥11) and the relative abundance of Capnodiales and Dothideales

increased with elevated CRIB-II score (6.0% <11 and 13.5%, ≥11; and 1.6% <11 and 10.8% ≥11,

respectfully). Clinical illness severity did not appear to alter fungal community composition

(PERMANOVA R^2 =0.0251 p =0.146 and RDA, variance =2.68 f =0.85 p =0.947, Fig. 4d).

Fig. 4. Fungal community structure is not determined by common perinatal factors.
a Host sex does not significantly alter fungal community composition. Principal coordinates analysis
(PCoA) of Bray-Curtis dissimilarity matrices, PERMANOVA R^2 =0.0209 p =0.529. Redundancy
analysis (RDA), variance =3.19 f =1.02 p =0.348. The subset displays a discriminant analysis of
principal components.
b Prenatal antibiotic exposure (PAE) does not significantly alter fungal community composition. PCoA
of Bray-Curtis dissimilarity matrices, PERMANOVA R^2 =0.202 p =0.598. RDA, variance =3.51 f =1.12
p =0.083.
c Mode of delivery does not significantly alter fungal community composition. PCoA of Bray-Curtis
dissimilarity matrices, PERMANOVA R^2 =0.202 p =0.598. RDA, variance =3.51 f =1.12 p =0.083.
d Illness severity does not significantly alter fungal community composition. PCoA of Bray-Curtis
dissimilarity matrices, PERMANOVA R^2 =0.0251 p =0.146. RDA, variance =2.68 f =0.85 p =0.947.
CRIB-II, Clinical Risk Index for Babies II.
For all analyses n = 71.

Fungi form complex interkingdom communities

We examined how fungi formed interkingdom microbial communities by the characterizing the

resulting community composition from the combined 16s and ITS sequencing results (n =71). For

most infants, fungal communities were a minor component their interkingdom microbiome, but a few

preterm infants had microbiomes that were dominated by fungi (Fig. 5 and Supplemental Fig. 1). This

may emphasize the importance of pioneer species to the ecological succession of the primordial gut.

In general, the interkingdom microbiome increased in complexity with increasing gestational age at

birth. While rarefaction analysis favored more microbial richness in term samples (Fig. 5a), alpha

diversity as quantified by the Shannon diversity index was not significantly altered (ANOVA f =2.2 p

=0.14, Fig 5b). PCoA of the combined microbial communities showed clear differential clustering for

with a binary cut off at 33 weeks' gestation (Bray-Curtis dissimilarity matrices, PERMANOVA with 999

permutations R^2 =0.103 p =0.000333, Fig. 5c). Discriminant analysis of principal components

suggests that the principal variables at the genus level are the prevalence of *Methylobacterium*,

Bacteroides and *Stereum*. The most prominent differences in abundance at the phylum level were in

Blasidiomycota, Bacteroidetes and Actinobacteria, all of which were more abundant in term born

infants. Acidobacteria, in contrast, were more prevalent in preterm infants (Supplemental Fig. 1b). At

the genus level, there were expansions in multiple genera with increasing gestational age. The most

prominent changes were in *Methylobacterium* and *Bacteroides* (Supplemental Fig. 1d). However,

there were some genera, in particular *Lactobacillus* and *Staphylococcus*, that were more abundant in

preterm infants, but at a much lower level. The expansion of the interkingdom microbiome with

increasing gestation is also supported by analysis of the core microbiome of term infants which

increased from 24 to 36 unique OTUs and added 7 unique phyla to the core preterm microbiome

(Supplemental Fig. 1e).

To understand interrelatedness in interkingdom ecology, we used network analysis. Compared to the

sparse inter-fungal networks, fungi and bacteria form complex interkingdom networks with numerous

interactions (Figure 5d). The expanding networks led to multiple unique nodes clearly separated by

gestational age at birth.

To characterize the influence of perinatal factors on interkingdom community structure, we used

PCoA and RDA to capture the impact of host sex, prenatal antibiotic exposure, mode of delivery and

illness severity. Similar to fungal community structure, the interkingdom structure was not significantly

altered by these common perinatal exposures as assessed by both constrained and unconstrained

analyses (Supplemental Fig. 2).

Fig 5. Fungi and bacteria form complex interkingdom microbial communities in human meconium.
a Sequencing rarefaction analysis of species richness.
b Alpha diversity quantified by the Shannon index, ANOVA f =2.2 p =0.14.
c Principal coordinates analysis of Bray-Curtis dissimilarity matrices, PERMANOVA with 999 permutations R^2 =0.103 p =0.000333. The subset displays a discriminant analysis of principal components.
d Network analysis developed using Spearman's correlations. Positive correlations with false discovery rate-adjusted, p-values <0.05 are presented as an edge.
e Heat map of differential distribution of taxa at the order level, ranked by gestation.
f Relative abundance of taxa at the phylum level. Sqrt(TSS), square root total sum normalization (Hellinger transformation).
For all analyses (n=71). Preterm samples are displayed in red and term-born samples in blue.

Key interkingdom community members predict gestational age of the host

To explore the distribution of microbial taxa with respect to gestational age at birth, we developed a machine learning model to predict the gestational age of a newborn based on the interkingdom composition of the microbiome. We developed our first machine learning model using only microbial taxa. Starting with 51 fungal and 209 bacterial taxa that were identified at least twice in the dataset, we utilized LASSO with 5-fold cross validation to identify potential significant predictors, leading to the identification of 9 fungal and 34 bacterial taxa clearly distributed by gestational age (Supplemental Fig. 6a). A support vector machine model reliably classified infants as preterm or term utilizing these taxa (Supplemental Fig. 6b).

Interkingdom community composition classifies preterm status

To understand the relative strength of key microbial taxa to function as predictors of preterm status relative to other clinical demographic data, we implemented a random forest model with 5-fold cross validation using the presence of 260 genus-level microbial taxa and select clinical demographics. After 100 independent permutations, average sensitivity analysis of the final random forest model provided an area under the curve (AUC) of 0.859 [95% confidence interval (CI) 0.857-0.862, Supplemental Fig. 6c]. We then performed a variable importance analysis on the random forest model and selected the top 10 variables to build a rarified random forest model using 5-fold cross validation. After 100 permutations, the average performance of the sensitivity analysis of the rarified random forest models had an AUC 0.867 (95% CI 0.861-0.871). Overall, the strongest predictors from these key taxa ran counter to the general trend and were more abundant in preterm than in term-born infants (Undefined genera within family Rikenellaceae, and the bacterial genera *Parabacteroides* and *Citrobacter*). With the exception of *Lactococcus*, the other key genera were more abundant in term-born infants. Of note, maternal body mass index and age also functioned as reliable predictors, which may be related to their known association with preterm birth[19].

We compared the random forest models to a variety of other classifiers. All other methods compared

provided classification accuracy < 71%, which was significantly lower than the accuracy of the random

forest classifiers (Supplemental Fig. 6d). During model development, we did not include clinical

variables such as prenatal steroid use, duration of rupture of membranes, antibiotic use and delivery

mode as predictors because such variables are closely associated with impending delivery and would

confound predictive modeling. However, we implemented Mann-Whitney-U (or Kruskal-Wallis) tests to

determine whether there were significant differences between the presence of these variables and the

10 key microbial taxa we previously identified during development of the random forest model. We

found no significant differences between antibiotics use, duration of membrane rupture or delivery

mode and presence of these 10 taxa. In contrast, there were statistically significant differences (p

<0.01) between use of prenatal steroids (none, 1 or 2 doses) and the abundance of 7 of these key

taxa (Supplemental Fig. 7a).

To quantify the performance of higher-order microbial taxa, we developed models exploring the

strength of different phylogenic levels to function as predictors of prematurity. We designed our

modeling approach in six levels to capture the influence of using increasingly more granular taxa

information on the classification accuracy. The accuracy of the model increased as the as the

phylogenic level of the taxa used decreased, highlighting the importance of the distribution of key tax

and their variation with gestational age (Fig. 5 and Supplemental Fig. 7b). In addition, when using

higher level microbial taxa as potential predictors, the mothers' age and body mass index were

relatively more important as predictors. These statistical models demonstrate the differential variation

of the interkingdom community composition and underscores the close relationship between the

microbial community composition and gestational age of the host at birth.

Fig. 6. Interkingdom microbial community composition accurately classifies preterm infants. The precision of rarified random forest classifier machine learning models developed using microbial taxa from each level of the phylogenic tree increases as the phylogenic level decreases, highlighting the importance of key microbial taxa. Darker shading indicates a more important predictor to the machine learning model. Blue represents bacteria, red represents fungi and black represents archaea and clinical demographics. Starting from the center and moving outward, concentric rings represent models developed utilizing taxa on the kingdom, phylum, class, order, family, and genus levels. The 10 most important predictors based on the genus-level model are highlighted in bold. Area under the curve statistics for each level are reported in Supplementary Fig. 6b.

Fungal and microbial organisms can be cultured from the primordial microbiome

To explore if our next generation sequencing-based analyses were potentially representative of live microorganisms as opposed to translocated microbial DNA, we performed anaerobic and aerobic culture-based assays for both fungi and bacteria on a subset of samples (n = 41) for which sufficient meconium was available (Supplemental Fig. 8). Aerobic fungi-selective culture media was positive in 32% of preterm and 29% of term infant samples (χ^2 p = 0.835). Aerobic bacteria-selective culture media was positive in 89% of preterm and 71% of term infant samples (χ^2 p = 0.128). Anaerobic fungi-selective culture media was positive in 50% of preterm and 21% of term-born infant samples (χ^2 p = 0.079). Anaerobic bacteria-selective culture media was positive in 69% of preterm and 88% of term infant samples (χ^2 p = 0.1543). Overall, bacterial cultures were diffusely positive, while fungal cultures were less likely to demonstrate growth. While contamination cannot be completely excluded, we also cannot eliminate the possibility of live fungi and bacteria in the primordial human gut.

The fetal gut contains fungal DNA

Considerable controversy persists as to whether the isolation of microbial DNA from the human placenta or newborn gut is representative of live microbial organisms[20]. While the similarities between microbes observed in the placenta and neonatal meconium suggest transfer of microbial DNA across the placenta, they do not represent direct observation of the fetal intestine. However, bacterial DNA has been identified in the fetal gut in a mouse model of normal pregnancy[21]. To directly investigate the fetal intestine for the presence of fungal DNA and live organisms we developed a mouse model of normal pregnancy. Using C57BL6J mice, we collected samples from the fetal gut and several maternal body sites on either estimated gestational day 17 (E17) or within 24 h of birth (P1). We used real time PCR to identify ITS biomass across all maternal and fetal samples. The relative fungal biomass was about 14-fold lower in the fetal intestine before birth than in the maternal amniotic fluid. However, fungal biomass was about 6-fold more abundant in the neonatal intestine than the amniotic

fluid (mean cycle threshold, C_t = 36.75 ±15.48 at E17 and C_t = 25.71 ±15.16 at P1 and C_t = 29.20

±6.58 in amniotic fluid).

Dectin-1 is a pattern recognition receptor that recognizes the fungal cell-wall β-glucan and is important

for detection of fungal pathogens in humans and mice[22]. To investigate if inhibited host sensing of

fungi altered the transplacental deposition of fungal biomass in the neonatal gut, we collected

samples from the fetal gut and multiple maternal body sites from dectin-1[-/-] mice on the C57Bl6J

background (B6.clec7a[tm1Gdb]). In the fetal intestine, dectin-1[-/-] mice had about a 6-fold increase in

fungal biomass over dectin-1 competent mice (C_t = 33.50 ± 14.81 versus C_t = 36.75 ±15.16).

However, after birth, the neonatal intestinal fungal biomass was lower on P1 (C_t = 29.33 ±18.01

versus C_t = 25.71 ±15.48) and remained a more significant proportion of the intestinal biomass. While

there are considerable differences between the placentas of humans and mice[23], these results

suggest that the intrauterine transfer of fungal DNA may be widespread across mammals. Impaired

host-sensing of fungi may alter both intrauterine deposition and post-natal colonization.

449
450   **Discussion**
451
452
Here we present the first description of the primordial fungal ecology of the neonatal gut using both

next-generation sequencing and culture-based techniques. Microeukaryotes, primarily fungi, were

present in the first-pass meconium samples collected within hours of birth in both preterm and term-

born newborns. In general, as quantified by amplicon sequencing of eukaryotic ITS rDNA, infantile

fungal communities developed greater taxonomic diversity as the gestational age at birth increased.

The structure of these communities was unaffected by perinatal factors, such as mode of delivery,

antibiotic exposure, infant sex, and illness severity, consistent with intrauterine formation of

meconium. Culture-based techniques indicate microbial DNA may originate from viable

microorganisms present in the neonatal gut.

Understanding the primary succession of microorganisms into the human gut is important because

the order in which these potential colonizers are introduced into the intestine influences the mature

community structure. Colonization experiments in mice suggest the timing of arrival of microbial

colonizers is the major factor in determining final community structure[24]; which helps explain why

other perinatal factors, such as perinatal antibiotic exposure, account for only limited variability in

mature community structure. Bacterial communities then undergo a rapid expansion during the first

several years of life[25], eventually establishing a host-specific core microbiome that resists further

perturbations[26]. Because misconfiguration of the core microbiome during this period likely contributes

to multiple host disease states, understanding the ecological processes driving microbiome

assemblage is crucial to developing therapeutic interventions[27]. Recent evidence suggests that the

mycobiome remains relatively constant during the first year of life[11,12], which underscores that the

primary succession of initial fungal colonizers may be even more important for final community

structure than the succession of bacterial species. In addition, fungi may also form a crucial nidus for

bacterial colonization[6]. Our evidence for the presence of commensal fungi in the primordial gut and

the development of these communities with gestational age demonstrates this process may begin

earlier that was previously thought - with implications that extend throughout life.

Meconium forms during early human gut development, eventually migrating into the colon around the

19th week of gestation[28]. Passage of meconium is a developmentally programmed event that usually

occurs within 48 hours after birth[28]. Once enteral feeding has been established environmental

microbes invade the gastrointestinal tract and are detectable in the first true stools. Until the

development of next-generation sequencing techniques, the presence of microorganisms in the first-

pass meconium was opaque using traditional culture-based techniques. Meconium, and therefore the

uterine environment, has been widely accepted to be sterile[9]. However, during the past decade

several researchers have demonstrated bacterial DNA can be isolated from meconium[3,8,29-34] and the

placenta[35-37], but if this microbial biomass is representative of live organisms remains controversial.

These studies have focused on the bacterial components of meconium, however, leaving fungi

undescribed.

The human amniotic fluid may also harbor a very low level of microbial organisms. Chu et al.[17] have

demonstrated that the bacterial components of the meconium microbiome are distinct and do not

resemble adult vaginal and skin communities like the predominant bacterial members of other

neonatal body sites. Our results show interkingdom microbial communities that are essentially

unaffected by perinatal exposures, with the possible exception of prenatal steroids. The

demonstration of live microorganisms in the newborn gut represents a monumental shift from the

sterile womb hypothesis that dominated perinatal biology for the last century[8,9]. The increasing

prevalence and the development of bacterial ecology with increasing gestational age suggests that

this is a natural, non-pathogenic phenomenon[3,33,34]. The human fetus regularly swallows and

replenishes the amniotic fluid throughout gestation, which suggests a mechanism for how rare

microbes in the amniotic fluid could become more concentrated within the primordial gut. Based on

these findings, we postulate that low abundant amniotic fluid bacterial and fungal biomass is

continuously collected during gestation and accumulates in the intestine. By the time of delivery,

meconium may contain a collective record of fetal microbial exposure in an analogous manner to how

meconium contains a record of transplacental illicit drug exposure[38]. This process would also expose

the fetal mucosal immune system to low levels of microbial products that may initiate host microbial

surveillance mechanisms required to process the influx of microbes into the gut that occurs during and

after delivery.

To directly observe the fetal environment, we utilized a mouse model of normal pregnancy. This work

highlights the presence of fungal biomass within the developing fetal gut. The implication of a steady

deposition of microbial biomass in the fetal intestine suggests that this is a natural process, which

raises the question of whether translocation of microbes is regulated. To gain insight into this process,

we asked if impairment of the host's recognition of fungi would alter the deposition of fungal biomass

within the fetal and neonatal gut. Indeed, we found that loss of the host fungal pattern recognition

receptor Dectin-1 resulted in more fungal biomass within the fetal intestine. While there are marked

differences between the placentas of humans and mice[21], this suggests that the deposition of fungal

biomass within the fetal gut may be a regulated process that requires host recognition of fungal

organisms.

Counter to the general trend of our results which support an apparently natural process of microbial

biomass accumulation in the neonatal intestine, our finding that some preterm infants are dominated

by fungal order Saccharomyces, and in particular the genus *Candida*, may support the alternate

hypothesis that in some instances preterm birth is a product of polymicrobial disease of the uterine

cavity[39]. After birth, systemic infection with *Candida* are a persistent threat during neonatal care.

Candida species, primarily *Candida albicans*, are opportunistic pathogens particularly associated with

bacterial dysbiosis[40]. *Candida* are significant commensals of the human gut[41], so the presence of

Candia within the gastrointestinal tract is understandable. In the mature gut, commensalism is

maintained by a tripartite interaction between *Candida*, other members of the microbiome, and host

immunity[41]. However, *Candida* are also the dominate fungal species associated with microbial

invasion of the amniotic cavity[42], and are detectable in the amniotic fluid by culture and culture-

independent techniques[43]. As quantified by culture-based techniques, *Candida* chorioamnionitis is a

rare but accepted cause of preterm birth[44]. Our study showed that preterm meconium samples were

more likely to contain fungi than term infants. Furthermore, the mycobiomes of preterm infants were

also more likely to be dominated by *Candida*, as opposed to the more diverse mycobiomes of infants

with a longer gestation. Regardless of these concerning findings, conclusively demonstrating that

Candida colonization is associated with preterm birth will require a clinical trial powered to test this

hypothesis.

Fungi are ubiquitous in the environment. In low biomass samples like meconium, the presences of

environmental fungi are difficult to separate from organisms originating within the samples. Because

microeukaryotic DNA is a relatively low proportion of the fecal DNA content, especially in meconium,

exacting extraction, purification and amplification techniques are required. A prominent reason that

investigation of the mycobiome has lagged behind investigation of the microbiome is a lack of

standardization in detection methods for fungal organisms[41]. Our choice to utilize amplification of the

ITS2 region, the read depth and the database used to align our amplicon reads, therefore, could all

influence the fungal species identified in our samples. Incidentally, the exacting DNA extraction

techniques we utilized may also explain the differences in bacterial community structure with prior

work because they increased the detectability of low biomass samples. We froze meconium samples

immediately after collection to prevent post-excretion microbial expansion that would alter the relative

composition of microbial communities, and passed negative control samples throughout the sample

collection, DNA extraction and amplicon sequencing process. However, the fact that perinatal

exposures known to shape the post-natal microbiome did not significantly alter community

composition in our samples argues that significant post-natal microbial colonization did not occur

before sample collection.

Our findings support the presence of fungal organisms within the neonatal gut. Taken together, our

sequencing and culture-based techniques support the presence of live fungal organisms in meconium

in addition to bacteria. Homeostasis of fungal commensals may represent an important aspect of

human biology present even before birth with functional implications throughout early life. Low

abundance colonization of the primordial gut with interkingdom ecologies appears to be a normal part

of healthy pregnancy that might represent a priming of the intestine for the exponential expansion of

the microbiome after birth. The domination of microbial communities by fungal order Saccromycetes

in some preterm infants, however, suggests a potential pathologic association with preterm birth.

566
567

Methods

Study patients

Written and informed consent was obtained from the infants' mothers. All investigations were carried

out in accordance with the declaration of Helsinki under The University of Tennessee Health Science

Center Institutional Review Board protocol 17-05311-XP. We prospectively enrolled either very low

birthweight infants (< 1500 g) preterm or term-born (estimated gestational age 37-41 weeks) infants

(Fig. 1a). We excluded any mother-infant dyads with the potential for immune deficiency, but

otherwise invited all mothers that meet our entrance criteria during the study period to participate.

Infant exposures

Data was collected regarding infant demographics and the following clinical exposures: prenatal and

postnatal antibiotic exposure, perinatal steroid exposure, delivery mode, and illness severity (the

CRIB-II[18]). Additional clinical characteristics were also abstracted from the medical record to

characterize potential co-variates with gestational age.

Sample collection

The first-pass meconium was collected during the first 48 h after birth. Samples were collected from

diapers using sterile spatulas, placed into sterile microcentrifuge tubes and frozen at -30C within

minutes of collection. Several times a week, batches of samples were collected and frozen at -80C

until further analysis could be performed. Infants that did not stool during the first 48 h or for which a

meconium sample was not collected were excluded from downstream analysis.

DNA extraction and Illumina MiSeq sequencing

Meconium samples were resuspended in 500 μL of TNES buffer containing 200 units of lyticase and

100 μL of 0.1/0.5 (50/50 Vol.) zirconia beads. Incubation was performed for 20 min at 37°C. Following

mechanical disruption using ultra-high-speed bead beating, 20 μg of proteinase K was added to all

samples, and they were incubated overnight at 55C with agitation. Total DNA was extracted using

601   chloroform isoamyl alcohol, and total DNA concentration per mg stool was determined by qRT-PCR.

602   Purified DNA samples were sent to the Argonne National Laboratory (Lemont, IL) for amplicon

603   sequencing using the NextGen Illumina MiSeq platform. Blank samples passed through the entire

604   collection, extraction and amplification process remained free of DNA amplification.

605
606   **Bioinformatics**
607
Sequencing data were processed and analyzed using QIIME (Quantitative Insights into Microbial

Ecology) 1.9.1. Sequences were first demultiplexed, then denoised and clustered into sequence

variants. For bacteria we rarified to a depth of 3,000 sequences. (In combined analysis after excluding

samples with a bacterial depth of < 3,000, we used a depth of 300 for further analysis).

Representative bacterial sequences were aligned via PyNAST, taxonomy assigned using the RDP

Classifier. Processed data was then imported into Calypso 8.84 for further analysis and data

visualization[45]. The Shannon Index was used to quantify alpha diversity (inter-sample)[46,47].

Unweighted UniFrac analysis was used to quantify beta diversity (intra-sample) [48], and the differences

were compared using PERMANOVA with 999 permutations. For fungi, sequences were aligned, and

taxonomy was assigned using the UNITE (dynamic setting) database[49]. Fungal OTUs were rarified at

a depth of 300 sequences for alpha diversity using Shannon or Chao1 and beta diversity using Bray-

Curtis dissimilarity or Jaccard abundance [50] indices. As with bacteria, beta diversity significance was

then assessed using PERMANOVA. To quantify relative abundance of taxa between groups, we

utilized ANOVA adjusted using the Bonferroni correction and FDR for multiple comparisons. We used

LEfSe to test for significance and perform high-dimensional biomarker identification[51]. Network

analysis was generated from Spearman's correlations. Positive correlations with an FDR-adjusted, *p*

<0.05 were presented as an edge.

Machine learning and variable selection

In the first phase, we developed a support vector machine utilizing 34 bacterial and 9 fungal taxa

identified utilizing an unbiassed LASSO approach with 5-fold cross validation and generated a

confusion matrix to characterize the performance of the model as a predictor of gestational age

In the second phase, we developed a random forest-based classification model to distinguish

between preterm and term-born newborns using maternal age, maternal body mass index, newborn

sex and 260 microbial taxa that were present in at least two samples in the data set. Random forest

classifiers are based on the arrangement of multiple decision trees built using a random sample of

predictors. The outcome of a random forest is based on a weighted average of the outcomes of

individual decision trees within the model. To identify and control for potential overfitting, we

implemented a 5-fold cross validation strategy. By repeating this process 5 times for each subject, we

obtained a single class prediction that is purely based on a classifier built on a distinct dataset. We

then repeated the development of the entire model 100 times. We also implemented a recursive

variable selection based on the out-of-bag predictor importance analysis. Here, variable importance

was assessed by randomly permitting values of a single predictor and quantifying the increment in the

out-of-bag error rate. Variables that caused no drop on the error rate were assumed to have no

influence in predicting outcome. We then selected the 10 most important variables and built another

100 permutations of our random forest model using 5-fold cross validation.

We also compared the performance of the random forest classifier we generated using this method to

other classifiers including logistic regression, discriminant analysis, k-nearest neighborhood, decision

trees, support vector machines and neural networks using AUC statistics. Finally, we calculated

commonly used classification accuracy statistics such as specificity, sensitivity, and positive predictive

value for the final random forest model to characterize the ability of the model to distinguish between

preterm and term-born babies.

In the final phase, we repeated this model development process for each phylogenic level to

understand the variable importance of different taxa. To achieve this, in level 1 (kingdom) we used

four demographic variables and 3 microbial kingdoms (bacteria, fungi, and archaea) features by

aggregating all bacterial and fungal taxa under their particular kingdom definition. Therefore, at level 1

(kingdom), we built a random forest classifier using a total of 6 features. The four demographics

variables included host sex (female: 0, male: 1), maternal self-reported race/ethnicity (Black: 0, other:

1), maternal age, and maternal BMI. Fungal and bacterial taxa were represented in six different levels

following, kingdom > phylum > class > order > family > genus. For level 2 (phylum) we aggregated

taxa data at 23 distinct phylum-level taxa resulting in total of 27 features and including the same four

demographics data. We repeated this process until we had reached the 6^{th} level (genus) where we

utilized a total of 265 variables (4 demographics and 261 taxa data at genus level). At each level, we

included only taxa that were observed in at least two different subjects to avoid overfitting. For each

level, we implemented five-fold cross validation. To avoid sampling bias, we repeated the cross-

validation after randomly shuffling the data for 100 times at each level. Classification performance was

summarized as AUC statistics.

674
675   **Microbial culture**
676
We prepared culture broth using either brain heart infusion (BHI) with 250 mg/L of fluconazole

(bacteria-selective) or yeast-extract-peptone-dextrose (YPD) with 50 mg/mL of chloramphenicol

(fungal-selective) under both aerobic and anaerobic conditions. Meconium samples were thawed and

diluted in 1 mL of PBS. Tubes containing 1 mL of culture broth were inoculated with 100 μL of room

temperature meconium diluted in 1 mL of culture broth. Samples were then incubated at 32°C under

either aerobic or anaerobic conditions. Bacterial samples were assessed by optical density after 48 h

and fungal samples after 120 h.

Statistical analysis

We compared demographic and clinical variables between infants with a gestational age < 33 weeks'

and > 33 weeks' using the χ^2 or two-tailed Welch t-test for parametric variables or the Wilcoxon rank-

sum test for non-parametric variables, as appropriate.

Animal model of pregnancy

All animal experiments were approved by the University of Tennessee Health Science Center

Institutional Animal Care and Use Committee. Timed-mating of adult C57BL6J mice or dectin-1$^{-/-}$ on

the C57Bl6J background (B6.clec7a^{tm1Gdb}, Jackson Labs stock #012337) at 8-10 weeks of age was

performed to obtain offspring samples at embryonic day 17 (E17) or post-partum day 1 (P1). After

euthanasia, swabs were collected from the maternal mouth, skin and genitourinary tract. The dam

was then soaked in exspor (chlorine dioxide disinfectant). Gravid uteri and intestines were removed

from prepped dams in a UV-light sterilized laminar flow hood by sterile gowned and gloved personnel

extensively trained in sterile technique (a maternal-fetal medicine specialist gynecologist and a

neonatologist) using sterile technique and instruments that were sterilized between animals. Utilizing

a second set of sterilized instruments, pups were removed, euthanized and placed in exspor then

passed to another operator who removed them in a second laminar flow hood. The body cavity was

opened with another set of sterile instruments and removed the intestine *en bloc* and placed them into

a sterile container. Samples of the placenta, amniotic fluid, small intestine and colonic contents were

also placed in sterile containers. All samples were snap frozen and stored at -80C until DNA

extraction.

Data Availability

Microbial sequence data is available at: [To be uploaded on acceptance]

Computer Code

Custom computer code is available upon request.

Disclosure

The authors declare no conflict of interest.

Author Contributions

KW, AT and JFP conceived of the study design; KW and JHP recruited the patients and collected the

meconium samples; KW and EM extracted clinical data from the medical record; CG and JFP

performed the DNA purification; KW, BP and JFP performed the culture-based analyses; KW and MA

performed the murine experiments, IK and OA performed the statistical modeling and developed the

machine learning models; KW, IK, OA and JFP performed the data quality control, statistical analysis

and prepared the figures; KW wrote the paper; KW, MA, JHP, OA, BP, AT and JFP edited the

manuscript; and all authors approved the final version.

Acknowledgements

The authors would like to thank the mothers and infants who contributed to the study and to our

research coordinators, Gail Camp and Nancy Ruch, for their assistance in identifying and recruiting

patients.

References

1. Romano-Keeler, J. & Weitkamp, J.-H. Maternal influences on fetal microbial colonization and immune development. *Pediatr. Res.* **77,** 189–195 (2015).
2. Mueller, N. T. *et al.* Delivery Mode and the Transition of Pioneering Gut-Microbiota Structure, Composition and Predicted Metabolic Function. *Genes (Basel)* **8,** 364 (2017).
3. Ardissone, A. N. *et al.* Meconium Microbiome Analysis Identifies Bacteria Correlated with Premature Birth. *PLoS ONE* **9,** e90784 (2014).
4. Dominguez-Bello, M. G. *et al.* Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci U S A* **107,** 11971–11975 (2010).
5. Olin, A. *et al.* Stereotypic Immune System Development in Newborn Children. *Cell* **174,** 1277–1292.e14 (2018).
6. Zhang, Y., Kastman, E. K., Guasto, J. S. & Wolfe, B. E. Fungal networks shape dynamics of bacterial dispersal and community assembly in cheese rind microbiomes. *Nat Commun* **9,** 336 (2018).
7. Arrieta, M.-C. *et al.* Associations between infant fungal and bacterial dysbiosis and childhood atopic wheeze in a nonindustrialized setting. *J. Allergy Clin. Immunol.* **142,** 424–434.e10 (2018).

8. Jiménez, E. *et al.* Is meconium from healthy newborns actually sterile? *Res. Microbiol.* **159,** 187–193 (2008).

9. Mackie, R. I., Sghir, A. & Gaskins, H. R. Developmental microbial ecology of the neonatal gastrointestinal tract. *Am. J. Clin. Nutr.* **69,** 1035S–45S (1999).

10. Limon, J. J., Skalski, J. H. & Underhill, D. M. Commensal Fungi in Health and Disease. *Cell Host Microbe* **22,** 156–165 (2017).

11. Ward, T. L. *et al.* Development of the Human Mycobiome over the First Month of Life and across Body Sites. *mSystems* **3,** e00140–17 (2018).

12. Schei, K. *et al.* Early gut mycobiota and mother-offspring transfer. *Microbiome* **5,** 107 (2017).

13. Chernikova, D. A. *et al.* The premature infant gut microbiome during the first 6 weeks of life differs based on gestational maturity at birth. *Pediatr. Res.* **84,** 71–79 (2018).

14. Strati, F. *et al.* Age and Gender Affect the Composition of Fungal Population of the Human Gastrointestinal Tract. *Front Microbiol* **7,** 1227 (2016).

15. Becattini, S., Taur, Y. & Pamer, E. G. Antibiotic-Induced Changes in the Intestinal Microbiota and Disease. *Trends in Molecular Medicine* **22,** 458–478 (2016).

16. Samonis, G. *et al.* Prospective evaluation of effects of broad-spectrum antibiotics on gastrointestinal yeast colonization of humans. *Antimicrob. Agents Chemother.* **37,** 51–53 (1993).

17. Chu, D. M. *et al.* Maturation of the infant microbiome community structure and function across multiple body sites and in relation to mode of delivery. *Nat. Med.* **23,** 314–326 (2017).

18. Parry, G., Tucker, J., Tarnow-Mordi, W.UK Neonatal Staffing Study Collaborative Group. CRIB II: an update of the clinical risk index for babies score. *The Lancet* **361,** 1789–1791 (2003).

19. Kong, L., Nilsson, I. A. K., Gissler, M. & Lavebratt, C. Associations of Maternal Diabetes and Body Mass Index With Offspring Birth Weight and Prematurity. *JAMA Pediatr* **173,** 371 (2019).

20. Perez-Muñoz, M. E., Arrieta, M.-C., Ramer-Tait, A. E. & Walter, J. A critical assessment of the "sterile womb" and "in utero colonization" hypotheses: implications for research on the pioneer infant microbiome. *Microbiome* **5,** 243 (2017).

21. Martinez, K. A. *et al.* Bacterial DNA is present in the fetal intestine and overlaps with that in the placenta in mice. *PLoS ONE* **13,** e0197439 (2018).

22. Schorey, J. S. & Lawrence, C. The pattern recognition receptor Dectin-1: from fungi to mycobacteria. *Curr Drug Targets* **9,** 123–129 (2008).

23. Schmidt, A., Morales-Prieto, D. M., Pastuschek, J., Fröhlich, K. & Markert, U. R. Only humans have human placentas: molecular differences between mice and humans. *J. Reprod. Immunol.* **108,** 65–71 (2015).

24. Martínez, I. *et al.* Experimental evaluation of the importance of colonization history in early-life gut microbiota assembly. *Elife* **7,** 1 (2018).

25. Arrieta, M.-C., Stiemsma, L. T., Amenyogbe, N., Brown, E. M. & Finlay, B. The Intestinal Microbiome in Early Life: Health and Disease. *Front. Immunol.* **5,** 355 (2014).

26. Martínez, I., Muller, C. E. & Walter, J. Long-term temporal analysis of the human fecal microbiota revealed a stable core of dominant bacterial species. *PLoS ONE* **8,** e69621 (2013).

27. Koskella, B., Hall, L. J. & Metcalf, C. J. E. The microbiome beyond the horizon of ecological and evolutionary theory. *Nat Ecol Evol* **1,** 1606–1615 (2017).

28. Ahanya, S. N., Lakshmanan, J., Morgan, B. L. G. & Ross, M. G. Meconium Passage in Utero: Mechanisms, Consequences, and Management. *Obstet Gynecol Surv* **60,** 45–56 (2005).

29. Rougé, C. *et al.* Investigation of the intestinal microbiota in preterm infants using different methods. *Anaerobe* **16,** 362–370 (2010).

30. Mshvildadze, M. *et al.* Intestinal microbial ecology in premature infants assessed with non-culture-based techniques. *J Pediatr* **156,** 20–25 (2010).

31. Gosalbes, M. J. *et al.* Meconium microbiota types dominated by lactic acid or enteric bacteria are differentially associated with maternal eczema and respiratory problems in infants. *Clinical & Experimental Allergy* **43,** 198–211 (2013).

32. Stewart, C. J. *et al.* The preterm gut microbiota: changes associated with necrotizing
enterocolitis and infection. *Acta Paediatr* **101,** 1121–1127 (2012).
33. Hansen, R. *et al.* First-Pass Meconium Samples from Healthy Term Vaginally-Delivered
Neonates: An Analysis of the Microbiota. *PLoS ONE* **10,** e0133320 (2015).
34. Tapiainen, T. *et al.* Maternal influence on the fetal microbiome in a population-based study of
the first-pass meconium. *Pediatr. Res.* **84,** 371–379 (2018).
35. Collado, M. C., Rautava, S., Aakko, J., Isolauri, E. & Salminen, S. Human gut colonisation may
be initiated in utero by distinct microbial communities in the placenta and amniotic fluid. *Sci Rep*
6, 23129 (2016).
36. Aagaard, K. *et al.* The placenta harbors a unique microbiome. *Sci Transl Med* **6,** 237ra65–
237ra65 (2014).
37. Zheng, J. *et al.* The Placental Microbiome Varies in Association with Low Birth Weight in Full-
Term Neonates. *Nutrients* **7,** 6924–6937 (2015).
38. Palmer, K. L. & Krasowski, M. D. Alternate Matrices: Meconium, Cord Tissue, Hair, and Oral
Fluid. *Methods Mol. Biol.* **1872,** 191–197 (2019).
39. Payne, M. S. & Bayatibojakhi, S. Exploring preterm birth as a polymicrobial disease: an
overview of the uterine microbiome. *Front. Immunol.* **5,** 595 (2014).
40. Iliev, I. D., Immunology, I. L. N. R.2017. Fungal dysbiosis: immunity and interactions at mucosal
barriers. *nature.com*
41. Neville, B. A., d'Enfert, C. & Bougnoux, M.-E. Candida albicans commensalism in the
gastrointestinal tract. *FEMS Yeast Res.* **15,** fov081 (2015).
42. DiGiulio, D. B. Diversity of microbes in amniotic fluid. *Semin Fetal Neonatal Med* **17,** 2–11
(2012).
43. DiGiulio, D. B. *et al.* Prevalence and Diversity of Microbes in the Amniotic Fluid, the Fetal
Inflammatory Response, and Pregnancy Outcome in Women with Preterm Pre-Labor Rupture
of Membranes. *American Journal of Reproductive Immunology* **64,** 38–57 (2010).
44. Maki, Y., Fujisaki, M., Sato, Y. & Sameshima, H. Candida Chorioamnionitis Leads to Preterm
Birth and Adverse Fetal-Neonatal Outcome. *Infect Dis Obstet Gynecol* **2017,** 9060138–11
(2017).
45. Zakrzewski, M. *et al.* Calypso: a user-friendly web-server for mining and visualizing
microbiome-environment interactions. *Bioinformatics* **33,** 782–783 (2017).
46. Caporaso, J. G. *et al.* QIIME allows analysis of high-throughput community sequencing data.
Nat. Methods **7,** 335–336 (2010).
47. Hughes, J. B., Hellmann, J. J., Ricketts, T. H. & Bohannan, B. J. Counting the uncountable:
statistical approaches to estimating microbial diversity. *Appl. Environ. Microbiol.* **67,** 4399–4406
(2001).
48. Lozupone, C., Lladser, M. E., Knights, D., Stombaugh, J. & Knight, R. UniFrac: an effective
distance metric for microbial community comparison. *The ISME journal* **5,** 169–172 (2011).
49. Nilsson, R. H. *et al.* The UNITE database for molecular identification of fungi: handling dark
taxa and parallel taxonomic classifications. *Nucleic Acids Res.* **5,** 1–D264 (2018).
50. Faith, J. J. *et al.* The long-term stability of the human gut microbiota. *Science* **341,** 1237439–
1237439 (2013).
51. Segata, N. *et al.* Metagenomic biomarker discovery and explanation. *Genome Biol.* **12,** R60
(2011).
858