1   # Predation strategies of the bacterium *Bdellovibrio*

2   # *bacteriovorus* result in overexploitation and bottlenecks

3   **J. Kimberley Summers[1][†][*] and Jan-Ulrich Kreft[1]**

4   [1]Institute of Microbiology and Infection & Centre for Computational Biology & School of Biosciences,
5   University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom
6
[†]Present address: School of Life Sciences, University of Warwick, Coventry, CV4 7AL, United Kingdom

ORCIDs:
Jan-Ulrich Kreft: 0000-0002-2351-224X
Kimberley Summers: 0000-0002-0092-7848

*Corresponding author

J. Kimberley Summers
Tel: +44 (0)757 9037167
E-mail address: kimberley.summers@warwick.ac.uk
17

Keywords: predator-prey interactions; generalist versus specialist; prey size; ordinary

differential equations; mathematical modelling; robust permanence; competition; fitness

Running title: Predation strategies of *Bdellovibrio*

Abstract

With increasing antimicrobial resistance, alternatives for treating infections or removing

resistant bacteria are urgently needed, such as the bacterial predator *Bdellovibrio*

bacteriovorus or bacteriophage. Therefore, we need to better understand microbial

predator-prey dynamics. We developed mass-action mathematical models of predation

for chemostats, which capture the low substrate concentration and slow growth typical for

intended application areas of the predators such as wastewater treatment, aquaculture

or the gut. Our model predicted that predator survival required a minimal prey size,

explaining why *Bdellovibrio* is much smaller than its prey. A too good predator (attack rate

too high, mortality too low) overexploited its prey leading to extinction (tragedy of the

commons). Surprisingly, a predator taking longer to produce more offspring outcompeted

a predator producing fewer offspring more rapidly (rate versus yield trade-off). Predation

was only efficient in a narrow region around optimal parameters. Moreover, extreme

oscillations under a wide range of conditions led to severe bottlenecks. A bacteriophage

outcompeted *Bdellovibrio* due to its higher burst size and faster life cycle. Together,

results suggest that *Bdellovibrio* would struggle to survive on a single prey, explaining

why it must be a generalist predator and suggesting it is better suited than phage to

environments with multiple prey.

Importance

The discovery of antibiotics led to a dramatic drop in deaths due to infectious disease.

Increasing levels of antimicrobial resistance, however, threaten to reverse this progress.

There is thus a need for alternatives, such as therapies based on phage and predatory

bacteria that kill bacteria regardless of whether they are pathogens or resistant to

antibiotics. To best exploit them, we need to better understand what determines their

effectiveness. By using a mathematical model to study bacterial predation in realistic slow

growth conditions, we found that the generalist predator *Bdellovibrio* is most effective

within a narrow range of conditions for each prey. For example, a minimum prey size is

required, and the predator should not be too good as this would result in over-exploitation

risking extinction. Together these findings give insights into the ecology of microbial

predation and help explain why *Bdellovibrio* needs to be a generalist predator.

51 **Introduction**

Predator-prey relationships are some of the oldest and most important interactions in

nature and occur at every level, from the smallest virus infecting a bacterium, to lions

attacking wildebeest. Predators are often keystone species in natural ecosystems (1).

Investigations into predator-prey dynamics in natural settings are, however, complicated

by an array of confounding factors (2). Microbes by contrast make attractive models for

studying predator-prey interactions in a controlled environment, with millions of individuals

in a drop of liquid that can go through many generations in days. A completely separate,

but increasingly important, reason for an interest in microbial predators is the antimicrobial

resistance crisis, with resistance to even last resort antibiotics such as colistin rising (3).

There is an urgent need for 'living antibiotics' as alternatives to antibiotics, such as phage

and bacterial predators.

The best-studied bacterial predator is *Bdellovibrio bacteriovorus*. It is a Gram-negative

bacterium that predates a wide range of other Gram-negative bacteria (4), regardless of

their antimicrobial resistance or pathogenicity. *Bdellovibrio* alternates between two

phases in its lifecycle. In the free-living attack phase, it does not grow but hunts prey. It

swims faster than most bacteria, at speeds of up to 160 µm s^{-1} or ~100 body lengths s^{-1}

(5). This requires a high metabolic rate and loss of viability within 10 hours if prey is not

encountered (6). Once a cell is encountered, its suitability is investigated for several

minutes (7). If suitable, it enters the prey's periplasm by creating and squeezing through

a pore in the outer membrane and peptidoglycan layer, shedding its flagellum in the

process (8). Once inside the periplasm, *Bdellovibrio* kills the prey and lets the cytoplasmic

nutrients leak into the periplasm (9). It also alters the prey's peptidoglycan, causing the

prey cell to round up into a "bdelloplast". In this bdelloplast phase, *Bdellovibrio* uses these

nutrients to grow into a long filament rather than dividing (5). When the nutrients have

been used up, this filament septates into as many new cells as resources allow for,

typically between 3 and 6 new predators per *E. coli* cell (10). If it were using normal binary

fission, it could only produce 2^n offspring, potentially forsaking prey resources if they

would only suffice for say 5 rather than 8 offspring. The new *Bdellovibrio* form flagella and

lyse the remains of the prey cell, allowing them to burst out in search of fresh prey.

Mathematical models explain how predator-prey interactions can generate stable

oscillations (11). Most models of microbial predator-prey interactions focussed on protists

or bacteriophage. Only a few models considered predatory bacteria, see Table S1 for an

overview of models and the review by Wilkinson (12). Varon & Zeigler (13) fitted a Lotka-

Volterra model to their experimental results, which led them to propose that a minimal

prey density is necessary for the survival of the predator. Crowley's (14) model tracked

substrate levels and used Monod kinetics for prey growth, it also included a delay between

prey attack and the production of new predators to reflect the ~3 h long bdelloplast phase.

The dynamics were destabilised by higher nutrient concentrations, especially at low

dilution rates, leading to extreme oscillations. Wilkinson (15) developed two *Bdellovibrio*

models, one based on a Holling type II functional response, but without specifically

modelling the bdelloplast stage. The other model did include a combined predator-prey

complex, but used a Holling type I functional response (15). Both models again showed

a destabilising effect of nutrient enrichment, albeit somewhat ameliorated by the presence

of a decoy species. Similarly, Hobley et al. (7), Baker et al. (16) and Said et al. (17)

included a predator-prey complex and a Holling type I functional response. Hobley,

Summers et al. (18) fitted a series of models, with a separate bdelloplast stage and

various predator functional responses, to data from batch culture predation by either

Bdellovibrio, a bacteriophage or both, and found a synergistic effect from dual predation.

Two further models have been developed by Hol et al. (19) and Dattner et al. (20), both

including spatial structure. These previous models focussed on the effects of dilution rate

and substrate inflow on the dynamics of the chemostat systems. However, the effects of

prey and predator characteristics such as prey size, attack kinetics and predation

efficiency have not been studied.

In this study, we developed a family of models, based on 'ingredients' from previous

models, to identify a unique combination of ingredients that generates realistic outcomes

from realistic model assumptions and to ask many novel questions. This model predicts

that *Bdellovibrio* has to be much smaller than its prey to survive, which is in agreement

with empirical evidence (21). We also found that high attack rates, large prey sizes and

low predator mortality, which might be expected to help the predator, are in fact

detrimental for predator survival. Instead, these parameters have optimal values that

maximize predator density. These optima occurred at the tipping points into oscillations

and were often very narrow relative to the natural variation of these parameters.

Moreover, we found that the system was prone to extreme oscillations in bacterial

densities, leading to bottlenecks that would result in stochastic extinction. Additionally,

Bdellovibrio would easily be outcompeted by a bacteriophage. Together, these three key

predictions (narrow optima, population bottlenecks and phage superiority) suggest that

Bdellovibrio would struggle to survive on a single prey species in a natural environment

full of phages, and where conditions are unlikely to be optimal, and if so, not for long. Our

findings are consistent with the fact that *Bdellovibrio* is a generalist predator, so we posit

that our model explains why *Bdellovibrio* is a generalist predator and why it has to be as

small as it is.

Model development

We developed models to investigate the effects of a consumer, the predatory bacterium

Bdellovibrio bacteriovorus or a bacteriophage, on a bacterial population under continuous

culture conditions in a chemostat. A chemostat captures the low substrate concentrations

and slow growth rates typical of the natural environment (22). Also, the prey on its own

will reach a steady state population density in the chemostat, which facilitates the

investigation of oscillations caused by predation. Oscillations could not be observed in a

batch culture model. We developed a family of models to investigate the effect of changes

in the structure of the model on the predator-prey dynamics. This structural sensitivity

analysis (23) identified Model 6 as the most realistic in terms of assumptions and

predictions (see SI text, Table S2 and Fig. S4), so we explored Model 6 further and

describe it in Fig. 1 and list its parameters and their values in Table 1. Model 6 included

a single abiotic resource (substrate S), single prey species (N) and a single obligate

predator (P). The prey species grew by consuming the substrate according to Monod

kinetics. The predator had a Holling type II functional response (predation rate

proportional to predator density but saturating at high predator density), a bdelloplast (for

Bdellovibrio) or infected cell (for bacteriophage) stage and mortality. The bdelloplast (B)

is a distinct stage in the *Bdellovibrio* life cycle that usually lasts for 2 to 4 hours. There are

distinct parallels between this bdelloplast stage and a bacteriophage infected cell as the

prey cell does not grow and replicate when infected by a bacteriophage or consumed by

Bdellovibrio and the predator or phage does not prey on or infect further prey. We

modelled the bdelloplast or infected cell stage as a separate entity to account for the delay

in producing offspring and the combining of the consumer and its prey. For simplicity,

these entities will be referred to as a bdelloplast from now on, but the principles apply

equally to an infected cell.

Results

Model implementation and validation showed the system to be brittle

We first ensured that the numerical results were reliable by testing all MatLab Ordinary

Differential Equation (ODE) solvers. Only the Runge-Kutta ode45 solver could correctly

handle all test cases including extreme oscillations (Fig. S1). It was also necessary to set

tolerances of this solver to very low values and to constrain variables to be non-negative

(Fig. S2). Secondly, we used the same test cases to compare simulations using biomass-

based units with those using particle-based units, required to obtain numerically stable

simulations with bacteriophage parameters. The results were in agreement (SI text and

Fig. S3). These unusual difficulties of obtaining correct numerical results highlight that the

system tends to undergo extremely rapid changes, followed by periods of stasis, and is

very sensitive to parameter settings.

Structural sensitivity analysis identified most appropriate model

To gain a better understanding of the impact of various modelling choices, we compared

a number of ordinary differential equation (ODE) and delay differential equation (DDE)

models (Table S2, Fig. S4). In all cases, the same set of standard conditions, based on

Bdellovibrio predating *E. coli* that are growing on glucose (0.05 mg ml^{-1}) and a dilution

rate of 0.0333 h^{-1} (equivalent to a 30 hour retention time) were used. We first ran the

simulation without predators until the prey had reached a steady state (same for all

models). (Model 1) Addition of a predator without a bdelloplast stage (or other form of

delay between prey killing and predator birth) gave rise to sustained, extreme oscillations.

(Model 2) Incorporating an explicit delay of 4 hours, approximately doubled the oscillatory

period from ~150 hours to ~300 hours. (Model 3) Adding mortality reduced the oscillatory

period to approximately 100 hours. (Models 4-6) Addition of an explicit bdelloplast stage

stabilised the system, resulting in a stable steady state of co-existing predator and prey,

regardless of the predator's Holling type functional response. (Model 4 vs. 6) With a

Holling type I predator functional response (Model 4), the final prey density was lower,

and the predator density higher, than with the saturating type II response (Model 6).

(Model 5) Constant input of prey to the system; this is akin to growing the prey on its own

in one chemostat that feeds into a second chemostat containing predator. This gave a

very similar response to the single chemostat with constant input of substrate. Model 6

was chosen as the most biologically appropriate model for all further work, for several

reasons. Firstly, the time required for *Bdellovibrio* to consume the contents of a prey cell,

convert these into new *Bdellovibrio* predators, septate and finally lyse the prey cell to

release new predators is about four hours (10). As the prey is killed very shortly after

penetration (26), there is a significant delay between prey killing and birth of new

predators, which is best modelled by treating the bdelloplast stage as a separate entity.

Secondly, *Bdellovibrio* has a high endogenous respiration rate and a correspondingly low

life span in the absence of suitable prey (6). Hence, it is appropriate to include predator

mortality in the model. Thirdly, the saturating Holling type II functional response results

from the 'handling time' of a predator (27). This would correspond to the minimum time

for a successful attack in a prey-saturated environment, i.e., the attachment and

penetration time of *Bdellovibrio* of ~10 minutes (28).

Dynamic regimes from steady states to extreme oscillations

Model 6 has 12 parameters and six possible dynamic regimes (Fig. 2a). In order to gain a better understanding of the factors determining which regimes were observed, we swept through a range of inflow substrate concentrations (S_0) and dilution rates and evaluated the steady state and its stability analytically (see model analysis in SI). As expected, at the highest dilution rates and lowest S_0, all biological species washed out. Reducing the dilution rate or increasing S_0 enabled survival of first the prey alone and then the predator. Further increases in S_0 destabilised the system, resulting first in damped and then sustained oscillations, before finally reaching a linearly unstable state. We ran simulations in each of the regimes and found that they agreed with the outcomes predicted from the model analysis (Fig. 2b-g), apart from giving sustained, extreme oscillations where the analytical results predicted a linearly unstable state that would correspond to washout (Fig. 2d).

Since the oscillations generated by our *Bdellovibrio* model had a much longer period, extremely abrupt rises and falls, and a strong asymmetry in wave shapes compared to the typical Lotka-Volterra models for animal populations, we compared our model with a protist predator model by Curds and Bazin (29) that generated similarly extreme, but shorter period, oscillations. The period of the protist model could be increased to that of the *Bdellovibrio* model just by replacing protist with *Bdellovibrio* kinetic parameters (SI text on Protist model, Fig. S6).

Dimensional analysis was then performed to deduce which combinations of parameters determine the qualitative behaviour of the system, yielding 7 independent parameter combinations. Essentially, rate parameters became relative to the dilution rate; inflow

substrate concentration relative to the prey's K-value and the predator's K-value replaced

by its ratio to the prey K-value (for details see SI text). These K-values are half-saturation

constants and a low K-value corresponds to high affinity and vice-versa. The next 7

sections show the effect of these 7 dimensionless parameter combinations that determine

qualitative behaviour.

Bdellovibrio has a minimal and optimal attack rate constant

To our knowledge, Varon & Zeigler (13) and Hobley, Summers et al. (18) are the only

studies of predation kinetics where experimental data were used to infer parameters of a

model. Varon & Zeigler (13) used a relative of *Bdellovibrio*, the marine strain BM4, and

Photobacterium leiognathi as prey. *P. leiognathi* can grow at rates of up to 0.2 h^{-1} (30),

however, nothing beyond the study by Varon and Zeigler (13) is known about the growth

kinetics of strain BM4. Hobley, Summers et al. (18) used the type strain of *Bdellovibrio*

bacteriovorus (HD100), with *E. coli* as the prey species, which are the species assumed

in our study. Predictions from the Hobley, Summers et al. study were used to calculate

default values for the attack kinetics. Since different predator and prey combinations may

have different kinetics, we investigated the effect of varying the attack rate constant (μ_P),

which in the Holling type II functional response corresponds to the catalytic rate constant

of an enzyme with Michaelis-Menten type saturation. The ratio of attack rate constant and

K-value ($K_{N,P}$) gives the initial slope of the Holling type II function, which determines the

predation rate at low prey densities. We expected that the faster the predator was at

locating and attacking prey, the more successful it would be, particularly given its high

mortality (half-life of 10 hours), implying that *Bdellovibrio*'s attack rate constant was limited

by intrinsic constraints preventing it from being any faster (at up to 160 µm s^{-1} it is already

one of the fastest swimming bacteria known, especially considering its small size).

Instead, we found that the highest attack rate constant was not the best, as a lower value

was optimal for the abundance of the predator, and conversely for the prey (Fig. 3). The

position and width of this optimum varied with dilution rate (Fig. 3c, d). Increasing levels

of S_0 narrowed the range of μ_P in which the predator could achieve near maximal density.

Below the optimal μ_P, there was a sharp drop to predator extinction. The optimal μ_P was

also the rate at which the system underwent a Hopf bifurcation (31) from a stable steady

state of co-existence into an oscillatory regime (Fig. 3e-g).   The finding that too fast

predation was not optimal can be understood by realising that the food source for the

predator, a living organism, is a renewable resource, if it is given time to regrow. Hence,

a too effective predator reduces regrowth of its prey and will starve.

Higher prey growth rate does not benefit prey

Since *Bdellovibrio* can prey on a wide range of Gram-negative bacteria, which have a

wide range of maximum specific growth rates (μ_N), not just *E. coli*, we investigated the

effects of prey growth rate. We expected that an increased prey growth rate would benefit

both predator and prey. Instead, increasing prey growth rate benefitted only the predator

and never the prey. At low inflow substrate concentration (S_0), populations were co-

existing in a stable steady state, where surprisingly prey growth rate had no effect on prey

and predator density (Fig. 4b, cf. phase diagram in Fig. 2a). At high S_0, populations were

co-existing in sustained oscillations, and increasing prey growth rate led to decreasing

prey abundance and then a sharp drop in prey abundance at the bifurcation point where

the system became stable (Fig. 4a, c). Overall, higher prey growth rate did not benefit the

prey.

Predators benefit from mortality

Mortality of *Bdellovibrio* is much higher than that of other bacteria. Therefore, we included

it in our model and swept death rates from 0 to 0.2 h^{-1} – substantially more than the 0.06

h^{-1} reported for *Bdellovibrio* (6). Surprisingly, predator death was beneficial for the

predator with an optimal death rate just above a critical mortality where oscillations were

replaced with stable co-existence (Fig. S7). Further increases in mortality caused

predator extinction. Increasing inflow substrate concentration (S_0) and dilution rate (D)

narrowed the predator peak, giving a narrow window for predator persistence (Fig. S7).

There is an optimal maturation rate for bdelloplasts

The bdelloplast stage, where the predator grows inside the prey periplasm with a certain

specific growth rate (maturation rate - k_P) until prey resources are exhausted and

offspring is produced, takes ~3 hours with *E. coli*. We found, as with most other

parameters, that there was a minimal maturation rate required for predator survival, and

an optimal maturation rate (Fig. S8). This optimal rate was just below a critical rate, above

which populations oscillated at higher inflow substrate concentration.

Highest affinity of the predator for its prey is not always optimal

The dimensional analysis identified that the system behaviour depends on the ratio of the

K-values $K_{N,P}$ and $K_{S,N}$ (SI text, so we swept through a range of values). At low inflow

substrate concentrations (S_0), populations did not cycle and the lowest $K_{N,P}$ was optimal

(Fig. S9a). At high S_0, in contrast, too low $K_{N,P}$ resulted in oscillations with reduced

average predator levels (Fig. S9b-e). Raising the $K_{N,P}$ resulted in a bifurcation from

extreme oscillations to a stable co-existence that benefited the predator at the expense

of the prey. The optimal $K_{N,P}$ for the predator was just above this critical $K_{N,P}$ (SI text).

Increasing prey productivity benefits predator until an optimum is

reached

285   Substrate inflow (S_0) determines prey productivity. Increasing prey productivity from 0 at

286   the outset benefits the predator much more than the prey, which remains at very low

287   levels, until a predator maximum is reached. Further increasing substrate inflow led to a

288   drop in predator and rise in prey (SI text, Fig. S10).

Bdelloplast burst size

The last of the 7 dimensionless parameter combinations is the burst size, which increases

with prey size. It is not surprising that there was a minimal burst size for predator survival,

but it was not expected that this minimal burst size (at higher dilution rates or inflow

substrate concentrations) was higher than the experimentally determined burst size for

E. coli of 3.5 (Fig. S11). We also found an optimal burst size, after which predator

abundance declined, i.e., too large prey was not optimal. The lower the dilution rate, the

lower the minimal and optimal burst size and the broader the optimum such that the

optimum was in the range corresponding to typical prey sizes (Fig. S11). Increasing the

burst size above the optimal value resulted in a bifurcation to extreme oscillations,

corresponding to a sharp rise in prey and drop in predator average densities, but only at

low S_0. At higher S_0, the optimum was very narrow, above values corresponding to typical

prey and oscillations did not occur (Fig. S11).

Bacteriophage outcompetes Bdellovibrio

We asked whether bacteriophages would outcompete *Bdellovibrio* on single prey populations, and if so, under which conditions and why. The bacteriophage was parameterised based on the well-studied T4 phage infecting *E. coli*. T4 caused oscillations and outcompeted *Bdellovibrio* (Fig. 5). Why did the phage win? There are three processes where the two predators differ. Firstly, the attack kinetics, where the phage had a higher attack rate constant (μ_P), but a higher K-value ($K_{N,P}$). Secondly, the kinetics of prey consumption, where the phage had a higher burst size ($Y_{P/B}$) and a faster maturation rate (k_P). Thirdly, the phage, unlike *Bdellovibrio*, did not have mortality. To find out which advantage(s) allowed the phage to win, we ran competitions where the phage kept the prey $K_{N,P}$ disadvantage and had one or more of the advantages. Increased burst size ($Y_{P/B}$) alone was sufficient for the phage to win (Fig. S12d). A combination of increased μ_P and reduced mortality was also sufficient (Fig. S12a, b, e). Increased k_P was insufficient even in the presence of either an increased μ_P or reduced mortality (Fig. S12c, f, g). Simulating the experiments of Williams et al. (32) where a bacterial predator (but no phage) responded upon addition of prey to a mesocosm, suggests that if suitable phage were present, they should have responded better to the prey addition (SI text, Fig. S13).

Trade-off between high rate versus high yield for predators

Since it takes time to convert prey resources into predator biomass, more complete exploitation of prey resources (higher yield or burst size) should come at the cost of a longer maturation time before offspring will emerge (lower maturation rate - k_P). The fast

324   but wasteful predator was assumed to convert bdelloplasts into offspring at a higher rate

325   (k_P one third higher), but had a lower burst size ($Y_{P/_B}$ halved) to reflect a reduced yield.

326   The high yield strategy outcompeted the high rate strategy (Fig. 6). Note that the phage

327   had both higher burst size and faster maturation rate, considering this, it is not surprising

328   that the phage won (cf. Fig. 5).

Co-existence of two predators on one prey

330   We assumed that a trade-off between attack rate constant (μ_P) and K-value ($K_{N,P}$) exists

331   because time has to be invested for finding prey as well as for binding and examining

332   potential prey. We found that this trade-off would allow coexistence of a 'fast' predator

333   with a higher μ_P and a 'high affinity' predator with lower $K_{N,P}$ on a single prey (Fig. S14).

Minimal and optimal prey cell sizes

Bacteria have a large range of sizes (and therefore biomass) (33). Clearly prey can be

physically too small to be entered and consumed. *Bdellovibrio* also needs to produce at

least two offspring per prey cell. Cells of *B. bacteriovorus* are around seven times smaller

than *E. coli* cells; this small size might enable *Bdellovibrio* to prey on cells with a wider

range of sizes. While consuming a larger prey cell will produce more offspring, it will take

longer. Also, larger prey cells will mean fewer prey are available if prey growth is limited

by the substrate entering the system as in our chemostat case and most environments. It

is not obvious whether the higher number of offspring per prey would offset the

disadvantages of longer maturation times and fewer cells to hunt. We found that there

was both a minimum prey to predator size ratio required for predator persistence and an

optimal value for maximal predator biomass (Fig. 7). Fat prey caused the system to

display extreme oscillations. The optimal prey size was just below the size causing these

oscillations. Increases in S_0 narrowed the optimum towards the minimal prey size (Fig.

7a-c). Increases in dilution rate also narrowed the peak, but increased the optimal prey

size (Fig. 7c, d).

Bottlenecks, permanence and paradox of enrichment

When sweeping parameters, it became clear that the system oscillated with long periods

of hundreds of hours and extreme amplitudes for many of the conditions tested, with

extremely low minimum values (< 1 x 10^{-30} mg dry mass ml^{-1}). Such bottlenecks would

result in the extinction of the predator (or both predator and prey) in any natural system

prone to extrinsic fluctuations and governed by stochastic processes. We therefore

examined how prey size (previous section) in combination with dilution rate and inflow

substrate concentration (cf. Fig. 2a) would affect robust predator persistence (union of

regions of stable co-existence and damped oscillations in Fig. 2a). Robust persistence is

known as permanence and means that population densities do not approach zero as the

boundary is a repeller (34). The prey size range allowing permanence narrowed with

increasing S_0 (Fig. S15). This effect of increased productivity (higher influx of resource for

prey) destabilising the system is known as the paradox of enrichment (35) and is typical

for predator-prey systems. Prey also had to be unrealistically large to enable permanence

at higher dilution rates and lower productivity.

Tragedy of the commons

Similarly to increasing productivity, the prey size range enabling permanence shrank with

increasing attack rate constant of the predator (Fig. S16). This is an example of the

tragedy of the commons, where overexploitation of a shared resource known as the

commons is to the detriment of the users of the resource, such as overfishing (36). Here,

a too effective predator that consumed its prey faster than it could regrow led to extinction

of the predator. Increases in the inflow substrate concentration (S_0) narrowed the prey

size range for permanence further whilst increases in dilution rate expanded the range

(Fig. S16).

374   *Global parameter sensitivity analysis*

To understand how much the system behaviour would change if the parameters were

different due to changes in substrate, prey or predator species, we conducted a global

sensitivity analysis (SI text and Figs. S17, S18). Substrate concentration was only

sensitive to the inflow substrate concentration. Prey density was not sensitive to substrate

concentration and prey growth kinetics, but sensitive to predator parameters. Predator

densities were sensitive to the attack rate constant, but also to the maximal specific prey

growth rate.

Discussion

Our results suggest that *Bdellovibrio* consuming a single prey species can only survive

permanently within a very narrow range of conditions. We therefore characterise this

predator-prey system as 'brittle'. For example, over a wide range of conditions, the system

is prone to extreme oscillations with periods of over a hundred hours and bacterial

densities dropping below 0.1 pg ml^{-1}. In a deterministic mathematical model, species

densities can eventually recover from even the smallest positive number but in a biological

system, there has to be at least a single cell left (~1 pg). More importantly, when a system

contains just a few cells over a long time, stochastic fluctuations will almost inevitably

result in the loss of those few cells, leading to local extinction. For most system

parameters, there was a narrow range of values that allowed the predator to reproduce

fast enough to avoid being washed out and not trigger these oscillations, and the optimal

value to maximise predator numbers occurred near the threshold triggering these

oscillations. Increasing nutrient concentrations narrowed this survival range. Previous

models of microbial predator-prey interactions in chemostats, including those with

Bdellovibrio as the predator, also showed a tendency to extreme oscillations, especially

for higher nutrient concentrations (14, 15, 29, 37). Only the group of Varon studied this

experimentally in chemostats with the *Bdellovibrio* like predator BM4 and *Photobacterium*

leiognathi as prey. They did observe oscillations, albeit less extreme ones than in our

model (38, 39). They also found that BM4 could survive in a chemostat with a prey density

of 2-5 x 10^4 CFU ml^{-1} (39), substantially less than the 7 x 10^5 CFU ml^{-1} predicted by their

model (13) and less still than the 4.4 x 10^6 CFU ml^{-1} minimum required for survival in our

model. In contrast, Keya and Alexander (40) found *Bdellovibrio* strain PF13, isolated from

soil, would only replicate in the presence of at least 3 x 10^7 CFU ml^{-1} of its *Rhizobium*

prey. Studies so far do not allow us to judge whether predictions are reasonable.

Given the brittle behaviour predicted by the model it is surprising that *Bdellovibrio* is

ubiquitous in non-marine environments, while *Bdellovibrio* like organisms are ubiquitous

in marine environments (41–44). This suggests that there must be other forces at work

stabilising population numbers. One possibility is that *Bdellovibrio* is a hotspot organism

targeting habitats of high prey density, or structured environments containing areas of

high prey density such as biofilms, and that these hotspots are the sources of *Bdellovibrio,*

and other habitats are sinks. Biofilms represent a lump of prey for *Bdellovibrio*. It

possesses the lytic enzymes needed to chew the extracellular matrix that holds cells

within the biofilm and can likely derive valuable nutrients from this (45). *Bdellovibrio* can

predate the metabolically inactive cells found deeper within biofilms and its gliding motility

allows it to move within the biofilm (46). *Bdellovibrio* is also highly motile, and contains

chemotaxis genes that might enable it to locate these prey rich areas (47) – so hotspots

that are only temporary can be located. Indeed, there is ample evidence that *Bdellovibrio*

is a hotspot organism. It moves towards regions rich in bacteria (48). Higher numbers are

found in sewage rather than in rivers (49, 50), and downstream sewage treatment plants

rather than upstream (51) or in sewage treated rather than untreated soils (52). The prey

rich rhizosphere sustains higher predator numbers than the bulk soil (53). Eutrophic lakes

support higher numbers than oligotrophic lakes (54). *Bdellovibrio* like organisms are more

abundant in marine sediments than the water column and much higher in oyster shell

biofilms (55). *Bdellovibrio* are likewise more common in trickling filter biofilms than in their

inflow (56).

However, the model predicts a narrower range of conditions for survival at the higher

nutrient concentrations expected in hotspots. Therefore, it is unlikely that these hotspots

would sustain *Bdellovibrio* if it were only predating a single prey species. Moreover, our

model also suggests that bacteriophage would outcompete *Bdellovibrio* under all

conditions tested. This is because phage, as specialist predators, have several big

advantages. Indeed, bacteriophage are far more numerous in nature than *Bdellovibrio*

(57, 58) (this comparison is reasonable assuming that *Bdellovibrio* in total can prey on

about half of all bacteria that phage in total can prey on). This is despite the fact that

phage have several disadvantages that we did not consider. First, half of the sequenced

bacterial genomes contain a CRISPR-CAS system providing adaptive immunity against

phage (59); restriction enzymes cutting up phage DNA are also common (60). Second,

bacteria can rapidly evolve resistance to phages (61), in contrast to *Bdellovibrio*, although

phenotypic plasticity of prey can afford temporary protection against *Bdellovibrio* (18, 44).

Third, phage require a metabolically active host cell to replicate whereas *Bdellovibrio* can

consume dormant prey (4). Fourth, phage do not benefit from chemotaxis as *Bdellovibrio*

likely does. Nevertheless, phage outnumber *Bdellovibrio*, suggesting that these four

factors can be overcome by the huge diversity of phage.

There are several factors which may promote the survival of BALOs in the natural

environment. Firstly, natural environments are spatially heterogeneous, facilitating spatial

refuges promoting the survival of prey species and by extension the survival of BALOs

(19, 20, 62). Secondly, characteristics of the environment can make predation less

efficient (giving prey a greater opportunity to recover) (63–66). Thirdly, seasonal changes

can result in a temporal refuge, with prey concentrations varying with temperature (67,

68). Other temporal fluctuations could create hotspots and refuges. Fourthly, other

predators, such as protists, can stabilize prey populations as they can prey on BALOs

(69) and can also permit the survival of prey species in a mixed microbial community (70).

Finally, prey diversity itself promotes the survival of a generalist predator, such as

Bdellovibrio, and the presence of *Bdellovibrio* has been shown to positively correlate with

alpha-diversity (68, 71–73). Overall, our results suggest that *Bdellovibrio* must prey on

several prey species and locate transient hotspots to survive. Indeed, all known

Bdellovibrio and like organisms have a wide prey range (4, 53, 74, 75).

One might expect that prey cells have to be large enough to give rise to two predator

offspring, although it has been reported that it is possible for *Bdellovibrio* to start

replication within one prey cell and complete this in a second prey (76). Surprisingly, our

model predicts that, in all conditions tested, *Bdellovibrio* needs prey that is at least 7 times

larger than itself. Indeed, this is about the difference in size between *Bdellovibrio* and *E.*

coli (21) and other typical prey are of similar size. However, the predicted optimal prey

size is considerably larger. Maybe *Bdellovibrio* cannot be much smaller than it already is,

or accessing diverse prey species avoids precarious oscillations.

Our model also predicts that a predator that kills its prey too efficiently (has a too high

attack rate, or too little mortality) will drive its prey to extinction and become extinct itself,

a tragedy of the commons similar to over-fishing (36). Indeed, *Bdellovibrio* has a much

higher mortality than other bacteria. While this may make over-exploitation of prey less

likely, the high mortality is likely caused by its high energy expenditure when swimming

fast and not feeding at the same time, also its small size prevents storage of energy

reserves (6).

In conclusion, our model results suggest that *Bdellovibrio* and like organisms are unlikely

to survive in most natural environments if they were preying only on a single prey species.

They would also be outcompeted by phage. In line with empirical evidence, *Bdellovibrio*

ought to be a generalist predator and would only thrive in prey density hotspots – which

it should be able to find by chemotaxis. For application as a living antibiotic to reduce the

abundance of pathogens or antimicrobial resistant bacteria in aquaculture or plant and

animal agriculture, *Bdellovibrio* would be expected to be more effective where multiple

prey species, not only the target species, are naturally available or added artificially.

482 Acknowledgments

483   We would like to thank Andrew Lovering (University of Birmingham) and Liz Sockett

484   (University of Nottingham) for helpful discussions of *Bdellovibrio* biology and Jamie Wood

485   (University of York) for suggestions on predator-prey modelling. KS was supported by a

486   Biotechnology and Biological Sciences Research Council (BBSRC) UK funded, Midlands

487   Integrative Biosciences Training Partnership (MIBTP) PhD studentship.

488 Competing Interests

489   The authors declare that they have no competing financial interests.

References

1.   Paine RT. 1969. A note on trophic complexity and community stability. Am Nat
103:91–93.
2.   Krebs CJ, Boonstra R, Stan B, Sinclair ARE. 2001. What drives the 10-year cycle
of Snowshoe hares? Bioscience 51:25–35.
3.   Arcilla MS, van Hattem JM, Matamaros S, Melles DC, Penders J, de Jong MD.
2016. Dissemination of the *mcr-1* colistin resistance gene. Lancet Infect Dis
16:147–149.
4.   Stolp H, Starr MP. 1963. *Bdellovibrio bacteriovorus* gen. et sp. n., a predatory,
ectoparasitic, and bacteriolytic microorganism. Antonie Van Leeuwenhoek 29:217–
248.
5.   Lambert C, Morehouse KA, Chang C-Y, Sockett RE. 2006. *Bdellovibrio*: growth and
development during the predatory cycle. Curr Opin Microbiol 9:639–644.
6.   Hespell RB, Thomashow MF, Rittenberg SC. 1974. Changes in cell composition
and viability of *Bdellovibrio bacteriovorus* during starvation. Arch Microbiol 97:313–
327.
7.   Hobley L, King JR, Sockett RE. 2006. *Bdellovibrio* predation in the presence of
decoys: Three-way bacterial interactions revealed by mathematical and
experimental analyses. Appl Environ Microbiol 72:6757–6765.
8.   Starr MP, Baigent NL. 1966. Parasitic interaction of *Bdellovibrio bacteriovorus* with
other bacteria. J Bacteriol 91:2006–2017.
9.   Romo AJ, Ruby EG, Saier MHJ. 1992. Effect of *Bdellovibrio bacteriovorus* infection
on the phosphoenolpyruvate:sugar phosphotransferase system in *Escherichia coli*
: evidence for activation of cytoplasmic proteolysis. Res Microbiol Paris 143:5–14.
10. Seidler RJ, Starr MP. 1969. Factors affecting the intracellular parasitic growth of
Bdellovibrio bacteriovorus developing within *Escherichia coli*. J Bacteriol 97:912–
23.
11. Berryman AA. 1992. The origins and evolution of predator-prey theory. Ecology
73:1530–1535.
12. Wilkinson MHF. 2006. Mathematical modelling of predatory prokaryotes, p. 94–130.
In Jurkevitch, E (ed.), Predatory Prokaryotes.
13. Varon M, Zeigler BP. 1978. Bacterial predator-prey interaction at low prey density.
Appl Environ Microbiol 36:11–17.
14. Crowley PH, Straley SC, Craig RJ, Culin JD, Fu YT, Hayden TL, Robinson TA,
Straley JP. 1980. A model of prey bacteria, predator bacteria, and bacteriophage
in continuous culture. J Theor Biol 86:377–400.
15. Wilkinson MHF. 2001. Predation in the presence of decoys: an inhibitory factor on
pathogen control by bacteriophages or *Bdellovibrios* in dense and diverse
ecosystems. J Theor Biol 208:27–36.
16. Baker M, Negus D, Raghunathan D, Radford PM, Moore C, Clark G, Diggle M,
Tyson J, Twycross J, Sockett RE. 2017. Measuring and modelling the response of
Klebsiella pneumoniae KPC prey to *Bdellovibrio bacteriovorus* predation, in human
serum and defined buffer. Sci Rep 7:1–18.
17. Said N, Chatzinotas A, Schmidt M. 2019. Have an ion on it: The life-cycle of
Bdellovibrio bacteriovorus viewed by helium-ion microscopy. Adv Biosyst

3:1800250.

18. Hobley L, Summers JK, Till R, Milner DS, Atterbury RJ, Stroud A, Capeness MJ,
Gray S, Leidenroth A, Lambert C, Connerton I, Twycross J, Baker M, Tyson J, Kreft
J-U, Sockett RE. 2020. Dual predation by bacteriophage and *Bdellovibrio* can
eradicate *E. coli* prey in situations where single predation cannot. J Bacteriol 202:1–
32.

19. Hol FJH, Rotem O, Jurkevitch E, Dekker C, Koster DA. 2016. Bacterial predator–
prey dynamics in microscale patchy landscapes. Proc R Soc B Biol Sci 283:1–9.

20. Dattner I, Miller E, Petrenko M, Kadouri DE, Jurkevitch E, Huppert A. 2017.
Modelling and parameter inference of predator–prey dynamics in heterogeneous
environments using the direct integral approach. J R Soc Interface 14.

21. Cover WH, Martinez RJ, Rittenberg SC. 1984. Permeability of the boundary layers
of *Bdellovibrio bacteriovorus 109J* and its bdelloplasts to small hydrophilic
molecules. J Bacteriol 157:385–390.

22. Jannasch HW. 1969. Estimations of bacterial growth rates in natural waters. J
Bacteriol 99:156–160.

23. Hellweger FL, Clegg RJ, Clark JR, Plugge CM, Kreft J-U. 2016. Advancing
microbial sciences by individual-based modelling. Nat Rev Microbiol 14:461–471.

24. Kreft J-U, Booth G, Wimpenny JWT. 1998. BacSim, a simulator for individual-based
modelling of bacterial colony growth. Microbiology 144:3275–3287.

25. Kubitschek HE, Friske JA. 1986. Determination of bacterial cell volume with the
Coulter Counter. J Bacteriol 168:1466–7.

26. Rittenberg SC, Shilo M. 1970. Early host damage in the infection cycle of
Bdellovibrio bacteriovorus. J Bacteriol 102:149–60.

27. Jeschke JM, Kopp M, Tollrian R. 2002. Predator functional responses:
Discriminating between handling and digesting prey. Ecol Monogr 72:95–112.

28. Hespell RB. 1976. Glycolytic and tricarboxylic acid cycle enzyme activities during
intraperiplasmic growth of *Bdellovibrio bacteriovorus* on *Escherichia coli*. J Bacteriol
128:677–80.

29. Curds CR, Bazin MJ. 1980. Protozoan predation in batch and continuous cultures,
p. 115–176. *In* Droop, MR, Jannasch, HW (eds.), Advances in Aquatic
Microbiology. Academic Press.

30. Dunlap P V. 1985. Osmotic control of luminescence and growth in *Photobacterium
leiognathi* from ponyfish light organs. Arch Microbiol 141:44–50.

31. Hopf E. 1948. A mathematical example displaying features of turbulence. Commun
Pure Appl Math 1:303–322.

32. Williams HN, Lymperopoulou DS, Athar R, Chauhan A, Dickerson TL, Chen H,
Laws EA, Berhane T-K, Flowers ARA, Bradley N, Young S, Blackwood D, Murray
J, Mustapha O, Blackwell C, Tung Y, Noble RT, Williams HN, Florida A, King SML,
Blvd J. 2016. *Halobacteriovorax*, an underestimated predator on bacteria: potential
impact relative to viruses on bacterial mortality. ISME J 10:491–499.

33. Schulz HN, Barker B, Jørgensen BB. 2001. Big bacteria. Molecules 55:105–37.

34. Jansen VAA, Sigmund K. 1998. Shaken not stirred: on permanence in ecological
communities. Theor Popul Biol 54:195–201.

35. Rosenzweig ML. 1971. Paradox of enrichment: destabilization of exploitation
ecosystems in ecological time. Science 171:385–7.

36. Hardin G. 1968. The tragedy of the commons. The population problem has no
technical solution; it requires a fundamental extension in morality. Science
162:1243–8.
37. Nisbet RM, Cunningham A, Gurney WS. 1983. Endogenous metabolism and the
stability of microbial prey-predator systems. Biotechnol Bioeng 25:301–306.
38. Varon M. 1979. Selection of predation-resistant bacteria in continuous culture.
Nature 277:386–388.
39. Varon M, Fine M, Stein A. 1984. The maintenance of *Bdellovibrio* at low prey
density. Microb Ecol 10:95–98.
40. Keya SO, Alexander M. 1974. Regulation of parasitism by host density: the
Bdellovibrio-Rhizobium interrelationship. Soil Biol Biochem 7:231–237.
41. Shilo M, Bruff B, Shilo BYM, Bruff B, Bacteriology D, Calqorrnia U. 1965. Lysis of
Gram-negative   bacteria   by   host-independent   ectoparasitic   *Bdellovibrio*
bacteriovorus isolates. J Gen Microbiol 40:317–28.
42. Williams HN. 1987. The recovery of high numbers of *Bdellovibrios* from the surface
water micro layer. Can J Microbiol 33.
43. Williams HN, Schoeffield AJ, Guether D, Kelley J, Shah D, Falkler WA. 1995.
Recovery of *bdellovibrios* from submerged surfaces and other aquatic habitats.
Microb Ecol 29:39–48.
44. Shemesh Y, Jurkevitch E. 2004. Plastic phenotypic resistance to predation by
Bdellovibrio and like organisms in bacterial prey. Environ Microbiol 6:12–18.
45. Im H, Dwidar M, Mitchell RJ. 2018. *Bdellovibrio bacteriovorus HD100*, a predator
of Gram-negative bacteria, benefits energetically from *Staphylococcus aureus*
biofilms without predation. ISME J 12:2090–2095.
46. Lambert C, Fenton AK, Hobley L, Sockett RE. 2011. Predatory *Bdellovibrio* bacteria
use gliding motility to scout for prey on surfaces. J Bacteriol 193:3139–41.
47. Hobley L, Lerner TR, Williams LE, Lambert C, Till R, Milner DS, Basford SM,
Capeness MJ, Fenton AK, Atterbury RJ, Harris MATS, Sockett RE. 2012. Genome
analysis of a simultaneously predatory and prey-independent, novel *Bdellovibrio*
bacteriovorus from the River Tiber, supports *in silico* predictions of both ancient and
recent lateral gene transfer from diverse bacteria. BMC Genomics 13:670.
48. Straley SC, Conti SF. 1977. Chemotaxis by *Bdellovibrio bacteriovorus* toward prey.
J Bacteriol 132:628–40.
49. Dias FF, Bhat J V. 1965. Microbial ecology of activated sludge. II. bacteriophages,
Bdellovibrio, coliforms, and other organisms. Appl Microbiol 13:257–261.
50. Seliavko A V., Lambina VA. 1985. Distribution of microorganisms lysing
Pseudomonas aeruginosa and *Staphylococcus aureus* in the system of sewage
waters. Nauchnye Doki Vyss Shkoly Biol Nauk 4:84–8.
51. Staples DG, Fry JC. 1973. Factors which influence the enumeration of *Bdellovibrio*
bacteriovorus in sewage and river water. J Appl Bacteriol 36:1–11.
52. Klein DA, Casida LEJ. 1967. Occurrence and enumeration of *Bdellovibrio*
bacteriovorus in soil capable of parasitizing *Escherichia coli* and indigenous soil
bacteria. Can J Microbiol 13:1235–41.
53. Jurkevitch E, Minz D, Ramati B, Barel G. 2000. Prey range characterization,
ribotyping, and diversity of soil and rhizosphere *Bdellovibrio spp.* isolated on
phytopathogenic bacteria. Appl Environ Microbiol 66:2365–71.

54.   Chauhan A, Cherrier J, Williams HN. 2009. Impact of sideways and bottom-up
628   control factors on bacterial community succession over a tidal cycle. Proc Natl Acad
629   Sci U S A 106:4301–6.
630   55.   Williams HN, Kelley JI, Baer ML, Turng B-F. 1995. The association of *Bdellovibrios*
631   with surfaces in the aquatic environment. Can J Biochem 41:1142–1147.
632   56.   Fry JC, Staples DG. 1976. Distribution of *Bdellovibrio bacteriovorus* in sewage
633   works, river water, and sediments. Appl Environ Microbiol 31:469–474.
634   57.   Kutter   EM,   Sulakvelidze   A.   2004.   Bacteriophages: Biology   and
635   applicationsBacteriophages Biology and Applications. Taylor & Francis Group,
636   Boca Raton.
637   58.   Stolp H. 1973. The *Bdellovibrios*: Bacterial parasites of bacteria. Annu Rev
638   Phytopathol 11:53–76.
639   59.   Koonin E V., Wolf YI. 2015. Evolution of the CRISPR-Cas adaptive immunity
640   systems in prokaryotes: models and observations on virus–host coevolution. Mol
641   Biosyst 11:20–27.
642   60.   Bayliss CD, Callaghan MJ, Moxon ER. 2006. High allelic diversity in the
643   methyltransferase gene of a phase variable type III restriction-modification system
644   has implications for the fitness of *Haemophilus influenzae*. Nucleic Acids Res
645   34:4046–59.
646   61.   Luria SE, Delbrück M. 1943. Mutations of bacteria from virus sensitivity to virus
647   resistance. Genetics 28:491–511.
648   62.   Petrenko M, Friedman SP, Fluss R, Pasternak Z, Huppert A, Jurkevitch E. 2019.
649   Spatial heterogeneity stabilizes predator–prey interactions at the microscale while
650   patch connectivity controls their outcome. Environ Microbiol 00.
651   63.   Monnappa AK, Dwidar M, Mitchell RJ. 2013. Application of bacterial predation to
652   mitigate recombinant bacterial populations and their DNA. Soil Biol Biochem
653   57:427–435.
654   64.   Im H, Son S, Mitchell RJ, Ghim C-M. 2017. Serum albumin and osmolality inhibit
655   *Bdellovibrio bacteriovorus* predation in human serum. Sci Rep 7:5896.
656   65.   Sathyamoorthy R, Maoz A, Pasternak Z, Im H, Huppert A, Kadouri DE, Jurkevitch
657   E. 2019. Bacterial predation under changing viscosities. Environ Microbiol 1462-
658   2920.14696.
659   66.   Im H, Kwon H, Cho G, Kwon J, Choi SY, Mitchell RJ. 2019. Viscosity has
660   dichotomous effects on *Bdellovibrio bacteriovorus* HD100 predation. Environ
661   Microbiol 21:4675–4684.
662   67.   Ottaviani D, Pieralisi S, Chierichetti S, Rocchegiani E, Hattab J, Mosca F, Tiscar
663   PG, Leoni F, Angelico G. 2020. *Vibrio parahaemolyticus* control in mussels by a
664   *Halobacteriovorax* isolated from the Adriatic sea, Italy. Food Microbiol 92:103600.
665   68.   Ezzedine JA, Jacas L, Desdevises Y, Jacquet S. 2020. *Bdellovibrio* and like
666   organisms in lake Geneva : An unseen elephant in the room? Front Microbiol 11:1–
667   14.
668   69.   Johnke J, Boenigk J, Harms H, Chatzinotas A. 2016. Killing the killer: Predation
669   between protists and predatory bacteria. FEMS Microbiol Lett 364.
670   70.   Johnke J, Baron M, Leeuw M de, Kushmaro A, Jurkevitch E, Harms H, Chatzinotas
671   A. 2017. A generalist protist predator enables coexistence in multitrophic predator-
672   prey systems containing a phage and the bacterial predator *Bdellovibrio*. Front Ecol

Evol 5:124.
71. Johnke J, Fraune S, Bosch T, Hentschel U, Schulenburg H. 2019. *Bdellovibrio* and
like organisms are predictors of microbiome diversity in distinct host groups. bioRxiv
1–20.
72. Cao Q, Najnine F, Han H, Wu B, Cai J. 2020. BALOs improved gut microbiota
health in postlarval shrimp (*Litopenaeus vannamei*) after being subjected to salinity
reduction treatment. Front Microbiol 11:1–22.
73. Johnke J, Dirksen P, Schulenburg H. 2020. Community assembly of the native *C.*
elegans microbiome is influenced by time, substrate and individual bacterial taxa.
Environ Microbiol 22:1265–1279.
74. Feng S, Tan CH, Cohen Y, Rice SA. 2016. Isolation of *Bdellovibrio bacteriovorus*
from a tropical wastewater treatment plant and predation of mixed species biofilms
assembled by the native community members. Environ Microbiol
https://doi.org/10.1111/1462-2920.13384.
75. Dashiff A, Junka RA, Libera M, Kadouri DE. 2010. Predation of human pathogens
by the predatory bacteria *Micavibrio aeruginosavorus* and *Bdellovibrio*
bacteriovorus. J Appl Microbiol 110:431–444.
76. Makowski Ł, Trojanowski D, Till R, Lambert C, Lowry RC, Sockett RE, Zakrzewska-
Czerwińska J. 2019. Dynamics of chromosome replication and its relationship to
predatory attack lifestyles in *Bdellovibrio bacteriovorus*. bioRxiv 85.
77. Bauer A, Forchhammer K. 2021. Bacterial Predation on Cyanobacteria. Microb
Physiol 1–10.
695

696   **Table 1** Baseline parameters for the principal model (Model 6) and range over which

697   global sensitivity analysis was performed. All yields are expressed in the form $Y_{A/B}$, which

698   is the yield of consumer A per resource B consumed. The half-saturation constant, or K-

699   value, of consumer B for resource A is expressed as $K_{A,B}$.

Parameter	Units	Value	Minimum	Maximum	Source
Inflow substrate concentration (S_0)	mg ml^{-1}	5 x 10^{-3}	5 x 10^{-3}	2.5	Assumed to be ~2 × $K_{S,N}$
Prey maximum specific growth rate (μ_N)	h^{-1}	1.23	¾ ln(2)	3 ln(2)	(24)
Predator attack rate constant (μ_P)	mg bdelloplast mg predator^{-1} h^{-1}	0.38	0.3	6	(18)
Predator K-value $(K_{N,P})$	mg prey ml^{-1}	8.6 x 10^{-4}	1.5 x 10^{-6}	6.5 x 10^{-3}	(18)
Yield of bdelloplasts from predators $(Y_{B/P})$	mg bdelloplast mg predator^{-1}	8	5	20	Relative sizes of *E. coli* and *Bdellovibrio* cells (4, 25)
Bdelloplast maturation rate (k_P)	mg predator mg bdelloplast^{-1} h^{-1}	0.109	0.075	0.3	(10)
Predator mortality rate (m)	h^{-1}	0.06	0.03	0.09	(6)
Dilution rate (D)	h^{-1}	Always 0.05			
Prey K-value $(K_{S,N})$	mg ml^{-1}	Always 2.34 x 10^{-3}			(24)
Yield of prey from substrate $(Y_{N/S})$	mg dry mass prey mg substrate^{-1}	Always 0.4444			(24)
Yield of bdelloplasts from prey $(Y_{B/N})$	mg bdelloplast^{-1} mg prey^{-1}	Always $\frac{Y_{B/P}}{Y_{B/P}-1}$			Relative sizes of *E. coli* and *Bdellovibrio* cells (4, 25)
Yield of predators from bdelloplasts $(Y_{P/B})$	mg predator mg bdelloplast^{-1}	Always 0.438			Relative sizes of *E. coli* and *Bdellovibrio* cells and burst size (4, 10, 25)

FIG 1 Principal model used to track predator and prey densities under chemostat
conditions (Model 6, see Table S2). Substrate is consumed by prey to fuel growth. Prey
and predators combine to form a bdelloplast. The bdelloplast matures to give new
predators. Predators have mortality. This is the model used unless stated otherwise.

FIG 2 Dynamic regimes of Model 6. **a** Analytically calculated regimes, as depending on

the inflow substrate concentration S_0 and dilution rate D. **b-g** Simulations at S_0 = 0.25 mg

ml^{-1} and increasing dilution rates, indicated by white crosses in **a**. **b** Damped oscillations

that ended in steady state co-existence. **c** Damped oscillations of much shorter period

than **b**. **d** Amplifying oscillations that ended in sustained, extreme oscillations. **e** The

analytically predicted linearly unstable region gave sustained, extreme oscillations in the

numerical simulations. **f** Stable co-existence of predator and prey. **g** Predator extinction.

FIG 3 Minimal and optimal attack rate constant (μ_P). The average population densities

and substrate concentrations, oscillatory periods and phase shifts strongly depend on μ_P,

shown at increasing inflow substrate concentrations (S_0) and two dilution rates (cf. phase

diagram in Fig. 2a). Top row shows concentrations at steady state or averaged over one

oscillatory cycle. Bottom row shows the oscillatory period (blue, left axis) and phase shifts

(green, right axis) from substrate peak to peak of prey (solid line), free *Bdellovibrio*

(dashed line) or bdelloplast (dotted line). Note that oscillations start above the optimal μ_P.

In order to obtain accurate simulation results at all parameter values, the absolute

tolerance of the ode45 solver had to be reduced from 1 x 10^{-9} to 1 x 10^{-12}.

FIG 4 Increasing the maximal specific growth rate of the prey (μ_N) leads to a sharp drop

in prey density and stabilizes the system at high inflow substrate concentrations (S_0). The

system was more stable at low S_0 (cf. phase diagram in Fig. 2a). Top row shows

concentrations at steady state or averaged over one oscillatory cycle. Bottom row shows

the oscillatory period (blue, left axis) and phase shifts (green, right axis) from substrate

peak to peak of prey (solid line), free *Bdellovibrio* (dashed line) or bdelloplast (dotted line).

Note that oscillations occur at the higher S_0 and below a critical μ_N. Prey density is much

higher in the oscillatory regimes, where it decreases with increasing μ_N. Since prey

density will of course be 0 if μ_N is zero, there is an optimal μ_N.

FIG 5 T4 phage caused oscillations and outcompeted *Bdellovibrio* at two very different inflow substrate concentrations. Dilution rate was 0.02 h^{-1}. Note that units had to be changed to particle densities from the biomass densities used in the other sections. Panel c shows a zoomed in version of a peak in panel b, the substrate concentration appears to be constant on the scale needed to show the other variables.

FIG 6 A slow, but economical (efficient resource use, therefore high $Y_{P/_B}$) predator

outcompetes a fast (high k_P) but wasteful predator at different inflow substrate

concentrations (S_0). Dilution rate = 0.02 h^{-1}, (a) S_0 = 0.005 mg ml^{-1} and (b) S_0 = 0.25 mg

ml^{-1}. Panel (c) shows a zoomed in version of a peak of panel (b).

FIG 7 Minimal and optimal prey biomass, relative to predator biomass (0.028 pg dry

biomass predator cell^{-1}). Predator density showed a broad optimum at low inflow

substrate concentrations (S_0) that became narrower at higher S_0 and dilution rate. Too

large prey caused oscillations. Top row shows concentrations at steady state or averaged

over one oscillatory cycle. Bottom row shows the oscillatory period (blue, left axis) and

phase shifts (green, right axis) from substrate peak to peak of prey (solid line), free

Bdellovibrio (dashed line) or bdelloplast (dotted line). The vertical orange line, at a

prey:predator ratio of 7, is the biomass ratio between an average *E. coli* and *Bdellovibrio*

and similar to most prey used in laboratory studies or for isolating *Bdellovibrio* from the

environment.