Pollen tube-triggered accumulation of NORTIA at the filiform apparatus facilitates

fertilization in *Arabidopsis thaliana*

Jing Yuan[1,2*], Yan Ju[1,2*], Daniel S. Jones[4,5], Weiwei Zhang[2,3], Noel Lucca[1,2], Christopher J.

Staiger[1,2,3], and Sharon A. Kessler[1,2,**]

[1]Department of Botany and Plant Pathology, Purdue University, West Lafayette, Indiana USA

[2]Purdue Center for Plant Biology, Purdue University, West Lafayette, Indiana USA

[3]Department of Biological Sciences, Purdue University, West Lafayette, Indiana USA

[4]Biology Department, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina

USA (current address)

[5]Department of Microbiology and Plant Biology, University of Oklahoma, Norman, Oklahoma

USA

*These authors contributed equally

** Corresponding author

Email: sakessler@purdue.edu

Abstract

During gamete delivery in *Arabidopsis thaliana*, intercellular communication between the

attracted pollen tube and the receptive synergid cell leads to subcellular events in both cells

culminating in the rupture of the tip-growing pollen tube and release of the sperm cells to

achieve double fertilization. Live imaging of pollen tube reception revealed dynamic subcellular

changes that occur in the female synergid cells. Pollen tube arrival triggers the trafficking of

NORTIA (NTA) MLO protein from Golgi-associated compartments and the accumulation of

endosomes at or near the synergid filiform apparatus, a membrane-rich region that acts as the

site of communication between the pollen tube and synergids. Domain swaps and site-directed

mutagenesis reveal that NTA's C-terminal cytoplasmic tail with its calmodulin-binding domain

influences the subcellular localization and function of NTA in pollen tube reception and that

accumulation of NTA at the filiform apparatus is necessary and sufficient for MLO function in

pollen tube reception.

Key words

MLO, pollination, gametophyte, synergid, pollen tube reception, calmodulin binding domain,

pollen tube

Introduction

Intercellular communication is central to the proper development and maintenance of all

multicellular organisms. During this communication, signals from one cell are perceived by

receptors in another cell and translated into various subcellular responses. These responses

include signal transduction cascades leading to transcription of other genes, calcium signaling,

and trafficking of proteins to different organelles or regions of the cell. A well-studied example of

signal-induced protein trafficking in plant development is the redistribution of the PIN polar auxin

transporters to different sides of the cell during important developmental events such as embryo

patterning, leaf initiation and lateral root initiation [1-3]. In plants, most intercellular

communication occurs between cells that are genetically identical and connected by adjoining

cell walls. One exception is pollination, in which pollen (the male gametophyte) is released from

an anther, transported to a receptive stigma, and produces a tip-growing pollen tube that grows

through the female tissues of the pistil and delivers the two sperm cells to the female

gametophyte (also known as the embryo sac, Figure 1A). The pollen tube's journey through the

pistil requires cell-to-cell interactions with the female that allows water and nutrient uptake and

enables the detection of cues important for guidance toward the female gametes [4].

In the model plant *Arabidopsis thaliana*, complex signaling events ranging from pollen landing

on the stigma to fusion of gametes occur over several hours. Most of our knowledge about the

signaling pathways involved along the pollen tube's journey through the female is limited to the

final stages of pollination and involve a highly specialized pair of female gametophyte cells

known as synergids. During female gametophyte development, meiosis followed by three

rounds of mitosis produce the egg cell and central cell along with 2 synergid cells flanking the

egg cell and 3 antipodal cells on the chalazal end of the embryo sac [5] (Figure 1A). The

synergid cells are accessory cells that control the behavior of the pollen tube during the final

stages of pollination. Before pollen tube arrival, they secrete cysteine-rich LURE peptides that

act as short-range pollen tube attractants that are recognized by receptor-like kinases in the tip

of the pollen tube to regulate the direction of pollen tube growth and guide the pollen tube to the

micropyle of the ovule [6-8]. After pollen tube arrival, the synergids communicate with the pollen

tube to induce changes that result in pollen tube rupture and delivery of the sperm cells [4, 9].

Thus, synergids are critical for ensuring that double fertilization can occur to produce seeds.

In Arabidopsis, live imaging has been used to examine the behavior of both the pollen tube and

the synergids during the process of pollen tube reception. A pollen tube follows the gradient of

LURE attractants, enters the micropyle of the ovule, and pauses its growth for 30 min to 1 h just

outside the receptive synergid [10-12]. During this pause in pollen tube growth, communication

occurs between the pollen tube and the synergids leading to subcellular changes and ultimately

the death of both the pollen tube and the receptive synergid. Cytoplasmic calcium oscillations

occur in both the tip of the pollen tube and in the 2 synergid cells during this communication

phase. Cytoplasmic calcium levels continue to increase in both cell types until the pollen tube

starts to grow again and bursts to release the sperm cells, a catastrophic event for both the

pollen tube and the receptive synergid, which also degenerates [10-12]. Mutations in genes that

regulate communication between the synergids and pollen tube during pollen tube reception

result in a pollen tube overgrowth phenotype in which the pollen tubes are attracted normally to

the ovules, but do not get the signal to burst and release the sperm cells. Presumably, synergid-

induced changes in the cell wall of the pollen tube tip do not occur in these mutants, therefore

the pollen tube continues to grow and coil inside the embryo sac. Synergid-expressed genes

that participate in pollen tube reception include the FERONIA (FER) receptor-like kinase, the

GPI-anchored protein LORELEI (LRE), and the Mildew Resistance Locus-O (MLO) protein

NORTIA (NTA, also known as AtMLO7) [12-17]. Mutations in all of these genes lead to the

pollen tube overgrowth phenotype due to disruption of the pollen tube-synergid communication

pathway.

FER and LRE are necessary for the calcium oscillations that occur in synergids in response to

pollen tube arrival [12]. In contrast, *nta-1* mutants have $[Ca^{2+}]_{cyto}$ oscillations at lower amplitudes,

indicating that NTA may participate in modulating Ca^{2+} fluxes in the synergids during

communication with the pollen tube and likely act downstream of FER and LRE [12]. Like all

members of the MLO gene family, NTA has seven membrane-spanning domains and a

predicted calmodulin-binding domain (CaMBD) in its C-terminal intracellular tail [18, 19].

Calmodulin (CaM) is a small protein that binds Ca^{2+} and is involved in signal transduction for

many cellular processes [20]. We previously showed that the C-terminal domain of NTA could

confer pollen tube reception function to the related MLO8 protein [21], but the significance of the

CaMBD in pollen tube reception remains an open question.

The subcellular localization of important pollen tube reception proteins is not always predictive

of their function in communicating with the pollen tube. As expected for early response proteins,

both FER and LRE are expressed in synergid cells where they localize in or near a specialized

region called the filiform apparatus, a membrane rich area located at the micropyle end of the

synergids [14-17, 22-24]. The filiform apparatus is thought to be important for the secretion of

attractant peptides and is the first site of interaction between the pollen tube and synergid cell

prior to pollen tube reception [25-27]. In contrast, before pollen tube arrival, NTA is localized to a

Golgi-associated compartment within the synergid cell and absent from the filiform apparatus

[21]. At the end of pollen tube reception, NTA protein is only detected at the filiform apparatus,

indicating that this protein changes its subcellular localization during pollen tube reception [13].

This suggests that pollen tube-triggered regulation of the synergid secretory system may be a

crucial subcellular response to pollen tube arrival and that NTA function may be related to its

subcellular distribution; however, the precise timing and significance of NTA's redistribution

remain unclear.

Here, we use a live-imaging system to further characterize synergid cellular dynamics during

pollen tube reception and to determine the timing and significance of the polar redistribution of

NTA to the filiform apparatus. To investigate the link between Ca^{2+} and MLO function in pollen

tube reception, we assayed the influence of the CaMBD on NTA's function and subcellular

distribution through C-terminal truncations and a point mutation disrupting the CaMBD. We

show that the polar redistribution of NTA is triggered by the approach of a pollen tube and is

facilitated by the CaMBD. While most subcellular compartments remain distributed throughout

the synergid cells during pollen tube reception, recycling endosomes respond to pollen tube

arrival by accumulating towards the filiform apparatus. Manipulation of NTA subcellular

localization with domain swaps and a Golgi retention signal revealed that filiform apparatus

localization is necessary and sufficient for MLO function in pollen tube reception.

Results

NTA dynamically accumulates at the filiform apparatus during pollen tube reception

Pollen tube reception requires synergid cells to recognize the approaching pollen tube and to

send signals back to the pollen tube that result in release of the sperm cells at the correct time

and place so that double fertilization can be completed. Based on static images, we previously

reported that NTA-GFP fusion protein localizes to a Golgi-associated compartment in synergids

prior to pollen tube attraction [21]. When imaged after pollen tube reception, NTA-GFP is

concentrated at the micropylar end of the synergid (in or near the filiform apparatus) [13]. NTA-

GFP does not accumulate at the filiform apparatus in *fer* ovules with pollen tube overgrowth,

suggesting that FER-mediated signaling during pollen tube reception triggers NTA-GFP

redistribution that in turn contributes to the interaction of the synergid with the pollen tube [13].

An alternative hypothesis is that pollen tube rupture triggers NTA-GFP redistribution and is a

symptom of pollen tube reception rather than an important contributor to the signaling pathway.

To distinguish between these two possibilities, we used a semi-*in vivo* pollination system

combined with time-lapse spinning disk confocal microscopy to determine the timing of NTA-

GFP redistribution during the pollen tube reception process. In the semi-*in vivo* system, pollen

tubes grow out of a cut style and are attracted to ovules arranged on pollen germination media

[28]. This system has previously been used to quantify and track pollen tube attraction to ovules

and to image $[Ca^{2+}]_{cyto}$ reporters during pollen tube reception [10, 12, 29-32]. To follow

subcellular changes in NTA-GFP protein localization before, during, and after pollen tube arrival,

we used pollen from plants expressing the pollen-specific *AUTOINHIBITED Ca^{2+}-*

ATPase9$_{pro}$::DsRed (*ACA9$_{pro}$::DsRed*) reporter and ovules expressing *NTA$_{pro}$::NTA-GFP* in the

semi-*in vivo* system. Approximately 4 h after pollination, pollen tubes emerged from the style

onto the media and were attracted to ovules (Figure 1B). Fluorescence images in both channels

were collected every 5 min from when a pollen tube approached an ovule until after the pollen

tube ruptured inside the ovule. In ~83% (n=93) of the ovules that attracted a pollen tube and

successfully burst to deliver the sperm cells, NTA-GFP accumulated at the filiform apparatus of

the synergids (Figures 1C-E). The rest of the ovules (~17%) attracted pollen tubes that stopped

growing in the micropyle and did not rupture. In these ovules, NTA-GFP did not accumulate at

the filiform apparatus. To exclude the possibility that prolonged imaging causes stress in

synergids which leads to filiform apparatus accumulation of NTA-GFP, we analyzed neighboring

ovules that did not attract pollen tubes but were imaged together with ovules that attracted

pollen tubes. NTA-GFP did not accumulate at the filiform apparatus in these ovules (n=103)

(Figure 1F). Likewise, ovules that were incubated on pollen germination media without a

pollinated pistil (n=133, Figures 1F and S1) and imaged over the same time frame did not

accumulate NTA-GFP at the filiform apparatus. These data indicate that pollen tube arrival is

necessary for NTA-GFP accumulation at the filiform apparatus rather than being retained in the

Golgi, and that this accumulation is not an artifact of the semi- *in vivo* imaging system. (Figure

1F).

Our semi-*in vivo* system also allowed us to determine the timing of NTA-GFP accumulation at

the filiform apparatus in relation to the position of the pollen tube as it approached the

synergids. We defined the 0 min timepoint as the time where the pollen tube just reached the

micropylar opening of the ovule (Figure 1C, ovules with yellow stars, Movies S1 and S2). In all

cases, the shift of NTA-GFP signal from the Golgi to the filiform apparatus also started from this

time point. During the following 30–50 min, pollen tubes grew through the micropyle region of

ovule and arrived at the filiform apparatus of the receptive synergid cell. During this timeframe,

three quarters to half of the NTA-GFP signal moved to the micropylar end of synergid cells,

indicating that the approach of the pollen tube triggers NTA-GFP accumulation at the filiform

apparatus. As reported in [12] and [10], the arriving pollen tubes paused their growth outside the

filiform apparatus for 30–50 min, presumably for cell-to-cell communication. During this period,

the NTA-GFP signal continued to shift toward the filiform apparatus. At 70–80 min after pollen

tube arrival at the micropyle, NTA-GFP signal was only detected at the filiform apparatus and

pollen tubes resumed growth and ruptured to release the sperm cells (Figures 1C, D and

Movies S1 and S2). Even though only one of the synergids receives the pollen tube, NTA-GFP

accumulated at the filiform apparatus in both synergid cells in response to pollen tube arrival,

similar to the activation of $[Ca^{2+}]_{cyto}$ oscillations in both synergids during pollen tube reception

reported in [12].

Golgi do not concentrate at the filiform apparatus during pollen tube reception

We previously determined that NTA is sequestered in a Golgi-associated compartment in

synergid cells that have not attracted a pollen tube [21]. Our live-imaging data suggest that

NTA-GFP is selectively moved out of the Golgi and transported to the filiform apparatus in

response to pollen tube arrival. However, it is possible that the observed NTA-GFP

accumulation at the filiform apparatus is a result of massive reorganization of subcellular

compartments. To distinguish between these possibilities, we investigated the behavior of the

Golgi in synergid cells during pollen tube reception. We used the semi-*in vivo* imaging system

described above with a synergid-expressed Golgi marker (Man49-mCherry) co-expressed with

NTA-GFP [33]. In all replicates, the Golgi marker was distributed throughout the synergids,

excluded from the filiform apparatus, and co-localized with NTA-GFP as reported previously

(Figure 2A, n=93). When a pollen tube approached the synergids, NTA-GFP accumulated at the

filiform apparatus of the synergids as observed previously (Figure 1), but the Golgi-mCherry

marker remained consistently distributed throughout the synergid cells and did not concentrate

near the filiform apparatus (Figure 2B). In order to examine the behavior of the Golgi during later

stages of pollen tube reception, we used the synergid-expressed Golgi marker line

(*LRE_{pro}::Man49-mCherry*) and pollen that was expressing GFP (*Lat52_{pro}::GFP*). In all cases, the

Golgi marker remained randomly distributed throughout the synergid cells, even after pollen

tube rupture (Figures 2C-D, Movies S3 and S4, n=106). These results indicate that the

accumulation of NTA-GFP at the filiform apparatus during pollen tube reception is not linked to

mass redistribution of Golgi.

Endosomes accumulate at the filiform apparatus during pollen tube reception

A signal from the arriving pollen tube seems to trigger the movement of NTA-GFP out of the

Golgi-associated compartments. It is possible that pollen tube arrival triggers other changes to

synergid subcellular organization. We previously reported the localization of synergid-expressed

markers for the ER, peroxisome, endosome and the trans-Golgi Network (TGN) in unfertilized

ovules *in vivo* using confocal laser scanning microscopy [33]. Before pollen tube arrival, SP-

mCherry-HDEL (an ER-associated marker), mCherry-SKL (a peroxisome-associated marker),

and mCherry-RabA1g (a recycling endosome-associated marker) were all distributed evenly

throughout synergid cells (Figures S2-S4 and 3A; [33]. The TGN-associated marker SYP61

exhibited two types of distribution patterns before pollen tube arrival: in type 1 synergids, the

marker accumulated near the filiform apparatus, whereas type 2 synergids displayed a punctate

distribution pattern throughout the cells (Figure S3; [33]. During pollen tube reception, no

change was seen in the TGN marker distribution in either type of synergids (Figure S3, Movies

S5 and S6). Likewise, the ER and peroxisome markers maintained a diffuse distribution

throughout the synergids and did not accumulate at the filiform apparatus (Figures S2 and S4,

Movies S7-10). In contrast, we detected a more dynamic behavior of the endosome marker

during pollen tube reception. Endosomes are membrane-bound compartments that are involved

in the endocytic membrane transport pathway from the plasma membrane to the vacuole.

Endosomes also transport molecules from the Golgi and either continue to the vacuole or

recycle back to the Golgi [34]. We previously reported that mCherry-RabA1g is distributed

throughout synergid cells and had some overlap with NTA-GFP in synergids of unpollinated

ovules [33]. Using the semi-*in vivo* system, we confirmed that before pollen tube arrival,

mCherry-RabA1g distributed throughout synergid cells (Figure 3A). Interestingly, as pollen

tubes approached, the endosome marker started to accumulate at the filiform apparatus of the

synergid cells (Figures 3A and B). By the time pollen tube reception was completed, most of the

endosome signal was concentrated at or near the filiform apparatus (Figures 3B-F and S5,

Movies S11 and S12). These results indicate that the RabA1g endosome compartments have a

distinct response to pollen tube arrival and may play a role in facilitating the intercellular

signaling pathway that occurs between the synergids and the pollen tube.

Pollen tube-independent targeting of NTA to the filiform apparatus is not toxic to the synergids

The selective targeting of NTA-GFP from the Golgi to the filiform apparatus during pollen tube

arrival (Figures 1 and 2) suggests that NTA trafficking to the pollen tube/synergid interface is

important for the intercellular communication process that occurs between the pollen tube and

synergids. In *nta-1* mutants, around 30% of ovules display pollen tube overgrowth and fail to

complete double fertilization, but the other 70% are fertilized normally [13]. This indicates that

NTA is not absolutely required for pollen tube reception, but may function as a modifier of the

signaling pathway. Since FER signaling in the synergids leads to cell death as pollen tube

reception is completed [17, 23, 24], NTA trafficking to the filiform apparatus could be a

mechanism to regulate this death and would thus require sequestration of "toxic" NTA in the

Golgi before pollen tube arrival. To test this hypothesis, we took advantage of sequence

variation leading to differential subcellular localization of MLO proteins to manipulate the

subcellular localization of NTA. When expressed in synergids, other proteins from the

Arabidopsis MLO family have different subcellular localization patterns, indicating that specific

sequences within the MLOs direct them to different parts of the secretory system [35]. MLO1-

GFP localizes at the filiform apparatus when it is ectopically expressed under control of the

synergid-specific *MYB98* promoter and cannot complement the *nta-1* pollen tube reception

phenotype [21], and Figure 4C and 4G). Domain swaps between different regions of NTA and

MLO1 revealed that the C-terminal cytoplasmic tail including the CaMBD of MLO1 (NTA-

MLO1[CTerm], see diagram in Figure 4D) was sufficient to direct the fusion protein to the filiform

apparatus of the synergids, in a pattern very similar to MLO1-GFP (Figure 4C). Quantification of

the GFP signal along the length of the synergids from the chalazal end to the filiform apparatus

in the NTA-GFP, NTA-MLO1[Cterm]-GFP, and MLO1-GFP confirmed that the MLO1 tail was

sufficient to cause NTA protein accumulation at the filiform apparatus (Figure 4A, 4C and 4D). In

MLO1-GFP and NTA-MLO1^{Cterm}-GFP ovules, the majority of GFP signal was detected in the 20-

40% of the micropylar pole of synergids and most of the signal overlapped with the diffuse FM4-

64 staining at the filiform apparatus (Figure 4F). In contrast, wild-type NTA-GFP is excluded

from the filiform apparatus (Figure 4A and [21].

The successful manipulation of NTA subcellular localization provided a tool for determining the

functional relevance of NTA redistribution. We transformed the NTA-MLO1^{CTerm}-GFP construct

into *nta-1* plants and used the percentage of unfertilized ovules as a measurement for the ability

of the fusion construct to complement the *nta-1* phenotype of unfertilized ovules caused by

pollen tube overgrowth [13]. Synergid-expression of NTA-MLO1^{CTerm}-GFP rescued the *nta-1*

pollen tube reception phenotype (Figure 4G), indicating that 1) the MLO1^{CTerm} domain is

functionally equivalent to the NTA^{CTerm} domain when the protein is localized at the filiform

apparatus and 2) premature targeting of NTA to the filiform apparatus is not toxic to synergid

cells.

Polar accumulation of NTA at the filiform apparatus is necessary for pollen tube

reception

Early targeting of NTA-MLO1^{CTerm}-GFP to the filiform apparatus was sufficient to complement

the *nta-1* pollen tube reception phenotype, but did not give an indication of whether filiform

apparatus targeting is necessary for NTA function. Since NTA accumulation is triggered by the

arrival of a growing pollen tube, drugs that block membrane protein trafficking are not

appropriate to inhibit NTA accumulation at the filiform apparatus because they would also inhibit

pollen tube growth. Therefore, we used a genetic approach to inhibit NTA trafficking to the

filiform apparatus by adding a Golgi retention signal (RNIKCD) to the C-terminus of our MLO-

GFP variants. This Golgi retention signal has been reported to cause Golgi-retention when

added to proteins that would normally be targeted to the trans-Golgi network and tonoplast [36].

To test if RNIKCD works on MLO proteins, we added RNIKCD to the C-terminus of MLO1

(MLO1-RNIKCD) and NTA (NTA-RNIKCD) and co-infiltrated tobacco leaves with these

constructs and the Golgi-mCherry marker (*35S::Man49-mCherry*). Both NTA and NTA-RNIKCD

co-localize with Golgi-mCherry marker in tobacco leaves (Figure S7A), as was previously

reported for NTA-GFP [21]. However, in contrast to MLO1-GFP which accumulates at the

plasma membrane and not in the Golgi, most of MLO1-RNIKCD was retained in the Golgi

(Figure S7A). We next added RNIKCD to MYB98$_{pro}$-driven NTA and NTA-MLO1^{CTerm} (NTA-

RNIKCD and NTA-MLO1^{CTerm}-RNIKCD) and transformed the constructs into *nta-1* and Golgi-

mCherry plants for functional and subcellular localization studies, respectively. In synergids of

emasculated ovules, both NTA-RNIKCD and NTA-MLO1^{CTerm}-RNIKCD fusion proteins co-

localized with the Golgi marker (Figure 4B, 4E and S7B), indicating that MLO fusion proteins

were mostly retained in the Golgi. Synergid-expressed NTA-RNIKCD and NTA-MLO1^{CTerm}-

RNIKCD could only partially rescue the *nta-1* pollen tube reception phenotype (Figure 4G),

indicating that filiform apparatus-localized NTA is both necessary and sufficient for pollen tube

reception.

Disrupting the NTA CaMBD compromises NTA's ability to accumulate at the filiform

apparatus during pollen tube reception

The dramatic differences in protein localization between NTA-GFP and NTA-MLO1^{CTerm}-GFP

revealed that the C-terminal cytoplasmic domain of NTA contains sequences that cause NTA to

be sequestered in the Golgi before pollen tube arrival and possibly also to allow for directed

transport to the filiform apparatus in response to signals from the arriving pollen tube. The C-

terminal cytoplasmic domains of MLO proteins contain a predicted CaMBD followed by an

unconserved tail of variable length [37-39]. A comparison of C-terminal domains (Figure 5A)

revealed that NTA and MLO1 share the conserved 20 amino acid 1-8-14 CaM binding motif of

hydrophobic residues interspersed with basic residues that define the MLO CaMBD [38], but

have different residues surrounding the conserved tryptophan that has been shown to

necessary for CaM binding in MLO proteins [38, 39]. In addition, NTA and MLO1 have different

length C-terminal tails after the CaMBD (Figure 5A). In order to determine if the C-terminal tail

after the CaMBD is necessary for NTA sequestration in the Golgi before pollen tube arrival, we

made a GFP fusion with NTA truncated just after the predicted CaMBD (NTA$^{\Delta481}$, Figure 5A)

expressed under control of the synergid-specific MYB98 promoter. This construct

complemented the *nta-1* pollen tube reception phenotype and localized in the Golgi before

pollen tube arrival (Figure 5C and 5E), revealing that the C-terminal tail after the CaMBD is not

necessary for NTA sequestration or function. We next tested whether disruption of the CaMBD

would affect NTA localization and/or function. A point mutation in the conserved tryptophan

necessary for CaM-binding in other CaMBDs [38, 40, 41] was introduced into our NTA-GFP

expression construct (NTA^{W458A}, Figure 5A) and transformed into the *nta-1* background. In

contrast to wild-type NTA and NTA$^{\Delta481}$, which result in an even higher fertility in *nta-1* mutants

than in the Ws controls, NTA^{W458A} only partially complemented the *nta-1* fertility phenotype

(Figure 5E). Before pollen tube arrival, NTA^{W458A} co-localizes with a Golgi marker in synergids

similar to wild-type NTA (Figure 5D and Figure S6). However, during pollen tube reception,

NTA^{W458A} displays different accumulation patterns that correlate with the ability of the pollen

tube to rupture. In our semi-*in vivo* system, ovules with normal pollen tube rupture had at least

partial NTA^{W458A} accumulation at the filiform apparatus, while ovules with pollen tube overgrowth

did not accumulate NTA^{W458A} at the filiform apparatus (Figure 6). These data reveal that an

intact CaMBD enhances NTA's redistribution to the filiform apparatus during pollen tube

reception and that NTA accumulation at the filiform apparatus promotes pollen tube rupture,

while the C-terminal tail after the CaMBD is dispensable for NTA function.

The influence of the CaMBD on NTA accumulation at the filiform apparatus could indicate a

connection with the calcium oscillations that occur in synergids during signaling with the arriving

pollen tube. The receptor-like kinase FER and the GPI-anchored protein LRE are filiform-

localized proteins that interact with each other and form a co-receptor for signals from the

arriving pollen tube [14, 16, 17]. *fer* and *lre* mutant ovules fail to initiate synergid calcium

oscillations at pollen tube arrival and have highly penetrant pollen tube overgrowth phenotypes

[12]. In *nta-1* mutant ovules, synergids have dampened calcium oscillations and around 30% of

ovules carrying the mutation display pollen tube overgrowth [12, 13]. In both *fer-1* and *lre-7*

backgrounds, NTA-GFP does not accumulate at the filiform apparatus in response to signals

from the arriving pollen tube (Figure S8 and [13]), placing NTA accumulation at the filiform

apparatus downstream of both FER and LRE and the calcium oscillations conditioned by these

proteins.

Discussion

Synergids respond to a signal from the approaching pollen tube

Successful pollination and production of seeds requires a series of signaling events between the

male gametophyte (pollen tube) and both sporophytic and gametophytic cells of the female. In

this study, we used live imaging to characterize dynamic subcellular changes that occur in the

synergid cells of the female gametophyte in response to the arrival of the pollen tube. We

showed that both the NTA protein and endosomes are actively mobilized to the filiform

apparatus where male-female communication occurs during pollen tube reception (Figure 7).

Disrupting the ability of NTA to pass through the secretory system by adding a Golgi retention

signal compromised its ability to participate in pollen tube reception, revealing that NTA

accumulation at the filiform apparatus is necessary for its function. Mutation of NTA's CaMBD

partially compromised NTA redistribution and function in pollen tube reception, revealing that

Ca^{2+} may play a role in the synergid response to the signal from the pollen tube.

The polar accumulation of the RabA1g endosome marker, but not a Golgi marker (Man49-GFP),

near the filiform apparatus during pollen tube reception suggests a change in synergid secretory

system behavior that is triggered by the approaching pollen tube. Our results with the ER, Golgi,

TGN, and peroxisome markers indicate that the mobilization of the RabA1g endosomes toward

the approaching pollen tube is not just a symptom of FER signaling triggering synergid cell

death that leads to mass disruption of subcellular compartments. Trans-Golgi Network/Early

endosomes (TGN/EEs) have been shown to be involved in the trafficking of both secretory and

endocytic cargo [42]. RabA1g is present in endosomes that are highly sensitive to Brefeldin A in

roots, suggesting that they could function as recycling endosomes [43]. While the resolution of

our live imaging system did not allow us to determine whether NTA completely co-localizes with

this compartment, it is tempting to speculate that the RabA1g endosomes are mediating the

polar movement of NTA to the filiform apparatus. Alternatively, these endosomes could be

transporting other signaling molecules either to or from the filiform apparatus.

Signal-mediated protein trafficking

Signal-mediated regulation of protein trafficking is an elegant mechanism to control the delivery

of molecules to the precise location where they are needed for critical signaling events that

occur over relatively short time frames. Selective protein targeting similar to NTA movement in

response to pollen tube arrival has also been observed during cell-to-cell communication

between the egg and sperm cells in Arabidopsis. After pollen tube reception and release of the

sperm cells, a signal from the sperm and/or the degenerated synergid cell causes the egg cell to

secrete EGG CELL 1 (EC1) peptides that have been stored in punctate compartments in the

egg cytoplasm toward the sperm cells. The sperm cells perceive the EC1 signal and, in turn,

mobilize the gamete fusogen HAPLESS2/GENERATIVE CELL SPECIFIC1 (HAP2/GCS1) from

a cytoplasmic compartment to the cell surface [44]. This controlled movement of proteins that

have already been translated and stored facilitates a quick response to activate the egg and

sperm for fertilization. Likewise, NTA mobilization to the filiform apparatus of the synergids as

the pollen tube arrives could play a role in sending a signal to the pollen tube that leads to the

mobilization of pollen tube proteins that allow the pollen tube to rupture and release the sperm

cells. For example, proteins that regulate the integrity of the tip of the pollen tube could be

quickly delivered after the "arrival" signal from the synergid is perceived. Recent work on the

role of the pollen-expressed ANXUR1 and 2 and BUDDHA PAPER SEAL1 and 2 receptor-like

kinases in pollen tube tip integrity support this hypothesis. During pollen tube growth through the

female tissues, RALF4 and RALF19 peptides that are secreted from pollen tubes act as ligands

for ANX1/2 and BPS1/2 to promote tip growth, while RALF34 secreted from the synergids

displaces RALF4 and 19 from the receptors leading to subcellular changes that result in pollen

tube rupture [45].

Calcium and NTA movement

Our study revealed that an intact CaMBD facilitates the trafficking of NTA from the Golgi to the

filiform apparatus in response to a stimulus from the approaching pollen tube. Our live imaging

studies and previous reports all suggest that the FER/LRE co-receptor is necessary for

perceiving this stimulus and that two other CrRLK1L genes, HERK1 and ANJEA are also

necessary for NTA accumulation at the filiform apparatus [13, 46]. Polar accumulation of NTA-

GFP at the filiform apparatus occurs in a similar time frame to the FER/LRE-dependent Ca^{2+}_{cyto}

spiking in the synergids during pollen tube reception [10, 12]. Subcellular Ca^{2+} spiking occurs in

plant responses to both biotic and abiotic external stimuli. Notably, cytosolic Ca^{2+} oscillations

occur during pollen tube-synergid interactions, egg-sperm interactions, and in biotrophic

interactions between plant cells and both beneficial and harmful microbes (reviewed in [47]). In

most cases, the mechanism for decoding Ca^{2+} spikes into a cellular response is not known, but

Ca^{2+}-binding proteins such as CaM and CaM-like proteins could play a role in relaying Ca^{2+}

signals [48]. In *nta-1* mutants, the cytosolic Ca^{2+} oscillations still occur, but at a lower magnitude

than in wild-type synergids, suggesting that NTA could be involved in modulating Ca^{2+} flux [12].

A recent study of the pollen tube-expressed MLO5 and MLO9 genes revealed a link between

MLO proteins and recruitment of the calcium channel CNGC18 to control Ca^{2+} flux during

directional growth [49]. Several CNGC genes are expressed in synergids and could possibly

interact with filiform apparatus-localized NTA to regulate the flow of Ca^{2+} ions into or out of the

apoplast near the filiform apparatus.

Calcium has also been linked to regulation of endomembrane trafficking (reviewed in [50]. In

animals, CaM plays a role in regulating vesicle tethering and fusion [51], and in plants CaM-like

proteins are associated with endosomal populations [52]. Thus, it is possible that the CaMBD in

NTA is critical for the precise targeting of NTA in response to pollen tube arrival. Another

mechanism that has been proposed for Ca^{2+} regulation of protein targeting is that electrostatic

interactions between Ca^{2+} and anionic phospholipids in specific domains of the plasma

membrane regulate vesicle fusion and differential recruitment of proteins to these domains [53,

54].

An interesting parallel between powdery mildew infection and pollen tube reception is that in

both cases, an MLO protein accumulates at the site of interaction with a tip-growing cell. During

powdery mildew infection, active transport of proteins and lipids to penetration site leads to

membrane remodeling and establishment of the extrahaustorial membrane (EHM) which

separates the plant cytoplasm from the invading fungal hyphae [55]. This specialized membrane

is enriched in the anionic phospholipid phosphatidylinositol 4,5 bisphosphate ($PI(4,5)P_2$).

Depletion of $PI(4,5)P_2$ in *phosphatidylinositol 4-phosphate 5-kinase 1/2* double mutants (*pip5k1*

pip5k2) resulted in powdery mildew resistance associated with a failure of MLO2 to accumulate

at the EHM in early stages of fungal penetration [56]. Although the composition of the filiform

apparatus and its relationship with the arriving pollen tube has not been established, it is

possible that similar membrane modification occurs in the filiform apparatus during signaling

with the pollen tube and is linked to NTA accumulation at pollen tube arrival.

Cell death and pollen tube reception

MLOs were first discovered in barley as powdery mildew resistance genes [57]. *mlo* mutants in

both monocots and dicots are resistant to powdery mildew infection, indicating that the MLO

proteins are required for infection. These mutants also have ectopic cell death, indicating that

one function of MLO proteins is to negatively regulate cell death [58]. Pollen tube reception is

catastrophic for both the pollen tube and the receptive synergid cell: both cells die as a result of

successful male-female signaling and delivery of the male gametes. The timing of synergid

degeneration remains under debate, with some studies suggesting that it occurs upon pollen

tube arrival at the synergid and others concluding that it occurs concurrently with pollen tube

rupture [31, 59, 60]. Our live imaging experiments with both NTA-GFP and the subcellular

compartment reporters suggest that the synergid cells are still alive and regulating their

secretory systems up to the point of pollen tube rupture. Given the function of the "powdery

mildew" members of the MLO gene family in regulating cell death [61], it is possible that one

role of NTA is to prevent early degeneration of the synergids. In both pollen tube reception and

powdery mildew infection, an MLO protein might be needed at the site of penetration to stabilize

the cell and prevent precocious cell death. Our result that premature delivery of NTA (in the

NTA-MLO1^{CTerm} domain swap construct) to the filiform apparatus of the cell does not disrupt

pollen tube reception is consistent with this hypothesis, since other signaling processes

occurring in the synergids during communication with the arriving pollen tube would likely not be

triggered by simply moving NTA to the filiform apparatus in the absence of a pollen tube.

In this study, we showed that signals from an approaching pollen tube trigger the movement of

NTA out of the Golgi and to the filiform apparatus and that this redistribution is necessary for

pollen tube reception. Future work will focus on determining the mechanism through which NTA

polarly accumulates at the filiform apparatus and on identifying the signals from the pollen tube

that lead to important subcellular changes in the synergids during pollen tube reception.

Materials and methods

Plant materials and growth conditions

Arabidopsis thaliana lines expressing *NTA$_{pro}$::NTA-GFP*, *MYB98$_{pro}$::NTA-GFP*, *MYB98$_{pro}$::MLO1-GFP*, Golgi-associated marker (Man49-mCherry), ER-associated marker (SP-mCherry-HDEL), TGN-associated marker (SYP61-mCherry), peroxisome-associated marker (Peroxisome-mCherry) and recycling endosome-associated marker (mCherry-RabA1g), were generated and described in previous publications [13, 14, 21, 33]. Pollen tube marker lines, *ACA9::DsRed* [62] and *Lat52::GFP* [28] were generously provided by Dr. Aurelien Boisson-Dernier and Dr. Ravi Palanivelu. *fer-1* seeds were provided by Dr. Ueli Grossniklaus and *lre-7* seeds were provided by Dr. Ravi Palanivelu. Seeds were sterilized and plated on ½-strength Murashige and Skoog (MS) plates. All plates were sealed and stratified at 4°C for two days, and then transferred to the growth chamber (long day conditions, 16 h of light and 8 h of dark at 22°C) for germination and growth. After 5-7 days, seedlings were transplanted to soil. Seeds from transformed lines were sterilized and plated on ½ MS plate with 20 mg/L hygromycin for selection of transgenic seedlings, which were then transplanted to soil and grown in long days.

Live Imaging of Pollination Using a Semi-*in vivo* Pollen Tube Guidance Assay

The semi-*in vivo* system of pollen tube reception was modified from [28]. Approximately 150 μL of pollen germination media (5 mM KCl, 1 mM MgSO$_4$, 0.01% (w/v) H$_3$BO$_3$, 5 mM CaCl$_2$, 20% sucrose, 1.5% agarose, and adjusted pH to 7.5 with KOH) was poured into a Glass Bottom Culture Petri Dish (MatTek Corporation, P35G-1.0-20-C) and spread out using a pipette. Pistils were emasculated and 2 d later were hand pollinated with *ACA9::DsRed* or *Lat52::GFP* pollen and returned to the growth chamber. Approximately 1 h after pollination, pistils were removed from plants and placed on double sided tape on a glass slide. Stigmas were cut using single-sided razor blade and placed on pollen germination media using forceps. 8-10 ovules were arranged around the cut style and the petri dish was returned to the growth chamber. After 4-6

h, pollen tubes grew through the stigma and style and emerged near the ovules. Imaging

commenced when the pollen tubes approached ovules. Time-lapse images were acquired at 5

min intervals by spinning disk confocal microscopy using an Andor Revolution XD platform with

a Yokogawa CSU-X1-A1 scanner unit mounted on an Olympus IX-83 microscope and a 20X/0.5

NA objective (Olympus). An Andor iXon Ultra 897BV EMCCD camera was used to capture GFP

fluorescence (488-nm excitation) and red fluorescent protein (dsRed or mCherry) fluorescence

(561-nm excitation).

For each experimental condition, at least 60 ovules were imaged over the time course from

pollen tube approaching the ovule to completion of pollen tube reception (pollen tube rupture to

release the sperm cells). Neighboring ovules that did not attract pollen tubes were imaged at the

same time and served as controls for photoxicity.

Confocal laser scanning microscopy (CLSM)

CLSM to examine MLO variant subcellular localization was performed on ovules dissected 2 d

after emasculation. FM4-64 staining was performed according to the protocol described in [21].

CLSM was done using either a Nikon A1Rsi inverted confocal microscope according to [63] or a

Leica SP8 upright confocal microscope according to [21].

Quantification of fluorescence signal intensity

Two-channel images were adjusted for brightness and contrast using Fiji [64]. Then, they were

input to NIS-Elements software (Ver. 5.02) to measure the fluorescence signal intensity. A line

that spanned from the chalazal end to the filiform apparatus end of the synergid was used for

the signal intensity measurements. For a more accurate representation of the total area of the

synergid, signal intensities were recorded along the same length line at five parallel position

within the synergid cell. Finally, all the measurement data were output as Excel files. Graphs

and statistical analysis were performed with Prism software (www.graphpad.com).

Video processing

Images were filtered to remove the noise and cropped using Fiji (Version 2.0.0). QuickTime

Player (Version 10.5) was used for movie editing and time-lapse analyses.

Cloning and Generation of Transgenic Lines

PCR amplification with PHUSION High-Fidelity Polymerase (NEB, M0535S) or Q5 High-Fidelity

DNA Polymerase (NEB, M0419S) were used to generate the following constructs in this study.

Genes were amplified with primers that had attB1 and attB2 sites for recombination via BP

reaction into the Gateway-compatible entry vector pDONR207. Full-length NTA entry vectors

used in this study was generated as described previously [21] . NTA truncation was amplified

using NTA full-length entry vector as a template with forward primer NTA-FattB1 and the

reverse primer NTA481-RattB2 (See all primer sequences in Table S1). The NTA^{W458A} point

mutation was generated using the same *NTA* template and amplifying two fragments of *NTA*

with desired point mutations introduced into the primers: NTA-FattB1 + NTAW458A-R and

NTAW458A-F + NTA-RattB2. The two fragments were purified and pasted together with

overlaps using a PCR-pasting protocol. The NTA-MLO1^{CTerm} construct was generated using the

full-length entry vectors of NTA and MLO1 used in previous study [21] as templates and

amplifying two fragments of *NTA* and *MLO1* using the two pairs of primers: NTA-FattB1 +NTA-

R19 and MLO1-F + MLO1-RattB2. The two fragments were purified and pasted together with

overlaps using a PCR-pasting protocol. The NTA-GFP-RNIKCD construct was generated using

the full-length expression vector of MYB98$_{pro}$::NTA-GFP as template. Two separate fragments

of NTA and GFP-RNIKCD-1 were amplified with two pairs of primers: NTA-FattB1 + NTA-Gly5-

GFP-R and NTA-Gly5-GFP-F + GFP-RNIKCD-RattB2. The two fragments were purified and

pasted together with overlaps using a PCR-pasting protocol. The NTA-MLO1^{CTerm}-GFP-RNIKCD

construct was generated using the full-length expression vector of MYB98$_{pro}$::NTA-MLO1^{CTerm}-

GFP as template and two fragments of NTA-MLO1^{CTerm} and GFP-RNIKCD-2 were amplified

using the two pairs of primers: NTA-FattB1 + MLO1-Gly5-GFP-R and MLO1-Gly5-GFP-F +

GFP-RNIKCD-RattB2. The two fragments were purified and pasted together with overlaps using

a PCR-pasting protocol. The coding sequence from the truncation and the point mutation and

the fusion sequences were fully sequenced in entry vectors. NTA$^{\Delta481}$ and NTA^{W458A} entries were

then recombined via LR reactions into the pMDC83 backbone with the MYB98$_{pro}$ [65] to drive

expression of each NTA variant in synergid cells with a C-terminal GFP fusion and NTA-GFP-

RNIKCD and NTA-MLO1^{CTerm}-GFP-RNIKCD entries were then recombined via LR reactions into

pMDC32 plasmids with the MYB98$_{pro}$ for synergid expression and FER$_{pro}$ for tobacco transient

expression.

For *nta-1* complementation assay and co-localization analyses, expression vectors were

transformed into the *Agrobacterium tumefaciens* strain GV3101 and transformed into the *nta-1*

mutant background or the Col-0 background (for co-localization combinations, Col-0 stably

expressing the Golgi or TGN synergid secretory markers was used for transformation) via the

floral-dip method [66]. Stable transgenic lines were selected using their respective selections

described above. Homozygous T2 or T3 lines were used in *nta-1* complementation assay and

co-localization imaging in the synergid was done in a T1 or T2 analysis. The NTA$_{pro}$::NTA-GFP

construct described in [13] was introduced into the *lre-7* background by floral dip as described

above.

***nta-1* Complementation Assays**

2-4 independent insertion lines for each construct were taken to the T2 generation and

screened for homozygosity using fluorescence microscopy to ensure transgene expression in

synergids of every ovule. Unfertilized vs. fertilized ovule counts from self-pollinated flowers were

assessed in at least five plants of each insertion line and compared to *nta-1*, Wassilewskija (Ws-

2; wildtype), and the previously published full-length NTA (MYB98$_{pro}$::NTA-GFP in *nta-1*

background, [21]. Ovule counts were statistically analyzed using Prism with significance

determined using a Student's *t*-test. Comparisons of the NTA variants were made to full-length

NTA and the *nta-1* mutant.

Acknowledgements

We thank Patrick Day for technical assistance and Rachel Flynn and Thomas Davis for helpful

discussions and comments on the manuscript. This work was supported by funds from NSF

IOS-1733865 to SAK, Purdue University Start-up funds to SAK, and a grant from the Oklahoma

Center for the Advancement of Science and Technology #PS14-008 to SAK. Spinning disk

confocal microscopy in the Staiger laboratory was supported, in part, by an award from the

Office of Science at the US Department of Energy, Physical Biosciences Program, under

contract number DE-FG02-09ER15526 to CJS.

Author Contributions

JY, YJ, DSJ, and SAK conceived and designed the experiments. JY, YJ, DSJ, NL, and WZ

performed the experiments. JY, YJ, DSJ, and SAK analyzed the data. JY, YJ, DSJ, and SAK

wrote the manuscript, and all authors revised and approved the final manuscript.

Competing Interests

The authors declare no competing interests.

Figure Legends

Figure 1. NTA-GFP accumulates at the filiform apparatus as the pollen tube approaches.

(A) Diagram of a mature *Arabidopsis thaliana* ovule and embryo sac, modified from Jones et al,

2018. CC, Central Cell; Syn, Synergid Cells; EC, Egg Cell; An, Antipodal cells; FA, Filiform

Apparatus (B, C) Live imaging of pollen tube (PT) reception using NTA-GFP labeled synergids

(green signal) and *ACA9::DsRed* pollen tubes (magenta signal). (B) Polar NTA-GFP

accumulation at the filiform apparatus (FA) occurred in ovules that attracted a pollen tube

(ovules with yellow stars), while polar NTA-GFP accumulation did not occur in ovules without

pollen tube attraction (ovules with white stars). (C) Time-lapse imaging of NTA-GFP

accumulation during pollen tube reception. (D) Quantification of NTA-GFP signal along the

length of the synergid from chalazal end (0µm) to FA end (33µm) at 0min, 25min, 50min, 75min,

and 100min timepoints, respectively. (E) 6 more examples of the quantification of NTA-GFP

signal along the length of synergids after pollen tube arrival. (F) Quantification of the percentage

of ovules with different NTA accumulation patterns under the same imaging conditions. Bars=50

µm (B and C).

Figure 2. The Golgi marker is randomly distributed throughout synergids during pollen

tube reception. (A) NTA-GFP (green signal) and Golgi-mCherry signals (magenta signal) are

evenly distributed along the length of the synergid and co-localized within synergid cells before

pollen tube arrival. (B) After pollen tube arrival, NTA-GFP accumulated at the FA, but Golgi-

mCherry did not accumulate at the FA. (C) Live imaging of Golgi-mCherry during reception of

Lat52::GFP labeled pollen tubes. (D) Quantification of Golgi-mCherry signal along the length of

synergids shown in (C) before and after pollen tube arrival. Bars=30µm (A-B), 50µm (C).

Figure 3. RabA1g endosomes polarly accumulate toward the filiform apparatus during pollen tube reception. (A) RabA1g-mCherry endosome marker (magenta signal) accumulates at the FA region in response to pollen tube arrival (ovule with yellow star). The ovule with no pollen tube attraction (white star) serves as a negative control imaged under the same conditions. (B) Higher magnification of the micropylar region of starred ovule in (red box in panel A). (C) Timing of RabA1g-mCherry polar accumulation during pollen tube arrival. Bars=50µm. (D) Quantification of RabA1g-mCherry signal along the length of synergids during pollen tube reception. Synergid cell from chalazal end to filiform apparatus (FA) end was defined from 0 to 33 µm in length. (E) 6 more examples of the quantification of RabA1g-mCherry signal after pollen tube arrival. (F) Quantification of ovule percentage with RabA1g endosome marker throughout the synergids (gray bars) or with polar accumulation at or near the filiform apparatus (black bars).

Figure 4. Targeting of NTA to the filiform apparatus before pollen tube arrival is not toxic to synergid cells. (A-E) Localization patterns of MYB98 promoter driven MLO-GFP variants (green signal) in synergids of mature virgin ovules stained with FM4-64 (magenta signal) to reveal the outline of the synergid and the filiform apparatus (FA, diffuse magenta signal). Graphs in each panel show quantification of the GFP intensity of the MLO variants along the length of the synergids. Bars = 10 µm. (F) Percentage of ovules showing MLO-GFP signal throughout the synergids (100% of length), in the FA only (20% of length) and the region surrounding and including the FA (40% of length). (F) Scatter plot of unfertilized ovule percentages in homozygous plants of pMYB98::MLO-GFP in *nta-1* mutants to assess the ability of the MLO-GFP constructs to complement *nta-1*. WS, Wassilewskija. Significance was determined by a Student's *t*-test (****, $P < 0.0001$; *, $P = 0.0281$; and ns, $P = 0.2020$).

Figure 5. Analysis of NTA variants expressed in synergids of *nta-1*. (A) Alignment of MLO1

and NTA C-terminal domains after the seventh membrane span. Red bars highlight the CaMBD;

W458A and Δ481 indicate the point mutation in NTA and the deletion point, respectively ; a.a.,

amino acid. (B-D) NTA (variant)-GFP (green) distribution in synergid cells of unfertilized ovules

stained with FM4-64 (magenta signal). Bar = 10 μm. (E) Complementation analysis of NTA

variants in T2 plants homozygous for MYB98$_{pro}$::NTA(variant)-GFP constructs in *nta-1* mutants.

Adjusted P values from a Student's *t*-test are as follows: **** indicates $P < 0.0001$; *** indicates

P = 0.001 to 0.0001; and ns indicates $P > 0.05$.

664
Figure 6. A point mutation in the CaMBD (NTA^{W458A}) affects filiform apparatus

accumulation of NTA and pollen tube reception. (A) NTA^{W458A}-GFP has 3 different

localization patterns in response to PT arrival under semi *in-vivo* conditions. Bar=50 μm (B)

Quantification of GFP signal intensity in NTA^{W458A} synergids during pollen tube reception. (C)

Analysis of NTA^{W458A}-GFP distribution patterns in ovules with successful (PT bursting) and

unsuccessful (no PT or PT overgrowth (PT OG)) pollen tube reception.

671
Figure 7. Subcellular dynamics in synergids during pollen tube reception. (A) Pollination

using a semi-*in vivo* pollen tube guidance assay. (B) Before pollen tube arrival, NTA is in a

Golgi-associated compartment and RabA1g endosomes are distributed throughout the

synergids. (C) As a pollen tube arrives, NTA and RabA1g endosomes move toward the filiform

apparatus (FA). NTA accumulation is dependent on signaling from the FER receptor like kinase,

which acts in a complex with LRE. Abbreviations: CC, Central Cell; Syn, Synergid cells; EC,

Egg Cell; An, Antipodal cells; nuc, Nucleus; FA, Filiform Apparatus.

679
680
Supplemental Materials

Supplementary figure 1. NTA-GFP does not polarly accumulate at the filiform apparatus in

synergids without pollen tube attraction.

Supplementary figure 2. Peroxisomes do not exhibit polar accumulation at the filiform

apparatus during pollen tube reception.

Supplementary figure 3. Trans-Golgi marker distribution in synergids does not change in

response to pollen tube reception.

Supplementary figure 4. ER marker distribution in synergids does not change in response to

pollen tube reception.

Supplementary figure 5. Additional examples of the polar accumulation of RabA1g endosomes

during pollen tube reception.

Supplementary figure 6. NTA^{W458A} co-localizes with Golgi marker in synergid cells before pollen

tube arrival.

Supplementary figure 7. A C-terminal Golgi retention signal retains MLO1 and NTA variants in

the Golgi.

Supplementary figure 8. NTA-GFP polar accumulation requires FER and LRE.

Supplemental Movies

Supplementary movie 1. NTA-GFP (green signal) accumulates at the filiform apparatus as

ACA9::DsRed labeled pollen tube (magenta signal) approaches.

Supplementary movie 2. NTA-GFP (green signal) accumulates at the filiform apparatus during

pollen tube reception (GFP channel only, same movie as S1).

Supplementary movie 3. Golgi-mCherry signals (magenta signal) are evenly distributed along

the length of the synergid as Lat52::GFP labeled pollen tube (green signal) approaches.

Supplementary movie 4. Golgi-mCherry signals (magenta signal) are evenly distributed along

the length of the synergid during pollen tube reception (mCherry channel only, same movie as

S3).

Supplementary movie 5. The trans-Golgi marker SYP61-mCherry (magenta signal) is localized

toward the micropylar region of synergid cells both before and after pollen tube (green signal)

arrival.

Supplementary movie 6. The trans-Golgi marker SYP61-mCherry (magenta signal) is localized

toward the micropylar region of synergid cells during pollen tube reception (mCherry channel

only, same movie as S5).

Supplementary movie 7. Before and after pollen tube (green signal) arrival, the ER marker SP-

mCherry-HDEL (magenta signal) is distributed throughout synergid cells.

Supplementary movie 8. The ER marker SP-mCherry-HDEL (magenta signal) is distributed

throughout synergid cells (mCherry channel only, same movie as S7).

Supplementary movie 9. The peroxisome marker mCherry-SLK (magenta signal) does not

accumulate at the filiform apparatus after pollen tube (green signal) arrival.

Supplementary movie 10. The peroxisome marker mCherry-SLK (magenta signal) does not

accumulate at the filiform apparatus during pollen tube reception (mCherry channel only, same

movie as S9).

Supplementary movie 11. RabA1g-mCherry endosome marker (magenta signal) accumulates

at the filiform apparatus in response to pollen tube (green signal) arrival.

Supplementary movie 12. RabA1g-mCherry endosome marker (magenta signal) accumulates

at the filiform apparatus in response to pollen tube (green signal) arrival (mCherry channel only,

same movie as S11).

References

732   1.   Naramoto, S. (2017). Polar transport in plants mediated by membrane transporters:
733   focus on mechanisms of polar auxin transport. Curr Opin Plant Biol *40*, 8-14.
734   2.   Petrasek, J., Mravec, J., Bouchard, R., Blakeslee, J.J., Abas, M., Seifertova, D.,
735   Wisniewska, J., Tadele, Z., Kubes, M., Covanova, M., et al. (2006). PIN proteins perform a
736   rate-limiting function in cellular auxin efflux. Science *312*, 914-918.
737   3.   Salanenka, Y., Verstraeten, I., Lofke, C., Tabata, K., Naramoto, S., Glanc, M., and Friml, J.
738   (2018). Gibberellin DELLA signaling targets the retromer complex to redirect protein
739   trafficking to the plasma membrane. Proc Natl Acad Sci U S A *115*, 3716-3721.
740   4.   Johnson, M.A., Harper, J.F., and Palanivelu, R. (2019). A Fruitful Journey: Pollen Tube
741   Navigation from Germination to Fertilization. Annu Rev Plant Biol.
742   5.   Drews, G.N., and Yadegari, R. (2002). Development and function of the angiosperm
743   female gametophyte. Annu Rev Genet *36*, 99-124.
744   6.   Takeuchi, H., and Higashiyama, T. (2016). Tip-localized receptors control pollen tube
745   growth and LURE sensing in Arabidopsis. Nature *531*, 245-248.
746   7.   Wang, T., Liang, L., Xue, Y., Jia, P.F., Chen, W., Zhang, M.X., Wang, Y.C., Li, H.J., and Yang,
747   W.C. (2016). A receptor heteromer mediates the male perception of female attractants
748   in plants. Nature *531*, 241-244.
749   8.   Okuda, S., Tsutsui, H., Shiina, K., Sprunck, S., Takeuchi, H., Yui, R., Kasahara, R.D.,
750   Hamamura, Y., Mizukami, A., Susaki, D., et al. (2009). Defensin-like polypeptide LUREs
751   are pollen tube attractants secreted from synergid cells. Nature *458*, 357-361.
752   9.   Kessler, S.A., and Grossniklaus, U. (2011). She's the boss: signaling in pollen tube
753   reception. Curr Opin Plant Biol *14*, 622-627.
754   10.   Denninger, P., Bleckmann, A., Lausser, A., Vogler, F., Ott, T., Ehrhardt, D.W., Frommer,
755   W.B., Sprunck, S., Dresselhaus, T., and Grossmann, G. (2014). Male-female
756   communication triggers calcium signatures during fertilization in Arabidopsis. Nature
757   communications *5*, 4645.
758   11.   Iwano, M., Ngo, Q.A., Entani, T., Shiba, H., Nagai, T., Miyawaki, A., Isogai, A.,
759   Grossniklaus, U., and Takayama, S. (2012). Cytoplasmic Ca2+ changes dynamically during
760   the interaction of the pollen tube with synergid cells. Development *139*, 4202-4209.
761   12.   Ngo, Q.A., Vogler, H., Lituiev, D.S., Nestorova, A., and Grossniklaus, U. (2014). A Calcium
762   Dialog Mediated by the FERONIA Signal Transduction Pathway Controls Plant Sperm
763   Delivery. Dev Cell *29*, 491-500.
764   13.   Kessler, S.A., Shimosato-Asano, H., Keinath, N.F., Wuest, S.E., Ingram, G., Panstruga, R.,
765   and Grossniklaus, U. (2010). Conserved molecular components for pollen tube reception
766   and fungal invasion. Science *330*, 968-971.
767   14.   Liu, X., Castro, C., Wang, Y., Noble, J., Ponvert, N., Bundy, M., Hoel, C., Shpak, E., and
768   Palanivelu, R. (2016). The Role of LORELEI in Pollen Tube Reception at the Interface of
769   the Synergid Cell and Pollen Tube Requires the Modified Eight-Cysteine Motif and the
770   Receptor-Like Kinase FERONIA. Plant Cell *28*, 1035-1052.
771   15.   Capron, A., Gourgues, M., Neiva, L.S., Faure, J.E., Berger, F., Pagnussat, G., Krishnan, A.,
772   Alvarez-Mejia, C., Vielle-Calzada, J.P., Lee, Y.R., et al. (2008). Maternal Control of Male-

Gamete Delivery in Arabidopsis Involves a Putative GPI-Anchored Protein Encoded by
the LORELEI Gene. Plant Cell *20*, 3038-3049.

16.   Li, C., Yeh, F.L., Cheung, A.Y., Duan, Q., Kita, D., Liu, M.C., Maman, J., Luu, E.J., Wu, B.W.,
Gates, L., et al. (2015). Glycosylphosphatidylinositol-anchored proteins as chaperones
and co-receptors for FERONIA receptor kinase signaling in Arabidopsis. Elife *4*.

17.   Escobar-Restrepo, J.M., Huck, N., Kessler, S., Gagliardini, V., Gheyselinck, J., Yang, W.C.,
and Grossniklaus, U. (2007). The FERONIA receptor-like kinase mediates male-female
interactions during pollen tube reception. Science *317*, 656-660.

18.   Devoto, A., Hartmann, H.A., Piffanelli, P., Elliott, C., Simmons, C., Taramino, G., Goh, C.S.,
Cohen, F.E., Emerson, B.C., Schulze-Lefert, P., et al. (2003). Molecular phylogeny and
evolution of the plant-specific seven-transmembrane MLO family. J Mol Evol *56*, 77-88.

19.   Kusch, S., Pesch, L., and Panstruga, R. (2016). Comprehensive Phylogenetic Analysis
Sheds Light on the Diversity and Origin of the MLO Family of Integral Membrane
Proteins. Genome Biol Evol *8*, 878-895.

20.   Yang, T., and Poovaiah, B. (2003). Calcium/calmodulin-mediated signal network in
plants. Trends in plant science *8*, 505-512.

21.   Jones, D.S., Yuan, J., Smith, B.E., Willoughby, A.C., Kumimoto, E.L., and Kessler, S.A.
(2017). MILDEW RESISTANCE LOCUS O Function in Pollen Tube Reception Is Linked to Its
Oligomerization and Subcellular Distribution. Plant Physiol *175*, 172-185.

22.   Lindner, H., Kessler, S.A., Müller, L.M., Shimosato-Asano, H., Boisson-Dernier, A., and
Grossniklaus, U. (2015). TURAN and EVAN mediate pollen tube reception in Arabidopsis
synergids through protein glycosylation. PLoS Biol *13*, e1002139.

23.   Huck, N., Moore, J.M., Federer, M., and Grossniklaus, U. (2003). The Arabidopsis mutant
feronia disrupts the female gametophytic control of pollen tube reception.
Development *130*, 2149-2159.

24.   Rotman, N., Rozier, F., Boavida, L., Dumas, C., Berger, F., and Faure, J.E. (2003). Female
control of male gamete delivery during fertilization in Arabidopsis thaliana. Curr Biol *13*,
432-436.

25.   Mansfield, S.G., Briarty, L.G., and Erni, S. (1991). Early embryogenesis in Arabidopsis
thaliana. I. The mature embryo sac. Canadian Journal of Botany *69*, 447-460.

26.   Huang, B.-Q., and Russell, S.D. (1992). Female Germ Unit: Organization, Isolation, and
Function. *140*, 233-293.

27.   Leshem, Y., Johnson, C., and Sundaresan, V. (2013). Pollen tube entry into the synergid
cell of Arabidopsis is observed at a site distinct from the filiform apparatus. Plant
reproduction *26*, 93-99.

28.   Palanivelu, R., and Preuss, D. (2006). Distinct short-range ovule signals attract or repel
Arabidopsis thaliana pollen tubes in vitro. BMC Plant Biol *6*, 7.

29.   Hamamura, Y., Nishimaki, M., Takeuchi, H., Geitmann, A., Kurihara, D., and Higashiyama,
T. (2014). Live imaging of calcium spikes during double fertilization in Arabidopsis. Nat
Commun *5*, 4722.

30.   Hamamura, Y., Nagahara, S., and Higashiyama, T. (2012). Double fertilization on the
move. Curr Opin Plant Biol *15*, 70-77.

31.   Hamamura, Y., Saito, C., Awai, C., Kurihara, D., Miyawaki, A., Nakagawa, T., Kanaoka,
M.M., Sasaki, N., Nakano, A., Berger, F., et al. (2011). Live-cell imaging reveals the

dynamics of two sperm cells during double fertilization in Arabidopsis thaliana. Curr Biol
21, 497-502.

32.   Huang, J., Ju, Y., Wang, X., Zhang, Q., and Sodmergen (2015). A one-step rectification of
sperm cell targeting ensures the success of double fertilization. J Integr Plant Biol *57*,
496-503.

33.   Jones, D.S., Liu, X., Willoughby, A.C., Smith, B.E., Palanivelu, R., and Kessler, S.A. (2018).
Cellular distribution of secretory pathway markers in the haploid synergid cells of
Arabidopsis thaliana. Plant J *94*, 192-202.

34.   Stoorvogel, W., Strous, G.J., Geuze, H.J., Oorschot, V., and Schwartz, A.L. (1991). Late
endosomes derive from early endosomes by maturation. Cell *65*, 417-427.

35.   Jones, D.S., and Kessler, S.A. (2017). Cell type-dependent localization of MLO proteins.
Plant Signal Behav *12*.

36.   Gao, C., Yu, C.K., Qu, S., San, M.W., Li, K.Y., Lo, S.W., and Jiang, L. (2012). The Golgi-
localized Arabidopsis endomembrane protein12 contains both endoplasmic reticulum
export and Golgi retention signals at its C terminus. Plant Cell *24*, 2086-2104.

37.   Devoto, A., Piffanelli, P., Nilsson, I., Wallin, E., Panstruga, R., von Heijne, G., and Schulze-
Lefert, P. (1999). Topology, subcellular localization, and sequence diversity of the Mlo
family in plants. J Biol Chem *274*, 34993-35004.

38.   Kim, M.C., Lee, S.H., Kim, J.K., Chun, H.J., Choi, M.S., Chung, W.S., Moon, B.C., Kang,
C.H., Park, C.Y., Yoo, J.H., et al. (2002). Mlo, a modulator of plant defense and cell death,
is a novel calmodulin-binding protein. Isolation and characterization of a rice Mlo
homologue. J Biol Chem *277*, 19304-19314.

39.   Kim, M.C., Panstruga, R., Elliott, C., Muller, J., Devoto, A., Yoon, H.W., Park, H.C., Cho,
M.J., and Schulze-Lefert, P. (2002). Calmodulin interacts with MLO protein to regulate
defence against mildew in barley. Nature *416*, 447-451.

40.   Arazi, T., Baum, G., Snedden, W.A., Shelp, B.J., and Fromm, H. (1995). Molecular and
Biochemical-Analysis of Calmodulin Interactions with the Calmodulin-Binding Domain of
Plant Glutamate-Decarboxylase. Plant Physiology *108*, 551-561.

41.   Yamada, M., Miyawaki, A., Saito, K., Nakajima, T., Yamamoto-Hino, M., Ryo, Y., Furuichi,
T., and Mikoshiba, K. (1995). The calmodulin-binding domain in the mouse type 1
inositol 1,4,5-trisphosphate receptor. Biochem J *308 (Pt 1)*, 83-88.

42.   Viotti, C., Bubeck, J., Stierhof, Y.D., Krebs, M., Langhans, M., van den Berg, W., van
Dongen, W., Richter, S., Geldner, N., Takano, J., et al. (2010). Endocytic and secretory
traffic in Arabidopsis merge in the trans-Golgi network/early endosome, an independent
and highly dynamic organelle. Plant Cell *22*, 1344-1357.

43.   Geldner, N., Denervaud-Tendon, V., Hyman, D.L., Mayer, U., Stierhof, Y.D., and Chory, J.
(2009). Rapid, combinatorial analysis of membrane compartments in intact plants with a
multicolor marker set. Plant J *59*, 169-178.

44.   Sprunck, S., Rademacher, S., Vogler, F., Gheyselinck, J., Grossniklaus, U., and
Dresselhaus, T. (2012). Egg cell-secreted EC1 triggers sperm cell activation during double
fertilization. Science *338*, 1093-1097.

45.   Ge, Z., Bergonci, T., Zhao, Y., Zou, Y., Du, S., Liu, M.C., Luo, X., Ruan, H., Garcia-Valencia,
L.E., Zhong, S., et al. (2017). Arabidopsis pollen tube integrity and sperm release are
regulated by RALF-mediated signaling. Science *358*, 1596-1600.

46.   Galindo-Trigo, S., Blanco-Tourinan, N., DeFalco, T.A., Wells, E.S., Gray, J.E., Zipfel, C., and
Smith, L.M. (2020). CrRLK1L receptor-like kinases HERK1 and ANJEA are female
determinants of pollen tube reception. EMBO Rep *21*, e48466.

47.   Chen, J., Gutjahr, C., Bleckmann, A., and Dresselhaus, T. (2015). Calcium signaling during
reproduction and biotrophic fungal interactions in plants. Mol Plant *8*, 595-611.

48.   Chin, D., and Means, A.R. (2000). Calmodulin: a prototypical calcium sensor. Trends Cell
Biol *10*, 322-328.

49.   Meng, J.G., Liang, L., Jia, P.F., Wang, Y.C., Li, H.J., and Yang, W.C. (2020). Integration of
ovular signals and exocytosis of a Ca(2+) channel by MLOs in pollen tube guidance. Nat
Plants *6*, 143-153.

50.   Himschoot, E., Pleskot, R., Van Damme, D., and Vanneste, S. (2017). The ins and outs of
Ca(2+) in plant endomembrane trafficking. Curr Opin Plant Biol *40*, 131-137.

51.   Burgoyne, R.D., and Clague, M.J. (2003). Calcium and calmodulin in membrane fusion.
Biochim Biophys Acta *1641*, 137-143.

52.   Ruge, H., Flosdorff, S., Ebersberger, I., Chigri, F., and Vothknecht, U.C. (2016). The
calmodulin-like proteins AtCML4 and AtCML5 are single-pass membrane proteins
targeted to the endomembrane system by an N-terminal signal anchor sequence. J Exp
Bot *67*, 3985-3996.

53.   Platre, M.P., Noack, L.C., Doumane, M., Bayle, V., Simon, M.L.A., Maneta-Peyret, L.,
Fouillen, L., Stanislas, T., Armengot, L., Pejchar, P., et al. (2018). A Combinatorial Lipid
Code Shapes the Electrostatic Landscape of Plant Endomembranes. Dev Cell *45*, 465-480
e411.

54.   Simon, M.L., Platre, M.P., Marques-Bueno, M.M., Armengot, L., Stanislas, T., Bayle, V.,
Caillaud, M.C., and Jaillais, Y. (2016). A PtdIns(4)P-driven electrostatic field controls cell
membrane identity and signalling in plants. Nat Plants *2*, 16089.

55.   Huckelhoven, R., and Panstruga, R. (2011). Cell biology of the plant-powdery mildew
interaction. Curr Opin Plant Biol *14*, 738-746.

56.   Qin, L., Zhou, Z., Li, Q., Zhai, C., Liu, L., Quilichini, T.D., Gao, P., Kessler, S.A., Jaillais, Y.,
Datla, R., et al. (2020). Specific Recruitment of Phosphoinositide Species to the Plant-
Pathogen Interfacial Membrane Underlies Arabidopsis Susceptibility to Fungal Infection.
Plant Cell.

57.   Buschges, R., Hollricher, K., Panstruga, R., Simons, G., Wolter, M., Frijters, A., van
Daelen, R., van der Lee, T., Diergaarde, P., Groenendijk, J., et al. (1997). The barley Mlo
gene: a novel control element of plant pathogen resistance. Cell *88*, 695-705.

58.   Panstruga, R. (2005). Serpentine plant MLO proteins as entry portals for powdery
mildew fungi. Biochem Soc Trans *33*, 389-392.

59.   Leydon, A.R., Tsukamoto, T., Dunatunga, D., Qin, Y., Johnson, M.A., and Palanivelu, R.
(2015). Pollen Tube Discharge Completes the Process of Synergid Degeneration That Is
Initiated by Pollen Tube-Synergid Interaction in Arabidopsis. Plant Physiol *169*, 485-496.

60.   Sandaklie-Nikolova, L., Palanivelu, R., King, E.J., Copenhaver, G.P., and Drews, G.N.
(2007). Synergid cell death in Arabidopsis is triggered following direct interaction with
the pollen tube. Plant Physiol *144*, 1753-1762.

61.   Piffanelli, P., Zhou, F., Casais, C., Orme, J., Jarosch, B., Schaffrath, U., Collins, N.C.,
904   Panstruga, R., and Schulze-Lefert, P. (2002). The barley MLO modulator of defense and
905   cell death is responsive to biotic and abiotic stress stimuli. Plant Physiol *129*, 1076-1085.
906   62.   Boisson-Dernier, A., Frietsch, S., Kim, T.H., Dizon, M.B., and Schroeder, J.I. (2008). The
907   peroxin loss-of-function mutation abstinence by mutual consent disrupts male-female
908   gametophyte recognition. Curr Biol *18*, 63-68.
909   63.   Yuan, J., and Kessler, S.A. (2019). A genome-wide association study reveals a novel
910   regulator of ovule number and fertility in Arabidopsis thaliana. PLoS Genet *15*,
911   e1007934.
912   64.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
913   Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
914   platform for biological-image analysis. Nat Methods *9*, 676-682.
915   65.   Muller, L.M., Lindner, H., Pires, N.D., Gagliardini, V., and Grossniklaus, U. (2016). A
916   subunit of the oligosaccharyltransferase complex is required for interspecific
917   gametophyte recognition in Arabidopsis. Nat Commun *7*, 10826.
918   66.   Bent, A. (2006). Arabidopsis thaliana Floral Dip Transformation Method. In Methods in
919   Molecular Biology, Volume 343, K. Wang, ed. (Totowa, NJ: Humana Press Inc.).
920

Figure 1. NTA-GFP accumulates at the filiform apparatus as the pollen tube approaches. (A) Diagram of a mature *Arabidopsis thaliana* ovule and embryo sac, modified from Jones et al, 2018. CC, Central Cell; Syn, Synergid Cells; EC, Egg Cell; An, Antipodal cells; FA, Filiform Apparatus (B, C) Live imaging of pollen tube (PT) reception using NTA-GFP labeled synergids (green signal) and ACA9::DsRed pollen tubes (magenta signal). (B) Polar NTA-GFP accumulation at the filiform apparatus (FA) occurred in ovules that attracted a pollen tube (ovules with yellow stars), while polar NTA-GFP accumulation did not occur in ovules without pollen tube attraction (ovules with white stars). (C) Time-lapse imaging of NTA-GFP accumulation during pollen tube reception. (D) Quantification of NTA-GFP signal along the length of the synergid from chalazal end (0μm) to FA end (33μm) at 0min, 25min, 50min, 75min, and 100min timepoints, respectively. (E) 6 more examples of the quantification of NTA-GFP signal along the length of synergids after pollen tube arrival. (F) Quantification of the percentage of ovules with different NTA accumulation patterns under the same imaging conditions. Bars=50 μm (B and C).

Figure 2. A Golgi marker is randomly distributed throughout synergids during pollen tube reception. (A) NTA-GFP (green signal) and Golgi-mCherry signals (magenta signal) are evenly distributed along the length of the synergid and co-localized within synergid cells before pollen tube arrival. (B) After pollen tube arrival, NTA-GFP accumulated at the FA, but Golgi-mCherry did not accumulate at the FA. (C) Live imaging of Golgi-mCherry during reception of Lat52::GFP labeled pollen tubes. (D) Quantification of Golgi-mCherry signal along the length of synergids shown in (C) before and after pollen tube arrival. Bars=30μm (A-B), 50μm (C).

Figure 3. RabA1g endosomes polarly accumulate toward the filiform apparatus during pollen tube reception. (A) RabA1g-mCherry endosome marker (magenta signal) accumulates at the FA region in response to pollen tube arrival (ovule with yellow star). The ovule with no pollen tube attraction (white star) serves as a negative control imaged under the same conditions. (B) Higher magnification of the micropylar region of starred ovule in (red box in panel A). (C) Timing of RabA1g-mCherry polar accumulation during pollen tube arrival. Bars=50μm. (D) Quantification of RabA1g-mCherry signal along the length of synergids during pollen tube reception. Synergid cell from chalazal end to filiform apparatus (FA) end was defined from 0 to 33 μm in length. (E) 6 more examples of the quantification of RabA1g-mCherry signal after pollen tube arrival. (F) Quantification of ovule percentage with RabA1g endosome marker throughout the synergids (gray bars) or with polar accumulation at or near the filiform apparatus (black bars).

Figure 4. Targeting of NTA to the filiform apparatus before pollen tube arrival is not toxic to synergid cells. (A-E) Localization patterns of MYB98 promoter driven MLO-GFP variants (green signal) in synergids of mature virgin ovules stained with FM4-64 (magenta signal) to reveal the outline of the synergid and the filiform apparatus (FA, diffuse magenta signal). Graphs in each panel show quantification of the GFP intensity of the MLO variants along the length of the synergids. Bars = 10 µm. (F) Percentage of ovules showing MLO-GFP signal throughout the synergids (100% of length), in the FA only (20% of length) and the region surrounding and including the FA (40% of length). (F) Scatter plot of unfertilized ovule percentages in homozygous plants of pMYB98::MLO-GFP in *nta-1* mutants to assess the ability of the MLO-GFP constructs to

complement *nta-1*. WS, Wassilewskija. Significance was determined by a Student's *t*-test (****, *P* < 0.0001; *, *P* = 0.0281; and ns, *P* = 0.2020).

Figure 5. Disruption of the CaMBD affects NTA localization and function. (A) Alignment of MLO1 and NTA C-terminal domains after the seventh membrane span. Red bars highlight the CaMBD; W458A and Δ481 indicate the point mutation in NTA and the deletion point, respectively ; a.a., amino acid. (B-D) NTA (variant)-GFP (green) distribution in synergid cells of unfertilized ovules stained with FM4-64 (magenta signal). Bar = 10 μm. (E) Complementation analysis of NTA variants in T2 plants homozygous for MYB98$_{pro}$::NTA(variant)-GFP constructs in *nta-1* mutants. Adjusted P values from a Student's *t*-test are as follows: **** indicates *P* < 0.0001; *** indicates *P* = 0.001 to 0.0001; and ns indicates *P* > 0.05.

Figure 6. A point mutation in the CaMBD (NTA^{W458A}) affects filiform apparatus accumulation of NTA and pollen tube reception. (A) NTA^{W458A}-GFP has 3 different localization patterns in response to PT arrival under semi *in-vivo* conditions. Bar=50 μm (B) Quantification of GFP signal intensity in NTA^{W458A} synergids during pollen tube reception. (C) Analysis of NTA^{W458A}-GFP distribution patterns in ovules with successful (PT bursting) and unsuccessful (no PT or PT overgrowth (PT OG)) pollen tube reception.

Figure 7. Subcellular dynamics in synergids during pollen tube reception. (A) Pollination using a semi-*in vivo* pollen tube guidance assay. (B) Before pollen tube arrival, NTA is in a Golgi-associated compartment and RabA1g endosomes are distributed throughout the synergids. (C) As a pollen tube arrives, NTA and RabA1g endosomes move toward the filiform apparatus (FA). NTA accumulation is dependent on signaling from the FER receptor like kinase, which acts in a complex with LRE. Abbreviations: CC, Central Cell; Syn, Synergid cells; EC, Egg Cell; An, Antipodal cells; nuc, Nucleus; FA, Filiform Apparatus.

A
+0min +25min +50min +75min +100min

B

NTA-GFP intensity

+0min +25min +50min +75min +100min

Chalazal end *FA end*

Synergid cell length

Supplementary figure 1. NTA-GFP does not polarly accumulate at the filiform apparatus in synergids without pollen tube attraction. (A) NTA-GFP (green signal) distribution pattern in synergids at 0min, 25min, 50min, 75min, and 100min timepoints, respectively. (B)Signal intensity measurement of NTA-GFP intensity at 0min, 25min, 50min, 75min, and 100min timepoints, respectively. Bar=50µm.

Supplementary figure 2. Peroxisomes do not exhibit polar accumulation at the FA during pollen tube reception. (A) The peroxisome marker mCherry-SLK (magenta signal) does not accumulate to the FA region after pollen tube (green signal) arrival. (B) Quantification of mCherry-SLK signal before and after PT arrival in the ovule with the white star. Bars=50μm.

Supplementary figure 3. Trans-Golgi marker distribution in synergids does not change in response to pollen tube reception. (A) The trans-Golgi marker SYP61-mCherry (magenta signal) is concentrated toward the micropyle region of synergid cells both before and after pollen tube (green signal) arrival (ovules with yellow stars). (B) Quantification of SYP61-mCherry signal along the length of a synergid before and after pollen tube reception. Bars=50μm.

Supplementary figure 4. ER marker distribution in synergids does not change in response to pollen tube reception. (A) Before and after pollen tube (green signal) arrival, the ER marker SP-mCherry-HDEL (magenta signal) is distributed throughout synergid cells. (B) Quantification of SP-mCherry-HDEL signal along the length of synergids in ovule with white star before and after pollen tube reception. Bars=50μm.

Supplementary figure 5. **Additional examples of the polar accumulation of RabA1g endosomes during pollen tube reception.** Endosome marker accumulates near the filiform apparatus region as pollen tube approaches (ovules with yellow stars), but no polar accumulation was found in ovules that did not attract a pollen tube (ovules with white stars). Bars=50μm.

Supplementary figure 6. NTA^{W458A} co-localizes with Golgi marker in synergid cells before pollen tube arrival. Colocalization of NTA^{W458A}-GFP (green) with Golgi marker (*LRE::proMan49-mCherry*, magenta) (A) and the trans-Golgi network marker (TGN, *MYB98pro::SYP61:mCherry*, magenta) (B) in the synergid cell of unpollinated ovules. Bars = 10 μm.

Supplementary figure 7. **A C-terminal Golgi retention signal retains MLO1 and NTA variants in the Golgi.** (A) Colocalization of MLO-GFP (green) with the Golgi-mCherry marker (magenta) indicates that the Golgi retention signal RNIKCD partially retains MLO1 in the Golgi in transiently-transformed tobacco epidermis, Bars = 20 μm. (B) Colocalization of MLO-RNIKCD (green) with the Golgi-mCherry marker (magenta) indicates that the Golgi retention signal RNIKCD retains NTA-MLO1^{CTerm} in the Golgi in the synergid. Bars = 10 μm.

Supplementary figure 8. **NTA-GFP polar accumulation requires FER and LRE.** (A-B) In *fer* mutant synergids, NTA-GFP (green signal) distributes throughout synergids (arrows) with ACA9::DsRed pollen tube overgrowth (magenta signal). (C) Quantification of NTA-GFP signal along the length of *fer-1* synergids before and after pollen tube reception. (D-E) In *lre-7* ovules, NTA-GFP distributes throughout synergids (white arrow) with pollen tube overgrowth. (F) Quantification of NTA-GFP signal along the length of *lre-7* synergids before and after pollen tube reception. FA, Filiform Apparatus. Bars=50µm.

Supplementary table 1. List of primers used for cloning.

Primer Name	Primer Sequences (5' to 3')
NTA-FattB1	GGGGACAAGTTTGTACAAAAAAGCAGGCTTCACCATGATCACAAGAAGCAGGTGT
NTA481-RattB2	GGGGACCACTTTGTACAAGAAAGCTGGGTGTGAAACACCAAGTGTCTTGCT
NTAW458A-R	C**GC**CTTCTTCAATGCCTTTGCCAT
NTAW458A-F	GCATTGAAGAAG**GC**GCACAAAGACATCAAATTGAAGAAAG
NTA-RattB2	GGGGACCACTTTGTACAAGAAAGCTGGGTGAGAGTTGTGGAATTGCATCTC
NTA-R19	CTCATCAAACACTGCTTTCTTCATG
MLO1-F	GCAGTGTTTGATGAGAATGTGCAGGTTGGTCTTGTTG
MLO1-RattB2	GGGGACCACTTTGTACAAGAAAGCTGGGTGGTTGTTATGATCAGGTGTAATCTCA
NTA-Gly5-GFP-R	TTCTCCTTTACTCATTCCTCCTCCTCCACCAGAGTTGTGGAATTG
NTA-Gly5-GFP-F	CAATTCCACAACTCTGGTGGAGGAGGAGGAATGAGTAAAGGAGAA
GFP-RNIKCD-RattB2	GGGGACCACTTTGTACAAGAAAGCTGGGTGCTAGTCGCACTTGATGTTTCTTTTGTATAGTTC
MLO1-Gly5-GFP-R	TTCTCCTTTACTCATTCCTCCTCCTCCACCGTTGTTATGATCAGG
MLO1-Gly5-GFP-F	CCTGATCATAACAACGGTGGAGGAGGAGGAATGAGTAAAGGAGAA

Supplemental Movies

Supplementary movie 1. NTA-GFP (green signal) redistributes to filiform apparatus region as ACA9::DsRed labeled pollen tube (magenta signal) approaches.

Supplementary movie 2. NTA-GFP (green signal) redistributes to filiform apparatus region during pollen tube reception (GFP channel only, same movie as S1).

Movie S1

Movie S2

Supplementary movie 3. Golgi-mCherry signals (magenta signal) are evenly distributed along the length of the synergid as Lat52::GFP labeled pollen tube (green signal) approaches.

Supplementary movie 4. Golgi-mCherry signals (magenta signal) are evenly distributed along the length of the synergid during pollen tube reception (mCherry channel only, same movie as S3).

Movie S3

Movie S4

Supplementary movie 5. The trans-Golgi marker SYP61-mCherry (magenta signal) is localized toward the micropyle region of synergid cells both before and after pollen tube (green signal) arrival.

Supplementary movie 6. The trans-Golgi marker SYP61-mCherry (magenta signal) is localized toward the micropyle region of synergid cells during pollen tube reception (mCherry channel only, same movie as S5).

Movie S5   Movie S6

Supplementary movie 7. Before and after pollen tube (green signal) arrival, the ER marker SP-mCherry-HDEL (magenta signal) is distributed throughout synergid cells.

Supplementary movie 8. The ER marker SP-mCherry-HDEL (magenta signal) is distributed throughout synergid cells (mCherry channel only, same movie as S7).

Movie S7   Movie S8

Supplementary movie 9. The peroxisome marker mCherry-SLK (magenta signal) does not redistribute to the filiform apparatus region after pollen tube (green signal) arrival.

Supplementary movie 10. The peroxisome marker mCherry-SLK (magenta signal) does not redistribute to the filiform apparatus region during pollen tube reception (mCherry channel only, same movie as S9).

Movie S9   Movie S10

Supplementary movie 11. RabA1g-mCherry endosome marker (magenta signal) accumulates at the filiform apparatus region in response to pollen tube (green signal) arrival.

Supplementary movie 12. RabA1g-mCherry endosome marker (magenta signal) accumulates at the filiform apparatus region during pollen tube reception (mCherry channel only, same movie as S11).

Movie S11   Movie S12