Fisheries-induced evolution of the circadian system and collective personality traits

Valerio Sbragaglia[1,†], Jose Fernando López-Olmeda[2], Elena Frigato[3], Cristiano

Bertolucci[3], Robert Arlinghaus[1,4]

[1] Department of Biology and Ecology of Fishes, Leibniz-Institute of Freshwater Ecology and Inland
Fisheries, Berlin, Germany
[2] Department of Physiology, Faculty of Biology, Regional Campus of International Excellence "Campus
Mare Nostrum", University of Murcia, Murcia, Spain
[3] Department of Life Sciences and Biotechnology, University of Ferrara, Ferrara, Italy
[4] Division of Integrative Fisheries Management, Faculty of Life Sciences & Integrative Research Institute
on Transformations of Human-Environment Systems (IRI THESys), Humboldt-Universität zu Berlin, Berlin,
Germany
[†] Corresponding authors: valeriosbra@gmail.com

ABSTRACT

The circadian system is phylogenetically conserved across a wide range of taxa. It

orchestrates organismal biological processes and is linked to life history and risk-taking

behavior. Fisheries-induced evolution affects life history and may also alter behavioral

traits in exploited populations. Thus, intensive and selective harvesting may lead to the

evolution of the circadian system and thereby affect temporal organization of

physiological processes. We present experimental results based on zebrafish (*Danio*

rerio) selection lines exposed to either large or small size-selective mortality relative to

a control, simulating evolutionary responses to intensive fishing. We show that size-

selective harvest leads to evolution of life-history, risk-taking and shoaling behaviors,

and importantly, also shifts the molecular circadian clockwork. Evolutionary effects of

size-selective harvesting on risk-taking and shoaling were linked to daily activity

rhythms in the small, but not in the large, size-selective mortality scenario. Both small

and large size-selective mortality induced an evolutionary shift of the molecular

circadian clockwork in the same direction in the brain and liver. Interestingly, the shifts

in the molecular circadian clockwork disappeared in the clock output pathway,

resulting in similar transcription profile among size-selective scenarios. We reveal that

size-selective harvesting not only alters life-histories, but also leaves an evolutionary

legacy in the behavioral and physiological machinery of exploited fish populations.

These changes can affect catchability, yield, recovery and the way exploited fish

population use space and time and maybe difficult to be reversed even when

harvesting is halted.

KEYWORDS: boldness, shoaling, collective, self-feeding, life history, gene expression

Significance statement:

Harvesting constitutes a strong driver of life history evolution in wild populations, but

there is limited knowledge whether harvesting also leads to adaptive changes in the

underlying physiology and behavior of exploited populations. We use a multi-

generation zebrafish (*Danio rerio*) harvesting experiment simulating two different

fisheries scenarios and we show that size-selective mortality affect the circadian

system and collective fish personality traits. Such adaptations were present eight

generations after size-selective harvesting was stopped, indicating induced evolution

of the circadian system and behavioural traits in response to size-selective harvesting.

INTRODUCTION

Harvesting drives the evolution of animal populations on contemporary time scales by creating selection pressures that substantially depart from natural conditions (1-3). Unselective, yet elevated harvesting-induced mortality alone triggers the evolution of a fast life history characterized by elevated reproductive investment, reduced age and size at maturation, reduced post maturation growth and longevity (4, 5). The evolution of a fast life history is reinforced by harvesting mortality being size-selective (6). Life history evolution can alter correlated traits, such as physiological and behavioral traits (6-9). However, physiological and behavioral adaptations in response to harvesting are much less understood compared to harvesting-induced changes to life-histories (2, 8). Our study fills this gap using an experimental system simulating fisheries-induced evolution in response to size-selective harvest.

Fisheries-induced evolution can affect physiological and behavioral traits through at least two key mechanisms. The first relates to correlated selection responses of adaptive changes in life-histories (10). For example, the pace-of-life syndrome proposes that life-history traits, such as growth rate or maturation size, are systematically and genetically related to behavioural and physiological traits (11). Accordingly, life-history adaptations caused by fisheries can also be expected to indirectly affect the physiology and behavior of exploited fishes (5, 6, 9). Secondly, fisheries can also directly select on physiological and behavioural traits, e.g., because bold, aggressive, social or stress-resilient fishes are preferentially harvested with certain gears (12-16). Theory (6, 9) and experimental harvesting with model fish species (17-19) have indeed shown that both elevated and size-selective harvesting typical of most real fisheries foster evolutionary adaptation of behavioral traits. What is largely unknown is the underpinning mechanisms linking life history, physiology and behavior in response to harvesting (8, 20) and how such mechanisms contribute to shape the evolutionary trajectories in a multivariate selection environment (21).

One candidate pathway linking life-history, behavior and physiology is the circadian system. The circadian system is widely conserved phylogenetically (22) and controls daily rhythms of physiology and behavior (23). Thereby, the circadian clock fulfills an adaptive function helping the organism to anticipate and maintain synchronization to external 24-h environmental cycles (23-26). Functional circadian

clocks are essential for precise timing of activity, thereby helping the organism to avoid

or minimize risk of predation while maximizing foraging (27). Recent studies reported

an insofar unknown relationship of animal personality, particularly an animal's

tendency to take risks, and the circadian system (28, 29) suggesting genetic

correlations among traits. Zebrafish (*Danio rerio*) of different coping styles, commonly

referred to as proactive/reactive behavioural phenotypes (30), were found to be

systematically linked to different properties of circadian rhythms (29). Moreover, life

history changes correlate with the circadian clock (e.g., 31, 32). Accordingly,

evolutionary adaptations of the circadian system in exploited populations may occur in

response to both behavioral or life history adaptations. Here, we report a study

documenting evidence for evolutionary shifts in the circadian system in response to

intensive size-selective harvesting using a model system based on zebrafish size-

selected lines.

The proximate mechanisms governing fisheries-induced evolution are difficult

to disentangle in the wild because phenotypic changes are also affected by plasticity

(2). A powerful means to single out the cause-and-effect of selective harvesting is

conducting selection experiments with model fish species in the laboratory (33). We

use a long-term harvesting experimental system based on zebrafish selection lines

exposed to strong directional harvest selection (a 75% per-generation harvest rate)

acting on either large or small fish, relative to a control line harvested randomly with

respect to body size (17). Zebrafish were selected in a social environment over five

generations. Afterwards, selection was halted for eight further generations to remove

all "maternal" effects (i.e. a summary term for all non-genetic ecological effects on the

phenotype of the offspring), allowing clean comparisons among the lines at F_{13} in

terms of long-term outcomes of harvesting-induced evolution on behavioural traits

and the circadian system. The experimental zebrafish lines that we used in the present

study maintained key life-history adaptations until F_{13} that were already induced at F_9

(17; see also Fig. S1), indicating that harvesting-induced evolution of life-history did not

recover to pre-harvesting levels.

Previous results using the zebrafish model system have shown that the

selection lines differ evolutionarily in reproductive allocation, adult body size, risk-

taking behavior, trait variability, allele composition and transcriptomic profiles (17, 19,

34, 35). Moreover, size-selection affected single nucleotide polymorphisms occurring

within regulatory areas of genes related to the circadian system (17), in particular

guca1c that is involved in the light reception cascade in the retina and pineal gland (36)

and *tph2* that is involved in serotonin synthesis (37). These data collectively provided

strong evidence that size-selection over just five generations affected both genes and

phenotypes, suggesting that size-selection could also have left a molecular legacy in

the circadian system. In the present work, we test the hypothesis that risk taking

behavior and the circadian system share common pathways (29) and we predicted that

harvesting-induced evolution of risk-taking behavior has altered the circadian system

in the zebrafish selection lines. We also expected that possible changes in shoaling

behavior (i.e., sociability; 19) would affect the social synchronization of individual

circadian clocks (Bloch et al., 2013), with consequent changes in the collective

circadian output at the group level. Finally, we expected that changes of daily

behavioral rhythms would be mirrored by the underlying circadian molecular

clockwork in the central (i.e., brain; possibly associated with the control of swimming

activity) and peripheral (i.e., liver; possibly associated with the control of feeding

activity) oscillators, and by neuroendocrinological molecular markers related to energy

balance, appetite and growth.

RESULTS

Shoaling and risk-taking behavior

We first studied zebrafish shoaling behavior in a round experimental arena (see

Methods for details). We found significant ($p < 0.001$) repeatability of inter-individual

distance among shoal members (R = 0.70 [confidence interval: 0.49 – 0.85]) over time

(i.e., 150 and 190 days post fertilization, dpf; Fig. 1A). Moreover, we found that the

small-harvested line formed significantly (t_{71} = - 3.52; $p < 0.001$) more cohesive (i.e.,

smaller inter-individual distance) shoals than controls, yet no significant (t_{71} = 0.90; p =

0.373) differences existed among the large-harvested line and the controls (Fig. 1B).

Next, we studied zebrafish risk-taking behavior using a diving test in groups (see

Methods for details). We found significant ($p < 0.001$) repeatability for the time spent

by the shoals feeding at the surface (R = 0.62 [confidence interval: 0.39 – 0.79]) over

time (i.e., 230 and 240 dpf; Fig. 1C). The small-harvested line spent significantly (p <

0.01) more time feeding at the surface (i.e., a risky environment) than the control,

while the large-harvested line spent significantly ($p < 0.05$) less time than the control at

the risky surface (Fig. 1D). Hence, the large-harvested line was found to be significantly

shyer, while the small-harvested line was bolder than the control.

Daily and circadian activity rhythms

We recorded robust daily activity rhythms of swimming (Fig. 2A, B) and self-feeding

(Fig. 2C, D) behaviour in a separate group of zebrafish between 170 and 190 dpf (see

Methods for details; Fig. S2, S3). In all selection lines, the behavioral rhythms revealed

significant 24-hour periodicities during both light-dark (LD) trials (first LD trial, days 1-

15; second LD trial, days 22-36) and significant free-running rhythmicity under

constant dim light conditions, with periods close to 24-h (Fig. S3, Table 1, see

Supplement for details).

All the three zebrafish lines displayed the typical diurnal swimming activity

rhythms (Fig. 2A, B). The total swimming activity during the photophase was

significantly ($p < 0.001$) repeatable over the two LD trials (R = 0.78 [confidence interval:

0.53 − 0.92]; Fig. 3A), with the small-harvested line being significantly ($t_{30} = 3.52$; $p <$

0.01) more active than controls (Fig. 3A). The selection lines were also significantly ($p <$

0.05) repeatable in their early peaks of swimming activity (R = 0.58 [confidence

interval: 0.19 − 0.84]). In particular, the small-harvested line concentrated significantly

($t_{30} = 2.96$; $p < 0.01$) more of their daily activity during the first four hours of the

photophase relative to controls (Fig. 3B). By contrast, the late peaks of swimming

activity were not significantly ($p = 0.123$) repeatable traits (Fig. 3C), therefore this

behaviour was not considered as a stable behavioral trait and was thus not compared

statistically among the selection lines.

All lines displayed self-feeding activity rhythms during the last hours of the

scotophase, immediately prior light-on (Fig. 2C, D). The self-feeding activity was

significantly ($p < 0.05$) repeatable (R ranged from 0.45 to 0.52) over the two LD trials in

all the four consecutive hourly sections that were analyzed (from three hours before to

one hour after light on, Fig. 4). At the onset of the self-feeding events (06:00 - 08:00 h;

darkness) both size-selected lines were significantly ($p < 0.01$) more active than

controls (Fig. 4A, B). Subsequently, during the hour before light-on (08:00 - 09:00 h),

and the first hour after light-on (09:00 - 10:00 h), the small-harvested line, but not the

large-harvested line, was more active in demanding food than controls (Fig. 4C, D). To

obtain a more quantitative measure of feeding behavior, we measured the amount of

food demanded every two days normalized by the biomass of each fish group during

the first days of the first LD trial. Significant ($F_{2,33}$ = 5.161, p < 0.05) differences among

lines during the first two days were found. Specifically, the large-harvested fish

demanded more food per fish mass (0.80 ± 0.58 g/g) than controls (0.18 ± 0.36 g/g),

but these differences disappeared over time (Fig. S4).

Core clock genes

At the end of the behavioral activity trials we sacrificed the fish, under constant-

lighting conditions, to measure circadian transcript abundance with RT-qPCR (see

Methods for details). Core clock genes expression revealed significant differences in

their circadian oscillations among the zebrafish lines (Table 2). In the brain, the control

line showed a strong trend (p = 0.052) in the circadian rhythmicity of *arntl1a* and a

clear circadian oscillation of *clock1a* with a peak at the same Circadian Time (CT) of

arntl1a (CT0 for both genes; Table 2). By contrast, the small-harvested line lost

circadian rhythmicity of *arntl1a* and *clock1a* with respect to the control, while the

large-harvested line shifted its peaks of circadian expression for both genes (CT16

instead of CT0, Table 2 and Fig. 5). Results of *per1b* indicated significant circadian

patterns in all lines with main differences in peaks in both size-selected lines that

shifted from CT8 to CT12 relative to the control (Fig. 5, Table 2). In the liver, the small-

harvested fish showed similar circadian rhythmicity to control fish in terms of *arntl1a*,

while the large-harvested line revealed only a trend (p = 0.086) in the circadian

oscillation (Table 2). The control line did not show any circadian rhythmicity in terms of

clock1a and *per1b*, however, the size-selected lines revealed clear circadian

rhythmicity for these two genes with slightly different peaks (Table 2 and Fig. 5).

Light-inducible genes

In the brain, the control line showed no significant rhythmicity of *per2*, while the size-

selected lines oscillated with peaks at different circadian times (CT12 and CT8 for large-

and small-harvested lines, respectively; Table 2, Fig. 5). Results of *cry1a* indicated

significant circadian rhythmicity with different peaks in the size-selected lines relative to the control: the large-harvested line shifted the peak from CT8 to CT12, while the small-harvested line shifted from CT8 to CT16 (Table 2 and Fig. 5). In the liver, both size-selected lines lost circadian rhythmicity of *per2* with respect to the control. The small-harvested line lost rhythmicity for *cry1a*, while the large-harvested line showed the same peak compared to the control (Table 2 and Fig. 5).

Clock output genes

The circadian expression of clock output genes showed less peak shifts compared to the other two groups of genes that we investigated (Table 2). In the brain, all lines had a circadian rhythmicity of *tef1* and *dbpa* with peaks at the same circadian time. However, the large-harvested line showed a trend (p = 0.078) in the circadian rhythmicity of *dbpa* (Table 2 and Fig. 5). In the liver, both size-selected lines shifted the peak of the circadian oscillation of *tef1* from CT12 to CT8 compared to the control (Table 2 and Fig. 5). By contrast, *dbpa* did not show differences among the lines. The control line only showed a trend (p = 0.095) for the circadian rhythmicity of *dbpa* (Table 2 and Fig. 5).

Clock controlled genes related to growth and energy balance

The circadian expression of genes related to physiological processes, such as energy balance (*lepa*) and growth (*igf1*), also revealed significant differences between the size-selected lines and controls. The circadian oscillation of *lepa* expression in the brain was only significant in the size-selected lines, but not in the control, with peaks at the same circadian time (CT8). By contrast, we did not find significant circadian oscillation of *lepa* in the liver (Table 3, Fig. 6). Finally, the *igf1* circadian oscillation in the liver was significant only in the control line (Table 3, Fig. 6).

DISCUSSION

Using an experimental system simulating fisheries-induced evolution following size-selection, we reveal the evolutionary legacy left by five generations of size-selective harvesting on shoaling and risk-taking behaviors as well as on the circadian molecular and behavioral outputs. In particular, we demonstrate that individuals of the small-

harvested line (which is conceivable in recreational fisheries managed with a maximum

size limit or in commercial gears having an upper bounds on mesh sizes, e.g. gill nets)

took more risk during feeding at the surface, were more cohesive, were more active at

the beginning of the photophase and extended self-feeding activity during photophase

relative to the control. We also found the large-harvested line (which can be expected

in recreational fisheries managed with a minimum-length limit or in commercial

fisheries operating with a trawl) to be shyer and to initially demand more food per

biomass than controls, and there was also a trend for this line to be less cohesive than

control. The evolutionary effects of size-selective harvesting on risk taking, shoaling

and foraging seem to be linked to daily activity rhythms, particularly in the small-

harvested line. From a molecular perspective, we also found shifts of the clock genes

circadian expression in the brain and liver in the same direction for both size-selected

lines. Importantly, the molecular shifts of the clock genes disappeared in the clock

output pathway, suggesting the occurrence of a switch leading to overall similar daily

rhythms at the behavioral level in all three selection lines. This suggests that being

overall active during the day and look for food around sunrise has strong fitness value

for zebrafish, no matter what the molecular clock is signaling.

Collective personality traits and daily activity rhythms

The behavioural traits we studied represented stable and repeatable traits and thus

were indicative of zebrafish collective personality traits (38, 39). Importantly, we

demonstrated that shoaling and risk-taking behaviors were repeatable behavioral

traits over ontogenetic time. We also revealed daily swimming and self-feeding activity

rhythms (except for late peaks of swimming activity) to be repeatable across

experimental stages.

We documented behavioral adaptations in all the studied collective personality

traits, indicating that the evolutionary response to the disruptive size-selective

harvesting treatments allowed the small-harvested fish to elevate boldness to increase

their foraging efficiency, while the opposite was found in the large-harvested

treatment which showed increased shyness. These results agreed with predictions

from theory (6, 12). Feeding during the selection experiment occurred with clumped

food at the surface of the water during daylight (17), which can be considered a risky

environment for zebrafish (i.e. avian predation; 40). Therefore, bold individuals

spending more time at the surface, as the small-harvested line, were likely more

efficient at foraging. Our interpretation is additionally supported by the fact that the

small-harvested line formed more cohesive shoals relative to controls. Social fish

species, such as zebrafish, can forage more effectively in groups than solitary

individuals (e.g., 41). Therefore, being more cohesive could provide additional foraging

advantages in the small-, but not in the large-harvested line. By contrast, the large-

harvested line evolved increased shyness, most likely a response to the evolution of a

fast life-history (6). Thus, our findings are fully consistent with the idea that elevated

and positively size-selective harvesting selected for increased timidity/shyness (6, 12).

As regarding daily activity rhythms, the small-harvested line was found to be

more active during the photophase and concentrated swimming activity at the

beginning of the photophase. These findings again suggested the evolution of a

foraging-optimizing timing of activity, likely linked to boldness. Our finding are

supported by a recent study demonstrating that proactive zebrafish (i.e., risk-taking

individuals) are more active in the first hours of photophase compared to reactive ones

(29). Relatedly, the self-feeding activity events of the small-harvested line extended to

the hours before and after light-on. The timing of feeding is a key aspect for survival

and is traded-off against predation risk and food availability (24). The ecological

significance of zebrafish feeding during the last hours of darkness may be related to

avoidance of visual predators (e.g. birds; 40) and/or to storing energy to be prepared

for spawning that occurs at the beginning of photophase (42). However, the food

demand sensor was located close to the water surface (42; see also Fig. S2) and an

increase of self-feeding events towards light-on by the small-harvested line could

indicate an additional increase of foraging efficiency and safeguarding energy intake in

risky contexts. A further support for a link between boldness and daily activity rhythms

has been reported in a recent study that revealed artificial light at night affects risk-

taking behavior in the Trinidad guppy, *Poecilia reticulata* (43).

Overall, our results indicated an asymmetrical adaptation among behavior and

daily activity rhythms in the size-selective treatments as previously reported for life

history traits in other selection experiments (20, 21, 44). In fact, we demonstrated an

adaptive behavioral link between boldness and daily activity rhythms in the small-, but

not in the large-, harvested line. Fisheries-induced evolution of behavior likely has a
complex and multivariate nature (9, 21), thereby complicating generalizations related
to evolutionary correlations among traits (45-47). The somewhat asymmetrical
adaptation of behavior in the two size-selection lines did not extend to all traits that
we measured, which could be related to a functional limitation of the range of
evolution of our model species (21, 48).

Brain clock gene shift

Purely size-selective harvesting modified the circadian clock transcript oscillations in
the same direction in both size-selected lines relative to controls. This result was a
surprise. In fact, despite the differences documented in the timing of daily swimming
and self-feeding activity rhythms among size-selection lines, we did not expect shifts in
the underlying molecular clockwork because all lines revealed an overall diurnal
swimming activity, while the self-feeding activity was mainly performed within a range
of 4-5 hours (Fig. 2). Conversely to expectations, we found that peaks of clock gene
expressions in the brain were shifted up to eight hours in both size-selection lines with
respect to controls. Yet, the peak of the oscillations of the clock output genes (*tef1* and
dbpa) showed a strong consistency among the three lines. Both *tef1* and *dbpa* belong
to a group of clock-controlled genes that feedback onto the core clock (49), and *tef1* is
also involved in the molecular switching mechanism controlling for photoperiodic
responsiveness in mammals (50), suggesting that the shifts documented in the core
clock and light-inducible genes can be switched before the clock output pathway. The
occurrence of a switch in the circadian clock was also reinforced by the fact that the
size-selected lines at F_9 showed genetic changes in areas of genes associated with
serotonin synthesis (17). Serotonin is a key element in the synthesis of melatonin that
is rhythmically secreted by the pineal gland (51). Melatonin represents a major
rhythmic output of the fish circadian system controlling photoperiodic-dependent
functions and synchronization of biological processes (51, 52). Thus, the serotonergic
system could have played a role in the switch mechanism we document in response to
size-selective harvest (17).

Our results did not indicate a link between harvesting-induced evolution of risk-
taking or other behaviors and the circadian molecular machinery in the brain because

the size-selection lines should have shown opposing adaptations in the circadian

system mirroring the adaptations in collective personality traits. Instead, the circadian

molecular shifts were pointing into the same direction in the two size-selection lines

relative to controls. Recent studies on zebrafish demonstrated that proactive and

reactive phenotypes in zebrafish explain a large fraction of the variation of the brain

transcriptome including expression of genes related to the circadian system (28).

Moreover, proactive phenotypes were found to have more robust molecular (i.e., clock

genes daily expression), humoral (i.e., daily secretion of melatonin) and behavioral

(i.e., locomotor activity) oscillations of the circadian system (29). However, both of the

cited studies measured transcripts abundance under LD conditions; therefore the truly

endogenous nature of the circadian molecular clockwork could have been masked by

the LD cycle. In our study, we measured transcript oscillations under constant

conditions, thereby providing a truly endogenous effect of size-selective harvesting on

the molecular circadian clockwork, and in our work both size-selection lines that

differed strongly in boldness responded similarly in the underlying molecular circadian

clockwork.

A possible explanation for our results could be related to a link between

harvesting induced evolution of life-history and the circadian system. In particular,

both size selected lines were previously reported to show evolution of a reduced age

and size at maturation compared to controls (17). Indeed, clock genes associate with

quantitative trait loci related to life history traits in fish, such as growth rate and length

in *Oncorhynchus kisutch* (53) and developmental rate and spawning time in

Oncorhynchus mykiss (32, 54). In particular, the core clock gene *clock* (i.e., *clock1a*, one

of the genes used here to characterize the core clock molecular machinery) is mainly

mapped in regions related to life history traits in salmonid fishes (32, 55). Therefore,

the possibility of a common molecular pathway or a pleiotropic effect of the brain core

clock machinery exist that is associated with the maturation schedule that evolved in

parallel directions in both size-selection lines according to previous studies (17).

Liver clock gene shift

The size-selected lines showed a shift in the peak of oscillations of some clock genes in

the liver, following a similar pattern to that observed in the brain. Moreover, the size-

selected lines did not reveal circadian rhythmicity of the light-inducible genes. Only the

large-harvested line indicated a circadian oscillation of *cry1a*. However, the clock

output genes had consistent peaks of circadian oscillations. The mechanisms governing

the synchronization of the liver's peripheral oscillator in fishes are not fully

understood, both LD and feeding play an important role in maintaining fitness (26, 56).

This could be the reason why we were not able to detect a clear effect of size-selective

harvesting in the liver circadian system as sampling was performed under constant dim

light to avoid a masking effect of the light-darkness cycle (57). The serotonergic system

also plays an important role in feeding processes and other behaviors in fish (57-59),

reinforcing the idea proposed above of its role as a mediator in the molecular switch of

the circadian clockwork.

We observed differences in the genes related to growth and energy balance

among the size-selection lines. The *lepa* expression in the brain displayed significant

circadian variations in both size-selected lines compared with the absence of

rhythmicity in the control group. Moreover, both size-selected lines showed a

disruption of the *igf1* circadian rhythm in the liver. Results on both genes indicated a

modification of growth and reproductive allocation processes, lipid and protein

metabolism and energy balance (59), as previously reported for both size selected lines

at the phenotypical level at F_9 (17). Modifications of the insulin growth factors

pathways in the zebrafish's skeleton muscle have also been documented in previous

size-selection experiments with zebrafish in other labs (20), supporting the idea that

this pathway could have a major role in mediating growth processes in response to

size-selection. Phenotypically, the large-harvested line also demanded more food

compared to the control at the beginning of the experimental phase, and earlier work

has revealed changes in genes related to growth-related processes and energy release

in the liver at F_{11} (35). Therefore, it is plausible that size-selective harvest has

evolutionarily altered the allocation of energy as well as the behavioral mechanisms of

energy acquisition.

Our interpretation of altered energy allocation caused by harvesting is also

supported by the results of the Lester biphasic growth model (60) that describes life-

time growth patterns and energy allocation to soma vs. reproduction in the zebrafish

selection lines (Fig. S1). The growth trajectories indicate harvest-induced changes in

energy allocation patterns at F_9 and F_{13} (Fig. S1), with the large-harvested line

investing early and intensively into reproduction, and the small-harvested one

investing early but at low intensity into gonad production and more intensively in

somatic growth (17). Note the terminal size attained by the large-harvested line was

previously found to be smaller than the other two lines (17; see also Fig. S1), while the

terminal size of the small-harvested and control lines was found to be similar. The

same pattern was revealed at F_{13}, suggesting evolutionary fixation and lack of

phenotypic rebound eight generations after selection was halted. The phenotypic data

suggests the evolution of a mechanism unrelated to energy allocation through which

the small-harvested line is able to attain similar terminal size to the control fish. This

mechanism, as elaborated above, could be mediated by a correlated response of

foraging efficiency in risky contexts.

Collectively, our results indicate the occurrence of a switch downstream the

circadian clock both in central (brain) and peripheral (liver) oscillators, in response to

five generations of size-selection in zebrafish. This is the first documented evidence

that classical harvesting patterns of fisheries can leave a legacy in the molecular

circadian system of evolving fish populations. There are three mechanisms used to

explain circadian system switches (61): *i)* altered properties of the circadian clock

leading to different activity patterns; *ii)* identical properties of the circadian clock

switching in the output pathway, leading to different activity patterns; and *iii)* identical

properties of the circadian clock subjected to masking that ultimately determine the

activity patterns. Our results suggest the presence of a further mechanism where

altered properties of the circadian clock are switched in the molecular output

pathway, leading to overall similar daily activity patterns. The role of the clock output

in determining diurnal and nocturnal phenotypes is also supported by a recent study

with the fruit fly *Drosophila melanogaster* (62), where ten generations of artificial

selection for diurnal and nocturnal phenotypes were sufficient to obtain highly

divergent strains where most differentially expressed genes were associated with the

clock output (62). The switching mechanism we describe here could be confounded by

the effect of size-selective harvesting on social interactions. In particular a possible

alteration of shoaling behavior could alter the social synchronization of circadian clocks

of the shoal members (63). Our results indicate that shoaling behavior (measured as

inter-individual distance among shoal members) was significantly altered in the small-

harvested line and also showed a trend to be different, in a different direction, in the

large-harvested line, suggesting that a social-mediated mechanism is not likely the

cause of the switching described for both size-selected lines in the present study.

Conclusions

The main impacts of human activities on biological rhythms of wildlife are related to

diurnal disturbance as documented in an increase of nocturnality in mammalian

species in response to human presence (64). Another important human-induced

change of activity rhythms is represented by artificial light at night that has been

demonstrated to have a profound impact on circadian rhythms in a wide range of taxa

(65). Using a selection experiment simulating fisheries-induced evolution, our work

adds to this literature by showing that size-selective harvesting as is typical in many

fisheries or under specific harvest regulations can substantially shift the circadian

molecular clockwork, yet without strong changes in the daily behavioral activity

rhythms of our model species. Moreover, the changes of risk-taking and shoaling

behaviors provide empirical evidence for size-selection to leave a strong behavioural

legacy, supporting recent theory (6). Our work shows that the evolutionary impact of

fisheries far extends from life-history traits to behavioral traits and to the circadian

system, providing advancements in the understanding of the molecular architecture

controlling the evolutionary trajectories of populations exposed to a multivariate

anthropogenic selection. Evolutionary changes as the ones documented here can have

important consequences for organism performance, social groups, population

dynamics, recovery, and ultimately fisheries quality (12, 13, 66).

MATERIAL AND METHODS

Selection lines

Our study system consists of wild-collected zebrafish (*Danio rerio*) from West Bengal in

India (for more details see 67). The parental wild-collected population was

experimentally harvested by exposing them to strong directional selection (a 75% per-

generation harvest rate) for either large body size (large-harvested) or small body size

(small-harvested). A third line was harvested randomly with respect to size and used as

control (17). Each selection line was replicated twice for a total of six selection lines.

Size-selective harvesting treatments were applied during the first five generations (17)

and then harvesting halted for eight generations until F_{13}. At F_9, the large-harvested

line evolved a smaller adult length and weight and higher relative fecundity compared

to controls (17). By contrast, the small-harvested line showed reduced reproductive

investment and no change in adult length compared to the control line (17). Both size

selected line evolved different maturation schedules by maturing smaller and at

younger age than the control line (for more details see 17) and such differences are

still detected at F_{13} (the generation used here) as indicated by results of a Lester

biphasic growth model and associated life-history traits (See Fig. S1). These changes

indicated a harvest-induced change in energy allocation patterns and no recovery to

pre-harvesting levels until F_{13}, with the large-harvested line investing early and

intensively into reproduction, and the small-harvested line investing early but at low

intensity into gonad production and more intensively in somatic growth. Moreover,

small-harvested juveniles at F_{10} were significantly more explorative and bolder than

controls, suggesting evolution of personality (17). In summary, the genetically diverged

selection lines differed significantly in their reproductive investment, adult body size,

juvenile behavior, trait variability, allele composition and gene expression (17, 34, 35).

Zebrafish selection lines for the present experiment where from F_{13}. They were

reared in groups and under *ad libitum* feeding and maintained under the following

conditions: water temperature at 26 ± 0.5 °C; photoperiod at 12:12 h LD cycle (lights

on/off at 07:00 and 19:00, respectively); fed three times per day with dry food

(TetraMin, Tetra) mainly in the first part of the light hours (at 09:00; 11:00 and 13:00).

Fish were reared and manipulated following the guidelines of the European Union

(2010/63/EU) and the Spanish legislation (RD 1201/2005 and Law 32/2007). The

experimental protocols were approved by the Spanish National Committee and the

Committee of the University of Murcia on Ethics and Animal Welfare.

Shoaling and risk-taking behavior experiments

Zebrafish is a social species that forms shoals (40), and both risk-taking behavior and

circadian rhythms are sensitive to social modulation (68, 69). Therefore, we used

zebrafish shoals for testing our hypothesis because it is the appropiate reflection of the

phenotypes in the original environment that the selection experiment operated on.

Shoaling behavior was recorded two consecutive times when zebrafish were

150 and 190 dpf (at about the same age used for the circadian experiment, see below).

We randomly selected 6 groups of 8 fishes for each of the six selection line (12 groups

for each treatment: large-harvested, small-harvested and control line). The groups

were transferred to 3-L rearing boxes at 30 dpf and maintained under the conditions

explained above. The standard length of the experimental fish groups at 170 dpf was

within the following range: large-harvested (1.9 – 2.7 cm); control (2.1 – 2.8 cm); small-

harvested (2.1 – 2.7 cm). Shoals behavior was measured in a white round arena

(diameter of 60 cm) with 10 cm of water as reported in other studies that have tested

shoaling behavior in zebrafish (70). The arena was placed on a table behind a white

curtain to avoid any kind of disturbance to the fish during the experimental trials. On

top of the arena (at about 1 m) we placed a webcam (Logitech C920; resolution: 1920 x

1080 pixels; frame rate: 30 frames per second) that was connected to a computer

were video recordings were stored. Behavior was measured between 09:00 and 13:00

h. We started the experimental trials two hours after lights on to avoid measuring

behavior when mating usually occurs (first hours after lights on; see 40). The fish group

was introduced in the experimental arena by gently pouring them together with the

water from the rearing boxes and left undisturbed for 25 min. Then, inter-individual

distance was measured for five min as reported in other studies (70).

We used the same groups of zebrafish to perform an additionally behavioral

test on the time spent feeding at the surface of the water. We repeated trials two

times (i.e., at 230 and 240 dpf). The groups of eight zebrafish were moved from the 3-

liter rearing box in a new experimental tank (width x length x height = 10 x 30 x 25 cm)

with 22 cm of water. The experimental tank was placed on a table behind a white

curtain. On the side of the experimental tank (at about 50 cm) we placed a webcam (as

described above). Zebrafish were introduced in the experimental tank and left

undisturbed for 3 min before starting the experimental trial. That time was sufficient

to remove the diving response usually seen in these methodological approach (71).

Then, a small quantity of food was added at the surface and at the same time we

started video recording for 30 s (sufficient time for zebrafish to consume almost all the

food) to quantify the cumulative time each of the eight fish spent feeding at the

surface of the experimental tank (one third of the water column = at about 7 cm). All

video recordings were analyzed with automated behavioral tracking (EthoVision XT 9,

Noldus Information Technologies Inc.; www.noldus.com).

Circadian experimental set up and design

Activity rhythms were recorded in a parallel experiment with a new set of fish when

they were at about 170 dpf. We randomly selected 3 groups of 15 fishes for each of

the six selection line (6 groups and 90 fishes for each treatment: large-harvested,

small-harvested and control line). The mass of the experimental fish groups was within

the following range: large-harvested (3.6 – 5.7 g); control (3.3 – 6.5 g); small-harvested

(4.6 – 6.1 g). Each 15-fish group was then placed in a 8-L experimental aquarium

(30×13×23 cm; see Fig. S2) equipped with aeration and mechanical and biological

filters. Light was provided by white LEDs (LED Flex-strip 1043-W, Solbright, RAYTE,

Murcia, Spain), with a light intensity of 435 lux at the water surface. The photoperiod

was set at a 12:12 h LD cycle, and a heater (75 W Magictherm; Prodac, China) kept the

water temperature at 26 ± 0.5 °C. The lights were set to turn on and off at 09:00 h and

21:00 h, respectively.

The experimental set up we used was previously designed to record at the same

time swimming (as a proxy of locomotor activity) and self-feeding activity by a group of

zebrafish (Fig. S2; see also 42). An infrared photocell (E3S-AD62 Omron, Kyoto, Japan)

was installed in the middle of the longer side of each aquarium, 15 cm below the water

surface. The sensitivity of this photocell was adjusted to detect fish at any distance

from the sensor to cover the full length of the aquarium (30 cm) and measure the

swimming activity. The self-feeding system consisted of another photocell located in

the upper corner of the aquarium, 2.5 cm below the water surface and 1.5 cm from

the side. The sensitivity of the photocell was adjusted to detect fish approaching within

a range of 1 cm independently of the size range used in the experiment. The photocell

was connected to a commercial feeder (Eheim, Deizisau, Germany) and when zebrafish

interrupted the infrared light beam, the feeder turned on and at about 0.2% g

flakes/biomass dropped into the aquarium. The feeder was placed in the same area, 10

cm from the sensor (for more details see 42). Swimming and self-feeding activities

were recorded and stored in a computer every 10 min. Before starting experimental

trials, fishes were acclimated to laboratory conditions and to the use of the self-feeder

for 15 days.

After the above mentioned acclimation period we recorded the swimming and

self-feeding activity for 38 days (from at about 170 to 210 dpf). The experimental trial

was subdivided in four consecutive phases: (first LD trial: 15 days) 12:12 h LD cycle to

investigate daily swimming and self-feeding rhythms; (constant conditions: 7 days)

constant dim light (3.5 lux at the water surface) to investigate the endogenous free-

running rhythms of swimming and self-feeding activity; (second LD trial: 15 days) 12:12

h LD cycle to investigate their resynchronization; (final constant conditions) constant

dim light to ascertain the endogenous expression patterns of clock genes. Moreover,

during the first 6 days of the first LD trial we measured the daily amount of food

demanded normalized by the mass of the fish group to have more quantitative

information on the self-feeding behavior of the fishes.

Gene expression analysis

The first group of genes was composed by genes related to the core transcriptional-

translational feedback loop driving circadian oscillation in vertebrates (*per1b, clock1a,*

arntl1a). The second group involved light-inducible genes (*per2, cry1a*). The third

group were genes related to the circadian clock output (*dbpa, tef1*), and the fourth

group was composed of circadian clock controlled genes (*lepa*, and *igf1*) related to

growth and energy balance. *lepa* (*lepa*) was measured both in brain and liver and *igf1*

only in liver. The gene *lepa* acts as satiation signal in teleosts and has implications in

energy balance and glucose homeostasis, it is mainly secreted in the liver with a daily

rhythm of expression (57, 72). The gene *lepa* is also expressed in the teleost brain

where expression increases after a meal (73). The gene *igf1* is involved in growth

processes and signals lipid metabolism and is secreted with a daily rhythm of

expression (57).

Gene expression was measured at the end of the activity rhythms experiment.

Samples were collected throughout a 24-hour cycle at six different time points

(circadian time, CT: 0, 4, 8, 12, 16 and 20). Fish were exposed for 24 h to the second

constant dim light phase. We sampled a total of five fishes for each time point and

each treatment. Fishes were killed on ice and then the whole brain and liver were

dissected and put in RNAlater at -80°C. Total RNA was isolated using a trizol reagent

(Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions. The amount,

quality and composition of isolated RNA were analyzed by Nanodrop ND-1000

(Thermo Fisher Scientific Inc., Wilmington, USA). DNase-treated RNA was used to

perform cDNA synthesis in a final volume of 20 µl using the iScript cDNA synthesis kit

(Biorad, USA). The reaction was performed at 46°C for 20 min, followed by a 1-min

inactivation step at 95°C.

cDNA was PCR-amplified with the StepOnePlus Real- Time PCR System (Applied

Biosystems, Foster City, CA, USA) using SsoAdvanced™ Universal SYBR® Green

Supermix (Biorad, USA). The thermal cycling conditions were as follows: 30 s of

denaturation at 95°C, followed by 40 cycles of a 15-s denaturation step at 95°C, and

then by an annealing-elongation step for 30 s at 60°C. After amplification, a melting

curve analysis was performed to confirm amplicon specificity. All the samples were run

in triplicate.

Gene-specific primers are indicated in Table S1. The relative expression levels of

each sample were calculated by the $2^{-\Delta\Delta CT}$ method (74), using geometric mean of three

housekeeping genes (*βactin*, *GADPH* and *EF1α*) in both tissues.

Statistical analysis

Inter-individual distance and time spent feeding at the surface of the water were

transformed by finding the exponent (lambda), which made the values of the response

variable as normally distributed as possible with a simple power transformation. Next,

we calculated the adjusted repeatability (selection lines as fixed effect; 75) across age

for inter-individual distance among group members (150 and 190 dpf) and for time

spent feeding at the surface (230 and 240 dpf). Finally, we modelled the behavioral

measures using linear mixed-effects models (76) with selection lines as fixed effect and

shoal nested within trials as random intercepts. Model fitting was assessed by checking

the plot of the residuals vs. the fitted values.

Time series of swimming and self-feeding activity were visually inspected by

plotting actograms and then using a Chi-square periodogram (77) to scan for the

presence of significant ($p < 0.05$) periodicity in the range 8-28 h and percentage of

variance (%V) explained by each period, reported as a measure of rhythms' robustness

(78). Waveform analysis (24-h based) was carried out in both LD trials, and raw time

series values (i.e. number of infrared light beam interruptions) were transformed in %

of maximum to standardize the data. The Midline Estimating Statistic Of Rhythm

(MESOR) was computed and represented as a horizontal threshold in waveform plots

(78).

As regarding swimming activity, we estimated total activity for each group by

calculating the area under the waveform curve during both LD trials (78). Next, early

and late peaks of activity were calculated as percentage of activity during the first

(early) and last (late) four hours of photophase. As regarding feeding activity, we

estimated feeding events for each group by calculating the area under the waveform

curve during four different hourly sections: *i*) from 06:00 to 07:00; *ii*) from 07:00 to

08:00; *iii*) from 08:00 to 09:00 (i.e., the hour before lights on); *iv*) from 09:00 to 10:00

(i.e., the hour after lights on). We calculated the adjusted repeatability (selection lines

as fixed effect) of these behavioral traits over the two LD trials as described above. We

lambda transformed them as above and modelled the data using a linear mixed-effects

models (76) with selection lines as fixed effect and shoals nested within LD trials as

random intercepts. Finally, we modelled the amount of food demanded by using a

linear model with selection lines and days of experiment (1-2, 3-4, 5-6) as fixed effects.

Model fitting was assessed by checking the plot of the residuals vs. the fitted values.

Periodicity in gene expression was checked using RAIN, a robust non-parametric

method for the detection of rhythms in biological data that can detect arbitrary

waveforms and has been widely applied in the measurement of circadian transcripts

abundance (79).

Periodogram and waveform analysis were performed using the software

Eltemps (www.el-temps.com). The rest of analyses were implemented using R 3.3.1

(https://www.R-project.org/) and the following additional packages: "rain" (79);

"rcompanion" (https://CRAN.R-project.org/package=rcompanion), "rptR" (80), and

"nlme" (81). In all cases, we used a 95% confidence interval.

Acknowledgments

We are grateful to Silva Uusi-Heikkilä for her contribution in the establishment of the

selection lines ($F_1 - F_9$) and for sharing unpublished data, Andrew E. Honsey for

modelling the Lester biphasic growth trajectories at F_{13}, the technicians at the IGB for

animal care support and Juan Fernando Paredes and José Antonio Oliver for helping

during sampling for molecular analysis. We are also grateful to Horacio de la Iglesia,

Achim Kramer and Carl H. Johnson for scientific discussion on preliminary results of the

experiments presented here. VS was supported by a Leibniz-DAAD postdoctoral

research fellowship (n. 91632699). CB was supported by University of Ferrara research

grants (FAR2018 and FIR2018). JFL-O was supported by a "*Ramón y Cajal*" research

fellowship (RYC-2016-20959) granted by the Spanish Ministry of Economy and

Competitiveness.

Author contributions

V.S., J.F.L., C.B. and R.A. designed research; V.S., J.F.L. and E.F. performed research;

V.S. analyzed data; V.S., J.F.L., E.F., C.B., and R.A. interpreted results; V.S. and R.A

wrote the paper with contributions from all the co-authors.

REFERENCES

1.   Allendorf FW & Hard JJ (2009) Human-induced evolution caused by unnatural
selection through harvest of wild animals. *Proc. Natl. Acad. Sci. USA*
106(Supplement 1):9987-9994.

2.   Heino M, Díaz Pauli B, & Dieckmann U (2015) Fisheries-induced evolution.
Annu. Rev. Ecol., Evol. Syst. 46(1):461-480.

3.   Jørgensen C*, et al.* (2007) Ecology: managing evolving fish stocks. *Science*
318:1247-1248.

4.   Dieckmann U & Heino M (2007) Probabilistic maturation reaction norms: their
history, strengths, and limitations. *Mar. Ecol. Prog. Ser.* 335:253-269.

5.   Jørgensen C & Holt RE (2013) Natural mortality: Its ecology, how it shapes fish
life histories, and why it may be increased by fishing. *J. Sea Res.* 75:8-18.

6.   Andersen KH, Marty L, & Arlinghaus R (2018) Evolution of boldness and life
history in response to selective harvesting. *Can. J. Fish. Aquat. Sci.* 75(2):271-
281.

7.   Uusi-Heikkilä S, Wolter C, Klefoth T, & Arlinghaus R (2008) A behavioral
perspective on fishing-induced evolution. *Trends Ecol. Evol.* 23(8):419-421.

8.   Hollins J*, et al.* (2018) A physiological perspective on fisheries-induced
evolution. *Evol. Appl.* 11(5):561-576.

9.   Claireaux M, Jorgensen C, & Enberg K (2018) Evolutionary effects of fishing gear
on foraging behavior and life-history traits. *Ecol. Evol.* 8(22):10711-10721.

10.   Biro PA & Stamps JA (2008) Are animal personality traits linked to life-history
productivity? *Trends Ecol. Evol.* 23(7):361-368.

11.   Réale D*, et al.* (2010) Personality and the emergence of the pace-of-life
syndrome concept at the population level. *Philos. Trans. R. Soc. Lond. B: Biol.*
Sci. 365(1560):4051-4063.

12.   Arlinghaus R*, et al.* (2017) Passive gear-induced timidity syndrome in wild fish
populations and its potential ecological and managerial implications. *Fish Fish.*
18(2):360-373.

13.   Diaz Pauli B & Sih A (2017) Behavioural responses to human-induced change:
Why fishing should not be ignored. *Evol. Appl.* 10(3):231-240.

14. Koeck B, *et al.* (2018) Angling selects against active and stress-resilient
phenotypes in rainbow trout. *Can. J. Fish. Aquat. Sci.* (ja).

15. Louison MJ, Jeffrey JD, Suski CD, & Stein JA (2018) Sociable bluegill, *Lepomis*
macrochirus, are selectively captured via recreational angling. *Anim. Behav.*
142:129-137.

16. Killen SS, Nati JJH, & Suski CD (2015) Vulnerability of individual fish to capture
by trawling is influenced by capacity for anaerobic metabolism. *Proc. R. Soc.*
Lond. B 282(1813).

17. Uusi-Heikkilä S, *et al.* (2015) The evolutionary legacy of size-selective harvesting
extends from genes to populations. *Evol. Appl.* 8(6):597-620.

18. Walsh MR, Munch SB, Chiba S, & Conover DO (2006) Maladaptive changes in
multiple traits caused by fishing: impediments to population recovery. *Ecol.*
Lett. 9(2):142-148.

19. Sbragaglia V, *et al.* (2019) Experimental size-selective harvesting affects
behavioral types of a social fish. *Trans. Am. Fish. Soc.* 0.

20. Amaral IP & Johnston IA (2012) Experimental selection for body size at age
modifies early life-history traits and muscle gene expression in adult zebrafish.
J. Exp. Biol. 215(Pt 22):3895-3904.

21. Renneville C, *et al.* (2018) Anthropogenic selection along directions of most
evolutionary resistance. *bioRxiv*:498683.

22. Panda S, Hogenesch JB, & Kay SA (2002) Circadian rhythms from flies to human.
Nature 417(6886):329-335.

23. Dunlap JC, Loros JJ, & DeCoursey PJ (2004) *Chronobiology: biological*
timekeeping (Sinauer Associates, Sunderland Massachusetts).

24. Kronfeld-Schor N & Dayan T (2003) Partitioning of time as an ecological
resource. *Annu. Rev. Ecol., Evol. Syst.* 34(1):153-181.

25. Cowan M, Azpeleta C, & López-Olmeda JF (2017) Rhythms in the endocrine
system of fish: a review. *J. Comp. Physiol., B*.

26. Idda ML, *et al.* (2012) Circadian clocks: lessons from fish. *Prog. Brain Res.*
199:41-57.

27. DeCoursey PJ, Walker JK, & Smith SA (2000) A circadian pacemaker in free-
living chipmunks: essential for survival? *J. Comp. Physiol. A* 186(2):169-180.

28.   Rey S, Boltana S, Vargas R, Roher N, & Mackenzie S (2013) Combining animal
738   personalities with transcriptomics resolves individual variation within a wild-
739   type zebrafish population and identifies underpinning molecular differences in
740   brain function. *Mol. Ecol.* 22(24):6100-6115.

741   29.   Tudorache C*, et al.* (2018) Biological clock function is linked to proactive and
742   reactive personality types. *BMC Biol.* 16(1):148.

743   30.   Koolhaas J*, et al.* (1999) Coping styles in animals: current status in behavior and
744   stress-physiology. *Neurosci. Biobehav. Rev.* 23(7):925-935.

745   31.   Nikhil KL & Sharma VK (2017) On the origin and implications of circadian
746   timekeeping: an evolutionary perspective. *Biological Timekeeping: Clocks,*
747   *Rhythms and Behaviour*, ed Kumar V (Springer India, New Delhi), pp 81-129.

748   32.   Leder EH, Danzmann RG, & Ferguson MM (2006) The candidate gene, *clock*,
749   localizes to a strong spawning time quantitative trait locus region in rainbow
750   trout. *J. Hered.* 97(1):74-80.

751   33.   Diaz Pauli B & Heino M (2014) What can selection experiments teach us about
752   fisheries-induced evolution? *Biol. J. Linn. Soc.* 111(3):485-503.

753   34.   Uusi-Heikkilä S*, et al.* (2016) Altered trait variability in response to size-selective
754   mortality. *Biol. Lett.* 12(9).

755   35.   Uusi-Heikkilä S, Savilammi T, Leder E, Arlinghaus R, & Primmer CR (2017) Rapid,
756   broad-scale gene expression evolution in experimentally harvested fish
757   populations. *Mol. Ecol.* 26(15):3954-3967.

758   36.   Weger BD*, et al.* (2011) The light responsive transcriptome of the zebrafish:
759   function and regulation. *PloS one* 6(2):e17080.

760   37.   Zhang X, Beaulieu J-M, Sotnikova TD, Gainetdinov RR, & Caron MG (2004)
761   Tryptophan hydroxylase-2 controls brain serotonin synthesis. *Science*
762   305(5681):217-217.

763   38.   Jolles JW, Laskowski KL, Boogert NJ, & Manica A (2018) Repeatable group
764   differences in the collective behaviour of stickleback shoals across ecological
765   contexts. *Proc. R. Soc. Lond. B* 285(1872):20172629.

766   39.   Bengston SE & Jandt JM (2014) The development of collective personality: the
767   ontogenetic drivers of behavioral variation across groups. *Front. Ecol. Evol.*
768   2(81).

40.   Spence R, Gerlach G, Lawrence C, & Smith C (2008) The behaviour and ecology
of the zebrafish, *Danio rerio*. *Biol. Rev. Camb. Philos. Soc.* 83(1):13-34.

41.   Pitcher TJ (1986) Functions of shoaling behaviour in teleosts. *The behaviour of*
teleost fishes, (Springer), pp 294-337.

42.   del Pozo A, Sanchez-Ferez JA, & Sanchez-Vazquez FJ (2011) Circadian rhythms
of self-feeding and locomotor activity in zebrafish (*Danio Rerio*). *Chronobiol. Int.*
28(1):39-47.

43.   Kurvers RHJM*, et al.* (2018) Artificial light at night affects emergence from a
refuge and space use in guppies. *Sci. Rep.* 8(1):14131.

44.   van Wijk SJ*, et al.* (2013) Experimental harvesting of fish populations drives
genetically based shifts in body size and maturation. *Front. Ecol. Environ.*
11(4):181-187.

45.   Polverino G, Santostefano F, Díaz-Gil C, & Mehner T (2018) Ecological
conditions drive pace-of-life syndromes by shaping relationships between life
history, physiology and behaviour in two populations of Eastern mosquitofish.
Sci. Rep. 8(1):14673.

46.   Royauté R, Berdal MA, Garrison CR, & Dochtermann NA (2018) Paceless life? A
meta-analysis of the pace-of-life syndrome hypothesis. *Behav. Ecol. Sociobiol.*
72(3):64.

47.   Wright J, Bolstad GH, Araya-Ajoy YG, & Dingemanse NJ (2018) Life-history
evolution under fluctuating density-dependent selection and the adaptive
alignment of pace-of-life syndromes. *Biol. Rev. Camb. Philos. Soc.*

48.   Arnold SJ (1992) Constraints on phenotypic evolution. *Am. Nat.* 140:S85-S107.

49.   Vatine G*, et al.* (2009) Light directs zebrafish *period2* expression via conserved
D and E boxes. *PLoS Biol.* 7(10):e1000223.

50.   Dardente H*, et al.* (2010) A molecular switch for photoperiod responsiveness in
mammals. *Curr. Biol.* 20(24):2193-2198.

51.   Lima-Cabello E*, et al.* (2014) A review of the melatonin functions in zebrafish
physiology. *J. Pineal Res.* 57(1):1-9.

52.   Falcón J, Migaud H, Muñoz-Cueto JA, & Carrillo M (2010) Current knowledge on
the melatonin system in teleost fish. *Gen. Comp. Endocrinol.* 165(3):469-482.

53.   McClelland EK, O'Malley KG, & Naish KA (2010) Clock genes localize to
quantitative trait loci for stage-specific growth in juvenile coho salmon,
Oncorhynchus kisutch. *J. Hered.* 101(5):628-632.

54.   Nichols KM*, et al.* (2007) Quantitative trait loci × maternal cytoplasmic
environment interaction for development rate in *Oncorhynchus mykiss*.
Genetics 175(1):335-347.

55.   Paibomesai MI, Moghadam HK, Ferguson MM, & Danzmann RG (2010) Clock
genes and their genomic distributions in three species of salmonid fishes:
Associations with genes regulating sexual maturation and cell cycling. *BMC
Research Notes* 3(1):215.

56.   Lopez-Olmeda JF, Tartaglione EV, de la Iglesia HO, & Sanchez-Vazquez FJ (2010)
Feeding entrainment of food-anticipatory activity and *per1* expression in the
brain and liver of zebrafish under different lighting and feeding conditions.
Chronobiol. Int. 27(7):1380-1400.

57.   Paredes JF, López-Olmeda JF, Martínez FJ, & Sánchez-Vázquez FJ (2015) Daily
rhythms of lipid metabolic gene expression in zebrafish liver: Response to
light/dark and feeding cycles. *Chronobiol. Int.* 32(10):1438-1448.

58.   Lillesaar C (2011) The serotonergic system in fish. *J. Chem. Neuroanat.*
41(4):294-308.

59.   Piccinetti CC*, et al.* (2013) Melatonin and peripheral circuitries: Insights on
appetite and metabolism in *Danio rerio*. *Zebrafish* 10(3):275-282.

60.   Lester NP, Shuter BJ, & Abrams PA (2004) Interpreting the von Bertalanffy
model of somatic growth in fishes: the cost of reproduction. *Proc. R. Soc. Lond.
B* 271(1548):1625.

61.   Hut RA, Kronfeld-Schor N, van der Vinne V, & De la Iglesia H (2012) In search of
a temporal niche: environmental factors. *Prog. Brain Res.* 199:281-304.

62.   Pegoraro M*, et al.* (2018) The genetic basis of diurnal preference in *Drosophila
melanogaster*. *bioRxiv*:380733.

63.   Bloch G, Herzog ED, Levine JD, & Schwartz WJ (2013) Socially synchronized
circadian oscillators. *Proc. R. Soc. Lond. B* 280(1765):20130035.

64.   Gaynor KM, Hojnowski CE, Carter NH, & Brashares JS (2018) The influence of
human disturbance on wildlife nocturnality. *Science* 360(6394):1232-1235.

65. Gaston KJ, Visser ME, & Hölker F (2015) The biological impacts of artificial light at night: the research challenge. *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 370(1667).

66. Laugen AT*, et al.* (2014) Evolutionary impact assessment: accounting for evolutionary consequences of fishing in an ecosystem approach to fisheries management. *Fish Fish.* 15(1):65-96.

67. Uusi-Heikkila S, Wolter C, Meinelt T, & Arlinghaus R (2010) Size-dependent reproductive success of wild zebrafish *Danio rerio* in the laboratory. *J. Fish Biol.* 77(3):552-569.

68. Castillo-Ruiz A, Paul MJ, & Schwartz WJ (2012) In search of a temporal niche: social interactions. *Prog. Brain Res.* 199:267-280.

69. Jolles JW, Boogert NJ, Sridhar VH, Couzin ID, & Manica A (2017) Consistent individual differences drive collective behavior and group functioning of schooling fish. *Curr. Biol.* 27(18):2862-2868. e2867.

70. Nunes AR, Ruhl N, Winberg S, & Oliveira RF (2017) Social phenotypes in zebrafish. *The rights and wrongs of zebrafish: Behavioral phenotyping of zebrafish*, ed V. KA (Springer International Publishing), pp 95-130.

71. Kalueff AV (2017) *The rights and wrongs of zebrafish: Behavioral phenotyping of zebrafish* (Springer).

72. Rønnestad I*, et al.* (2017) Appetite-controlling endocrine systems in teleosts. *Front. Endocrinol.* 8(73).

73. Yuan X*, et al.* (2016) Leptin expression in mandarin fish *Siniperca chuatsi* (Basilewsky): Regulation by postprandial and short-term fasting treatment. *Comp. Biochem. Physiol., A: Mol. Integr. Physiol.* 194:8-18.

74. Livak KJ & Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25(4):402-408.

75. Nakagawa S & Schielzeth H (2010) Repeatability for Gaussian and non-Gaussian data: a practical guide for biologists. *Biol. Rev. Camb. Philos. Soc.* 85(4):935-956.

76. Pinheiro J, Bates D, DebRoy S, & Sarkar D (2007) Linear and nonlinear mixed effects models. *R package version* 3:57.

77.   Sokolove PG & Bushell WN (1978) The chi square periodogram: its utility for analysis of circadian rhythms. *J. Theor. Biol.* 72(1):131-160.

78.   Refinetti R (2006) *Circadian physiology (2nd ed)* (Boca Raton, Fla, CRC Press).

79.   Thaben PF & Westermark PO (2014) Detecting rhythms in time series with RAIN. *J. Biol. Rhythms* 29(6):391-400.

80.   Stoffel MA, Nakagawa S, & Schielzeth H (2017) rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods Ecol. Evol.* 8(11):1639-1644.

81.   Pinheiro J*, et al.* (2017) Package 'nlme'. *Linear and Nonlinear Mixed Effects Models, version*:3-1.

Table 1 - The output of the periodogram analysis for swimming and self-feeding activity rhythms during the three steps of the experiment (first light-darkness trial: LD1; constant dim light: LL; and second light-darkness trial: LD2) according to the three selection lines (large-harvested: LH, small-harvested: SH and control: RH). Periods (T) or free running periods (*tau*) are averaged according to the number of groups (N; the proportion indicates the number of groups with significant rhythms) and the robustness of periodicities are expressed as percentage of variance explained (%V) together with the midline estimating statistic of rhythm (MESOR). The standard errors are reported between parentheses.

Phenotype	Experimental Phase	Line	N	T or *tau* (se)	%V (se)	MESOR (se)
Swimming	LD1	LH	6/6	24.00 (0.00)	47.41 (4.54)	0.070 (0.006)
		RH	6/6	24.00 (0.02)	29.74 (10.38)	0.103 (0.011)
		SH	5/6	24.02 (0.02)	52.48 (10.13)	0.116 (0.012)
	LL1	LH	4/6	20.88 (1.70)	29.43 (11.23)	-
		RH	2/6	26.33 (0.00)	30.29 (8.59)	-
		SH	2/6	23.42 (0.33)	23.54 (1.89)	-
	LD2	LH	6/6	24.03 (0.02)	52.45 (8.54)	0.081 (0.017)
		RH	6/6	24.00 (0.03)	51.01 (14.30)	0.066 (0.006)
		SH	5/6	24.00 (0.00)	56.96 (20.82)	0.108 (0.018)
Self-feeding	LD1	LH	6/6	24.00 (0.00)	46.44 (7.45)	0.018 (0.003)
		RH	6/6	23.97 (0.02)	38.89 (18.18)	0.016 (0.009)
		SH	6/6	24.01 (0.03)	43.21 (6.35)	0.026 (0.005)
	LL1	LH	4/6	23.56 (0.52)	28.59 (6.73)	-
		RH	3/6	22.56 (1.31)	18.86 (3.43)	-
		SH	1/6	16.83 (-)	16.03 (-)	-
	LD2	LH	6/6	24.03 (0.02)	46.07 (11.67)	0.013 (0.003)
		RH	6/6	24.01 (0.01)	32.29 (7.92)	0.008 (0.003)
		SH	6/6	24.00 (0.02)	37.87 (10.17)	0.016 (0.003)

Table 2 - The output of the analysis with RAIN (a robust non-parametric method for
the detection of rhythms in biological data) regarding the circadian oscillation of
transcript abundance of the genes related to the circadian system. Gene expression
was measured at the end of the experiment in constant dim light. Transcripts are
subdivided according to their role in the circadian system for each selection lines
(large-harvested: LH, small-harvested: SH and control: RH) and in each tissue (brain
and liver). The timing of the peak of the circadian oscillation (CT0, CT4, CT8, CT12, CT16
or CT20) is reported together with the p value (N is between 4 and 5). Significant
circadian rhythmicity is reported in bold.

Clock mechanism	Gene	Line	Brain		Liver	
			Peak	p value	Peak	p value
Core clock	arntl1a	LH	**16**	**< 0.01**	16	0.086
		RH	0	0.052	**0**	**< 0.01**
		SH	16	0.682	**0**	**< 0.01**
	clock1a	LH	**16**	**< 0.01**	**0**	**< 0.001**
		RH	**0**	**< 0.05**	0	0.472
		SH	20	0.279	**20**	**< 0.001**
	per1b	LH	**12**	**< 0.001**	**16**	**< 0.01**
		RH	**8**	**< 0.01**	8	0.878
		SH	**12**	**< 0.001**	**12**	**< 0.05**
Light-inducible	per2	LH	**12**	**< 0.001**	0	0.112
		RH	16	0.117	**16**	**< 0.05**
		SH	**8**	**< 0.05**	20	0.136
	cry1a	LH	**12**	**< 0.001**	**16**	**< 0.01**
		RH	**8**	**< 0.001**	**16**	**< 0.01**
		SH	**16**	**< 0.001**	20	0.120
Clock output	tef1	LH	**8**	**< 0.05**	**8**	**< 0.01**
		RH	**8**	**< 0.001**	**12**	**< 0.001**
		SH	**8**	**< 0.05**	**8**	**< 0.05**
	dbpa	LH	8	0.078	**8**	**< 0.05**
		RH	**8**	**< 0.05**	8	0.095
		SH	**8**	**< 0.001**	**8**	**< 0.05**

Table 3 - The output of the analysis with RAIN (a robust non-parametric method for
the detection of rhythms in biological data) regarding the circadian oscillation of
transcript abundance of the genes related to neuroendocrinological aspects of energy
balance, appetite and growth. Gene expression was measured at the end of the
experiment in constant dim light. Transcripts are subdivided according to their
function, for each selection lines (large-harvested: LH, small-harvested: SH and control:
RH) and in the corresponding tissue (brain and liver). The timing of the peak of the
circadian oscillation (CT0, CT4, CT8, CT12, CT16 or CT20) is reported together with the
p value (N is between 4 and 5). Significant circadian rhythmicity is reported in bold.
920
921

Gene	Function	Tissue	Line	Peak	*p* value
lepa	Appetite Food intake	Brain	LH	**8**	**<0.05**
			RH	8	0.456
			SH	**8**	**<0.01**
		Liver	LH	20	0.580
			RH	12	0.422
			SH	12	0.876
igf1	Growth	Liver	LH	4	0.227
			RH	**4**	**<0.01**
			SH	4	0.134

 FIGURE CAPTIONS

Figure 1 – Group behavioral traits of Zebrafish shoals composed of eight members

measured from 190 to 240 dpf during five-minute video recordings. Inter individual

distance (A) was measured at 190 and 210 dpf, while time spent feeding at the surface

(B) was measured at 230 and 240 dpf. Trial values are reported on the left with

repeatability score and confidence intervals, while boxplots are reported on the right.

The different colors represent the selection lines: red for large-harvested (LH; N = 12

for each trial) grey for control (RH; N = 11) and blue for small-harvested (SH; N = 12).

Boxplots represent the median (bold centerline), the 25^{th} (top of the box) and 75^{th}

percentile (bottom of the box). Significant differences are indicated by horizontal lines

(*: *p* value < 0.05; **: *p* value < 0.01; ***: *p* value < 0.001).

Figure 2 - Mean waveforms (time scale 24 h) for swimming (A, B) and self-feeding (C,

D) activity of Zebrafish shoals are expressed as percentage of the maximum during the

first (LD1: days 1-15) and second (LD2: days 22-36) light-darkness trial. Each point

represents the 10-min binned mean across all the experimental days in LD1 and LD2

for all the groups. The different colors represent the selection lines: red for large-

harvested (LH; N = 6) grey for control (RH; N = 6) and blue for small-harvested (SH; N =

6). The horizontal lines represent the midline estimating statistic of rhythm as reported

in Table 1. The vertical lines represent the standard error (*N* between 75 and 90). Grey

shadowed areas represent the darkness hours (lights on is at zeitgeber time = 12;

experimental time = 09:00).

Figure 3 – Total swimming activity during photophase (area under the waveform

curve, AUC; A) together with early (B) and late (C) peaks of activity (percentage of

activity during the first and last four hours of photophase, respectively) measured

during the first (LD1: days 1-15) and second (LD2: days 22:36) light-darkness trial. LD1

and LD2 values are reported on the left side together with repeatability score and

confidence intervals (n.s. = repeatability is not significant). Boxplots are reported on

the right side. The different colors represent the selection lines: red for large-

harvested (LH; N = 6) grey for control (RH; N = 6) and blue for small-harvested (SH; N =

5). Boxplots represent the median (bold centerline), the 25th (top of the box) and 75th

percentile (bottom of the box). Significant differences are indicated by black horizontal

lines (**: *p* value < 0.01).

Figure 4 – Self-feeding activity (area under the waveform curve, AUC) during the last

three hours of scotophase (grey shadow: 06:00 - 07:00; 07:00 - 08:00; 08:00 - 09:00)

and the first hour of photophase (09:00 - 10:00) measured during the first (LD1: days

1-15) and second (LD2: days 22-36) light-darkness trial. LD1 and LD2 values are

reported on the left side together with repeatability score and confidence intervals.

Boxplots are reported on the right side. The different colors represent the selection

lines: red for large-harvested (LH; N = 6) grey for control (RH; N = 6) and blue for small-

harvested (SH; N = 6). Boxplots represent the median (bold centerline), the 25th (top of

the box) and 75th percentile (bottom of the box). Significant differences are indicated

by black horizontal lines (**: *p* value < 0.01; ***: *p* value < 0.001).

Figure 5 - Mean transcript abundance of the genes related to the circadian system in

the brain (left) and liver (right) at different circadian time for the three selection lines

(large-harvested: LH, small-harvested: SH and control: RH; N between 4 and 5).

Figure 6 - Transcript abundance of the circadian clock controlled genes related to

growth and energy balance. Lines were used to better indicate the circadian oscillation

of the mean transcript abundance for the three selections lines (large-harvested, LH:

red; small-harvested, SH: blue; and control, RH: grey; N between 4 and 5). Boxplots

represent the median (bold centerline), the 25th (top of the box) and 75th percentile

(bottom of the box).

991   Fig. 1

993
994
995   Fig. 2
996

Fig. 3

Fig. 4

1005
1006

1007

1008   Fig. 5

1009

1010

1011

1012

1013   Fig. 6

1014

 SUPPLEMENTARY INFORMATION

SUPPLEMENTARY TEXT

Swimming and feeding activity rhythms

Visual inspection of actograms evidenced diurnal swimming activity (with some

evident bimodal peaks at lights on/off) while feeding activity indicated a main peak in

the last part of the darkness hours (Fig. S2).

The Chi-square periodogram analysis identified a significant 24-hours

periodicity in swimming activity during both light-dark (LD) trials. Interestingly during

the first LD trial the robustness of rhythms of the control line is lower (29.74±10.38

%V; Table 1) than the other two lines (large-harvested: 47.41±4.54 %V; small-

harvested: 52.48±10.13 %V, Table 1) in fact the control line showed a higher swimming

activity during darkness (Fig. 2). Such differences were no more evident during the

second LD trial when all lines have similar robust rhythms (large-harvested: 52.45±8.54

%V; control: 51.01±14.30 %V; small-harvested: 56.96±20.82 %V; Table 1) and the

control line did not show higher swimming activity during darkness (Fig. 2). Finally

swimming activity showed variable significant free running rhythmicity under constant

dim light (Table 1). The Chi-square periodogram also detected bimodality (peaks of

significant 12-hours periodicity) in swimming activity in both LD trials.

Self-feeding activity rhythms had a significant 24-hours periodicity during both

LD trials. Robustness was similar in all the lines during the first (large-harvested:

46.44±7.45 %V; control: 38.89±18.18 %V; small-harvested: 43.21±6.35 %V, Table 1)

and second (large-harvested: 46.07±11.67 %V; control: 32.29±7.92 %V; small-

harvested: 37.87±10.17 %V) LD trial. Finally under constant dim light the self-feeding

activity indicated variable free running rhythmicity (Table 1). The Chi-square

periodogram did not detect bimodality in self-feeding activity rhythms.

During the first LD trial bimodality was detected in 3 out of 6 groups in the

large-harvested line (robustness: 25.17±12.20 %V), in three out of six groups in the

control line (robustness: 8.50±1.73 %V) and in one out of five groups in small-

harvested line (robustness: 23.18 %V). During the second LD trial bimodality was still

present in two out of six groups in large-harvested line (robustness: 25.97±7.98 %V), in

four out of six groups in the control line (robustness: 21.79±3.82), and in two out of

five groups in small-harvested line (robustness: 9.17±0.97 %V).

SUPPLEMENTARY TABLE

1049

1050 **Table S1 –** Primers used for RT-qPCR together with the references used to design them. The function of each gene is reported together with the
1051 tissue (Brain: B; Liver: L) and the forward and reverse primers.

Function	Tissue	Genes	Forward	Reverse	Reference
Circadian clock: *Core clock*	B / L	*per1b*	CCGTCAGTTTCGCTTTTCTC	ATGTGCAGGCTGTAGATCCC	(1)
		clock1a	CTGGAGGATCAGCTGGGTAG	CACACACAGGCACAGACACA	(1)
		arntl1a	TAGAGCGCTGTTTGCTGATG	GACCCGTGGACTTCAGTGAC	(2)
Circadian clock: *Light-inducible*	B / L	*per2*	ATGTCGATGGCTTTAGGCAG	CGAGACATCCAGAAGGTGCT	(1)
		cry1a	TCCGCTGTGTGTACATCCTC	CAAACACTGCAGCAAAAACC	(1)
Circadian clock: *Clock output*	B / L	*dbpa*	GTCCAGGCCAATTTCTCAAA	CACAGCAGTCCTTCCCTCTC	(3)
		tef1	AAAGCTCTGCTTGAGTACCCTTT	ACTCCTCCAGATCCATGTACTCC	(3)
Growth	L	*igf1*	TCCGTCTCCTGTTCGCTAAATC	TTTGGGTCTCCAGCAAAAGC	(4)
Appetite	L/B	*lepa*	AACTGCAGGCAAAGACCATC	GCGGGAATCTCTGGATAATG	(4)
Housekeeping	B / L	*β-actin*	GCCTGACGGACAGGTCAT	ACCGCAAGATTCCATACCC	(1)
		gadph	GTGGAGTCTACTGGTGTCTTC	GTGCAGGAGGCATTGCTTACA	(5)
		ef1a	CCTTCGTCCCAATTTCAGG	CCTTGAACCAGCCCATGT	(6)

1052

REFERENCES USED IN THE SUPPLEMENTARY TABLE S1

1. Cavallari N*, et al.* (2011) A blind circadian clock in cavefish reveals that opsins mediate peripheral clock photoreception. *PLoS Biol.*
9(9):e1001142.

2. Weger M*, et al.* (2013) Real-time in vivo monitoring of circadian E-box enhancer activity: a robust and sensitive zebrafish reporter line
for developmental, chemical and neural biology of the circadian clock. *Dev. Biol.* 380(2):259-273.

3.   Mracek P*, et al.* (2012) Regulation of *per* and *cry* genes reveals a central role for the D-box enhancer in light-dependent gene

expression. *PLoS One* 7(12):e51278.

4.   Paredes JF, López-Olmeda JF, Martínez FJ, & Sánchez-Vázquez FJ (2015) Daily rhythms of lipid metabolic gene expression in zebrafish

liver: Response to light/dark and feeding cycles. *Chronobiol. Int.* 32(10):1438-1448.

5.   McCurley AT & Callard GV (2008) Characterization of housekeeping genes in zebrafish: male-female differences and effects of tissue

type, developmental stage and chemical treatment. *BMC Mol. Biol.* 9(1):102.

6.   Weger BD*, et al.* (2011) The light responsive transcriptome of the zebrafish: function and regulation. *PloS one* 6(2):e17080.

1065

 SUPPLEMENTARY FIGURES

1067

1068   **Figure S1** - Fits of the Lester biphasic growth model at generations F_9 and F_{13}. Colors

1069   represent the different selection lines: large-harvested (red), control (grey) and small-

1070   harvested (blue). Adapted from Uusi-Heikkilä et al. (2015) and Sbragaglia et al. (2019).

1071

1072

1073

Figure S2 – A schematic representation of the experimental aquarium used in this
study to measure swimming and self-feeding activity rhythms. Light illumination was
provided from above. An infrared sensor was placed in the central part of the
aquarium to measure swimming activity, while another infrared sensor was places to
one of the corners and connected to the nearby feeder (for more details see Del Pozo
et al., 2011).

1080

1081

1082
1083

Figure S3 - Six representative double plotted (time scale 48 h) actograms (one for each
selection line) representing locomotor and self-feeding activity over 38 days of
experiment (left: large-harvested line; middle: control line, right: small-harvested line).
Swimming and self-feeding activity of zebrafish are indicated by the height of black
areas that represent the number of infrared light beam interruptions/10 min.
Open/filled bars at top represent a 12-h light/12-h darkness photoperiod for the first
(LD1) and second (LD2) light-darkness trial, while during LL zebrafish shoals were
exposed to constant dim light.

1092

1093

1094

1095

1096

1097

1098

Figure S4 – Boxplots of the amount of food demanded (g) normalized on the weight of
the fish group (g) during the first 6 days in the first light-darkness trial (LD1: days 1-6).
The different colors represent the selection lines: red for large-harvested (LH; N = 6)
grey for control (RH; N = 6) and blue for small-harvested (SH; N = 6). Boxplots
represent the median (bold centerline), the 25th (top of the box) and 75th percentile
(bottom of the box). Significant differences are indicated by black horizontal lines (*: *p*
value < 0.05).