Title: Mosaic evolution, preadaptation, and the evolution of evolvability in apes

Author: Caroline Parins-Fukuchi*[1]

[1]Department of Geophysical Sciences, University of Chicago, Chicago, IL, USA

*Author for correspondence: parinsfukuchi@uchicago.edu

Acknowledgements

I wish to thank D Alvarado-Serrano, M Cosman, CW Dick, DC Fisher, M Foote, E

Greiner, D Jablonski, LM MacLatchy, H Marx, J Saulsbury, SA Smith, G Slater, N

Walker-Hale, M Webster for conversations that have greatly benefitted this work. I also

thank S Worthington for help accessing his skeletal measurements that comprised a large

portion of the character matrix analyzed here. The manuscript benefitted immensely from

careful reviews provided by P Wagner and PD Polly.

Abstract: A major goal in post-synthesis evolutionary biology has been to better understand how complex interactions between traits drive movement along and facilitate the formation of distinct evolutionary pathways. I present analyses of a character matrix sampled across the haplorrhine skeleton that revealed several modules of characters displaying distinct patterns in macroevolutionary disparity. Comparison of these patterns to those in neurological development showed that early ape evolution was characterized by an intense regime of evolutionary and developmental flexibility. Shifting and reduced constraint in apes was met with episodic bursts in phenotypic innovation that built a wide array of functional diversity over a foundation of shared developmental and anatomical structure. Shifts in modularity drove dramatic evolutionary changes across the ape body plan in two distinct ways: 1) an episode of relaxed integration early in hominoid evolution coincided with bursts in evolutionary rate across multiple character suites; 2) the formation of two new trait modules along the branch leading to chimps and humans preceded rapid and dramatic evolutionary shifts in the carpus and pelvis. Changes to the structure of evolutionary mosaicism may correspond to enhanced evolvability that has a 'preadaptive' effect by catalyzing later episodes of dramatic morphological remodeling.

Keywords: integration, evolvability, preadaptation, modularity, hominoidea

Introduction

The formation of new species and phenotypes is often hypothesized to be limited

by the availability of unoccupied 'adaptive zones', classically represented as peaks on a

fitness landscape (Wright 1932, Simpson 1944). Frequently referred to as 'ecological

opportunity' (Losos 2010), the opening of new adaptive zones through shifts in biotic

interactions and the environment is often invoked to explain large evolutionary radiations

(Rainey and Travisano 1998, Yoder et al., 2010, Wagner and Harmon et al 2012, but see

Slater 2015). The role of ecological processes in generating morphological novelty can be

contrasted with an increasing focus on 'constructional' factors that include functional

interactions and architectural constraints (Seilacher 1970, 1991). While character

evolution may sometimes reflect simple optimization processes when natural selection

operates independently on separate traits, the evolutionary pathways realized in nature are

also constrained by the complex interactions between functionally dependent traits and

the limitations of an organism's developmental architecture. When placed into a broader

evolutionary literature, these limiting factors might be coarsely partitioned into

developmental (Gould and Lewontin 1979, Alberch 1980, Olson 2012) and functional

(Charlesworth et al., 1982, Maynard-Smith et al., 1985, Arnold 1992) constraints.

Developmental and functional constraints limit the capacity for a clade to evolve

novel phenotypes. Together, these factors might be said to determine the level of

'constructional opportunity' available at a given time. Contrasted to ecological

opportunity, constructional opportunity reflects how the structure of developmental and

functional interactions between characters limits the 'evolvability' of a clade—its

capacity to evolve novel phenotypes. While ecological opportunity is often evoked as a

limiting factor on a clade's phenotypic diversification, constructional opportunity, as

measured by the overall level of constraint and internal integration between phenotypes,

may also limit the range of evolutionary modifications available to a population faced

with ecological pressure.

Evolutionary constraints are shaped by shared developmental pathways,

multivariate selection, and underlying genetic processes such as epistasis and pleiotropy.

Such interactions between traits and genes often create a common tendency to drive the

formation of modules of characters that are internally integrated in their evolutionary

trajectories. Such integration is typically identified by calculating numerical correlations

in their variance (Olson and Miller 1958). Integration between traits constrains the

combination of phenotypes accessible to each. For example, for two traits that are

strongly positively integrated, it is unlikely to observe an individual that possesses

phenotypes for each trait that lie at opposing ends of the spectrum of variation. As long as

this correlation exists, the evolution of the organism is constrained in the sense that only

certain combinations of phenotypes are accessible. At the population level, integration

can be caused by multivariate selection or physical/constructional linkages such as

pleiotropy or allometry (Cheverud 1984). However, it is generally not possible to

distinguish between these in macroevolutionary study systems.

Phenotypic integration generally occurs in a modular fashion (Wagner et al.

2007). Instead of covarying uniformly across the entire organism, integration often

manifests as a patchwork of modules of internally integrated traits that are free to evolve

largely independently (Goswami et al. 2014, 2015). Such modular structures lead to the

pattern of mosaic evolution when separate modules evolve at different rates as lineages

diverge (Felice and Goswami 2017, Parins-Fukuchi, in press). When analyzed

phylogenetically, these patterns result in 'mosaic disparity'—a scenario where individual

modules of traits each display a unique pattern in phylogenetic disparity across lineages

(Fig. 1).

The related concepts of mosaicism, constraint, and integration can help to explain

the emergence of phenotypic novelty across clades. For example, when ecological

opportunity is abundant, evolutionary radiations can be driven primarily by the release of

constraints (Wagner et al. 2003). In addition, bursts in evolutionary rate can also coincide

with the break-up of existing modules and the subsequent formation of a new modular

structure (Wagner 2018). Characterizing the role of integration and mosaic evolution in

shaping diversity across clades will ultimately contribute to a deeper understanding of

phenotypic evolution at the constructional level. This may improve our understanding of

evolutionary processes more generally by revealing the extent to which fluctuations in

constructional opportunity contribute to macroevolutionary patterns alongside ecological

opportunity.

In this study, I sought to determine the extent to which the evolutionary

trajectories blazed by ape species were shaped by early shifts in the structure of

integration and the timing of development. To address these questions, I reconstructed the

mosaic macroevolutionary patterns in morphological, neurological, and developmental

phenotypes across living and fossil great apes throughout the Miocene. To help place

these empirical results in a theoretical context, I also present a set of simple simulations

to generate a theoretical expectation for the patterns in disparity expected when

Markovian diffusion is constrained by trait integration imposed by both functional and

structural/developmental interactions. Taken together, these analyses suggest the

capability for shifting patterns in constraint imposed by integration to create

constructional opportunities that generate repeated and dramatic episodes in

morphological innovation.

Apes have been extensively studied in the context of both developmental

constraint and environmental adaptation. Young et al. (2010), found that apes display

weak integration between the fore- and hindlimb in comparison to cercopithecoids.

Reduced integration can confer greater functional flexibility, and so may have facilitated

the proliferation of the diverse locomotor function across living and fossil apes. Changes

in developmental timing have also been important in ape evolution. Developmental

heterochrony appears to have driven the emergence of many key aspects of hominin

morphology in the cranium and post-cranium (Gould 1977, Berge 1998). This long-

standing interest in the developmental basis of ape evolution has not excluded

environmental hypotheses. Environmental fluctuations throughout the Miocene and

Pliocene have also frequently formed the basis for adaptive hypotheses across apes

(Andrews 1992, Ruff 1994, Michel et al., 2014). The body plans of hominins, and

undoubtedly apes in general, have been shaped through modifications to separate suites

of traits occurring at different times (Holloway 1973, McHenry 1975). These diverse

threads suggest that a multiplicity of causes, perhaps including both ecological

opportunity and constructional factors, has shaped the evolution of hominoid body plans.

As a result, apes are an excellent exemplar taxon in which to pluralistically examine the

mosaic patterns in constraint and innovation that shape body plans in living and fossil

vertebrates.

Methods

Morphological data

I gathered a dataset of 149 quantitative morphological traits from the literature

(Lewton 2010, Worthington 2012) spanning the cranium, forelimb, and pelvis. These

traits were sampled across 10 extant taxa. I also collected fossil cranial and forelimb traits

from Worthington (2012), retaining all taxa with at least 25% matrix occupancy. Taxa

were all studied at the generic level, with each extant genus represented by a single

exemplar species. Each morphological trait represented either a dimensionless ratio or

geometric mean, depending on how the author of each dataset handled intraspecific

variation and body mass correction. I scaled the traits to display an empirical variance of

1 across all taxa. This simplified the analyses by reducing the complexity of the dataset,

while retaining the same comparative information. Since the original traits were

dimensionless, both the unscaled and the scaled datasets would facilitate examination of

relative, rather than absolute, evolutionary rates, but the scaled traits improve Markov-

chain Monte Carlo (MCMC) mixing and simplify the identification of mosaic suites.

Neurological data

To directly study the evolutionary patterns in related developmental and

phenotypic traits, I also gathered neurological data from the literature (Capellini et al.,

2010, Boddy et al., 2012). For neurological phenotypes, I collected encephalization

quotients (EQ) estimated in 76 primate species by Boddy et al. (2012). EQ is calculated

by fitting a nonlinear model to brain size and body size and taking the deviations

(residuals) from the best fit curve and so measures the enlargement of brain size while

controlling for allometry with body size. I compared EQ patterns to those in a postpartum

encephalization development (ED) metric calculated as:

$ED = EQ_{adult} - EQ_{neonatal}$ (1)

ED is therefore intended to reflect the amount of brain tissue grown by members of each

species following birth relative to their body size. The ED metric was scaled to unit

variance across all sampled taxa. Neonatal brain and body mass data representing 24

primate species were acquired from Capellini et al. (2010).

Fossil placement and divergence time estimation

Since the evolutionary relationships between the extant taxa represented in the

morphological dataset have all been extensively studied and are well-resolved, I used

these as a fixed scaffolding to place the fossil taxa. Fossil placements were inferred from

the continuous trait dataset using the cophymaru program (Parins-Fukuchi 2018). I used

the 'binary weights' procedure to filter out reliable traits from ones likely to mislead by

using the extant tree as a fixed point of reference. Markov-chain Monte Carlo (MCMC)

simulations were run for 1,000,000 generations and checked manually for convergence.

The posterior tree sample was summarized as the maximum clade credibility tree using

sumtrees.py (Sukumaran and Holder 2010).

For downstream comparative analyses, I also estimated divergence times on the

full 19-taxon tree. I downloaded cytochrome B sequences from Genbank representing

each of the 10 extant taxa. Molecular dating was performed using Beast version 2.4.8

(Bouckaert et al. 2014) using the fossilized birth-death (FBD) prior. The topology was

fixed to the summary tree generated during the fossil placement step. Temporal

occurrence ranges for the fossil taxa were assembled from the literature. These fossil and

extant occurrence ranges were used in the dating analysis to infer the diversification and

sampling rate parameters used in the FBD prior.

Measuring integration throughout the hominoid radiation

As an evaluation of the overall strength of integration in the skeletal traits across

the haplorrhine phylogeny, I estimated ancestral states for each character under Brownian

motion along a dated tree of nine extant haplorrhine taxa (*Pan paniscus* was removed

because it was missing pelvic measurements). I then estimated the strength with which

the reconstructed phenotypes at each internal node were correlated with the values at

each descendant node or tip. The correlations indicated how strongly the reconstructed

evolutionary changes in skeletal traits were integrated along each branch in the tree.

Stronger correlations corresponded to tighter overall integration. The Python script that I

developed for the reconstruction of ancestral states and calculation of character

correlations is available in the data supplement.

Evolutionary rates across mosaic morphological suites

I used the *greedo* program to recover mosaic patterns in disparity from the

continuous trait dataset (Parins-Fukuchi 2019). *Greedo* is a phylogenetic

clustering/mixture model estimation approach that uses the Akaike information criterion

(AIC) to iteratively merge and split clusters of traits to find a best-supported set of

character suites based on shared patterns in disparity (i.e., where integrated traits within

the same module are similarly disparate across taxa), where each is represented by a tree

with branch lengths (Fig. 1). Traits that display the highest improvement in log-likelihood

when assigned to a separate cluster are prioritized during the splitting steps to first

identify likely sources of heterogeneity in the dataset. During the merging step, only

clusters that show an improved AIC score when placed under the same model are joined.

This process recovers an estimate of the number of suites, the membership of each trait,

and a tree with branch lengths scaled to units of disparity for each suite. Fossil taxa were

not included in the mosaic analyses, because they were too fragmentary in their sampling

to inform disparity patterns when the traits were split into suites. I performed many runs

of the *greedo* procedure to avoid presenting results obtained from a suboptimal peak on

the likelihood surface. Since several of the top clusterings yielded close AIC support, I

performed an additional model-averaging step using Akaike weights to summarize the

results as a graph where each trait occupies a node where each link is weighted to reflect

the total weighted AIC support across all the models. Comparative analyses were

performed using a single summary clustering that resulted from application of the

Markov clustering (MCL) algorithm to the support network (Dongen 2000).

To examine the anatomical composition of the mosaic suites of characters, I

performed a statistical enrichment procedure to identify skeletal regions that were over

and under-represented in each suite relative to the proportions for each present in the

entire dataset. The proportion of the 149-trait dataset represented by each skeletal region

was used to generate a set of 'expected' values. I then compared these to the observed

proportions within each suite to examine trends in the deviations of the proportions

occupied by each region. I did not assess the statistical significance of these deviations,

because 1) my aim was to examine the general patterns in the composition of each

character suite rather than to determine absolute significance, and 2) the small sample

size of the dataset generated expected chi-square test statistic values that were nearly

entirely <5, making the test inappropriate. To examine the relative contributions of

separate anatomical regions in shaping the macroevolutionary patterns reconstructed in

each suite, I also performed a principal component analysis (PCA) on the traits contained

within each suite. I then transformed the loadings to calculate the total variance

contributed by each trait across all axes of the PCA, summing them to produce the

contribution of each anatomical region.

I transformed the branch lengths for each mosaic suite, which were scaled to

reflect disparity (total variance, v, accumulated over time t), to rates (v per unit t) using

the results from the divergence time analysis. I also calculated evolutionary rates

averaged over the entire morphological dataset while including the fossil taxa to coarsely

reconstruct evolutionary tempo throughout the Miocene. Although this step missed

valuable information recovered by the mosaic analyses, including the fossil taxa in this

manner enabled finer resolution into the coarse patterns in evolutionary tempo throughout

the Miocene.

Evolutionary rates in neurological data

I estimated macroevolutionary rates across primates in ED and EQ using BAMM

version 2.5.0 (Rabosky 2014). Since the trees that I constructed for the morphological

analysis contained only a fraction of the taxa present in the neurological datasets, I used

the dated primate supertree packaged in BAMM that was originally sourced from Mos

and Mooers (2006) and pruned the tips to match the taxa present in each dataset. MCMC

simulations were run until the estimated sample size (ESS) well exceeded 200 for each

parameter. Results were presented by plotting the mean rate estimated along each branch

and the maximum *a posteriori* configuration of rate shifts using BAMMtools version

2.5.0 in R (Rabosky et al., 2014).

Lineage diversification rate analyses

I performed analyses of origination and extinction rates in hominoid and

cercopithecoid fossil records to examine the correspondence between lineage

diversification and the patterns in disparification emphasized in the analyses of

morphological rates and mosaicism. Fossil occurrence data spanning primates were

acquired from the Paleobiology Database (paleobiodb.org) on July 31, 2019. This dataset

was partitioned into two subsets: one encompassing Hominoidea and the other

representing Cercopithecoidea. Rates of origination and extinction were inferred from

each subset separately using PyRate (Silvestero et al. 2014). The inferential model was

constrained to contain uniform origination and extinction rates across time, thus ignoring

the possibility of diversification shifts within each clade. Preservation was constrained to

being time-homogeneous and was assumed to be uniform within each clade. While

ignoring potential information regarding diversification rate heterogeneity through time,

this simplified model matches classic paleobiological work that assumes simple birth-

death-sampling models and was therefore appropriate for the comparison of average

diversification rates in sister clades while avoiding possible model overfitting that may

obfuscate the straightforward test sought here.

Simulated Markovian diffusion

To examine the expected effect of constructional opportunity stemming from both

developmental and functional constraint on phenotypic evolution, I designed a set of

evolutionary simulations based on a simple Markovian diffusion. In this system,

quantitative traits belonging to a single population of organisms evolve stochastically

along a fitness landscape. Each generation, values for each trait are proposed. Values that

increase the overall fitness of the population are accepted. To mimic the effect of drift,

values that decrease fitness are also accepted with some (user-specified) probability.

Constraint stemming from developmental integration caused by pleiotropy can be

mimicked by proposing values for traits that covary, while functional constraints are

modeled by evolving traits along a shared, multivariate adaptive landscape, modelled

here using a mixture of multivariate Gaussian distributions. In a completely

unconstrained system, each trait evolves according to its own univariate landscape, with

its value drawn independently of all others in each generation. Performing simulations in

this way facilitated an illustration of the potential for functional and developmental

integration to constrain evolutionary rates in an adaptive landscape. Although

distinguishing between these modes of integration is generally difficult or impossible in

paleontological study systems, the simulations performed here provide insight by testing

1) the potential for varying systems of integration to catalyze or constrain adaptive

change and 2) the extent to which functional and developmental integration generate

overlapping and distinct patterns in evolutionary rate and disparity. When placed in a

macroevolutionary context, the simulations here should be thought of as a demonstration

of the possible microevolutionary processes that may have driven the higher-level

patterns in evolutionary rate occurring along a single phylogenetic branch in the mosaic

analyses. The script used to generate these simulations is available in the data

supplement.

Results and Discussion

Strength of integration throughout the haplorrhine radiation

Calculating character correlations from the skeletal data along each branch of the

extant haplorrhine phylogeny revealed a single episode of reduced integration at the root

of Hominoidea, where the correlation decreased from 0.12 at the ancestor of catarrhines,

to -0.04 at the root of hominoids (Fig. 2). This suggests that the reduction in integration

across apes between the fore- and hindlimbs revealed by Young and colleagues (2009)

extended across the skeleton along the branch leading to the stem hominoids. Such

episodic relaxed integration was not detected in the other haplorrine lineages, with

cercopithecoids (0.2-0.33 correlation) and platyrrhines (0.16-0.31 correlation) displaying

higher integration than hominoids (-0.04-0.25 correlation) overall. However, it is possible

that more complete sampling of taxa and characters across both of these clades may

reveal changes in modularity and integration not detected here.

Composition of mosaic suites

The analysis of mosaic disparity recovered five suites of traits (Fig 3b). The

cranium was the largest contributor of both traits and variance across all suites except for

C4, which was represented by the pelvis alone (Fig. 3b and 3c). This reflects the higher

sampling of cranial traits in the dataset. While the cranium has previously been shown to

display substantial modularity (Felice and Goswami 2017), the results here suggest that

individual cranial modules may result from the formation of broader evolutionary

complexes shared with the post-cranium. The suites were distinct in their composition

across postcranial anatomy: C0 was represented strongly by the wrist, ulna, and humerus;

C1 was represented postcranially by the scapula; C2 by carpal and pelvic traits. The

cranium was represented even more strongly than expected in C3 and contributed nearly

half of the variance (Fig. 3b).

Rates of morphological evolution

319   The phylogenetic rate calculations for each suite revealed a strong pattern of

evolutionary mosaicism (Fig. 3a). Three of the suites, C0, C1, and C3, displayed the

highest rates after the divergence of the great apes from hylobatids, but before the

divergence of gorillas from humans and chimpanzees. The two suites that did not

experience the shared great ape rate increase, C2 and C4, experienced large bursts in

Homo. This finding is consistent with general knowledge of human evolution, as both

suites are dominated by cranial traits and pelvic traits related to birthing and locomotor

function (Table S1).

327   The rate analyses (Fig. 3a) show that humans have experienced major

macroevolutionary bursts. However, the unique characteristics of the human body plan

are built over an anatomical structure shared by all great apes that was shaped in the

Miocene during similarly dramatic episodes. For example, although suite C0 increased in

evolutionary rate along the human branch, its evolution was shaped by a sustained period

of elevated rate between 20 and 10 million years ago. This suggests that much of the

'groundwork' underlying the derivation of humans' unique features may predate the

divergence between our lineage and chimpanzees'. The mosaic analysis also

demonstrates that substantial visible phenotypic novelty can result from the evolutionary

remodeling of a small subset of the anatomy. The most elevated evolutionary rates in

humans were observed in suites C2, C3, and C4, which cumulatively comprise fewer than

one-third of the traits sampled in the matrix (Table S1). C4, the smallest suite (10 traits)

displayed by far the largest increase in rate along the *Homo* lineage. Perhaps notably, C4

also was the most static character suite throughout the earlier stages in the hominoid

radiation. The mosaic pattern recovered here shows that the uniqueness of the human

body plan was shaped by dramatic evolutionary shifts in only a small number of traits.

While the large amount of missing data among the fossil taxa made an additional

mosaic analysis infeasible, their inclusion revealed an otherwise hidden shift in

evolutionary rate that occurred at the root of Hominoidea (approximately 30 million years

ago). The fossil data also recapitulated the burst in evolutionary rate during the mid-

Miocene that was displayed by C0 and C1 in the mosaic analysis (Fig. 3 and 4). When

averaging over all the traits and including fossil taxa, these two episodes are the most

dramatic macroevolutionary events when tracing the evolutionary lineage leading to

Homo, suggesting the importance of early shifts in the ape body plan in shaping the

functional morphologies of living taxa.

Several of the major rate shifts that occurred throughout the great ape clade

corresponded to increased evolvability conferred by relaxed integration among early

apes. However, comparison of the human-specific evolutionary rates in cranial, pelvic,

and neurological traits to the phylogenetic character correlations in Figure 2 shows that

the elevated morphological rates in humans were immediately preceded by an *increase* in

the strength of integration. This illustrates an alternative means through which shifts in

systems of modularity promote rapid and dramatic evolutionary remodeling of

morphology. The suites that showed the most dramatic rate increases in humans, C2 and

C4, were evolutionarily inactive prior to the most recent common ancestor (MRCA) of

chimps and humans. This suggests that, while early ape evolution was characterized by

relaxed integration leading to rapid evolutionary change in existing modules, human

morphological evolution was shaped by the formation of two small modules of characters

from the pelvis and carpus that underwent an initial rate increase following their

formation along the branch leading to chimps and humans and then a more dramatic rate

increase in hominins. Although the overall structure of integration was well-defined

along the chimp-human MRCA branch, the formation of *new*, smaller modules

contributed to dramatic changes across the body plan in humans. The occurrence of these

initial shifts in modularity and rate along the chimp-human MRCA shows that

chimpanzees share the modified structure of integration that facilitated the evolution of

key human traits. This result is surprising given the typical assumption that chimpanzees

are morphologically more similar to other great apes than are humans.

Rates of neurological evolution

The great apes differentiated rapidly from other catarrhines in both EQ and ED.

Homo displays the highest rate of both EQ and ED evolution. However, the human rate

shift in both traits occurred as part of an older trend of rapid neurological evolution in

African apes. For EQ, humans experienced a substantial increase in evolutionary rate

relative to the rest of the African apes, despite the statistical evidence for a shared rate

shift at the root of the clade (Fig. 5b). The shared macroevolutionary regime shared by all

African apes is most apparent in encephalization development (Fig. 5a), where the

estimated evolutionary rate in *Homo* increased only slightly after splitting from *Pan* (Fig.

5c). It appears that the increased encephalization developed throughout the post-natal

period in humans reflects a general trend among great apes in the evolutionary plasticity

of neurological development. As a result, the ability to develop a relatively large mass of

brain tissue does not itself appear to be a strong limiting factor in the evolution of large

encephalization in humans. Instead, the morphological analyses suggest that pelvic traits

(C4 in Fig. 1) demanded a more dramatic alteration in macroevolutionary regime,

requiring both the formation of a new suite of characters and massive burst in rate.

Structural and ecological opportunity in ape evolution

The elevated evolutionary rates experienced by apes early in their divergence

correspond to a general relaxation of constraint in early Miocene stem and ancestral

hominoids. The reduction in integration across the skeleton and burst in the evolutionary

rate of neurological development that occur at the base of the hominoid clade correspond

in timing to the elevated rates in morphological evolution observed in most of the skeletal

characters examined here (Fig. 3 and 4). Freed from previous functional and

developmental limitations, ape body plans would have rapidly diversified when placed in

the context of the repeated environmental fluctuations that occurred throughout the mid

and late Miocene (Michel et al. 2014, Hunt 2016) before becoming functionally canalized

through development and stabilizing selection. This pattern is consistent with a scenario

in which morphological features evolve by stochastic diffusion on a Simpsonian adaptive

landscape that is alternately dampened and released by relaxed and shifting patterns in

character integration.

Rapid and frequent environmental fluctuations throughout the Miocene suggest a

possible abundance of available adaptive zones during the evolution of both

cercopithecoids and hominoids. Paleoecological information extracted from deposits

containing key hominoid taxa are so variable that environmental turnover often outpaces

the effects of time-averaging (Michel et al. 2014). Early Miocene ape and monkey

species evolved in overlapping environmental conditions and shared similar

ecomorphological and dietary habits. However, as their lineages diverged, apes evolved a

diverse set of locomotor suites, while evolution in locomotor features has remained

comparatively static across old world monkeys (Hunt 2016). The patterns in

developmental and mosaic morphological evolution revealed here suggest that the

substantial differences between the two taxa in phenotypic disparity were shaped more by

differences in patterns in integration and constraint rather than from the ecological

opportunities available to either. While hominoids and cercopithecoids inhabited a

similar range of habitats throughout their evolution, and so likely experienced a similar

abundance of ecological opportunity, greater constructional opportunity is a

distinguishing feature that correlates with hominoids' vast disparity in skeletal

morphology.

The scenario entertained here for apes is consistent with recent work that suggests

that the initial burst in phenotypic disparity that often accompanies the origin of a new

lineage can result from an early relaxation of constraints through the dissolution of

integrated modules that is followed by reformation of character suites that lead to rapid,

correlated evolutionary changes between constituent characters (Wagner 2018). In the

cited work, the author distinguishes this 'breakup-relinkage' model with one where

evolutionary rates are elevated by a relaxation of constraint alone. The 'breakup-

relinkage' model is directly supported by the phylogenetic integration pattern, which

show that an initial decrease in integration at the root of hominoids was followed by an

increase at the following node (Fig. 2). Therefore, the recanalization of functional

variation into a set of suites with distinct functional and developmental properties early in

hominoid variation is likely equally important to the initial relaxation in constraint in

having driven the remarkable morphological disparity observed across hominoid species.

"Preadaptation" and developmental enablers in apes

Bursts in phenotypic change and diversification are often preceded by

evolutionary enablers that facilitate the construction of more diverse morphological forms

or fulfill a necessary condition for the later emergence of a more derived trait. Such

patterns have been suggested to stem from higher-level processes variously referred to as

"preadaptation" (Bock 1959), "exaptation" (Gould and Vrba 1982), and "developmental

enablement" (Donoghue 2005). The evolution of developmental, morphological, and

behavioral traits is often hierarchical through functional inter-dependencies and

temporally autocorrelated, with changes in traits at relatively lower levels 'setting the

stage' for higher-level traits by providing the developmental or structural conditions

necessary for their emergence. The stepwise macroevolutionary bursts observed here

show that hominoid evolution has been defined by sequential, episodic releases in

constraint that likely set the stage for later innovations that compounded upon earlier

rearrangements in the structure of developmental and functional integration.

The related concepts of preadaptation, exaptation, and developmental enablement

all focus on identifying the origin of single characters that facilitate the later emergence

of more derived states. However, the patterns presented here suggest that such processes

may often be better characterized as reflecting larger changes in developmental and

functional character-linkage systems rather than by the sequential derivation of individual

characters. The dramatic bursts in evolutionary rate that occur in the largest character

suites coincide with or immediately follow the reduction in integration that occurred at

the root of hominoids. This suggests that a general shift in functional and/or

developmental integration occurred near this time. This shift precedes and, although

speculative, may have resulted in a general predisposition across apes that facilitated the

later functional divergence of later species into an unusually broad range of habits.

Therefore, rather than stemming from a single "preadaptive" character, the morphological

disparity of apes and the emergence of features derived in single lineages (such as

bipedalism in humans) may be better explained by a shift in developmental and

functional integration patterns earlier in ape evolution.

Enhanced evolvability conferred by shifts in the structure of integration

contributed to the emergence of key human traits. The apparent formation of suites C2

and C4, represented disproportionately by the cranium and pelvis, along the branch

leading to humans and chimpanzees coincided with an initial burst in evolutionary rate.

Both of these events immediately preceded the dramatic bursts in *Homo* (Fig. 3).

Likewise, EQ and ED have been evolving rapidly throughout the ape radiation, with ED

experiencing an initial increase in evolutionary rate before the splitting of gorillas,

chimps, and humans followed by more dramatic upticks of rate in humans in both traits

(Fig. 5c). While distinct in its details, this pattern in chimps and humans provides further

evidence for the preadaptive potency of shifts in systems of modularity and integration in

driving dramatic anatomical rearrangements. It also illustrates that the broad range of

possible evolutionary pathways opened by either an overall reduction in integration (at

the hominoid root) or the formation of new modules (in chimps and humans), rather than

movement toward a specific set of phenotypes, is the means through which such

predispositions facilitate evolutionary remodeling.

It is possible that the initial bursts in ED and suite C2 and C3 evolution, coupled

with the reduced integration in the MRCA of chimp and humans, may reflect a set of

structural rearrangements that facilitated, at least in part, the later emergence of human-

specific neurological and cognitive features. This scenario echoes the results of Rice

(1998), who demonstrated that humans and chimpanzees, to the exclusion of gorillas,

both inherited a shared 'additional' stage of brain development from our most recent

common ancestor. It is therefore possible that, in addition to sharing a similar derived

structure of modularity, chimpazees possess more of the neurological and cranial

groundwork that contributed to the evolution of humans' particular mode of cognition

than has previously been appreciated.

The patterns uncovered here suggest that the concepts of preadapation and

exaptation may sometimes be more compellingly applied by shifting focus from

individual features to patterns in integration and modularity across broad suites of

characters. Although the concept is essentially unchanged as applied to integration

patterns, it may be more accurate in the example here to refer instead to an 'evolutionary

predisposition' given the lack of evidence for a specific sequence of selective regimes

(Smith et al. 2018). This semantic argument is likely also valid in more traditional

examples of single trait exaptations or preadaptations. And so, a generalization from

preadaptation or exaptation to evolutionary predisposition that 1) incorporates integration

patterns and 2) shifts emphasis from adaptation, which makes assumptions about the

(often unknowable) set of historical microevolutionary processes, to the sequence of

observable states in both characters and integration patterns can provide a stronger

conceptual basis for future work that considers contingency and sequential order in the

evolutionary pathways of complex traits. Although this reconceptualization might

superficially appear to de-emphasize evolutionary process in favor of pattern, I would

instead argue that constraining the scale of inquiry and incorporating a broader range of

biological complexity can facilitate the reconstruction of elusive higher-level

macroevolutionary processes that shape organismal diversification and disparification

over deep timescales. This formulation also avoids the conceptual and practical

frustrations that occur when attempting to invoke the concepts of preadaptation and

exaptation to explain evolutionary history over deep timescales by reducing dependency

on knowledge of lower-level population processes.

Diversification and disparification in haplorrhine evolution

Cercopithecoids have diversified in species number at over twice the rate as

hominoids over the same time scale (Table 1). One limitation of these analyses stems

from cercopithecoids' somewhat poor fossil record as compared to hominoids.

Nevertheless, the difference in preservation rates estimated in each clade (~3 in apes

compared to ~2.6 in cercopithecoids) should ameliorate some of the potential effect of

this bias on parameter estimates. The general pattern of higher net diversification in

cercopithecoids is also consistent with previous neontological results (Purvis et al. 1995),

suggesting that the paleontological estimates presented here are adequately robust to

these effects. This difference in lineage diversification rates between both taxa

demonstrates that increased morphological disparification is, overall, not correlated with

faster lineage diversification in haplorrhine primates. While the evolutionary

predisposition conferred upon apes by their relaxed integration corresponds to an increase

in morphological disparity, this effect does not extend to lineage diversity.

On the surface, the overall similarity in the environments, dietary niches, and

geographical ranges of hominoids and cercopithecoids suggests that cercopithecoids have

a selective advantage in the species-sorting dynamics between the two clades. However,

the long-term persistence of hominoids and their remarkable innovation in locomotor

function relative to cercopithecoids throughout this timespan suggests an equilibrium in

their relative abundances (Van Valen 1975, Chesson 2000). Although cercopithecoids

experienced less dramatic innovation in postcranial morphology, they display a broader

range of derived digestive physiologies, including both intestinal and dental features

associated with food processing (Hunt 2016). Given these observations, it is possible that

the differential in diversification rate between cercopithecoids and hominoids stems from

differences in the timescales of their life histories and other factors not directly related to

interspecific competitive ability. If this is the case, hominoids may achieve steady

persistence through higher evolvability conferred by reduced integration and increased

developmental flexibility when faced with changing environments, while cercopithecoids

may do so through a combination of their higher net diversification rates and capability to

exploit a range of dietary resources through their digestive physiologies. Such differences

would be consistent with an equilibrium model of coexistence rather than a simple

species selection model that predicts the ultimate extinction of hominoids through

competitive exclusion by cercopithecoids. Nevertheless, more rigorous comparison of

each of these attributes in both lineages is needed to further constrain the range of

possible explanatory factors.

Constrained Markovian diffusion and entropy

The patterns in character disparity shown here evoke a Markovian diffusion that is

dampened by constraint. The tightly integrated nature of vertebrate body plans suggests

major constraints in their ability to fill morphospace. However, the pattern displayed by

apes suggests that the structure of integration may sometimes shift, allowing diffusion

into new areas. Morphological evolution might then be conceived as a multivariate

diffusion in morphospace where movement is directionally biased due to fitness

differences modelled by a traditional Simpsonian adaptive landscape and certain regions

are rendered inaccessible by structural interactions between characters. Although this

diffusive model, including the feedback links between phenotype, development, and

environment, has been considered previously (Fisher 1986), its behavior has not been

well explored in this context.

The Monte Carlo simulations presented here provide a theoretical illustration of

the population-level dynamics stemming from one possible set of integration scenarios

and their potential effects on the evolutionary disparity/rate along a single branch in

Figures 3 or 4. The results shed light on the empirical analyses by demonstrating the

overlapping and distinct empirical patterns that can result from varying levels of

developmental and functional integration in a simple closed system in the absence of

environmental fluctuation (Fig. 6). As is expected from quantitative genetic theory

(Cheverud 1984, Maynard-Smith et al. 1985), developmental and functional constraints

can generate similar patterns (Fig. 6a and 6b). In all these cases, the overall entropy

observed in the system over time remains low, with the system tending to become stuck

or moving away very slowly from a suboptimal location (Fig 4c and 4e) or crawling

slowly in concert along a gradient (Fig 6a and 6b). However, when both types of

constraint are released, the system displays high entropy, with each trait able to

independently jump between peaks (Fig. 6d). Such unconstrained movement would be

expected to result in higher phenotypic disparity over macroevolutionary timescales by

freeing individual lineages to generate novel character combinations when exposed to

distinct and changing environments.

The patterns in increased evolutionary rate and entropy across traits displayed by

the less constrained simulated diffusions provide mechanistic explanations that are

consistent with the bursts in evolutionary rate encountered throughout hominoid

evolution. Although it is not possible in the empirical example to distinguish between

functional and developmental integration as is done in the simulations, the general pattern

of reduced integration that gives way to increased evolutionary rate is consistent between

the two. Alternation between the varying strengths of integration as explored in the

simulations would likely generate substantial novelty during periods of relaxed

integration. Episodes of reduced developmental constraint (Figs. 2 and 5a) may have

generated an initial burst of constructional opportunity that lead to rapid morphological

diversification as ape populations were exposed to highly variable environments

throughout the mid-Miocene. If the processes modelled in the simulations drove the

patterns revealed in the empirical analyses, this sequence of events would represent true

preadaptation in the sense that relaxed integration leads to increased mean population

fitness. Under the evolutionary models explored here and elsewhere (Wagner 2018),

stabilizing effects such as developmental canalization (Waddington 1959) or functional

covariance (Cheverud 1984) would then be expected to re-constrain the newly divergent

phenotypes after an initial burst in disparification. This dampened diffusive model was

explored in early work in theoretical morphology that focused on the constrained filling

of morphospace (Raup 1968).

The simulated diffusions and empirical pattern in evolutionary predisposition

revealed here hint at the source of teleological concepts in human evolution. If the

morphological and neurological traits can be assumed to have followed a multivariate

Markovian diffusion that is at least qualitatively similar to the simulations in Figure 6

(especially Fig. 6a), the pattern in evolutionary rates along the path leading to humans

may paint a misleading view that such changes have followed a progressive trend leading

to a human-defined apex. However, the mosaic analysis here suggests that humans'

biological uniqueness can be attributed to a relatively small number of anatomical

rearrangements built over a longer diffusion into an area of developmental-morphospace

occupied by all apes that is characterized by increased structural opportunity. Instead of

treating humans as exceptional, this view suggests that the structural opportunities that

emerged early in ape evolution may have freed the ancestors of currently extant taxa to

blaze unique trajectories along a complex, multivariate adaptive landscape that would

have been otherwise inaccessible. Structurally, all ape species might therefore be viewed

as equally progressive in their evolution, with species-level differences in form and

function shaped by stochastic differences in separate realized evolutionary paths along a

shared adaptive landscape. Alternatively, separate taxa may have forged new and unique

adaptive landscapes following speciation events as structural differences between newly

isolated species drove the creation of new functional niches early in hominoid evolution.

The derived structure of integration and neurological development shared by chimps and

humans highlights the importance of contingency in human evolution—chimps' sharing

of much of the same underlying constructional architecture shows how this derived

predisposition allowed, but did not determine, the emergence of humans' unique suite of

phenotypes.

The empirical and simulated analyses presented here evoke a picture of

morphological evolution that involves episodic diffusion across adaptive zones.

However, the empirical and theoretical analyses allow for alternating and interacting

roles for both ecological and constructional opportunity. This view can supplement

existing conceptions of adaptive radiation and key innovation by providing an alternative

to ecological opportunity as a lone driving factor in the disparification of form. Instead,

phenotypic disparity produced through diffusion into a set of abundant adaptive zones

can be directed and limited by the availability of structural opportunity. Nature surely

merges these simplified extremes when shaping patterns in biodiversity (Seilacher 1991),

and so further integrative study of the interplay between ecological and constructional

factors will be critical in developing a pluralistic understanding of the complex patterns

and processes that have shaped the vast diversity across the tree of life.

References

Alberch P. 1980 Ontogenesis and morphological diversification. American Zoologist 20,

653-667.

Andrews P. 1992 Evolution and environment in the Hominoidea. Nature 360, 641–646.

(doi:10.1038/360641a0)

Arnold SJ. 1992 Constraints on phenotypic evolution. American Naturalist, 140(Supp):

S85-S107.

Berge C. 1998 Heterochronic processes in human evolution: An ontogenetic analysis of

the hominid pelvis. American Journal of Physical Anthropology 105, 441–459.

(doi:10.1002/(sici)1096-8644(199804)105:4<441::aid-ajpa4>3.0.co;2-r)

Bock J. 1959 Preadaptation and multiple evolutionary pathways. Evolution 13, 194-211.

Boddy AM, McGowen MR, Sherwood CC, Grossman LI, Goodman M, Wildman DE.

2012 Comparative analysis of encephalization in mammals reveals relaxed

constraints on anthropoid primate and cetacean brain scaling. Journal of

Evolutionary Biology 25, 981–994. (doi:10.1111/j.1420-9101.2012.02491.x)

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard MA, Rambaut

A, Drummond AJ. 2014 BEAST 2: A Software Platform for Bayesian

Evolutionary Analysis. PLoS Computational Biology 10, e1003537.

(doi:10.1371/journal.pcbi.1003537)

Capellini I, Venditti C, Barton, RA. 2010 Phylogeny and metabolic scaling in mammals.

Ecology, 91, 2783-2793.

Charlesworth B, Lande R, Slatkin M. 1982 A Neo-Darwinian Commentary on

Macroevolution. Evolution 36, 474–498. (doi:10.1111/j.1558-

5646.1982.tb05068.x)

Chesson P. 2000 Mechanisms of maintenance of species diversity. Annual Review of

Ecology and Systematics, 31, 343-366.

Cheverud JM. 1984 Quantitative genetics and developmental constraints on evolution by

selection. Journal of Theoretical Biology 110, 155–171. (doi:10.1016/s0022-

5193(84)80050-8)

Crouch NMA, Ricklefs RE. 2019 Speciation Rate Is Independent of the Rate of

Evolution of Morphological Size, Shape, and Absolute Morphological

Specialization in a Large Clade of Birds. The American Naturalist 193, E78–E91.

(doi:10.1086/701630)

Donoghue MJ, Ree RH. 2000 Homoplasy and Developmental Constraint: A Model and

an Example from Plants. American Zoologist 40, 759–769.

(doi:10.1093/icb/40.5.759)

Donoghue MJ. 2005 Key innovations, convergence, and success: macroevolutionary

lessons from plant phylogeny. Paleobiology 31, 77-93. (doi: 10.1666/0094-

8373(2005)031[0077:KICASM]2.0.CO;2)

Felice RN, Goswami A. 2017 Developmental origins of mosaic evolution in the avian

cranium. Proceedings of the National Academy of Sciences 115, 555–560.

(doi:10.1073/pnas.1716437115)

Felsenstein J. 1985 Phylogenies and the comparative method. The American Naturalist,

125, 1-15.

Fisher, DC. 1986 Progress in organismal design. Pp. 99-117 in: Patterns and Processes in

the History of Life. Raup DM and Jablonski D, eds. Springer, Berlin, Heidelberg.

Goswami A, Binder WJ, Meachen J, O'Keefe FR. 2015 The fossil record of phenotypic

integration and modularity: A deep-time perspective on developmental and

evolutionary dynamics. Proceedings of the National Academy of Sciences 112,

4891–4896. (doi:10.1073/pnas.1403667112)

Goswami A, Smaers JB, Soligo C, Polly PD. 2014 The macroevolutionary consequences

of phenotypic integration: from development to deep time. Philosophical

Transactions of the Royal Society B: Biological Sciences 369, 20130254–

20130254. (doi:10.1098/rstb.2013.0254)

Gould SJ, Vrba ES. 1982 Exaptation—a Missing Term in the Science of Form.

Paleobiology 8, 4–15. (doi:10.1017/s0094837300004310)

Gould SJ and Lewontin RC. 1979 The spandrels of San Marco and the Panglossian

paradigm: a critique of the adaptationist programme. Proceedings of the Royal

Society of London. Series B. Biological Sciences 205, 581–598.

(doi:10.1098/rspb.1979.0086)

Gould, SJ. 1977 Ontogeny and Phylogeny. Harvard University Press, Cambridge, USA.

Harmon LJ. 2003 Tempo and Mode of Evolutionary Radiation in Iguanian Lizards.

Science 301, 961–964. (doi:10.1126/science.1084786)

Holloway RL. 1973 Endocranial volumes of early African hominids, and the role of the

brain in human mosaic evolution. Journal of Human Evolution 2, 449–459.

(doi:10.1016/0047-2484(73)90123-1)

Hunt KD. 2016 Why are there apes? Evidence for the co-evolution of ape and monkey

ecomorphology. Journal of Anatomy 228, 630–685. (doi:10.1111/joa.12454)

Lewton KL. 2010 Locomotor function and the evolution of the primate pelvis. (Doctoral

dissertation, Arizona State University).

Losos JB. 2010 Adaptive Radiation, Ecological Opportunity, and Evolutionary

Determinism. The American Naturalist 175, 623–639. (doi:10.1086/652433)

Mahler DL, Ingram T, Revell LJ, Losos JB. 2013 Exceptional Convergence on the

Macroevolutionary Landscape in Island Lizard Radiations. Science 341, 292–295.

(doi:10.1126/science.1232392)

McHenry H. 1975 Fossils and the mosaic nature of human evolution. Science 190, 425–

431. (doi:10.1126/science.809842)

Michel LA et al. 2014 Remnants of an ancient forest provide ecological context for Early

Miocene fossil apes. Nature Communications 5. (doi:10.1038/ncomms4236)

Rice, SH. 2002. The role of heterochrony in primate brain evolution. In: Human

evolution through developmental change. N. Minugh-Purvis & K.J. McNamara,

eds. Johns Hopkins Univ. Press, Baltimore.

Seilacher A. 1991 Self-Organizing Mechanisms in Morphogenesis and Evolution. In

Constructional Morphology and Evolution, pp. 251–271. Springer Berlin

Heidelberg. (doi:10.1007/978-3-642-76156-0_17)

Silvestro D, Salamin N, Schnitzler J. 2014 PyRate: a new program to estimate speciation

and extinction rates from incomplete fossil data. Methods in Ecology and

Evolution 5, 1126-1131.

Smith JM, Burian R, Kauffman S, Alberch P, Campbell J, Goodwin B, Lande R, Raup D,

Wolpert L. 1985 Developmental Constraints and Evolution: A Perspective from

the Mountain Lake Conference on Development and Evolution. The Quarterly

Review of Biology 60, 265–287. (doi:10.1086/414425)

Smith SA, Brown JW, Yang Y, Bruenn R, Drummond CP, Brockington SF, Walker JF,

Last N, Douglas N, Moore MJ. 2018 Disparity, Diversity, and Duplications in the

Caryophyllales. New Phytologist 217, 836-854.

Olson ME. 2012 The developmental renaissance in adaptationism. Trends in Ecology and

Evolution 27, 278–287. (doi:10.1016/j.tree.2011.12.005)

Olson EC and Miller RL. 1958. Morphological Integration. University of Chicago Press.

Parins-Fukuchi C. 2018 Bayesian placement of fossils on phylogenies using quantitative

morphometric data. Evolution 72, 1801–1814. (doi:10.1111/evo.13516)

Parins-Fukuchi C. In press. Detecting mosaic patterns in macroevolutionary disparity.

American Naturalist.

Purvis A, Nee S, Harvey PH. 1995 Macroevolutionary inferences from primate

phylogeny. Proceedings of the Royal Society of London, Series B: Biological

Sciences 260, 329-333.

Rabosky DL, Santini F, Eastman J, Smith SA, Sidlauskas B, Chang J, Alfaro ME. 2013

Rates of speciation and morphological evolution are correlated across the largest

vertebrate radiation. Nature Communications 4. (doi:10.1038/ncomms2958)

Rabosky DL. 2014 Automatic Detection of Key Innovations, Rate Shifts, and Diversity-

Dependence on Phylogenetic Trees. PLoS One 9, e89543.

(doi:10.1371/journal.pone.0089543)

Rabosky DL, Grundler M, Anderson C, Title P, Shi JJ, Brown JW, Huang H, Larson JG.

2014 BAMMtools: an R package for the analysis of evolutionary dynamics on

phylogenetic trees. Methods in Ecology and Evolution 5, 701–707.

(doi:10.1111/2041-210x.12199)

Rainey PB, Travisano M. 1998 Adaptive radiation in a heterogeneous environment.

Nature 394, 69–72. (doi:10.1038/27900)

Raup DM. 1968 Theoretical morphology of echinoid growth. Journal of Paleontology 42,

50-63.

Ruff CB. 1994 Morphological adaptation to climate in modern and fossil hominids.

American Journal of Physical Anthropology 37, 65–107.

(doi:10.1002/ajpa.1330370605)

Simpson GG. 1944 Tempo and Mode in Evolution. Columbia University Press. New

York.

Slater GJ, Friscia AR. 2019 Hierarchy in adaptive radiation: A case study using the

Carnivora (Mammalia). Evolution 73, 524–539. (doi:10.1111/evo.13689)

Slater GJ. 2015 Iterative adaptive radiations of fossil canids show no evidence for

diversity-dependent trait evolution. Proceedings of the National Academy of

Sciences 112, 4897–4902. (doi:10.1073/pnas.1403666111)

Sukumaran J, Holder MT. 2010 DendroPy: a Python library for phylogenetic computing.

Bioinformatics 26, 1569–1571. (doi:10.1093/bioinformatics/btq228)

Stebbins GL. 1983 Mosaic evolution: An integrating principle for the modern synthesis.

Experientia 39, 823–834. (doi:10.1007/bf01990398)

Stebbins GL. 1984 Mosaic evolution, mosaic selection and angiosperm phylogeny.

Botanical Journal of the Linnean Society 88, 149–164. (doi:10.1111/j.1095-

8339.1984.tb01568.x)

Van Valen L. 1975 Group selection, sex, and fossils. Evolution, 29, 87-94.

Vos RA. 2006 Inferring large phylogenies: the big tree problem (Doctoral dissertation,

Simon Fraser University)

Waddington CH. 1959 Canalization of Development and Genetic Assimilation of

Acquired Characters. Nature 183, 1654–1655. (doi:10.1038/1831654a0)

Wagner CE, Harmon LJ, Seehausen O. 2012 Ecological opportunity and sexual selection

together predict adaptive radiation. Nature 487, 366–369.

(doi:10.1038/nature11144)

Wagner GP, Pavlicev M, Cheverud JM. 2007 The road to modularity. Nature Reviews

Genetics 8, 921-931.

Wagner GP, Amemiya C, Ruddle F. 2003 Hox cluster duplications and the opportunity

for evolutionary novelties. Proceedings of the National Academy of Sciences 100,

14603–14606. (doi:10.1073/pnas.2536656100)

Wagner GP. 1988 The influence of variation and of developmental constraints on the rate

of multivariate phenotypic evolution. Journal of Evolutionary Biology 1, 45–66.

(doi:10.1046/j.1420-9101.1988.1010045.x)

Wagner PJ. 2018 Early bursts of disparity and the reorganization of character integration.

Proceedings of the Royal Society B: Biological Sciences 285, 20181604.

(doi:10.1098/rspb.2018.1604)

Worthington S. 2012 New approaches to late Miocene hominoid systematics: Ranking

morphological characters by phylogenetic signal (Doctoral dissertation, New

York University).

Wright, S. 1932 The roles of mutation, inbreeding, crossbreeding and selection in

evolution. Proc. 6th Int. Cong. Genet. 1: 356–366.

Yoder JB, et al. 2010 Ecological opportunity and the origin of adaptive radiations.

Journal of Evolutionary Biology 23, 1581–1596. (doi:10.1111/j.1420-

9101.2010.02029.x)

Young NM, Wagner GP, Hallgrímsson B. 2010 Development and the evolvability of

human limbs. Proceedings of the National Academy of Sciences 107, 3400–3405.

(doi:10.1073/pnas.0911856107)

810 **Tables**

Clade	Origination	Extinction	Net diversification	Preservation
Hominoidea	0.395	0.338	0.057	3.121
Cercopithecoidea	0.251	0.128	0.123	2.551

812 *Table 1*. Lineage diversification and preservation rate parameters estimated in apes

813 (Hominoidea) and Old World monkeys (Cercopithecoids). Parameters were summarized

814 as the mean values from four million Markov-Chain Monte Carlo generations after

815 discarding 20% as burn-in.

829 **Figures**

Figure 1. The mosaic inference procedure starts by modelling all characters under a

unified phylogenetic model, in which branch lengths represent morphological disparity.

The algorithm then splits traits with divergent patterns in disparity into their own suites

and then greedily merges these suite fragments until the AIC score ceases to improve.

The splitting and merging steps are then alternated until the AIC scores of the final

merged models cease to improve. The best clustering achieved from this procedure

represents the best-supported heterogeneous model of character integration.

Figure 2. Strength of integration across haplorrhine evolution. Branch labels correspond

to the strength and polarity of correlation in the reconstructed changes along each branch.

Higher correlations in either direction (closer to either 1 or -1) indicate stronger

evolutionary integration, while lower correlations (closer to 0) indicate weaker

integration.

Figure 3. A) Mosaic evolutionary rates calculated while walking back from *Homo* to the

most recent common ancestor of New World monkeys, Old World Monkeys, and apes.

B) Variance contributed to each suite by each skeletal region, estimated from a principal

component analysis. C) Skeletal regions over-represented in each mosaic evolutionary

suite. Values above zero correspond to regions that occupy a higher proportion of their

clusters than expected.

Figure 4. Branch colors correspond to rates of morphological evolution across all

characters along individual branches. Phylogenetic positions of fossil genera reflect

placements on extant reference tree.

857

Figure 5. Rate shifts in A) encephalization development, and B) encephalization quotient.

C) Branch-specific evolutionary rates when tracing the lineage from *Homo* to the root.

Figure 6. Monte Carlo simulations of integrated and atomized systems of three

continuous traits. Each box is a simulated diffusion between three traits under A)

moderate developmental linkage and shared selection, B) no developmental linkage and

shared selection, C) no developmental linkage and selective independence, D) high

developmental linkage and selective independence, and E) high developmental linkage

and shared selection.

suite	skeletal region	trait
0	Ulna	Olecranon length
0	Ulna	Middle shaft shape
0	Ulna	Olecranon shape
0	Ulna	Distal shape
0	Ulna	Styloid process length
0	Ulna	Proximal shaft shape
0	Ulna	Coronoid
0	Ulna	Trochlea notch shape
0	Humerus	Head P D shape
0	Humerus	Trochlea
0	Humerus	Intertubercular angle
0	Humerus	Lat. trochlea keel dev.
0	Humerus	Bicipital groove shape
0	Humerus	Humeral Head Size
0	Humerus	Head torsion
0	Humerus	Trochlea sup. notch angle
0	Humerus	Zona conoidea dist. depth
0	Humerus	Grt. tuberosity proj.
0	Carpus	Capitate head M L width
0	Carpus	Capitate M C 3 shape
0	Carpus	CAPCM4
0	Carpus	Hamate DP depth COR
0	Carpus	Capitate head A P / P D
0	Carpus	Capitate dorsal nonarticular

0 Carpus	Capitate M C 3 depth
0 Carpus	Lunate dist.
0 Carpus	Hamate triquetral facet angle
0 Carpus	Capitate M C 2 facets
0 Cranium	Grt. palatine foramen position
0 Cranium	Zygomatic root origin elevation
0 Cranium	Basicranial length
0 Cranium	Piriform aperture shape
0 Cranium	Postglenoid projection
0 Cranium	Basicranium vs. palate
0 Cranium	Bregma position
0 Cranium	Lacrimal fossa & orbital floor
0 Cranium	Facial projection
0 Cranium	Zygomax. tubercle A P width
0 Cranium	Basicranium vs. nuchal
0 Cranium	Nasoalveolar clivus length
0 Cranium	Head M L shape
0 Cranium	Zygomatic
0 Cranium	Face height vs. depth
0 Cranium	Piriform max. M L elevation
0 Cranium	Neurocranial shape
0 Cranium	Asterion angle
0 Radius	Lunate facet shape
0 Radius	Lunate facet angle
0 Radius	Neck shape
0 Radius	Scapholunate facet areas
0 Radius	Lunate facet dorsal proj.

0 Radius	Distal facets area
0 Radius	Total length
0 Radius	Radial neck length
0 Radius	Shaft curvature M L
0 Radius	Head articular surface P D
0 Scapula	Trapezius insert angle
0 Scapula	Coraco
0 Scapula	Inf. glenoid cor
0 Scapula	Acromian size
0 Scapula	Inferior angle
0 Scapula	Sup. notch shape
0 Scapula	Spinatus ratio
0 Scapula	Glenoid vent. bar angle
0 Pelvis	pubic symphysis length
1 Ulna	Ulna head height
1 Ulna	Semilunar keel angle
1 Ulna	Radial notch shape
1 Ulna	Distal shaft shape
1 Humerus	Dist. articular shape
1 Humerus	Trochlea dist. waisting
1 Humerus	Med. trochlea keel dev.
1 Humerus	Olecranon fossa depth
1 Humerus	Olecranon fossa shape
1 Carpus	Capitate P D length
1 Carpus	Hamate M C 4
1 Carpus	Lunate radial facet M L
1 Carpus	Lunate triquetral facet shape

1 Cranium	Piriform base M L width
1 Cranium	M1 shape
1 Cranium	Lsr. palatine foramen
1 Cranium	Molar crown areas
1 Cranium	Entoglenoid width
1 Cranium	Entoglenoid projection
1 Cranium	Nasal bone M L width
1 Cranium	Orbital margin M L width
1 Cranium	Orbit area
1 Cranium	Interorbital width
1 Radius	Distal surface shape
1 Radius	Styloid process length
1 Radius	Radial Head Shape
1 Manubrium	Manubrium shape
1 Scapula	Scapula Body shape
1 Scapula	Trapezius med. insert
1 Scapula	Superior angle
1 Scapula	Subscapular depth
1 Scapula	Vertebral border length
1 Scapula	Glenoid fossa shape
1 Scapula	Coracoid length
1 Pelvis	illium length
1 Pelvis	bi
1 Pelvis	anteroposterior pelvic diameter
1 Pelvis	pelvic length
2 Humerus	Epicondyles projection
2 Humerus	Olecranon perforation

2 Ulna	Shaft curvature A P
2 Carpus	Capitate M C 3 waisting
2 Carpus	Capitate waisting
2 Carpus	Lunate PD Length
2 Carpus	Hamate hamulus P D proj.
2 Carpus	Capitate M C 2 margin
2 Carpus	Capitate M C 2 orientation
2 Carpus	Lunate dist. facet shape
2 Cranium	Orbit shape
2 Cranium	Pterion post. config
2 Cranium	Palate shape
2 Cranium	Lacrimal fossa visibility
2 Cranium	Palatine width
2 Cranium	Postorbital constriction
2 Cranium	Palate depth
2 Radius	Head depth P D
2 Scapula	Glenoid fossa depth
2 Pelvis	illium width
2 Pelvis	ischium length
2 Pelvis	AIIS to hip joint
2 Pelvis	inferior pubic ramus length
2 Pelvis	acetabulum diameter
2 Pelvis	superior pubic ramus length
3 Scapula	Spinoglenoid notch depth
3 Scapula	Infraglenoid tuberosity M L
3 Pelvis	lower illiac height
3 Pelvis	inferior pubic ramus cross

3 Ulna	Radial notch depth
3 Cranium	Temporal fossa width
3 Cranium	Face vs. nasal height
3 Cranium	Face height vs. width
3 Cranium	Palate width
3 Cranium	Entoglenoid lat. pterygoid
3 Humerus	Medial trochlea keel & medial epicon
4 Pelvis	ischium cross
4 Pelvis	superior pubic ramus cross
4 Pelvis	upper illiac height
4 Pelvis	lower ilium cross
4 Pelvis	ischium dorsal projection
4 Pelvis	pubic symphysis
4 Pelvis	auricular surface area
4 Pelvis	biacetabular diameter
4 Pelvis	maximum transverse diameter
4 Pelvis	sacrum width

874 **Table S1:** List of traits and their skeletal regions that comprise each of the mosaic suites detected
875 from the skeletal data.